Preface

Technological advance in endoscopic diagnosis and therapy have resulted in remarkable progress in medical engineering. This issue examines photodynamic therapy (PDT) as a special topic. PDT differs significantly from conventional cancer therapeutic modalities. The concept of PDT was initially reported by Jesionek and Tappeiner in 1903. Further research on PDT was made by Dougherty who used laser light and hematoporphyrin derivative in the 1970's, and clinical trials began in 1980. The number of patients treated by PDT throughout the world is estimated to be more than 5,000. PDT is indicated not only in malignant diseases but also non-malignant diseases. PDT can be used for cancer of the lung, esophagus, stomach, colon, bile duct, skin, cervix, genital organs, urinary bladder, brain, breast and eye.

As malignant diseases tend to develop in older persons and also have a tendency toward multiple incidence, conventional therapies are not always indicated for these patients. PDT is probably applicable in some of those patients.

Since PDT does not damage organ function, and is a minimally invasive procedure, it has a good level of toleration and allows rapid recovery. PDT yields a high rate of complete response in early stage cancer and can eliminate the necessity for surgery in certain patients.

Since recent studies on photodynamic therapy were started only 30 years ago, there are still a number of unsolved problems. However it is clear that PDT has become firmly established as one of the new therapeutic strategies for cancer.

This issue provides an overview of the clinical experiences with PDT for cancers of various organs and attempts to define the indications and limitations of modality.

Harubumi Kato, M.D., Ph.D.
Editor-in-Chief