

# ON THE ORTHOGONALIZATION OF ARBITRARY BOOLEAN FORMULAE

RENATO BRUNI

*Received 15 October 2002 and in revised form 19 March 2004*

The orthogonal conjunctive normal form of a Boolean function is a conjunctive normal form in which any two clauses contain at least a pair of complementary literals. Orthogonal disjunctive normal form is defined similarly. Orthogonalization is the process of transforming the normal form of a Boolean function to orthogonal normal form. The problem is of great relevance in several applications, for example, in the reliability theory. Moreover, such problem is strongly connected with the well-known propositional satisfiability problem. Therefore, important complexity issues are involved. A general procedure for transforming an arbitrary CNF or DNF to an orthogonal one is proposed. Such procedure is tested on randomly generated Boolean formulae.

## 1. Introduction

Let  $\mathbb{B} = \{0, 1\}$ , or, equivalently,  $\{\text{true}, \text{false}\}$ . A Boolean function of  $n$  Boolean variables  $x_i \in \mathbb{B}$  is a function  $f(x_1, \dots, x_n)$  from the Boolean hypercube  $\mathbb{B}^n$  to the Boolean set  $\mathbb{B}$ . We assume that the reader is familiar with the basic concepts of Boolean algebra (see, e.g., [11, 19]). A Boolean function can be represented in several manners. The most commonly used one is by means of a Boolean (or propositional) formula  $\mathcal{F}$  in *conjunctive* (CNF) or *disjunctive* (DNF) normal form. Both normal forms are widely used, the choice often depends on the applicative context. Orthogonal conjunctive normal form (OCNF) is a CNF in which any two clauses contain at least a pair of complementary literals. Orthogonal disjunctive normal form (ODNF) is defined similarly. The orthogonal form is of great relevance in solving several difficult problems, for example, in the reliability theory. One of the fundamental issues in reliability is to compute the probability  $p$  that a complex system is in *operating* state (and not in *failed* state, see, e.g., [3, 4]). The state of the system depends on the state  $x_i$  (operating or failed) of its  $i$ th component, for  $i = 1, \dots, n$ . Such relationship is usually described by means of a Boolean function  $g(x_1, \dots, x_n)$ , so that knowing the state of the components, the state of the system is immediately computable. Since the probabilities  $p_r$  of each component to be in operating state are generally known,  $p$  can also be computed by using  $g$ . However, such computation may be difficult [16]. For systems where  $g$  is expressed by a Boolean formula in ODNF, this probability is very easily

computed by summing the probabilities associated to all individual terms, since any two terms correspond to pairwise incompatible events. Another important application is in the field of mathematical statistics, where the orthogonality property is needed for assuring independence among statistical variables, and in particular in the analysis of variance, where it is used to separate the variation inside each group from the variation among the groups themselves [18].

A classical problem is therefore to derive the orthogonal form, or disjoint products form, of a Boolean function. A reliability algorithm based on the computation of the orthogonal form is proposed, for instance, in [14]. In general, however, computation of the orthogonal forms is a difficult problem [6]. One of the few interesting classes of formulae for which this can be performed efficiently is the class of *shellable* DNF, introduced by Ball and Provan [2]. It has been recently proved by Boros et al. [6] that every positive Boolean function (i.e., a Boolean function that can be written as a DNF that has no negative literals) can be represented by a shellable DNF. However, the complexity of recognizing shellable DNF is not known, and testing the lexicoexchange property (a strengthening of shellability) is NP-complete [6].

A procedure to transform a generic normal form formula into an orthogonal normal form, also briefly called orthogonal form, is described here. Such operation is called *orthogonalization*. The proposed procedure is applicable to both CNF and DNF. Therefore, in Section 2, we introduce a unified notation for normal forms in order to represent both CNF and DNF. A basic procedure to orthogonalize a generic formula is described in Section 4. During the above process, the size of the formula tends to exponentially increase. This is not surprising, since in Section 3 we show that an NP-complete problem like propositional satisfiability [7, 10, 13] becomes easy for formulae in orthogonal form. (This can be related to a procedure for solving satisfiability by counting the number of possible solutions proposed in [12].) Hence, the NP complexity [9] can be seen as being *absorbed* by the orthogonalization process. Improvements on the above basic procedure, with the aim of minimizing the size of the formula both in the final result and during the computation, are then presented in Section 5.

## 2. Notation and equivalence of problems

A Boolean CNF formula is the logic conjunction ( $\wedge$ ) of  $m$  *clauses*, each clause being the logic disjunction ( $\vee$ ) of *literals*, each literal being either a positive ( $x_i$ ) or a negative ( $\neg x_i$ ) Boolean (or propositional) variable. By denoting with  $P_j \subseteq \{1, \dots, n\}$  the set of positive variables of the  $j$ th clause, and with  $N_j \subseteq \{1, \dots, n\}$  the set of negative variables of the same clause, we have

$$\bigwedge_{j=1, \dots, m} \left( \bigvee_{i \in P_j} x_i \vee \bigvee_{i \in N_j} \neg x_i \right). \quad (2.1)$$

Conversely, a Boolean DNF formula is the logic disjunction of  $m$  *terms*, each term being the logic conjunction of literals, defined as above. By denoting with  $P_j \subseteq \{1, \dots, n\}$  the set of positive variables of the  $j$ th term, and with  $N_j \subseteq \{1, \dots, n\}$  the set of negative variables

of same term, we have

$$\bigvee_{j=1,\dots,m} \left( \bigwedge_{i \in P_j} x_i \wedge \bigwedge_{i \in N_j} \neg x_i \right). \quad (2.2)$$

The proposed procedure will apply to both CNF and DNF. Therefore, a notation which can represent both forms is needed. Clauses and terms can be viewed as pairs of sets  $(P_j, N_j)$  of literals plus a logical operator connecting all such literals. Such structures will be called *monomials*, and be denoted by  $m_j$ . The Boolean function expressed by a single monomial  $m_j$  will be denoted by  $m_j(x_1, \dots, x_n)$ . A CNF or DNF formula  $\mathcal{F}$  can now be viewed as a collection of monomials. An *external* operator is applied between monomials, and will be indicated here with the symbol  $\perp$ , and an *internal* operator is applied between literals of the same monomial, and will be indicated here with the symbol  $\top$ . Both CNF and DNF are therefore representable as follows:

$$\perp_{j=1,\dots,m} \left( \top_{i \in P_j} x_i \top_{i \in N_j} \neg x_i \right). \quad (2.3)$$

Clearly,  $\perp$  means  $\wedge$  when considering CNF, and  $\vee$  when considering DNF, and vice versa holds for  $\top$ . Given a so defined monomial  $m_j$ , let the set  $T_j \subseteq \mathbb{B}^n$  where  $m_j$  has value 1 be the set of *true points* of  $m_j$ :

$$T_j = \{(x_1, \dots, x_n) \in \mathbb{B}^n : m_j(x_1, \dots, x_n) = 1\}, \quad (2.4)$$

and let the set  $F_j \subseteq \mathbb{B}^n$  (the complement of  $T_j$  with respect to  $\mathbb{B}^n$ ) where  $m_j$  has value 0 be the set of *false points* of  $m_j$ :

$$F_j = \{(x_1, \dots, x_n) \in \mathbb{B}^n : m_j(x_1, \dots, x_n) = 0\}. \quad (2.5)$$

Given now a generic Boolean formula  $\mathcal{F}$ , let the global set of true points be  $T = \{(x_1, \dots, x_n) \in \mathbb{B}^n : f(x_1, \dots, x_n) = 1\}$ , and let the global set of false points be  $F = \{(x_1, \dots, x_n) \in \mathbb{B}^n : f(x_1, \dots, x_n) = 0\}$ . When  $\mathcal{F}$  has the structure of normal form (CNF or DNF), the following relations hold.

LEMMA 2.1. *In the case of CNF, the sets  $T$  and  $F$  are given by*

$$T = \bigcap_{j=1}^n T_j, \quad F = \bigcup_{j=1}^n F_j. \quad (2.6)$$

LEMMA 2.2. *In the case of DNF, the sets  $T$  and  $F$  are given by*

$$T = \bigcup_{j=1}^n T_j, \quad F = \bigcap_{j=1}^n F_j. \quad (2.7)$$

Note that  $T$  and  $F$  are not immediately computable from  $\mathcal{F}$ , nor are their cardinalities. Besides the fact that their cardinality can be exponential in the number of variables, even expressing such sets in some *compressed* but *usable* form appears difficult. In fact, knowing the set  $T$  (or equivalently  $F$ ) would give the solution of an NP-complete problem, namely the *propositional satisfiability* problem (see, e.g., [10]). Also, knowing the cardinality  $|T|$  (or equivalently  $|F|$ ) would give the solution of the decision version of the propositional satisfiability problem, which is still NP-complete. Moreover, this theoretically means that every problem in NP can be polynomially reduced to the problem of finding this cardinality [9].

On the contrary, the sets  $F_j$  for CNF and  $T_j$  for DNF are immediately computable and expressible in compressed form (see below). However, in the case of a generic CNF or DNF, such sets are not disjoint, but can overlap each other: it can be  $T_j \cap T_k \neq \emptyset$  or  $F_j \cap F_k \neq \emptyset$  for some  $j, k \in \{1, \dots, m\}$ . Due to the above reason, in order to find, respectively, the cardinalities  $|T|$  and  $|F|$ , it would be necessary to identify, respectively, all the  $T_j$  and all the  $F_j$ . Since the number of points in  $T_j$  and  $F_j$  can be exponential in the number of variables, the approach of identifying all the  $T_j$  and all the  $F_j$  has exponential worst-case time complexity. This is not surprising. On the other hand, if all the  $T_j$  (resp., all the  $F_j$ ) would be pairwise disjoint sets, in order to find the cardinality  $|T|$  (resp.,  $|F|$ ) it would suffice to know the cardinalities of the  $T_j$  (resp.,  $F_j$ ), and sum them. Such cardinalities are, in fact, trivially computable. In order to proceed with our notation unification, dissimilarities between *true* and *false* sets should be overcome. Consider again the satisfiability problem. It consists in finding if, in the Boolean hypercube  $\mathbb{B}^n$ , there is at least one true point for all clauses (for DNF formulae), or at least one false point for all terms (for DNF formulae). Altogether, false points are bad for CNF, while true points are bad for DNF. We will now call the set of such bad points  $B$ , with the convention that  $B = F$  for CNF, and  $B = T$  for DNF. Moreover, every monomial  $m_j$  has its set of bad points  $B_j$  of the Boolean hypercube  $\mathbb{B}^n$ , with the convention that  $B_j = F_j$  for CNF, and  $B_j = T_j$  for DNF. (More intuitively, every  $m_j$  *forbids* a set of points: in the case of CNF, the  $j$ th clause forbids its  $F_j$ , while in the case of DNF, the  $j$ th term forbids its  $T_j$ .) Conversely, we will call  $G$  the set of good points, with the convention that  $G = T$  for CNF, and  $G = F$  for DNF. Therefore, every monomial  $m_j$  has its set of good points  $G_j$ , with  $G_j = T_j$  for CNF, and  $G_j = F_j$  for DNF. Sets  $B_j$  and  $G_j$  on  $\mathbb{B}^n$  are directly obtainable by the structure of  $m_j$ . In the case of CNF,  $B_j$  (in implicit form) is given by a vector of length  $n$ , called *pattern*, having 0 for each variable appearing positive in  $m_j$ , 1 for each variable appearing negative in  $m_j$ , and  $*$  (*do not care*) for each variable not appearing in  $m_j$ . Expanding every  $*$  with both 0 and 1 gives all the points of  $B_j$  explicitly. Clearly,  $G_j$  is given by  $\mathbb{B}^n \setminus B_j$ . In the case of DNF,  $B_j$  (in implicit form) is given by a pattern having 1 for each variable appearing positive in  $m_j$ , 0 for each variable appearing negative in  $m_j$ , and  $*$  for each variable not appearing in  $m_j$ . Explicit expression of all points of  $B_j$  and  $G_j$  are obtainable as above. Pattern notation can be unified by using symbol “+” for 1 in case of CNF, for 0 in the case of DNF, and symbol “-” for 0 in the case of CNF, for 1 in the case of DNF (cf. Table 2.1).

*Example 2.3.* Suppose  $n = 5$ . Given monomial  $(x_1 \top \neg x_3 \top x_4)$ , the pattern for the set of its bad points is  $\{-, *, +, -, *\}$ , corresponding to  $\{0, *, 1, 0, *\}$  in the case of CNF, to  $\{1, *, 0, 1, *\}$  in the case of DNF.

Table 2.1. Conventions used in the unified notation for CNF and DNF.

Form	External operator	Internal operator	Bad point	Good point	Pattern	
CNF	$\wedge$	$\vee$	$F$	$T$	0	1
DNF	$\vee$	$\wedge$	$T$	$F$	1	0
Unified	$\perp$	$\top$	$B$	$G$	-	+

The cardinalities of the above  $B_j$  and  $G_j$  are easily computable, as follows.

LEMMA 2.4. *Let  $n$  be the number of variables, and let  $l(m_j)$  be the number of distinct literals appearing in  $m_j$ . The cardinalities of the above introduced  $B_j$  and  $G_j$  are  $|B_j| = 2^{n-l(m_j)}$ , and  $|G_j| = 2^n - |B_j| = 2^n - 2^{n-l(m_j)}$ .*

We denote with  $(\phi)$  the empty monomial, that is, the monomial  $m_\phi$  which is an empty set of literals. According to Lemma 2.4,  $B_{(\phi)} = \mathbb{B}^n$ , hence  $(\phi)$  has only bad points. Finally, we denote with  $\phi$  the empty formula, that is, the formula  $\mathcal{F}_\phi$  which is an empty set of monomials. By definition,  $\phi$  has only good points, so  $G_\phi = \mathbb{B}^n$ .

### 3. The orthogonal form

A Boolean formula (in unified notation) is in *orthogonal* normal form when every pair of monomials  $m_j$  and  $m_k$  contains at least one Boolean variable  $x_i$  (not necessarily the same  $i$  for all the couples of indexes  $(j, k)$ ) as a positive instance ( $x_i$ ) in one of them (e.g.,  $m_j$ ) and as a negative instance ( $\neg x_i$ ) in the other (e.g.,  $m_k$ ):

$$m_j = (\cdots \top x_i \top \cdots), \quad m_k = (\cdots \top \neg x_i \top \cdots) \quad \forall j, k \in \{1, \dots, m\}. \quad (3.1)$$

The above situation for  $m_j$  and  $m_k$  is variously expressed in literature: the above monomials are said to be *orthogonal* [6], or to *clash* [8] on  $x_i$ , or to *resolve* [17] on  $x_i$ , or also to *hit* [1] on  $x_i$ .

THEOREM 3.1. *A Boolean formula is in orthogonal normal form if and only if the above defined sets  $B_j$  are pairwise disjoint.*

The above theorem clearly particularizes for CNF as follows:

$$\begin{aligned} F_j \cap F_k &= \phi \quad \forall j, k \in \{1, \dots, m\}, \\ (T_j \cap T_k \text{ can be } \neq \phi \text{ for some } j, k \in \{1, \dots, m\}), \end{aligned} \quad (3.2)$$

and for DNF as follows:

$$\begin{aligned} T_j \cap T_k &= \phi \quad \forall j, k \in \{1, \dots, m\}, \\ (F_j \cap F_k \text{ can be } \neq \phi \text{ for some } j, k \in \{1, \dots, m\}). \end{aligned} \quad (3.3)$$

*Proof.* We first prove that orthogonal form implies  $B_j \cap B_k = \phi$  for all  $j, k \in \{1, \dots, m\}$ . If two monomials  $m_j$  and  $m_k$  clash on at least one variable  $x_c$ , the corresponding  $B_j$  and  $B_k$  are defined by two patterns which, respectively, have - and + in at least position  $c$ , hence they define two sets  $B_j$  and  $B_k$  which cannot have any common point. We now

prove that  $B_j \cap B_k = \emptyset$  for all  $j, k \in \{1, \dots, m\}$  implies orthogonal form. Since  $B_j$  and  $B_k$  are disjoint, the patterns corresponding to them must contain in at least one position  $c$ , respectively,  $+$  and  $-$  (or  $-$  and  $+$ ). This is because any other combination ( $+$  and  $+$ ,  $+$  and  $*$ , etc.) would contradict the hypothesis of  $B_j$  and  $B_k$  disjoint. Therefore, by letting  $x_c$  be the variable corresponding to position  $c$ , monomials  $m_j$  and  $m_k$  corresponding to such patterns must both contain  $x_c$  and clash on it. Finally, since we assumed that every pair of sets  $B_j, B_k$  has empty intersection, every pair of monomials  $m_j, m_k$  is orthogonal.  $\square$

Since the orthogonal form is a necessary and sufficient condition for having all the  $B_j$  pairwise disjoint, it is a condition for trivially solving the problem of computing  $|B|$ , which implies trivially solving the propositional satisfiability problem, with the above implications on all problems in NP.

*Example 3.2.* Suppose we are interested in checking satisfiability of

$$\begin{aligned} & (\neg x_1 \top \neg x_2 \top x_3 \top x_4 \top x_5) \perp (\neg x_1 \top \neg x_2 \top x_3 \top x_4 \top x_5) \perp (x_2 \top x_3 \top x_4 \top x_5) \\ & \perp (x_3 \top \neg x_4 \top x_5) \perp (x_3 \top x_4 \top \neg x_5) \perp (x_3 \top \neg x_4 \top \neg x_5) \perp (\neg x_3). \end{aligned} \quad (3.4)$$

In our terms, we need to check whether the global  $B$  covers the whole  $\mathbb{B}^5$ . There are many different and very efficient techniques to solve the satisfiability problem (see [10]). In practical cases, however, without imposing restrictions on the structure of the formula (Horn, quadratic, etc.), they have worst-case exponential time complexity. On the other hand, computing the above defined sets  $B_j$ , and their cardinalities, is straightforward:

$$\begin{aligned} (x_1 \top \neg x_2 \top x_3 \top x_4 \top x_5) &\longrightarrow B_1 = \{-, +, -, -, -\} \mid |B_1| = 1, \\ (\neg x_1 \top \neg x_2 \top x_3 \top x_4 \top x_5) &\longrightarrow B_2 = \{+, +, -, -, -\} \mid |B_2| = 1, \\ (x_2 \top x_3 \top x_4 \top x_5) &\longrightarrow B_3 = \{*, -, -, -, -\} \mid |B_3| = 2, \\ (x_3 \top \neg x_4 \top x_5) &\longrightarrow B_4 = \{*, *, -, +, -\} \mid |B_4| = 4, \\ (x_3 \top x_4 \top \neg x_5) &\longrightarrow B_5 = \{*, *, -, -, +\} \mid |B_5| = 4, \\ (x_3 \top \neg x_4 \top \neg x_5) &\longrightarrow B_6 = \{*, *, -, +, +\} \mid |B_6| = 4, \\ (\neg x_3) &\longrightarrow B_7 = \{*, *, +, *, *\} \mid |B_7| = 16. \end{aligned} \quad (3.5)$$

By computing the union of all the  $B_j$ , we have that  $B$  actually covers  $\mathbb{B}^5$  (see Figure 3.1 which reports the case of a CNF). Hence, the given formula is unsatisfiable. Since the number of points of such union is exponential (in the worst case) in the number of variables, this procedure has exponential time complexity. On the contrary, one could observe that the formula is orthogonal, hence the  $B_j$  are pairwise disjoint. On this basis, trivially,  $|B| = |B_1| + |B_2| + |B_3| + |B_4| + |B_5| + |B_6| + |B_7| = 32$ . This suffices to say that  $B$  covers  $\mathbb{B}^5$ , whose cardinality is  $2^5 = 32$ , and so the given formula is unsatisfiable. Altogether, by using the fact that the given formula is in orthogonal form, one can very easily solve the satisfiability problem.




Figure 3.1. The sets  $B_j$  of Example 3.2 on the Boolean hypercube  $\mathbb{B}^5$  in the case of a CNE.

#### 4. Basic orthogonalization procedure

In order to present a procedure for the orthogonalization of a generic Boolean formula, we first need to define an operation which will be called *multiplication*, denote it with  $\diamond$ , and apply it to a pair of monomials  $m_j$  and  $m_k$ . The result of such multiplication is a new monomial containing all the literals of  $m_j$  and  $m_k$  (but without repeated ones) when the two monomials are not orthogonal, and the empty formula  $\phi$  (i.e., a formula for which there are only good points, cf. Section 2) when they are orthogonal:

$$\begin{aligned}
 m_j \diamond m_k &= \left( \bigwedge_{i \in P_j} x_i \top \bigwedge_{i \in N_j} \neg x_i \right) \diamond \left( \bigwedge_{i \in P_k} x_i \top \bigwedge_{i \in N_k} \neg x_i \right) \\
 &= \begin{cases} \phi & \text{if } m_j \text{ and } m_k \text{ are orthogonal,} \\ \left( \bigwedge_{i \in (P_j \cup P_k)} x_i \top \bigwedge_{i \in (N_j \cup N_k)} \neg x_i \right) & \text{otherwise.} \end{cases} \quad (4.1)
 \end{aligned}$$

**THEOREM 4.1.** *Consider any two monomials  $m_j$  and  $m_k$  with their corresponding sets  $B_j$ ,  $G_j$ ,  $B_k$ , and  $G_k$ . Let  $m_l = m_j \diamond m_k$  be their product. The set of the bad points for  $m_l$  is  $B_l = B_j \cap B_k$ , while the set of good points is  $G_l = G_j \cup G_k$ .*

*Proof.* Given a generic monomial  $m_j$ , by adding literals to  $m_j$  the set  $B_j$  can in general only be reduced (this means decreasing the false set for CNE, decreasing the true set for DNF), and therefore the set  $G_j$  increased. Monomial  $m_l$  can be seen as adding literals to  $m_j$ , so  $B_l \subseteq B_j$ , and can also be seen as adding literals to  $m_k$ , so  $B_l \subseteq B_k$ . Therefore,  $B_l \subseteq B_j \cap B_k$ . Moreover, any point  $x \in B_j \cap B_k$  is a bad point for  $m_l$ , hence  $x \in B_l$ . This proves  $B_l = B_j \cap B_k$ , and consequentially  $G_l = G_j \cup G_k$ . Coherently, when  $m_j$  and  $m_k$  are orthogonal, the result of the multiplication, by definition, is the empty formula  $\phi$ , the sets  $B_j$  and  $B_k$  are disjoint by Theorem 3.1, their intersection is empty, and so is the set  $B_\phi$  by definition.  $\square$

Given an arbitrary monomial  $m_j = (x_h \top x_{h+1} \top \cdots \top x_i)$ , its negation  $\neg(m_j)$  is easily computable (by De Morgan's laws [19]) as the following set of monomials connected by our external operator:  $(\neg x_h) \perp (\neg x_{h+1}) \perp \cdots \perp (\neg x_i) = \neg(m_j)$ . However, the expression for  $\neg(m_j)$  is not unique. One could, in fact, consider a negation which is in orthogonal form, namely the *orthogonal negation*  $\neg^o(m_j)$  of  $m_j$ . Such negation  $\neg^o(m_j)$  is composed of  $k$  monomials  $\sigma_1^j \perp \sigma_2^j \perp \cdots \perp \sigma_k^j$ , the first of which contains the negation of the first variable, the second contains the first variable and the negation of the second one, and so on, as follows:

$$(\neg x_h) \perp (x_h \top \neg x_{h+1}) \perp \cdots \perp (x_h \top x_{h+1} \top \cdots \top \neg x_i). \quad (4.2)$$

*Example 4.2.* The orthogonal negation of  $m = (x_1 \top x_2 \top \neg x_3)$  is

$$\neg^o(m) = \sigma_1^m \perp \sigma_2^m \perp \sigma_3^m = (\neg x_1) \perp (x_1 \top \neg x_2) \perp (x_1 \top x_2 \top x_3). \quad (4.3)$$

We also define the multiplication of a monomial  $m_k$  by the negation  $\neg(m_j)$  of another monomial  $m_j$  as the set of monomials obtained by multiplying  $m_k$  by each of the monomials in  $\neg(m_j)$ . We denote this operation by  $m_k \diamond \neg(m_j)$ . Based on this, a basic orthogonalization operation can be developed. For clarity, we report the procedure without indicating negative variables. However, this does not cause any loss of generality, since negative variables can perfectly be present, and the negations will eventually appear in the result according to elementary rules of Boolean algebra.

*Basic orthogonalization operation.* Consider any two distinct monomials  $m_j$  and  $m_k$  not already orthogonal. Let  $C_{jk}$  be the (possibly empty) set of common literals between  $m_j$  and  $m_k$ , and  $D_j$  and  $D_k$  the (possibly empty) sets of literals, respectively, belonging only to  $m_j$  and only to  $m_k$ :

$$m_j = \left( \prod_{i \in D_j} x_i \top \prod_{i \in C_{jk}} x_i \right), \quad m_k = \left( \prod_{i \in D_k} x_i \top \prod_{i \in C_{jk}} x_i \right). \quad (4.4)$$

Note that, since they are not orthogonal, they cannot contain complementary literals:  $x_i \in m_j \Rightarrow \neg x_i \notin m_k$ . Choose one of the sets of different literals, for instance  $D_j$ , and consider the monomial  $m_d$  composed of all such literals. Compute now its orthogonal negation  $\neg^o(m_d) = \sigma_1^d \perp \sigma_2^d \perp \cdots \perp \sigma_j^d$ . We have that the (sub)formula  $m_j \perp m_k$  is equivalent (in the sense that they both represent the same Boolean function, we prove this throughout the rest of this section) to the following (sub)formula obtained by replacing  $m_k$  with  $m_k \diamond \neg^o(m_d)$ :

$$m_j \perp \sigma_1^d \diamond m_k \perp \sigma_2^d \diamond m_k \perp \cdots \perp \sigma_j^d \diamond m_k. \quad (4.5)$$

The essential point is that the obtained (sub) formula is now in orthogonal form. Hence, the (sub)formula composed of the two monomials  $m_j$  and  $m_k$  has been orthogonalized. Note that the number of monomials of the result is 1 plus the cardinality of the set of noncommon literals ( $D_j$ ) used. In order to obtain a smaller number of monomials, we




Figure 4.1. The partition of the Boolean hypercube  $\mathbb{B}^n$  determined by  $B_k$  and  $G_k$ .

always choose the set of noncommon literals of minimum cardinality. When one of these two sets is empty, this means that one of the monomials, say  $m_j$ , is a subset of the other  $m_k$ . Coherently, by choosing  $D_j$  for the above procedure, the result is only  $m_k$ . In fact, the Boolean (sub)formula  $m_j \perp m_k$  is equivalent, in this case, to the Boolean (sub)formula  $m_k$ . The following two theorems prove that replacing  $m_k$  with  $m_k \diamond \neg^o(m_d)$  produces an equivalent formula.

**THEOREM 4.3.** *Consider a monomial  $m_j$  and the negation  $\neg(m_k)$  of another monomial with their corresponding sets  $B_j$ ,  $G_j$ ,  $B_{-k}$ , and  $G_{-k}$ . The set of bad points for their product  $m_j \diamond \neg(m_k)$  is  $B_j \cap B_{-k}$ , while the set of good points is  $G_j \cup G_{-k}$ .*

*Proof.* Denote the set of bad points for the  $h$ th monomial of  $\neg(m_k)$  by  $B_{-k}^h$ , and denote the number of monomials composing  $\neg(m_k)$  by  $p$ . We clearly have  $B_{-k} = \bigcup_{h=1}^p B_{-k}^h$ . Moreover, by Theorem 4.1, for each single monomial product constituting  $m_j \diamond \neg(m_k)$ , we have that the corresponding set of bad points is  $B_j \cap B_{-k}^h$ . Therefore, the set of bad points of the entire  $m_j \diamond \neg(m_k)$  is  $\bigcup_{h=1}^p (B_j \cap B_{-k}^h)$ , which is  $B_j \cap B_{-k}$ . As a consequence, also the set of good points of the entire  $m_j \diamond \neg(m_k)$  is  $G_j \cup G_{-k}$ .  $\square$

**THEOREM 4.4.** *Consider an arbitrary Boolean formula  $\mathcal{F}$  in normal form representing the Boolean function  $f(x_1, \dots, x_n)$ . If an arbitrary monomial  $m_j \in \mathcal{F}$  is multiplied by the negation  $\neg(m_k)$  of another arbitrary monomial  $m_k \in \mathcal{F}$ , the new Boolean formula obtained  $\mathcal{F}'$  still represents the same  $f(x_1, \dots, x_n)$ .*

*Proof.* It is sufficient to prove that the sets  $B$  and  $G$  are the same for  $\mathcal{F}$  and  $\mathcal{F}'$ . As can be observed in Figure 4.1, monomial  $m_k$  determines in  $\mathbb{B}^n$  a partition in  $B_k$  and  $G_k$ . Its negation  $\neg(m_k)$  determines a partition  $B_{-k} = G_k$  and  $G_{-k} = B_k$ . Now multiply another monomial  $m_j$  by  $\neg(m_k)$ , to obtain new monomials  $m_j \diamond \neg(m_k)$ ; add  $m_j \diamond \neg(m_k)$  and remove  $m_j$  from the formula  $\mathcal{F}$ , thus obtaining  $\mathcal{F}'$ . The set  $G'_j$  corresponding to  $m_j \diamond \neg(m_k)$ , by Theorem 4.3, is  $G_j \cup G_{-k}$ , which is  $\supseteq G_j$ . So, the set of good points  $G$  for the formula  $\mathcal{F}$ , which is the intersection of all the  $G_j$ , cannot decrease. We now prove that  $G$  cannot increase. It could only increase in the area of  $G_{-k}$ , since  $G'_j = G_j \cup G_{-k}$ . However, all points of  $G_{-k}$  are forbidden by the fact that  $G \subseteq G_k$ . Hence,  $G$  is the same for  $\mathcal{F}$  and  $\mathcal{F}'$ , and therefore  $B$  also remains the same. The thesis follows.  $\square$

*Example 4.5.* Given the formula composed of two monomials  $m_1$  and  $m_2$ :

$$m_1 = (x_1 \top \neg x_2 \top x_5) \perp (\neg x_2 \top x_3 \top x_4) = m_2, \quad (4.6)$$

the defined sets of noncommon literals are

$$D_1 = (x_1 \top x_5), \quad D_2 = (x_3 \top x_4). \quad (4.7)$$

Their cardinality is the same. We choose  $D_1$ , and the orthogonal negation of the monomial corresponding to  $D_1$  is the following:

$$(\neg x_1) \perp (x_1 \top \neg x_5). \quad (4.8)$$

By using the orthogonalization operation, the above formula becomes

$$(x_1 \top \neg x_2 \top x_5) \perp ((\neg x_1) \diamond (\neg x_2 \top x_3 \top x_4)) \perp ((x_1 \top \neg x_5) \diamond (\neg x_2 \top x_3 \top x_4)) \quad (4.9)$$

which is the following orthogonal formula:

$$(x_1 \top \neg x_2 \top x_5) \perp (\neg x_1 \neg x_2 \top x_3 \top x_4) \perp (x_1 \top \neg x_2 \top x_3 \top x_4 \top \neg x_5). \quad (4.10)$$

**THEOREM 4.6.** *Given an arbitrary Boolean formula  $\mathcal{F}$  in normal form, representing the Boolean function  $f(x_1, \dots, x_n)$ , it is always possible to transform it into an orthogonal normal form  $\mathcal{O}$  still representing same  $f(x_1, \dots, x_n)$ .*

*Proof.* The (constructive) proof is given by the above orthogonalization operation, since that is a general procedure capable of orthogonalizing any two monomials. Define the orthogonalization of two monomials by means of procedure such as a *step*. Given therefore an arbitrary formula with  $m$  monomials, by iterating this orthogonalization operation to exhaustion until every pair of monomials is orthogonal, the orthogonal form is obtained in a finite number of steps, at most  $\binom{m}{2}$ .  $\square$

## 5. Improvements on the basic procedure

Unfortunately, by repeatedly applying above operation to exhaustion, the size of the formula tends to exponentially increase. As remarked above, this is not surprising, since the process of orthogonalization makes easy an NP-complete problem like satisfiability. Hence, the NP complexity [9] can be seen as being absorbed by the orthogonalization process; so it is unlikely that the orthogonalization process can be made inexpensive. However, improvements on the above basic procedure, with the aim of minimizing the size of the formula both in the final result and during the computation, are possible, as follows.

**5.1. Absorption of implied monomials.** Consider two generic monomials  $m_j$  and  $m_k$  appearing in the same formula  $\mathcal{F}$ , which represents the Boolean function  $f(x_1, \dots, x_n)$ , as follows:

$$m_j = \left( \prod_{i \in P_j} x_i \top \prod_{i \in N_j} \neg x_i \right), \quad m_k = \left( \prod_{i \in P_k} x_i \top \prod_{i \in N_k} \neg x_i \right). \quad (5.1)$$

If  $P_j \subseteq P_k$  and  $N_j \subseteq N_k$ , monomial  $m_k$  is logically implied by  $m_j$  [11, 19], and can therefore be removed from  $\mathcal{F}$  obtaining a smaller formula  $\mathcal{F}'$  still representing the same  $f(x_1, \dots, x_n)$ . This operation is applied in order to reduce the number of monomials in the formula.

**5.2. Synthesis resolution.** This operation is a special case of the general operation called resolution [5, 17] in the case of CNF, and consensus [15] in the case of DNF. Suppose a formula  $\mathcal{F}$  is given and that it represents the Boolean function  $f(x_1, \dots, x_n)$ . If  $\mathcal{F}$  contains two monomials, which are identical except for one literal  $x_s$  appearing positive in one monomial and negative in the other, as follows:

$$m_j = \left( x_s \top \prod_{i \in P_j} x_i \top \prod_{i \in N_j} \neg x_i \right), \quad m_k = \left( \prod_{i \in P_j} x_i \top \prod_{i \in N_j} \neg x_i \top \neg x_s \right), \quad (5.2)$$

their resolvent [17]  $m_r$ , reported below, can be added to  $\mathcal{F}$ , thus obtaining a new formula  $\mathcal{F}'$  which still represents the same Boolean function  $f(x_1, \dots, x_n)$ :

$$m_r = \left( \prod_{i \in P_j} x_i \top \prod_{i \in N_j} \neg x_i \right). \quad (5.3)$$

Moreover, their resolvent logically implies both its parents  $m_j$  and  $m_k$ , hence they can be removed from the formula, in order to obtain a new formula  $\mathcal{F}''$  which still represents the same Boolean function  $f(x_1, \dots, x_n)$ . This operation helps in reducing the number of monomials in the formula.

Finally, our aim being not to excessively increase the size of the formula, for each orthogonalization step  $t$ , we define the *quality*  $q_t$  of such step as the number  $o_t$  of clauses orthogonalized by such step divided by the number  $n_t$  of new clauses created by such step:  $q_t = o_t/n_t$  (it can be computed in advance). In our procedure, we set an initial quality limit  $q_{\text{limit}}$  in order to initially perform the most convenient basic orthogonalizations. During iterations, at the beginning of each phase of basic orthogonalization steps, if no steps respecting current limit are possible, current limit is decreased.

*Complete orthogonalization procedure.*

*Input:* an arbitrary Boolean formula  $\mathcal{F}$  in CNF or DNF.

*Output:* an equivalent Boolean formula  $\mathcal{F}'$  in OCNF or ODNE.

*Repeat*

- (a) If the current formula is orthogonal, *stop*.
- (b) Else if  $q_{\text{limit}}$  allows no basic orthogonalization steps, decrease  $q_{\text{limit}}$ .
- (c) Perform all basic orthogonalization steps of quality  $q \geq q_{\text{limit}}$ .
- (d) Perform all possible synthesis resolutions.
- (e) Perform all possible absorptions.

Table 6.1. Orthogonalization procedure on artificially generated CNF formulae.

Problem	$m_{\text{init}}$	$m_{\text{max}}$	$m_{\text{ortho}}$	Time
uf20-01	91	1443	130	1.93
uf20-02	91	912	100	1.53
uf20-03	91	859	132	1.59
uf20-04	91	861	134	1.09
uf20-05	91	341	31	0.11
uf20-06	91	723	40	0.58
uf20-07	91	506	85	0.49
uf20-08	91	895	136	1.67
uf20-09	91	1669	144	4.49
uf20-010	91	1068	128	1.72
uf20-011	91	418	130	0.35
uf20-012	91	1117	190	2.58
uf20-013	91	784	65	0.74
uf20-014	91	947	167	2.19
uf20-015	91	980	120	1.78
uf20-016	91	954	102	1.47
uf20-017	91	787	109	1.27
uf20-018	91	1530	105	2.44
uf20-019	91	861	70	1.09
uf20-020	91	1335	98	2.64
uf20-021	91	870	73	0.59
uf20-022	91	700	79	0.81
uf20-023	91	1575	211	3.40
uf20-024	91	837	171	1.63
uf20-025	91	935	82	1.31
uf20-026	91	836	59	1.05
uf20-027	91	888	63	0.78
uf20-028	91	740	93	1.03
uf20-029	91	618	66	0.70
uf20-030	91	509	90	0.61

## 6. Testing of the procedure

The algorithm was tested on artificially generated CNF formulae obtained from the SATLIB collection of the Darmstadt University of Technology. They represent 3-SAT problems. Table 6.1 reports the number of monomials of the original formula ( $m_{\text{init}}$ ),

the number of monomials in the orthogonalized formula produced ( $m_{\text{ortho}}$ ), the maximum number of monomials reached by the formula during the orthogonalization process ( $m_{\text{max}}$ ), and computational time in seconds on a PC Pentium IV 1.7 GHz. Such testing is intended solely as a study of the behavior of the orthogonalization procedure, since it currently does not constitute, from the practical point of view, a fast alternative for solving satisfiability problems.

It can be observed that the number of monomials in the orthogonalized formula generally increases, although not always. Moreover, intermediate formulae contain a much larger number of monomials. This turns out to be a general rule in performing similar operations. However, there are practical applications where the advantages of having the orthogonal form completely surmount the disadvantage of such size increase.

## 7. Conclusions

The orthogonal form of a Boolean formula has remarkable properties. Several difficult problems become easy when in orthogonal form. A general procedure for the orthogonalization of an arbitrary CNF or DNF is developed. A unified and coherent notation for representing at the same time CNF and DNF is therefore introduced. The procedure is proved to always produce the orthogonal form (OCNF or ODNF) in a finite number of steps. The problem is indeed computationally demanding. As predictable, in the initial phase of the procedure, the size of the formula tends to exponentially increase. On the other hand, the size of the formula decreases again when approaching the final phase. In spite of this size growth, orthogonalization appears to be the preferable way to solve some practical problems, for instance, in the field of reliability theory. Some computational complexity implications of the orthogonalization process are analyzed.

## Acknowledgments



The author wishes to thank Peter L. Hammer and Endre Boros for introducing the discussed problems to his attention.

## References

- [1] H. K. Büning and T. Lettmann, *Aussagenlogik: Deduktion und Algorithmen*, Leitfäden und Monographien der Informatik, B. G. Teubner, Stuttgart, 1994, reprinted in English.
- [2] M. O. Ball and J. S. Provan, *Disjoint products and efficient computation of reliability*, *Oper. Res.* **36** (1988), no. 5, 703–715.
- [3] R. E. Barlow, C. A. Clarotti, and F. Spizzichino, eds., *Reliability and decision making*, Chapman & Hall, London, 1993, Proceedings of the Conference Held at the University of Siena, Siena, 1990.
- [4] R. E. Barlow and F. Proschan, *Statistical Theory of Reliability and Life Testing*, International Series in Decision Processes. Series in Quantitative Methods for Decision Making, Holt, Rinehart and Winston, New York, 1975.
- [5] A. Blake, *Canonical expressions in Boolean algebra*, Ph.D. thesis, University of Chicago, Chicago, 1937.
- [6] E. Boros, Y. Crama, O. Ekin, P. L. Hammer, T. Ibaraki, and A. Kogan, *Boolean normal forms, shellability, and reliability computations*, *SIAM J. Discrete Math.* **13** (2000), no. 2, 212–226.

- [7] E. Boros, Y. Crama, P. L. Hammer, and M. Saks, *A complexity index for satisfiability problems*, SIAM J. Comput. **23** (1994), no. 1, 45–49.
- [8] V. Chvátal, *Resolution search*, Discrete Appl. Math. **73** (1997), no. 1, 81–99.
- [9] M. R. Garey and D. S. Johnson, *Computers and Intractability*, A Series of Books in the Mathematical Sciences, W. H. Freeman and Company, California, 1979.
- [10] J. Gu, P. W. Purdom, J. Franco, and B. W. Wah, *Algorithms for the satisfiability (SAT) problem: a survey*, Satisfiability Problem: Theory and Applications (Piscataway, NJ, 1996), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 35, American Mathematical Society, Rhode Island, 1997, pp. 19–151.
- [11] P. L. Hammer and S. Rudeanu, *Boolean Methods in Operations Research and Related Areas*, Econometrics and Operations Research, vol. 7, Springer-Verlag, New York, 1968.
- [12] K. Iwama, *CNF-satisfiability test by counting and polynomial average time*, SIAM J. Comput. **18** (1989), no. 2, 385–391.
- [13] D. W. Loveland, *Automated Theorem Proving: a Logical Basis*, Fundamental Studies in Computer Science, vol. 6, North-Holland, Amsterdam, 1978.
- [14] J. S. Provan, *Boolean decomposition schemes and the complexity of reliability computations*, Reliability of Computer and Communication Networks (New Brunswick, NJ, 1989), DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 5, American Mathematical Society, Rhode Island, 1991, pp. 213–228.
- [15] W. V. Quine, *A way to simplify truth functions*, Amer. Math. Monthly **62** (1955), 627–631.
- [16] K. G. Ramamurthy, *Coherent Structures and Simple Games*, Theory and Decision Library. Series C: Game Theory, Mathematical Programming and Operations Research, vol. 6, Kluwer Academic, Dordrecht, 1990.
- [17] J. A. Robinson, *A machine-oriented logic based on the resolution principle*, J. Assoc. Comput. Mach. **12** (1965), 23–41.
- [18] H. Scheffé, *The Analysis of Variance*, John Wiley & Sons, New York, 1959.
- [19] I. Wegener, *The Complexity of Boolean Functions*, Wiley-Teubner Series in Computer Science, John Wiley & Sons, Chichester, 1987.

Renato Bruni: Dipartimento di Informatica e Sistemistica, Università di Roma “La Sapienza,” Via Michelangelo Buonarroti 12, Roma 00185, Italy  
*E-mail address:* bruni@dis.uniroma1.it



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

