

Clinical Study

Exercise-Induced Bronchoconstriction and Exercise-Induced Respiratory Symptoms in Nurses

Jordan Minov,¹ Jovanka Karadzinska-Bislimovska,¹ Kristin Vasilevska,²
Snezana Risteska-Kuc,¹ Saso Stoleski,¹ and Dragan Mijakoski¹

¹Department of Cardiorespiratory Functional Diagnostics, Institute for Occupational Health of R. Macedonia—WHO Collaborating Center and GA2LEN Collaborating Center, II Makedonska Brigada 43, 1000 Skopje, Macedonia

²Institute for Epidemiology and Biostatistics, 1000 Skopje, Macedonia

Correspondence should be addressed to Jordan Minov, minovj@hotmail.com

Received 20 December 2010; Accepted 28 February 2011

Academic Editor: Donald H. Beezhold

Copyright © 2011 Jordan Minov et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In order to assess prevalence and characteristics of exercise-induced respiratory symptoms (EIRs) and exercise-induced bronchoconstriction (EIB) in health care workers, we performed a cross-sectional study including 48 female nurses from primary care settings and an equal number of female office workers studied as a control group. The evaluation of examined groups included completion of a questionnaire, skin prick tests to common inhalant allergens, spirometry, and exercise and histamine challenge. We found a similar prevalence of EIRs and EIB in both groups. EIB was closely related to asthma, atopy, family history of asthma, and positive histamine challenge in either group, while the association between EIB and daily smoking in nurses was of borderline statistical significance. Bronchial reaction to exercise was significantly higher in nurses than in controls with EIB. With the exception of exercise induced wheezing, EIRs were weakly associated with EIB in both groups with a large proportion of false positive results.

1. Introduction

Exercise-induced bronchoconstriction (EIB), also referred to as exercise-induced asthma (EIA), is a manifestation of bronchial hyperresponsiveness (BHR) that occurs in the majority of patients with current symptomatic asthma, especially in the patients with moderately to severely increased responsiveness [1–3]. The current thinking about mechanisms by which EIB develops emphasizes the loss of heat and/or water from the airways during exercise that leads to release of proinflammatory mediators [4]. Airborne particles and pollutants, as well as airborne allergens, are considered as stimulants that contribute to EIB [5]. A fish oil supplementation may have a protective effect on EIB, which is probably attributed to its anti-inflammatory properties [6].

Results from several studies indicated that BHR prevalence is higher in females than in males [7, 8]. The mechanisms responsible for a higher susceptibility of the airways in females to nonspecific stimuli include lower airway calibre, lower body weight, greater cholinergic irritability, and hormonal factors [9].

On the other side, data from the studies carried out in the last two decades suggest an increased risk for asthma among health care workers, yet only a few specific determinants have been elucidated [10–13]. As Delclos et al. [14] have suggested, the contribution of occupational exposures to respiratory impairment and asthma in health care professionals is not trivial, meriting both implementation of appropriate controls and further studies.

To our knowledge, so far, there is no study assessing exercise-induced respiratory symptoms (EIRs) and EIB in health care professionals. In the present study, we assessed effects of occupational exposure on EIRs and EIB among health care workers by comparison of their prevalence and characteristics between females working as nurses in primary care settings and female office workers.

2. Materials and Methods

2.1. Study Design and Setting. A cross-sectional survey was carried out in a university research laboratory, that is,

Department of Cardiorespiratory Functional Diagnostics at the Institute for Occupational Health of R. Macedonia, Skopje—WHO Collaborating Center for Occupational Health and GA²LEN Collaborating Center.

2.2. Subjects. We examined 48 females aged 24 to 51 years (mean age 37.8 ± 7.4) working as nurses in primary care settings with duration of employment 5 to 25 years (mean duration 14.7 ± 5.7).

The work shifts of the nurses lasted 8 hours per day, and their working tasks included completion of the medical documentation, assistance in medical interventions, administration of parenteral and aerosolized medications, and medical instruments cleaning. The workplace exposure included several types of cleaning products, disinfectants, adhesives, solvents, latex, and medications, some of which are in spray form. According to the classification of occupational muscular work, their work was classified as a light muscular work [15]. During the work shift, they use protective clothing, masks, and powdered latex gloves.

In addition, an equal number of female office workers matched to nurses as a group by age and smoking status were studied as a control group. According to the classification of occupational muscular work, their work was classified as a sedentary work.

In either group, there were no subjects in whom exercise challenge or histamine challenge were contraindicated [16, 17], nor there were subjects with the upper respiratory viral infection within three weeks before the challenge test was performed. None of the subjects took asthma medications or antihistamines at least one month before the challenge tests and skin-prick tests. Daily smokers were asked to refrain from smoking at least 3 hours before testing.

2.3. Questionnaire. The questionnaire was designed using the proposed model of the National Jewish Medical and Research Center, Denver, USA [18].

Subjects were considered having exercise-induced respiratory symptoms (EIRs) if one or more symptoms were reported: coughing during or after exercise, wheezing during or after exercise, inability to get deep breath after exercise, noisy breathing after exercise, and chest tightness after exercise.

Detailed smoking history, asthma diagnosed by physician, family history of asthma and allergic diseases (taking into account the first-degree relatives), accompanying disease, and medication use were also evaluated.

Classification of smoking status was done according to the World Health Organization (WHO) guidelines on definitions of smoking status [19].

Daily smoker was defined as a subject who smoked at the time of the survey at least once a day, except on days of religious fasting. In daily smokers, lifetime cigarette smoking and daily mean of cigarettes smoked were evaluated. Pack years smoked (one pack year denotes one year of smoking 20 cigarettes per day) were calculated according to the actual recommendations [20].

Ex-smoker was defined as a formerly daily smoker who no longer smokes.

Passive smoking or exposure to environmental tobacco smoke (ETS) was defined as the exposure of a person to tobacco-combustion products from smoking by others [21].

2.4. Skin-Prick Tests. Skin-prick tests (SPTs) to common inhalant allergens were performed in all subjects on the volar part of the forearm using commercial allergen extracts (Torlak, Serbia, and Montenegro) of birch (5000 PNU), grass mixed (5000 PNU), plantain (5000 PNU), fungi mixed (4000 PNU), *Dermatophagoides pteronyssinus* (3000 PNU), dog hair (4000 PNU), cat fur (4000 PNU), and feathers mixed (4000 PNU). All tests included positive (1 mg/mL histamine) and negative (0.9% saline) controls. Prick tests were considered positive if the mean wheal diameter 20 min after allergen application was at least 3 mm larger than the size of the negative control [22]. Atopy was defined as the presence of at least one positive SPT [23].

2.5. Spirometry. Spirometry, including measures of forced vital capacity (FVC), forced expiratory volume in one second (FEV₁), FEV₁/FVC ratio, and maximal expiratory flow at 50%, 25%, and 25–75% of FVC (MEF₅₀, MEF₂₅, and MEF_{25–75}, resp.), was performed in all subjects using spirometer Ganshorn SanoScope LF8 (Ganshorn Medizin Electronic GmbH, Germany) with recording the best result from three measurements the values of FEV₁ of which were within 5% of each other. The results of spirometry were expressed as percentages of the predicted values according to the European Community for Coal and Steel (ECCS) norms [24].

2.6. Histamine Challenge. The histamine challenge test was performed according to the actual European Respiratory Society (ERS)/American Thoracic Society (ATS) recommendations [16, 17]. Concentrations of 0.5, 1, 2, 4, and 8 mg/mL histamine (Torlak, Beograd) were prepared by dilution with buffered saline. The doses of aerosol generated by Pari LC nebulizer with output rate 0.17 mL/min were inhaled by mouthpiece. Subjects inhaled increasing concentrations of histamine using a tidal breathing method until FEV₁ fell by more than 20% of its base value (provocative concentration 20—PC₂₀) or the highest concentration was reached.

According to the ATS recommendations, bronchial hyperresponsiveness (BHR) was categorized as moderate to severe BHR (PC₂₀ < 1.0 mg/mL), mild BHR (PC₂₀ = 1.0–4.0 mg/mL), and borderline BHR (PC₂₀ > 4.0 mg/mL) [17]. The test was considered positive if PC₂₀ was equal or less than 4 mg/mL [16, 17].

2.7. Exercise Challenge Tests. The constant submaximal exercise challenge test (ECT) was performed in all subjects using cycle ergometer Hellige-dynavit Meditronic 40 (Hellige GmbH, Germany). ECT was conducted in an air-conditioned room with ambient temperature of 20–25°C and relative air humidity of 50% or less. According to the actual recommendations, subjects exercised

TABLE 1: Demographics of the study subjects.

Variable	Nurses (<i>n</i> = 48)	Office workers (<i>n</i> = 48)
Age (years)	37.8 ± 7.4	39.1 ± 9.2
BMI (kg/m ²)	25.4 ± 3.9	26.7 ± 4.3
Duration of employment (years)	14.7 ± 5.7	15.4 ± 7.8
Asthma diagnosed by physician	3 (6.3%)	2 (4.2%)
Family history of asthma	4 (8.3%)	4 (8.3%)
Family history of allergies	6 (12.5%)	8 (16.6%)
Daily smokers	14 (29.2%)	15 (31.2%)
Smoking experience (years)	17.7 ± 5.8	19.4 ± 7.9
Cigarettes per day	14.4 ± 6.9	16.8 ± 8.3
Pack years smoked	12.5 ± 3.1	13.4 ± 3.8
Daily smokers with less than 12 pack years smoked	6 (12.5%)	7 (14.6%)
Ex-smokers	5 (10.4%)	4 (8.3%)
Passive smokers	8 (16.6%)	10 (20.8%)

Numerical data are expressed as mean value with standard deviation and frequencies as number and percentage of study subjects with certain variable.

BMI: body mass index; kg: kilogram; m: meter.

8–10 min achieving 90% of predicted maximal heart rate ($HR_{max} = 220 - \text{age}$) in the last 4 min of exercise [16, 17]. Heart rate was monitored continuously throughout the exercise and for 5 minutes after its completion from a three-lead electrocardiographic configuration. The measurements of FEV₁ were performed before and 1, 3, 5, 7, 10, and 15 min after the exercise with inhaled bronchodilator (200 mcg salbutamol) application upon completion of the protocol.

The response to exercise was expressed as fall index FEV₁ ($100 \times [\text{pre-exercise FEV}_1 - \text{lowest postexercise FEV}_1] / \text{pre-exercise FEV}_1$). EIB was defined as fall index FEV₁ ≥ 10% [16].

2.8. Statistical Analysis. Continuous variables were expressed as mean values with standard deviation (SD) whereas the nominal variables as numbers and percentages. Analyses of the data involved testing the differences in prevalence, comparison of the means, and testing the association between EIRs and EIB and studied variables. Chi-square test was used for testing difference in the prevalence. Comparison of spirometric measurements and fall index FEV₁ values was performed by independent samples *t*-test. Chi-square test (or Fisher's exact test where appropriate) was used for testing association between EIRs and EIB and studied variables. A *P* value less than .05 was considered as statistically significant. Statistical analysis was performed using the Statistical Package for the Social Sciences (SPSS) version 11.0 for Windows.

3. Results

Demographic characteristics of the study subjects were similar in both examined groups (Table 1).

TABLE 2: Prevalence of EIRs in nurses and controls.

EIRs	Nurses (<i>n</i> = 48)	Office workers (<i>n</i> = 48)	<i>P</i> value*
Any exercise-induced respiratory symptom	19 (39.6%)	17 (35.4%)	.673
Cough	11 (22.9%)	13 (27.1%)	.637
Inability to get deep breath	14 (29.1%)	11 (22.9%)	.585
Wheezing	5 (10.4%)	4 (8.3%)	1.000
Chest tightness	9 (18.8%)	8 (16.7%)	.726
Noisy breathing	4 (8.3%)	6 (12.5%)	.740

Data are expressed as number and percentage of study subjects with certain variable.

EIRs: exercise-induced respiratory symptoms.

*Tested by Chi-square test (or Fisher's exact test where appropriate).

TABLE 3: Spirometric parameters in the study subjects.

Spirometric parameter	Nurses (<i>n</i> = 48)	Office workers (<i>n</i> = 48)	<i>P</i> value*
FVC (% pred)	88.9 ± 9.8	91.6 ± 10.4	.102
FEV ₁ (% pred)	84.3 ± 7.9	86.2 ± 9.6	.180
FEV ₁ /FVC%	76.1 ± 4.9	78.4 ± 5.8	.126
MEF ₅₀ (% pred)	64.8 ± 12.7	68.1 ± 9.8	.083
MEF ₂₅ (% pred)	60.8 ± 10.1	64.9 ± 8.9	.069
MEF ₂₅₋₇₅ (% pred)	72.8 ± 14.7	75.1 ± 10.9	.094

Data are expressed as mean value with standard deviation.

FVC: forced vital capacity; FEV₁: forced expiratory volume in one second; MEF₅₀, MEF₂₅, and MEF₂₅₋₇₅: maximal expiratory flow at 50%, 25%, and 25–75% of FVC, respectively; % pred: % of predicted value.

*Compared by independent samples *t* test.

Prevalence of EIRs, total and individual, was similar in both examined groups. Inability to get deep breath after exercise and cough during or after exercise was the most frequent EIRs in either group (Table 2).

EIRs were nonsignificantly associated with age and smoking in both examined groups. The association between EIRs in nurses and duration of employment was also nonsignificant.

Prevalence of subjects with positive SPT to common inhalant allergens was similar in both nurses and controls (33.3% versus 37.5%, *P* = .670; Chi-square test). Mite sensitization was the most important individual common allergen with no statistical difference between sensitized subjects in both groups (22.9% versus 25.0%, *P* = .811; Chi-square test).

Spirometric parameters were lower in nurses, but statistical significance was not found for any parameter (Table 3). Spirometric parameters were nonsignificantly lower in the subjects with asthma diagnosed by physician as compared to nonasthmatics in both nurses and controls.

Prevalence of overall subjects with BHR was nonsignificantly higher in nurses (12.5% versus 8.3%; *P* = .740),

TABLE 4: Characteristics of the ECT in nurses and controls.

BHR	Nurses (<i>n</i> = 48)	Office workers (<i>n</i> = 48)	<i>P</i> value*
Moderate to severe BHR	2 (4.2%)	1 (2.1%)	1.000
Mild BHR	1 (2.1%)	1 (2.1%)	1.000
Borderline BHR	3 (6.3%)	2 (4.2%)	1.000

Data are expressed as mean value with standard deviation.

BHR: bronchial hyperresponsiveness.

*Compared by Fisher's exact test.

whereas the prevalence of subjects with moderate to severe and mild BHR, that is, prevalence of subjects with positive histamine challenge, was similar in both examined groups (Table 4).

We found similar prevalence of EIB in both nurses and controls (8.3% versus 6.3%, $P = 1.000$; Fisher's exact test). The EIB severity, expressed as fall index FEV_1 , was significantly higher in nurses (28.1% versus 22.7%, $P = .033$; independent-samples *t*-test). Characteristics of the ECT performed in study subjects are shown Table 5.

EIB in both examined groups was significantly related to asthma diagnosed by physician, positive family history for asthma and allergies, and positive histamine challenge, whereas association with other variables was nonsignificant. Association between EIB and daily smoking in nurses was of borderline statistical significance ($P = .062$; Fisher's exact test), while association between EIB and pack years smoked (less or more than 12) was nonsignificant ($P = .097$; Fisher's exact test). These associations in controls were statistically nonsignificant.

Association between EIB and exercise-induced respiratory symptoms, with exception of exercise-induced wheezing in both nurses ($P = .037$; Fisher's exact test) and controls ($P = .034$; Fisher's exact test), was statistically nonsignificant. The frequency of false positive results was high in both nurses (84.3%) and controls (88.2%).

4. Discussion

According to the recent data, occupational exposures in health care professionals increase the risk of work-related asthma. Medical instruments cleaning, general cleaning, use of solvents/adhesives in patient care, use of powdered latex gloves, and aerosolized medication administration were identified as occupational risk factors associated with the development of asthma in nurses [10, 14, 25].

On the other hand, EIB is a common condition close related to asthma that is often unrecognized and uncontrolled leading affected subjects to avoid general and occupational physical activities and sports. We performed the present study on EIB among nurses in primary care settings as a continuum of our investigations on the effects of specific occupational exposures on the EIB occurrence and characteristics [26–28]. According to the results of several studies [11–13], the lowest risk of respiratory impairment and asthma was found in administrative workers, so this “unexposed” occupation was used as a control group.

TABLE 5: Characteristics of the ECT in nurses and controls.

Variable	Nurses (<i>n</i> = 48)	Controls (<i>n</i> = 48)
Exercise load (Watt)	102.1 ± 20.3	106.3 ± 16.1
Positive ECT	4 (8.3%)	3 (6.3%)
ΔFEV_1 in the subjects with EIB (%)	28.1 ± 3.4	22.7 ± 1.5
Time of EIB occurrence (minutes after exercise)	6.1 ± 2.1	6.9 ± 2.7

Numerical data are expressed as mean value with standard deviation and frequencies as number and percentage of study subjects with certain variable.

ECT: exercise challenge test; ΔFEV_1 : a fall in FEV_1 of pre-exercise value; FEV_1 : forced expiratory volume in 1 second; EIB: exercise-induced bronchoconstriction.

In the present study, both examined groups included subjects with similar demographic characteristics. In either group, there was a large proportion of daily and passive smokers similar to its prevalence among females in R. Macedonia documented in our previous studies [29, 30]. The prevalence of ex-smokers in both groups was low, suggesting insufficient smoking cessation activities. The situation in the developed countries seems to be somewhat different. In the study conducted in 12 European countries as well as Australia and the USA, Janson et al. [31] reported that both active and passive smoking rates have declined since the early 1990s but indicated lower quitting rates and higher risk of passive smoking among people with fewer qualifications and less skilled occupation groups.

We found high prevalence of EIRs in both examined groups that is similar to the findings of several studies which investigated EIB in different subpopulations of both sexes [32, 33] as well as to the findings of our studies among workers with different occupational exposures [26–28]. The prevalence of atopy and the pattern of allergic sensitization to common aeroallergens in both examined groups was comparable to that we had previously observed among adults in R. Macedonia [34, 35]. All spirometric parameters were lower in nurses, but statistical significance was not achieved for any of them. The prevalence of BHR was nonsignificantly higher in nurses than in office workers that is similar to the findings obtained in our previous studies on BHR prevalence among workers with specific occupational exposures (herbal tea processors, cooks, and cleaners) and office workers as a control group [36, 37].

Several studies indicated that the occurrence of EIB depends on degree of bronchial hyperresponsiveness (alias underlying chronic inflammation), exercise intensity, and ambient conditions [38, 39]. There are many studies about EIB occurrence in selected groups of general population (children, school children, adolescents, and recruits) as well as in recreative and elite athletes. On the contrary, there is a limited number of studies on EIB associated with specific workplace exposures. The EIB prevalence in elite athletes varies from 12% of basketball players to 55% of cross-country skiers [40, 41]. In the present study, we found similar EIB prevalence in both nurses and controls (8.3% and 6.3%, resp.). According to the results of our previous studies, the

EIB prevalence among workers with specific occupational exposures ranged from 6.4% in herbal tea processors, 6.9% in bakers, 7.1% in agricultural workers, 8.9% in textile workers, to 9.3% in agricultural workers. Bronchial reaction to exercise in the subjects with EIB was significantly higher in nurses than in controls. Significantly higher bronchial reaction to exercise in comparison to office workers we also found in ECT-positive female cleaners, whereas the difference in mean fall index FEV₁ did not differ significantly between workers exposed to organic dusts and office workers with EIB [26–28]. This difference may be due to the presence of the study subjects of both sexes among workers exposed to organic dusts as well as to the different occupational exposures (i.e., dominant exposure to chemical compounds in cleaners and nurses).

We found significant association between EIB and asthma, family history of asthma, and atopy in both examined groups. Contribution of genetic factor in the EIB development is confirmed in a number of studies [39, 42, 43]. We also found a significant association between positive ECT and positive histamine challenge in both examined groups. Data from the studies which compared results of two bronchial challenge types are somewhat inconsistent. Some authors reported significant association between the results of exercise and histamine challenge [26–28, 44]. On the contrary, other authors reported a weak association, which was explained by different pathomechanisms of BHR to histamine and EIB [45]. Correlation between EIB and daily smoking was nonsignificant in controls, whereas in nurses, it was of borderline statistical significance. A similar finding, suggesting possible interaction of tobacco smoke and occupational exposure in EIB development, was obtained in our previous study on EIB in female cleaners [28].

In the present study, there was no positive association in both examined groups between EIB and overall and individual EIRs with exception of exercise-induced wheezing. This finding, confirmed in a number of studies, was not unexpected, as the EIRs may be triggered by many conditions and diseases other than BHR (e.g., physical unfit, medical side effect of angiotensin-converting enzyme inhibitors and beta-blockers, anxiety, vocal cord dysfunction, arterial hypertension, gastroesophageal reflux, etc.) [26–28, 46–48].

The present study has some limitations. First, relatively smaller number of the subjects in the study groups could have certain implications on the data obtained and its interpretation. Second, we did not perform SPT to workplace allergens (e.g., latex), so we could not document relationship between sensitization to workplace allergens and EIB. Third, environmental measurements were not performed, so we could not document the effect of the type and the level of exposure on EIB. The strength of the study is the extensive examination of lung function in the study subjects with the possibility for comparison of the results of different tests.

5. Conclusions

In conclusion, in a cross-sectional study including nurses and office workers, we found a similar prevalence of EIRs

and EIB in both examined groups. EIB was strongly related to asthma in both nurses and controls. In addition, EIB was closely related to atopy, family history of asthma, and positive histamine challenge in either group as well as to daily smoking in nurses. Bronchial reaction to exercise in ECT-positive nurses was significantly higher than in ECT-positive controls. EIRs were weakly associated with EIB in both examined groups, with a large proportion of false positive results. Our study confirms the need of regular medical examinations in order to identify affected workers and to implement adequate preventive measures.

References

- [1] P. Palange, V. Brusasco, L. Delgado, and S. Del Giacco, "Exercise and airway physiology: interactions with immune and allergic responses," *European Respiratory Monograph*, vol. 10, no. 33, pp. 10–18, 2005.
- [2] C. L. Hermansen and J. T. Kirchner, "Identifying exercise-induced bronchospasm: treatment hinges on distinguishing it from chronic asthma," *Postgraduate Medicine*, vol. 115, no. 6, pp. 15–25, 2004.
- [3] J. Macan, T. Klepac, I. Busljeta, D. Plavec, and B. Kanceljak-Macan, "Exercise-induced bronchospasm and its prevention," *Lijecnicki Vjesnik*, vol. 122, no. 9–10, pp. 239–245, 2000.
- [4] K. W. Rundell, B. A. Spiering, J. M. Baumann, and T. M. Evans, "Bronchoconstriction provoked by exercise in a high-particulate-matter environment is attenuated by montelukast," *Inhalation Toxicology*, vol. 17, no. 2, pp. 99–105, 2005.
- [5] W. W. Storms, "Exercise-induced asthma: diagnosis and treatment for the recreational or elite athlete," *Medicine and Science in Sports and Exercise*, vol. 31, no. 1, pp. S33–S38, 1999.
- [6] T. D. Mickleborough, M. R. Lindley, A. A. Ionescu, and A. D. Fly, "Protective effect of fish oil supplementation on exercise-induced bronchoconstriction in asthma," *Chest*, vol. 129, no. 1, pp. 39–49, 2006.
- [7] B. Leynaert, J. Bousquet, C. Henry, R. Liard, and F. Neukirch, "Is bronchial hyperresponsiveness more frequent in women than in men?: a population-based study," *American Journal of Respiratory and Critical Care Medicine*, vol. 156, no. 5, pp. 1413–1420, 1997.
- [8] P. Paoletti, L. Carrozzi, G. Viegi et al., "Distribution of bronchial responsiveness in a general population: effect of sex, age, smoking, and level of pulmonary function," *American Journal of Respiratory and Critical Care Medicine*, vol. 151, no. 6, pp. 1770–1779, 1995.
- [9] G. Wassmer, R. A. Jörres, J. Heinrich, M. Wjst, P. Reitmeir, and H. E. Wichmann, "The association between baseline lung function and bronchial responsiveness to methacholine," *European Journal of Medical Research*, vol. 2, no. 2, pp. 47–54, 1997.
- [10] A. A. Arif, G. L. Delclos, and C. Serra, "Occupational exposures and asthma among nursing professionals," *Occupational and Environmental Medicine*, vol. 66, no. 4, pp. 274–278, 2009.
- [11] T. P. Ng, C. Y. Hong, L. G. Goh, M. L. Wong, K. T. C. Koh, and S. L. Ling, "Risks of asthma associated with occupations in a community-based case-control study," *American Journal of Industrial Medicine*, vol. 25, no. 5, pp. 709–718, 1994.
- [12] M. Kogevinas, J. M. Antó, J. B. Soriano, A. Tobias, and P. Burney, "The risk of asthma attributable to occupational

- exposures: a population- based study in Spain,” *American Journal of Respiratory and Critical Care Medicine*, vol. 154, no. 1, pp. 137–143, 1996.
- [13] D. Fishwick, N. Pearce, W. D’Souza et al., “Occupational asthma in New Zealanders: a population based study,” *Occupational and Environmental Medicine*, vol. 54, no. 5, pp. 301–306, 1997.
- [14] G. L. Delclos, D. Gimeno, A. A. Arif et al., “Occupational risk factors and asthma among health care professionals,” *American Journal of Respiratory and Critical Care Medicine*, vol. 175, no. 7, pp. 667–675, 2007.
- [15] J. R. BROWN and G. P. CROWDEN, “Energy expenditure ranges and muscular work grades,” *British Journal of Industrial Medicine*, vol. 20, pp. 277–283, 1963.
- [16] P. J. Sterk, L. M. Fabbri, P. H. Quanjer et al., “Airway responsiveness. Standardized challenge testing with pharmacological, physical and sensitizing stimuli in adults. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society,” *The European respiratory journal. Supplement*, vol. 16, no. 6, pp. 53–83, 1993.
- [17] American Thoracic Society, “Guidelines for methacholine and exercise challenge testing-1999,” *American Journal of Respiratory and Critical Care Medicine*, vol. 161, no. 1, pp. 309–329, 2000.
- [18] D. Parks, “Exercise-Induced Asthma. National Jewish Medical and Research Center,” <http://library.nationaljewish.org/MSU/10n5MSU-EIA>.
- [19] World Health Organization, *Guidelines for Controlling and Monitoring the Tobacco Epidemic*, WHO, Geneva, Switzerland, 1998.
- [20] “Smoking Pack-Years,” <http://smokingpackyears.com>.
- [21] US Department of Health and Human Services, “The health consequences of smoking: chronic obstructive pulmonary disease. A report of the Surgeon General,” Tech. Rep. 84-50 205, US Department of Health and Human Services, Public Health Service, Office of the Assistant for Health, Office of Smoking and Health. DHHS, 1984.
- [22] The European Academy of Allergology and Clinical Immunology, “Position paper: allergen standardization and skin tests,” *Allergy*, vol. 48, no. 14, pp. 48–82, 1993.
- [23] A. J. Frew, “Allergic basis of asthma,” *European Respiratory Monthly*, vol. 23, no. 8, pp. 74–83, 2003.
- [24] P. H. H. Quajner, “Standardization of lung function tests-1993 update. Report Working Party for the European Community for Steel and Coal. Official Statement of the European Respiratory Society,” *European Respiratory Journal*, vol. 16, no. 1, pp. 1–100, 1993.
- [25] M. Crippa, L. Balbiani, A. Baruffini et al., “Consensus Document. Update on latex exposure and use of gloves in Italian health care settings,” *Medicina del Lavoro*, vol. 99, no. 5, pp. 387–399, 2008.
- [26] J. Minov, J. Karadžinska-Bislimovska, S. Risteska-Kuc, and S. Stoleski, “Exercise-induced bronchoconstriction and exercise-induced respiratory symptoms in workers exposed to tea dust,” *Arhiv za Higijenu Rada I Toksikologiju*, vol. 56, no. 4, pp. 317–326, 2005.
- [27] J. Minov, J. Karadžinska-Bislimovska, K. Vasilevska, S. Risteska-Kuč, and S. Stoleski, “Exercise-induced bronchoconstriction in textile and agricultural workers and in bakers,” *Arhiv za Higijenu Rada I Toksikologiju*, vol. 57, no. 4, pp. 379–386, 2006.
- [28] J. Minov, J. Karadžinska-Bislimovska, S. Risteska-Kuc, Z. Stojanovski, and S. Stoleski, “Exercise-induced bronchoconstriction in female cleaners: effect of smoking,” *Nauka Pulmologia*, vol. 1, pp. 41–44, 2007.
- [29] J. Minov, J. Karadžinska-Bislimovska, K. Vasilevska, S. Risteska-Kuc, and S. Stoleski, “Exposure to environmental tobacco smoke in the workplace in Macedonia: where are we now?” *Arhiv za Higijenu Rada I Toksikologiju*, vol. 59, no. 2, pp. 103–109, 2008.
- [30] J. Minov, J. Karadžinska-Bislimovska, K. Vasilevska, and S. Stoleski, “Smoking status in exposed and unexposed workers,” *Makedonski Medicinski Pregled*, vol. 60, no. 68, p. 128, 2006.
- [31] C. Janson, N. Künzli, R. de Marco et al., “Changes in active and passive smoking in the European Community Respiratory Health Survey,” *European Respiratory Journal*, vol. 27, no. 3, pp. 517–524, 2006.
- [32] D. Gautrin, H. Ghezzi, and J. L. Malo, “Rhinconjunctivitis, bronchial responsiveness, and atopy as determinants for incident non-work-related asthma symptoms in apprentices exposed to high-molecular-weight allergens,” *Allergy*, vol. 58, no. 7, pp. 608–615, 2003.
- [33] M. Capão-Filipe, A. Moreira, L. Delgado, J. Rodrigues, and M. Vaz, “Exercise-induced bronchoconstriction and respiratory symptoms in elite athletes,” *Allergy*, vol. 58, no. 11, p. 1196, 2003.
- [34] J. Minov, V. Cvetanov, J. Karadžinska-Bislimovska, N. Ezova, S. Milkovska, and S. Risteska-Kuc, “Epidemiological characteristics of bronchial asthma in R. Macedonia,” *Makedonski Medicinski Pregled*, vol. 56, p. 156, 2003.
- [35] J. Karadžinska-Bislimovska, V. Cvetanov, J. Petrovska, S. Todorov, and S. Risteska-Kuc, “Respiratory symptoms and positive skin prick test in a prospective asthma study in Republic of Macedonia (initial results),” *European Respiratory Journal*, vol. 14, p. 78, 1999.
- [36] J. Minov, J. Karadžinska-Bislimovska, K. Vasilevska, S. Risteska-Kuc, and S. Stoleski, “Bronchial hyperresponsiveness in workers exposed to tea dust: host versus environmental factors,” *Allergy Hypersensitivity Asthma*, vol. 3, no. 1, pp. 33–39, 2005.
- [37] J. Karadžinska-Bislimovska, J. Minov, S. Risteska-Kuc, S. Stoleski, and D. Mijakoski, “Bronchial hyperresponsiveness in women cooks and cleaners,” *Arhiv za Higijenu Rada I Toksikologiju*, vol. 58, no. 2, pp. 223–231, 2007.
- [38] S. O’Sullivan, A. Roquet, B. Dahlén et al., “Evidence for mast cell activation during exercise-induced bronchoconstriction,” *European Respiratory Journal*, vol. 12, no. 2, pp. 345–350, 1998.
- [39] P. M. O’Byrne, “Leukotriene bronchoconstriction induced by allergen and exercise,” *American Journal of Respiratory and Critical Care Medicine*, vol. 161, no. 2, pp. S68–S72, 2000.
- [40] I. Helenius and T. Haahtela, “Allergy and asthma in elite summer sport athletes,” *Journal of Allergy and Clinical Immunology*, vol. 106, no. 3, pp. 444–452, 2000.
- [41] R. T. Thole, R. E. Sallis, A. L. Rubin, and G. N. Smith, “Exercise-induced bronchospasm prevalence in collegiate cross-country runners,” *Medicine and Science in Sports and Exercise*, vol. 33, no. 10, pp. 1641–1646, 2001.
- [42] R. A. Tan and S. L. Spector, “Exercise-induced asthma,” *Sports Medicine*, vol. 25, no. 1, pp. 1–6, 1998.
- [43] Y. I. Koh and I. S. Choi, “Blood eosinophil counts for the prediction of the severity of exercise-induced bronchospasm in asthma,” *Respiratory Medicine*, vol. 96, no. 2, pp. 120–125, 2002.

- [44] T. Klepac-Pulanić, J. Macan, D. Plavec, and B. Kanceljak-Macan, "Exercise and allergic diseases," *Arhiv za Higijenu Rada I Toksikologiju*, vol. 55, no. 2-3, pp. 197–204, 2004.
- [45] G. F. Joos, B. O'Connor, S. D. Anderson et al., "Indirect airway challenges," *European Respiratory Journal*, vol. 21, no. 6, pp. 1050–1068, 2003.
- [46] K. C. Ringsberg, H. Wetterqvist, O. Löwhagen, and T. Sivik, "Physical capacity and dyspnea in patients with asthma-like symptoms but negative asthma tests," *Allergy*, vol. 52, no. 5, pp. 532–540, 1997.
- [47] P. Weiner, N. Konson, A. Sternberg, D. Zamir, and Z. Fireman, "Is gastro-oesophageal reflux a factor in exercise-induced asthma?" *Respiratory Medicine*, vol. 92, no. 8, pp. 1071–1075, 1998.
- [48] O. Löwhagen, M. Arvidsson, P. Bjärneman, and N. Jörgensen, "Exercise-induced respiratory symptoms are not always asthma," *Respiratory Medicine*, vol. 93, no. 10, pp. 734–738, 1999.



Hindawi
Submit your manuscripts at
<http://www.hindawi.com>

