Research Article

Organic Solid Acid/ NaNO_2 : An Efficient System for the Oxidation of Urazoles and Bis-Urazoles under Mild and Heterogeneous Conditions

Gholamabbas Chehardoli,¹ Mohammad Ali Zolfigol,²
Mohsen Rasouli,³ and Shadpour Mallakpour⁴

¹ School of Pharmacy, Hamedan University of Medical Sciences, Hamedan 65178, Iran

² Faculty of Chemistry, Bu-Ali Sina University, Hamedan 65174, Iran

³ Department of Chemistry, Faculty of Sciences, Payame Noor University, Hamedan 65417, Iran

⁴ Organic Polymer Chemistry Research Laboratory, College of Chemistry, Isfahan University of Technology, Isfahan 51167, Iran

Correspondence should be addressed to Gholamabbas Chehardoli, cheh1002@gmail.com

Received 12 October 2011; Revised 25 November 2011; Accepted 26 November 2011

Academic Editor: Dipakranjan Mal

Copyright © 2011 Gholamabbas Chehardoli et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Various organic solid acids/ NaNO_2 were used as effective oxidizing agents for the oxidation of urazoles and bis-urazoles to their corresponding triazolinodiones under mild and heterogeneous conditions at room temperature with good-to-excellent yields.

1. Introduction

4-Substituted-1,2,4-triazole-3,5-diones (TADs) have been used both as substrates and reagents in various organic reactions, such as Diels-Alder-reaction, ene reactions, [2+2] cycloadditions, dehydrogenation reactions, electrophilic aromatic substitution, condensation of dicarbonyl compounds, and oxidation of alcohols to aldehydes and ketones [1–4]. Very recently aromatization of 1,4-dihydropyridines and pyrazolines and oxidation of thiols with TADs as well have been reported [5–7]. The unusual reactivity of TADs makes them interesting and also hard to prepare and purify [8]. It is interesting to note that, 4-phenyl-1,2,4-triazoline-3,5-dione is an extremely reactive dienophile and enophile which is at least 1000 times more reactive than tetracyanoethylene in the Diels-Alder reaction with 2-chlorobutadiene and 2000 times more reactive than maleic anhydride [1–4]. All known methods for the preparation of these compounds (1,2,4-triazolidine-3,5-diones) require oxidation of the corresponding urazoles.

Although a variety of reagents are capable of efficient oxidations of urazoles to TADs, this transformation is not easy because these compounds are very sensitive to the oxidizing

agents and reaction conditions. In addition, most of the reported reagents produce byproducts which either destroy the sensitive triazolinodiones, or are difficult to remove from the reaction product. Another major drawback of the older procedures is their use of reagents which are either highly toxic or impose serious disposal problems (or both) [9–13].

It has been demonstrated that application of heterogeneous systems, for the above reported oxidations, has many advantages over their liquid phase counterparts such as simple experimental procedures, mild reaction conditions and minimization of chemical waste materials [14, 18].

Polymer-supported sulfonic acid {(I) sulfonated polystyrenes (Dowex-50, Amberlite IR-112, Permutit-Q)} [10], sulfamic acid (II) [10], and isocyanuric acid (III) [10] are three stable and suitable organic solid acids that are commercially available. Easy workup and the stability of these reagents make them safe and convenient source of proton in comparison to liquid acids. These reagents are transformed during reaction into easily removable products. There are several reports that demonstrate the use of these reagents for various organic transformations under mild conditions (Scheme 1) [15].

TABLE 1: Oxidation of urazoles (1) and bis-urazoles (3) to their corresponding triazolinediones (2, 4) with either one of the Polymer-supported sulfonic acid (I), sulfamic acid (II) or isocyanuric acid (III) in the presence of NaNO₂ in dichloromethane at room temperature.

Entry	Urazole	Product	Reagent/substrate (mmol)			Time (h)			Yields ^a (%)		
			I	II	III ^b	I	II	III	I	II	III
1	1a	2a	2	0.2	0.75	1	2	0.0	100 ^c	100 ^c	100 ^c
2	1b	2b	2	0.2	0.75	1	2	0.0	100 ^c	100 ^c	100 ^c
3	1c	2c	2	0.2	0.75	1	2	0.0	90	72	91
4	1d	2d	2	0.2	0.75	1	2	0.0	99	99	90
5	1e	2e	2	0.2	0.75	1	2	0.5	90	85	90
6	1f	2f	2	0.2	1	1	2	0.5	91	80	95
7	1g	2g	2	0.2	1	1	2	0.5	94	83	89
8	1h	2h	2	0.2	1.25	1	2	0.5	90	92	80
9	1i	2i	2	0.2	1	1	2	0.5	89	81	100 ^{c,d}
10	1j	2j	2	0.2	1	1	2	0.5	93	75	70
11	1k	2k	2	0.2	1	1	2.5	0.5	98	94	96
12	1l	2l	2	0.2	1.25	1	2.5	0.5	87	80	84
13	3a	4a	4	0.4	2	2	4	0.5	85	82	94
14	3b	4b	4	0.4	2	2.5	5	0.5	90	99	93

^a Isolated yields. ^b In the presence of a few drops of water. ^c Conversion. ^d 4-Nitrophenyl triazolinedione is very labile. Therefore, it was destructed in the course of passing through a pad of silica gel.

SCHEME 1

2. Results and Discussion

In continuation of our studies on the use of *solid acids* in organic transformations [16, 17], we found that polymer-supported sulfonic acid (I), sulfamic acid (II) and isocyanuric acid (III) in the presence of NaNO₂ could be used as oxidizing agents for the oxidation of urazoles and bis-urazoles under mild and heterogeneous conditions (Scheme 2).

Herein, we wish to report a simple, inexpensive, and convenient method for the effective oxidation of urazoles (1) and bis-urazoles (3) to their corresponding triazolinediones (2, 4) by using the described reagents I–III.

A good range of urazoles (1) and bis-urazoles 3 were subjected to the oxidation reaction in the presence of the solid acids (I–III)/NaNO₂ in dichloromethane. All oxidation reactions were performed under mild and completely heterogeneous conditions, at room temperature with good to excellent yields (Table 1).

As reported above the oxidation reactions are heterogeneous because urazoles and bis-urazoles ((1, 3) as a white solids) are insoluble in dichloromethane whereas all of the triazolinediones and bis-triazolinediones ((2, 4), red, pink or brown solids) are very well soluble in dichloromethane. According to our previously reported results with other

Urazole	R ₁	R ₂
1a	H	Me
1b	H	Et
1c	Na	n-Pr
1d	H	n-Bu
1e	H	Cyclohexyl
1f	H	Ph
1g	H	4-Cl-Ph
1h	H	3,4-Cl ₂ -Ph
1i	H	4-NO ₂ -Ph
1j	H	4-OMe-Ph
1k	H	4- <i>t</i> -Bu-Ph
1l	H	4-Naphtyl
3a	Na	-(CH ₂) ₆ -
3b	H	

SCHEME 2

reagents [14, 16–18], the following mechanism for the oxidation reaction via *in situ* generation of NO⁺ may be suggested (Scheme 3).

Although Nafion-H is a very useful solid acid [10], unfortunately, several attempts for the oxidation of urazoles by Nafion-H have failed. Therefore, this reagent is not practical and suitable for this purpose.

In summary, we have described a practical, efficient, and convenient method for the oxidation of urazoles and bis-

SCHEME 3

urazoles. We suggest that these systems could be used for the oxidation of a wide variety of urazole derivatives under mild and safe conditions.

3. Experimental Section

3.1. General. Chemicals were purchased from Fluka, Merck and Aldrich chemical companies. Yields refer to isolated pure products. The oxidation products were characterized by comparison of their spectral (IR and $^1\text{H-NMR}$) and physical data with authentic samples which were produced by other reported procedures.

3.2. Oxidation of 4, 4'-(4, 4'-Diphenylmethylene)-bis-urazole (3b) to bis(p-3, 5-Dioxo-1, 2, 4-triazoline-4-ylphenyl) Methane (4b): A Typical Procedure. A mixture of compound **3b** (0.366 g, 1 mmol), **I-III** (see Table 1), and NaNO_2 (1 mmol) in CH_2Cl_2 (10 mL) was stirred for the specified time in Table 1. Then the reaction mixture was filtered and washed with CH_2Cl_2 (2×10 mL). Dichloromethane was removed by water bath ($40\text{--}50^\circ\text{C}$) and simple distillation. A crystalline pink solid (**4b**) is obtained in good-to-excellent yields. Mp = $182\text{--}185^\circ\text{C}$, and {lit. Mp = 185°C [14, 18]}.

Acknowledgments

This paper is extracted from the M. S. thesis of Mr. M. Rasouli. The authors acknowledge financial support for this work from the research affairs of Hamedan University of Medical Sciences, Hamedan, Iran and partial support of this work by the Research Affairs Office of Bu-Ali Sina University (Grant no. 32-1716 entitled development of chemical methods, reagent and molecules.) and also Center of Excellence in Development of Chemical Method (CEDCM) Hamedan, Iran.

References

- [1] G. Chehardoli, "Triazolinediones (TADs)," *Synlett*, no. 13, pp. 2154–2155, 2006.
- [2] S. E. Mallakpour, G. B. Butler, H. Aghabozorg, and G. J. Palenik, "Ene reaction of (S)-(-)-4-(a-Methylbenzyl)-1,2,4-triazoline-3,5-dione with Propylene, X-ray diffraction analysis of a single crystal of the brominated adduct," *Macromolecules*, vol. 18, pp. 342–347, 1985.
- [3] S. E. Mallakpour and G. B. Butler, "Alternating copolymers via diels-alder and ene reactions," in *Advanced in Polymer Synthesis*, B. M. Culbertson and J. E. McGrath, Eds., vol. 31 of *Polymer Science and Technology Series*, pp. 1–25, Plenum Press, New York, NY, USA, 1985.
- [4] S. E. Mallakpour and G. B. Butler, "Uncatalyzed polymerization of bistriazolinediones with electron-rich aromatic compounds via electrophilic aromatic substitution," *Journal of Polymer Science A*, vol. 27, no. 1, pp. 217–235, 1989.
- [5] M. A. Zolfigol, A. G. Choghamarani, M. Shahamirian et al., "4-Phenyl-1,2,4-triazole-3,5-dione as a novel and reusable reagent for the aromatization of 1,4-dihydropyridines under mild conditions," *Tetrahedron Letters*, vol. 46, no. 33, pp. 5581–5584, 2005.
- [6] M. A. Zolfigol, D. Azarifar, S. Mallakpour et al., "4-(p-Chloro)phenyl-1,2,4-triazole-3,5-dione as a novel and reusable reagent for the oxidation of 1,3,5-trisubstituted pyrazolines under mild conditions," *Tetrahedron Letters*, vol. 47, no. 5, pp. 833–836, 2006.
- [7] A. Christoforou, G. Nicolaou, and Y. Elemes, "N-Phenyltriazolinedione as an efficient, selective, and reusable reagent for the oxidation of thiols to disulfides," *Tetrahedron Letters*, vol. 47, no. 52, pp. 9211–9213, 2006.
- [8] J. C. Stickler and W. H. Pirkle, "An improved synthesis of 1,2,4-triazoline-3,5-diones," *Journal of Organic Chemistry*, vol. 31, no. 10, pp. 3444–3445, 1966.
- [9] G. Read and N. R. Richardson, "Synthetic studies on 4,5-dihydro-3H-1,2,4-triazole-3,5-diones bearing fluorogenic residues at N-4," *Journal of the Chemical Society—Perkin Transactions 1*, no. 2, pp. 167–174, 1996.
- [10] V. P. Arya and S. Shenoy, "Synthesis of novel heterocycles from 4-substituted 1,2,4-triazolidine-3,5-diones," *Indian Journal of Chemistry B*, vol. 11, pp. 883–886, 1976.
- [11] H. Warnho and K. Wald, "A Convenient preparation of 4-aryl-1,2,4-triazoline-3,5-diones," *Organic Preparations and Procedures International*, vol. 7, pp. 251–253, 1975.
- [12] S. E. Mallakpour, "A new method for the oxidation of 4-phenylurazole to 4-phenyltriazolinedione," *Journal of Chemical Education*, vol. 69, no. 3, pp. 238–241, 1992.
- [13] S. E. Mallakpour and M. A. Zolfigol, "A convenient method for preparation and Isolation of 4-n-Propyl-1,2,4-triazoline-3,5-Diane," *Iranian Journal of Science and Technology-Science*, vol. 4, pp. 199–205, 1993.
- [14] M. A. Zolfigol, M. Bagherzadeh, S. Mallakpour et al., "Mild and heterogeneous oxidation of urazoles to their corresponding triazolinediones via in situ generation Cl^+ using silica sulfuric acid/ KClO_3 or silica chloride/oxone system," *Catalysis Communications*, vol. 8, no. 3, pp. 256–260, 2007.
- [15] P. Gogoi, "Resin-supported sulfonic acid," *Synlett*, no. 14, pp. 2263–2264, 2005.
- [16] P. Salehi, M. A. Zolfigol, F. Shirini, and M. Baghbanzadeh, "Silica sulfuric acid and silica chloride as efficient reagents for organic reactions," *Current Organic Chemistry*, vol. 10, no. 17, pp. 2171–2189, 2006.
- [17] F. Shirini, M. A. Zolfigol, P. Salehi, and M. Abedini, "Applications of some metal hydrogen sulfates in organic transformations," *Current Organic Chemistry*, vol. 12, no. 3, pp. 183–202, 2008.
- [18] M. A. Zolfigol, G. Chehardoli, E. Ghaemi et al., "N-bromo reagent mediated oxidation of urazoles to their corresponding triazolinediones under mild and heterogeneous conditions," *Monatshefte fur Chemie*, vol. 139, no. 3, pp. 261–265, 2008.