Review Article

Metal Preferences of Zinc-Binding Motif on Metalloproteases

Kayoko M. Fukasawa,¹ Toshiyuki Hata,² Yukio Ono,² and Junzo Hirose²

¹ Department of Hard Tissue Research, Graduate School of Oral Medicine, Matsumoto Dental University, Shiojiri, Nagano 399-0781, Japan

² Faculty of Pharmacy and Pharmaceutical Science, Fukuyama University, Gakuen-cho, Fukuyama 729-0292, Japan

Correspondence should be addressed to Kayoko M. Fukasawa, kmf@po.mdu.ac.jp

Received 4 October 2010; Revised 25 January 2011; Accepted 7 March 2011

Academic Editor: Shandar Ahmad

Copyright © 2011 Kayoko M. Fukasawa et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Almost all naturally occurring metalloproteases are monozinc enzymes. The zinc in any number of zinc metalloproteases has been substituted by some other divalent cation. Almost all Co(II)- or Mn(II)-substituted enzymes maintain the catalytic activity of their zinc counterparts. However, in the case of Cu(II) substitution of zinc proteases, a great number of enzymes are not active, for example, thermolysin, carboxypeptidase A, endopeptidase from *Lactococcus lactis*, or aminopeptidase B, while some do have catalytic activity, for example, astacin (37%) and DPP III (100%). Based on structural studies of various metal-substituted enzymes, for example, thermolysin, astacin, aminopeptidase B, dipeptidyl peptidase (DPP) III, and del-DPP III, the metal coordination geometries of both active and inactive Cu(II)-substituted enzymes are shown to be the same as those of the wild-type Zn(II) enzymes. Therefore, the enzyme activity of a copper-ion-substituted zinc metalloprotease may depend on the flexibility of catalytic domain.

1. Introduction

Proteolytic enzymes are recognized by their catalytic type, that is, aspartic, cysteine, metallo, serine, threonine, and others as yet unclassified. The largest number of proteolytic enzymes are classified as metalloproteases [1]. Almost all metalloproteases contain one or two zinc ions, and several enzymes contain one or two cobalt or manganese ions. The HExxH motif forming an α -helix is well conserved in many monozinc enzymes as the active site in which the two histidine residues coordinate with the zinc ion [2]. Some other monozinc proteases have different zinc-binding motifs, for example, HxxE(D)-aa_n-H in the carboxypeptidase family or HxD-aa₁₂-H-aa₁₂-H in the matrix metalloprotease family [2]. Dipeptidyl peptidase (DPP) III also has a unique zinc-binding motif, which was classified as family M49 in 1999 by MEROPS (peptidase database) after rat DPP III had been cloned and the HELLGH motif of DPP III was identified as an active site coordinated with a zinc ion [3, 4]. Although the motif HELLGH could not be found in any other metalloproteases, it exists in three kinds of monooxygenases

(tyrosine, phenylalanine, and tryptophan hydroxylases) as an iron-binding site, as revealed by a search of the NBRF-PIR protein sequence database.

Zinc atoms in several zinc metalloproteases, for example, astacin [5], carboxypeptidase A [6], and thermolysin [7, 8], have been substituted by other divalent cations to probe the role of the metal for catalysis and structure. Some of these enzymes, for example, DPP III and astacin, were shown to have high metal substitution tolerance by metal substitution studies [9]. However, it is difficult to determine the relationship between the metal tolerance and the metal coordination structure of zinc metalloproteases.

Here, we show the metal coordination structure of the unique zinc-binding motif of DPP III, in which the zinc-binding motif is stabilized by several hydrogen bonds with acidic amino acid residues surrounding the zinc-binding motif, in order to clarify the relationship between the metal tolerance and the structure of the zinc-binding domain. The metal tolerances of both DPP III and del-DPP III, whose active site converts into a normal zinc-binding motif (HExxH), are shown here and compared with those

TABLE 1: The zinc coordination residues and the residues that fix the coordination with hydrogen bonds.

Zinc metalloprotease	Coordination residues	Residues that form the hydrogen bond with the coordination residues	PDB no.
(1) Thermolysin type	(HExxH- aa _n -E)	α -helix-aa _n - α -helix	
Thermolysin	His ¹⁴² , His ¹⁴⁶ Glu ¹⁶⁶	Asp ¹⁷⁰ -2.8 Å-His ¹⁴² Asn ¹⁶⁵ -2.8 Å-His ¹⁴⁶	1KEI
Vibriolysin	His ³⁴⁵ , His ³⁴⁹ Glu ³⁶⁹	Asp ³⁷³ -2.8 Å-His ³⁴⁵ Asn ³⁶⁸ -2.8 Å-His ³⁴⁹	3NQX
Staphylococcus aureus metalloproteinase	His ¹⁴⁴ , His ¹⁴⁸ Glu ¹⁶⁸	Asn ¹⁶⁷ -2.8 Å-His ¹⁴⁸ Asp ¹⁷² -2.8 Å-His ¹⁴⁴	1BQB
Zinc aminopeptidase	His ²⁶⁵ , His ²⁶⁹ Glu ²⁸⁸	Phe ²⁷² (C=O)-2.9 Å-His ²⁶⁹	1Z1W
Leukotriene A4 hydrolase	His ²⁹⁵ , His ²⁹⁹ Glu ³¹⁸	Glu ³²⁵ -2.8 Å-His ²⁹⁵ Gly ³⁰³ (C=O)-2.6 Å-His ²⁹⁹	1SQM
Human thimet oligopeptidase	His ⁴⁷³ , His ⁴⁷⁷ Glu ⁵⁰²	Glu ⁵⁰⁹ -2.6 Å-His ⁴⁷³	1SQM
Human neutral endopeptidase (Neprilysin)	His ⁵⁸³ , His ⁵⁸⁷ Glu ⁶⁴⁶	Asp ⁶⁵⁰ -2.9 Å-His ⁵⁸³ Asp ⁵⁹⁰ -2.7 Å-His ⁵⁸⁷	1DMT
(2) Endopeptidase type		(HExxH-aa _n -E or D) α -helix-aa _n -random coil	
Peptidyl-Lys metalloendopeptidase	His ¹¹⁷ , His ¹²¹ Asp ¹³⁰	Asp ¹⁵⁴ -2.7 Å-His ¹¹⁷ Thr ¹²⁸ (C=O)-2.8 Å-His ¹²¹	1GE6
(3) Carboxypeptidase A type		β -sheet-aa _n -random coil	
Carboxypeptidase A	His ⁶⁹ , His ¹⁹⁶ Glu ⁷²	Asp ¹⁴² -2.7 Å-His ⁶⁹	1YME
Putative lysostaphin peptidase	His ²³² , His ³¹¹ Asp ²³⁶	Glu ³¹⁵ -2.6 Å-His ³¹¹ Gly ²¹⁶ (C=O)-2.8 Å-His ²³²	2GU1

reported for other metalloproteases [10]. Finally, we discuss the relationship between the catalytic activities and metal coordination structures of metal-substituted enzymes.

2. Identification of a Zinc-Binding Motif in DPP III

We start with the identification of the zinc-binding motif in DPP III, which will be used for further investigation of the relationship between the metal tolerance and the metal coordination structures of DPP III. The deduced amino acid sequences from cDNA for human, rat, and fruit fly DPP IIIs are 723–738 amino acids long and conserve the amino acid sequence HELLGH-aa₅₂-E [3, 11, 12], which resembles the HExxH-aa_n-E zinc-binding motif conserved in many metalloproteases, such as thermolysin [13] and leukotriene A₄ hydrolase [14]. Site-directed mutagenesis was performed on rat DPP III in order to testify that the HELLGH-aa₅₂-E is a zinc-binding domain. Site-directed mutagenesis studies have clearly shown that the H450Y, H455Y, and E508A mutants, which lack zinc ions, lose their catalytic activity [4]. Replacement of Glu⁴⁵¹ in these mutants with an alanine or an aspartic acid restores a mol of zinc ion per mol of protein but does not restore catalytic activity [4]. These results show that the H⁴⁵⁰ELLGH-aa₅₂-E⁵⁰⁸ motif is a catalytic domain of which His⁴⁵⁰, His⁴⁵⁵, and Glu⁵⁰⁸ are ligands of a zinc ion and of which Glu⁴⁵¹ is a catalytic amino acid residue, in the same way that the H¹⁴²ExxH-aa₁₉-E¹⁶⁶ motif of thermolysin is a catalytic domain of which His¹⁴², His¹⁴⁶, and Glu¹⁶⁶ are ligands of a zinc ion and Glu¹⁴³ is a catalytic amino acid residue.

FIGURE 1: The superimposition of the active sites of rat DPP III and thermolysin. Zinc ion is shown as a green sphere, and amino acid side chains are shown as sticks colored red for oxygen and blue for nitrogen. Metal coordinates in light blue and hydrogen bonds in yellow are indicated by dashed lines. Carbon atom and amino acid chain are shown colored white for thermolysin and magenta for DPP III. Metal coordination bonds are indicated by light blue dashed lines.

The 1.95-Å crystal structure of yeast DPP III representing a prototype for the M49 family of metalloproteases was resolved by Baral et al. [15]. It shows a novel protein fold with two domains forming a wide cleft containing the catalytic metal ion. However, the three-dimensional structure of zinc coordination (His⁴⁶⁰, His⁴⁶⁵, and Glu⁵¹⁷) and the catalytic active (Glu⁴⁶¹) residues are structurally conserved,

(a)

(b)

(c)

FIGURE 2: Three-dimensional structures of the catalytic domain models for thermolysin ((a): PDB 1KEI), peptidyl-Lys metalloendopeptidase ((b): PDB 1GE6), and carboxypeptidase A ((c): PDB 1YME). The zinc ion is shown as a green sphere, and amino acid side chains are shown as sticks colored red for oxygen, blue for nitrogen, and silver for carbon. Hydrogen bonds are indicated by dashed lines.

TABLE 2: Kinetic parameters for the hydrolysis of Arg-Arg-NA, zinc contents, and zinc dissociation constants of wild-type and mutated rat DPP IIIs.

Enzymes	$k_{\text{cat}}/K_M \times 10^{-4}$ ($\text{M}^{-1} \text{s}^{-1}$)	Zinc content (mol/mol of protein ^a)	Zinc dissociation constant (M) (K_d)
Wild-type	73.6 ± 6.9	1.02 ± 0.15	$(4.5 \pm 0.1) \times 10^{-13}$
E507D	22.8 ± 1.9	0.65 ± 0.07	$(1.0 \pm 0.2) \times 10^{-11}$
E507A	4.43 ± 0.41	0.29 ± 0.04	$(1.0 \pm 0.2) \times 10^{-8}$
E512D	21.0 ± 0.19	0.45 ± 0.06	$(1.4 \pm 0.1) \times 10^{-12}$
E512A	2.45 ± 0.28	0.08 ± 0.01	$(2.6 \pm 0.7) \times 10^{-9}$

^a Values are means $\pm$ SD of two separately prepared enzymes with duplicate determinations.

similar to those presented in many metalloproteases, such as thermolysin [13]. The HELGH motif and the third ligand (Glu⁵¹⁷) of DPP III construct a helix α_{14} and a helix α_{16} , respectively [15]. The 3D structure of DPP III is similar to that of thermolysin [13] or leukotriene A₄ hydrolase [16], the zinc-binding domain of which is constructed of two α -helices, for HEXxH (containing two zinc ligands) and xNEx (third ligand).

Figure 1 shows the superimposition of the active sites of rat DPP III and thermolysin. The helix α_{14} of DPP III has a slightly larger loop than that of thermolysin, and the glutamic acid on the motif comes close to zinc ion comparing with the glutamic acid on the normal helix of thermolysin [13, 17].

3. Stabilization of the Coordination between Ligands and Metal

In the 3D structural model of the zinc-binding domain of many zinc enzymes—neprilysin [18], thermolysin [13], carboxypeptidase A [19], leukotriene A₄ hydrolase [16], aspzincin [20], and DPP III [17]—the His, His, and Glu residues that coordinate with the zinc ion are engaged in hydrogen bonds with one or two acidic amino acid residues (Glu or Asp) or other carbonyl oxygen atoms (Table 1). 3D structural models of catalytic domains of thermolysin (PDB: 1KEI), peptidyl-Lys metalloendopeptidase (PDB: 1GE6), and carboxypeptidase A (PDB: 1YME) are shown in Figure 2. In thermolysin (a), the oxygen atoms of Asp¹⁶⁵ and Asp¹⁷⁰ are engaged in hydrogen bonding with the nitrogen atoms of His¹⁴⁶ and His¹⁴², respectively. Asp¹⁵⁴ and Thr¹²⁸ of peptidyl-Lys metalloendopeptidase (b) and Asp¹⁴² of carboxypeptidase A (c) are also engaged in hydrogen bonding with His¹¹⁷, His¹²¹, and His⁶⁹, respectively. It was proved through the mutational studies of rat DPP III that this network of hydrogen bonds close to the zinc-binding motif plays an important role in stabilizing the coordination of the zinc ion to the protein [17]. The hydrogen bonds surrounding the zinc-binding motif of rat DPP III are shown in Figure 3, and the kinetic parameters, zinc contents and zinc dissociation constants of the several mutants are shown in Table 2. The replacement of Glu⁵⁰⁷ and Glu⁵¹², the oxygen atoms of which bind with the nitrogen atoms of His⁴⁵⁵ and

FIGURE 3: Molecular modeling of the catalytic site of rat DPP III. The model was generated as a template of the human DPP III crystal structure [39]. The zinc ion is shown as a green sphere, and amino acid side chains are shown as sticks colored red for oxygen, blue for nitrogen, and white for carbon. Metal coordinates in light blue and hydrogen bonds in yellow are indicated by dashed lines.

His⁴⁵⁰, respectively, increases the dissociation constants by factors of $10\sim 10^5$ and correlatively reduces the zinc contents and enzyme activities. The hydrogen bonds between acidic amino acid residues and zinc ligands (His, His, and Glu) may stabilize the coordination of the zinc ion with the protein of the metalloprotease.

4. Metal Substitutions of Monozinc Metalloproteases

Almost all metalloproteases are monozinc enzymes. Some enzymes contain two zinc ions for catalytic domains, for example, human renal dipeptidase [36], and a few are dicobalt or dimanganese enzymes, for example, *Pyrococcus furiosus* methionine aminopeptidase [37] or *Escherichia coli* proline aminopeptidase [38], respectively.

The zinc in numerous zinc metalloproteases has been substituted by several divalent cations. The cobalt(II)- or manganese(II)-substituted enzymes showed nearly restored catalytic activity or even excess activity from apoenzyme, as seen in Table 3.

Gomis-Rüth et al. [5] demonstrated in the metal substitution studies of astacin that Cu(II)-astacin displays enzyme activity of about 37%, while Ni(II)- and Hg(II)-astacin were almost inactive. In the crystal structure of Cu(II)-astacin, the metal ion is pentacoordinated with His⁹², His⁹⁶, His¹⁰², Tyr¹⁴⁹, and H₂O, as in native Zn(II)-astacin or Co(II)-astacin; however, in the Ni(II)-astacin or Hg(II)-astacin, the metal ion is hexacoordinated with an additional solvent molecule or tetracoordinated with no ordered solvent molecule, respectively [5]. The restoration of catalytic activity in these substituted astacins was shown to be dependent on the metal coordination structure [5].

Meanwhile, almost all Cu(II)-substituted enzymes, such as thermolysin [7, 8], carboxypeptidase A [6], aminopeptidase B [22], or endopeptidase from *Lactococcus lactis* [30],

show only partial activation or very low activities. The reason why these Cu(II) enzymes do not demonstrate catalytic activities may be that the coordination geometry of Cu(II) is more rigid than that of Zn(II) or Co(II).

In the case of DPP III, Co²⁺-, Ni²⁺- and Cu²⁺-DPP IIIs showed comparable catalytic activities to Zn²⁺-DPP III; the kinetic parameters are shown in Table 4 [9]. DPP III shows high flexibility of the metal ion for the catalytic activity compared with thermolysin or aminopeptidase B. Thermolysin or aminopeptidase B is a subclass MA (E) metalloprotease containing an HExxH-aa_n-E motif, and the 3D structure of the active domain is very similar to that of DPP III described above. The zinc ion in a subclass MA (E) metalloprotease or DPP III is tetraordinated with three ligands (His, His, and Glu) and a water molecule. The metal-substituted (Co²⁺, Cu²⁺, or Ni²⁺) DPP III may have the same tetrahedral coordination structure as Zn²⁺-DPP III, so these enzymes are able to maintain the catalytic activity. The zinc in del-DPP III, whose active site converted into HExxH, was substituted with Co²⁺, Ni²⁺, or Cu²⁺ to investigate the grounds for activation of the Cu²⁺-DPP III [10]. The Co²⁺-del-DPP III and Ni²⁺-del-DPP III showed comparable catalytic activity to that of Zn²⁺-del-DPP III, while the Cu²⁺-del-DPP III showed no catalytic activity, as in the case of thermolysin or aminopeptidase B [8–10].

The EPR (electron paramagnetic resonance) parameters of various Cu²⁺-substituted metalloproteases are shown in Table 5. Each parameter is exactly alike between DPP III and thermolysin, aminopeptidase B, or del-DPP III [8–10, 22]. The results show that the Cu(II) coordination structures of the HExxH-aa_n-E and HExxxH-aa₅₂-E motifs are very similar.

In the superimposition of the 3D structure models of active sites of DPP III and del-DPP III, the α -helix of DPP III, which is abnormally composed of 5 amino acid residues per one turn of the α -helix, is a larger loop than that of del-DPP III, the same as the case for the superimposition of the active sites of DPP III and thermolysin (Figure 1). The coordination geometries of the two enzymes are similar, while the position of Glu⁴⁵¹, which is essential for the enzyme activity, is slightly closer to the copper ion in DPP III than in del-DPP III. The distances of oxygen atoms of the Glu⁴⁵¹ residues of del-DPP III and wild-type DPP III are 4.8 Å and 3.2 Å from the zinc ion, respectively. Zn(II) coordination geometry is flexible, so both wild-type and del-DPP III could have catalytic activity. However, the oxygen atom of Glu⁴⁵¹ in Cu(II)-del-DPP III is not able to bind to the oxygen atom of the water molecule that is coordinated with the copper ion because the Cu(II) coordination geometry is very rigid. Therefore, the catalytic activity of Cu(II)-del-DPP III was diminished.

Some other Cu(II)-substituted enzymes, for example, aminopeptidase Ey [21], vibriolysin [40], hycicolysin [32], and *Legionella* metalloendopeptidase [33], were shown to have enzyme activities. These enzymes are all classified in subclass MA (E), the same as thermolysin or aminopeptidase B. The metal coordination structures of these enzymes have not been shown in detail; however, the catalytic domain may be more flexible than that of thermolysin or aminopeptidase B, in the same way as Cu(II)-substituted DPP III.

TABLE 3: Reactivated Co(II) and Mn(II) enzymes substituted from apo-metalloproteases.

Clan	Subclan	Name of enzyme	Replaced Ion	Reference
MA	E	Aminopeptidase Ey	Co ²⁺ , Mn ²⁺	Tanaka and Ichishima [21]
MA	E	Aminopeptidase B	Co ²⁺	Hirose et al. [22]
MA	E	Saccharolysin	Co ²⁺ , Mn ²⁺	Achstetter et al. [23] and Büchler et al. [24]
MA	E	Lysyl aminopeptidase	Co ²⁺ , Mn ²⁺	Klein et al. [25]
MA	E	Oligopeptidase F	Co ²⁺ , Mn ²⁺	Yan et al. [26] and Monnet et al. [27]
MA	E	Mycolysin (Thermolysin)	Co ²⁺ , Mn ²⁺ Co ²⁺ (200%), Mn ²⁺ (10%)	Chang and Lee [28], and Holmquist and Vallee [29]
MA	E	Oligopeptidase O	Co ²⁺ Mn ²⁺ (50%)	Tan et al. [30] and Baankreis et al. [31]
MA	E	Hycolysin	Co ²⁺	Ayora and Götz [32]
ME	E	Legionella metalloendopeptidase	Mn ²⁺ (69%)	Dreyfus and Iglewski [33]
MA	A	Epralysin	Co ²⁺ (58%)	Diermayr et al. [34]
MA	M	Astacin	Co ²⁺ (140%)	Gomis-Rüth et al. [5]
MA	M	MEP ^a (<i>Gf</i> ^b MEP)	Co ²⁺ Mn ²⁺ (200%)	Nonaka et al. [35]
MA	M	Po ^c MEP	Co ²⁺ (80%) Mn ²⁺ (30%)	Nonaka et al. [35]

^aPeptidyl-Lys metalloprotease; ^b*Grifola frondosa*; ^c*Pleurotus ostreatus*.

TABLE 4: Kinetic parameters for the hydrolysis of Arg-Arg-NA and metal contents of various metal-DPP IIIs.

Enzyme	K_M ($\times 10^{-5}$ M)	k_{cat} (s ⁻¹)	k_{cat}/K_M ($\times 10^4$ M ⁻¹ s ⁻¹)	Metal content (mol/mol of protein)
Zn ²⁺ -DPP III	8.1 (± 1.0)	7.1 (± 0.2)	8.8	0.8 (± 0.1)
Co ²⁺ -DPP III	8.2 (± 0.9)	7.0 (± 0.1)	8.5	1.0 (± 0.1)
Cu ²⁺ -DPP III	9.9 (± 1.1)	10.1 (± 0.3)	10.2	1.1 (± 0.1)

TABLE 5: EPR parameters of Cu²⁺ proteases.

	$g_{\perp}$	$g_{\parallel}$	$A_{\parallel}$ ($\times 10^{-4}$ cm ⁻¹)
Cu ²⁺ -DPP III ^a	2.06	2.27	167
Cu ²⁺ -del-DPP III ^b	2.06	2.27	161
Cu ²⁺ -thermolysin ^c	2.06	2.26	163
Cu ²⁺ -aminopeptidase B ^d	2.06	2.27	157
Cu ²⁺ -carboxypeptidase A ^e	2.05	2.33	115

References ^a[9], ^b[10], ^c[8], ^d[22], and ^e[6].

5. Conclusions

In this paper, we compared metal flexibility with the geometry of metal coordination of metalloproteases, to investigate why DPP III shows metal tolerance. Metal substitution of Zn(II) by Co(II) or Mn(II) on metalloproteases generally maintains catalytic activity, because the metal coordination geometries of Zn(II), Co(II), and Mn(II) are flexible. Most

Cu(II)-substituted enzymes could not restore the catalytic activities, because the Cu(II) coordination geometry is very rigid. However, Cu(II)-substituted DPP III showed the same catalytic activity as that of Zn(II)-DPP III. We then studied the metal flexibilities and metal coordination geometries of many metalloproteases, especially DPP III and del-DPP III, but we could not prove a relation between the metal flexibility and the metal coordination geometry. The metal tolerance of DPP III might depend on the flexibility of the metal-binding motif, not on the metal coordination geometry. By comparison of the 3D structure of active sites of DPP III and del-DPP III, both coordination geometries are seen to be similar, while the positions of catalytic amino acid residues (Glu) on those zinc-binding motifs are slightly different. We conclude that the catalytic site of Cu(II)-DPP III could be flexible enough to form the catalytic complex, with substrate and H₂O.

References

- [1] N. D. Rawlings and A. J. Barrett, "Introduction: metalloproteases and their clans," in *Handbook of Proteolytic Enzyme*, A. J. Barrett, N. D. Rawlings, and J. F. Woessner, Eds., pp. 231–268, Academic Press, San Diego, Calif, USA, 2nd edition, 2004.
- [2] A. David, "Catalytic mechanisms for metalloproteases," in *Handbook of Proteolytic Enzyme*, A. J. Barrett, N. D. Rawlings, and J. F. Woessner, Eds., pp. 268–289, Academic Press, San Diego, Calif, USA, 2nd edition, 1998.
- [3] K. Fukasawa, K. M. Fukasawa, M. Kanai, S. Fujii, J. Hirose, and M. Harada, "Dipeptidyl peptidase III is a zinc metalloexopeptidase. Molecular cloning and expression," *Biochemical Journal*, vol. 329, no. 2, pp. 275–282, 1998.
- [4] K. Fukasawa, K. M. Fukasawa, H. Iwamoto, J. Hirose, and M. Harada, "The HELLGH motif of rat liver dipeptidyl peptidase III is involved in zinc coordination and the catalytic activity

- of the enzyme," *Biochemistry*, vol. 38, no. 26, pp. 8299–8303, 1999.
- [5] F. X. Gomis-Rüth, F. Grams, I. Yiallourous et al., "Crystal structures, spectroscopic features, and catalytic properties of cobalt(II), copper(II), nickel(II), and mercury(II) derivatives of the zinc endopeptidase astacin. A correlation of structure and proteolytic activity," *Journal of Biological Chemistry*, vol. 269, no. 25, pp. 17111–17117, 1994.
  - [6] R. C. Rosenberg, C. A. Root, P. K. Bernstein, and H. B. Gray, "Spectral studies of copper(II) carboxypeptidase A and related model complexes," *Journal of the American Chemical Society*, vol. 97, no. 8, pp. 2092–2096, 1975.
  - [7] A. F. Monzingo and B. W. Matthews, "Binding of N-carboxymethyl dipeptide inhibitors to thermolysin determined by X-ray crystallography: a novel class of transition-state analogues for zinc peptidases," *Biochemistry*, vol. 23, no. 24, pp. 5724–5729, 1984.
  - [8] I. Bertini, G. Canti, H. Kozłowski, and A. Scozzafava, "Spectroscopic characterization of copper(II) thermolysin," *Journal of the Chemical Society Dalton Transactions*, no. 8, pp. 1270–1273, 1979.
  - [9] J. Hirose, H. Iwamoto, I. Nagao et al., "Characterization of the metal-substituted dipeptidyl peptidase III (rat liver)," *Biochemistry*, vol. 40, no. 39, pp. 11860–11865, 2001.
  - [10] J. Hirose, H. Kamigakiuchi, H. Iwamoto et al., "The metal-binding motif of dipeptidyl peptidase III directly influences the enzyme activity in the copper derivative of dipeptidyl peptidase III," *Archives of Biochemistry and Biophysics*, vol. 431, no. 1, pp. 1–8, 2004.
  - [11] C. Mazzocco, K. M. Fukasawa, A. A. Raymond, and J. Puiroux, "Purification, partial sequencing and characterization of an insect membrane dipeptidyl aminopeptidase that degrades the insect neuropeptide proctolin," *European Journal of Biochemistry*, vol. 268, no. 18, pp. 4940–4949, 2001.
  - [12] "Homo sapiens dipeptidyl-peptidase 3 (DPP3), transcript variant 1, mRNA," NCBI Reference Sequence: NM 005700.3.
  - [13] B. W. Matthews, "Structural basis of the action of thermolysin and related zinc peptidases," *Accounts of Chemical Research*, vol. 21, no. 9, pp. 333–340, 1988.
  - [14] J. F. Medina, A. Wetterholm, O. Rådmark et al., "Leukotriene A4 hydrolase: determination of the three zinc-binding ligands by site-directed mutagenesis and zinc analysis," in *Proceedings of the National Academy of Sciences of the United States of America*, vol. 88, no. 17, pp. 7620–7624, 1991.
  - [15] P. K. Baral, N. Jajčanin-Jozić, S. Deller, P. Macheroux, M. Abramić, and K. Gruber, "The first structure of dipeptidyl-peptidase III provides insight into the catalytic mechanism and mode of substrate binding," *Journal of Biological Chemistry*, vol. 283, no. 32, pp. 22316–22324, 2008.
  - [16] M. M. G. M. Thunnissen, P. Nordlund, and J. Z. Haeggström, "Crystal structure of human leukotriene A4 hydrolase, a bifunctional enzyme in inflammation," *Nature Structural Biology*, vol. 8, no. 2, pp. 131–135, 2001.
  - [17] K. M. Fukasawa, J. Hirise, T. Hata, and Y. Ono, "In rat dipeptidyl peptidase III, His568 is essential for catalysis, and Glu507 or Glu512 stabilizes the coordination bond between His455 or His450 and zinc ion," *Biochimica et Biophysica Acta*, vol. 1804, pp. 2063–2069, 2010.
  - [18] C. Oefner, A. D'Arcy, M. Hennig, F. K. Winkler, and G. E. Dale, "Structure of human neutral endopeptidase (neprilysin) complexed with phosphoramidon," *Journal of Molecular Biology*, vol. 296, no. 2, pp. 341–349, 2000.
  - [19] D. C. Rees, M. Lewis, and W. N. Lipscomb, "Refined crystal structure of carboxypeptidase A at 1.54 Å resolution," *Journal of Molecular Biology*, vol. 168, no. 2, pp. 367–387, 1983.
  - [20] T. Hori, T. Kumasaka, M. Yamamoto et al., "Structure of a new 'aspzincin' metalloendopeptidase from *Grifola frondosa*: implications for the catalytic mechanism and substrate specificity based on several different crystal forms," *Acta Crystallographica Section D*, vol. 57, no. 3, pp. 361–368, 2001.
  - [21] T. Tanaka and E. Ichishima, "Molecular properties of aminopeptidase Ey as a zinc-metalloenzyme," *International Journal of Biochemistry*, vol. 25, no. 11, pp. 1681–1688, 1993.
  - [22] J. Hirose, T. Ohsaki, N. Nishimoto et al., "Characterization of the metal-binding site in aminopeptidase B," *Biological and Pharmaceutical Bulletin*, vol. 29, no. 12, pp. 2378–2382, 2006.
  - [23] T. Achstetter, C. Ehmann, and D. H. Wolf, "Proteinase yscD. Purification and characterization of a new yeast peptidase," *Journal of Biological Chemistry*, vol. 260, no. 8, pp. 4585–4590, 1985.
  - [24] M. Büchler, U. Tislar, and D. H. Wolf, "Proteinase yscD (oligopeptidase yscD). Structure, function and relationship of the yeast enzyme with mammalian thimet oligopeptidase (metalloendopeptidase, EP 24.15)," *European Journal of Biochemistry*, vol. 219, no. 1–2, pp. 627–639, 1994.
  - [25] J. R. Klein, U. Klein, M. Schad, and R. Plapp, "Cloning, DNA sequence analysis and partial characterization of pepN, a lysyl aminopeptidase from *Lactobacillus delbrueckii* ssp. *lactis* DSM7290," *European Journal of Biochemistry*, vol. 260, pp. 4585–4590, 1993.
  - [26] T. R. Yan, N. Azuma, S. Kaminogawa, and K. Yamauchi, "Purification and characterization of substrate-size-recognizing metalloendopeptidase from *Streptococcus cremoris* H61," *Applied and Environmental Microbiology*, vol. 53, pp. 2296–2302, 1987.
  - [27] V. Monnet, M. Nardi, A. Chopin, M. C. Chopin, and J. C. Gripon, "Biochemical and genetic characterization of PepF, an oligopeptidase from *Lactococcus lactis*," *Journal of Biological Chemistry*, vol. 269, no. 51, pp. 32070–32076, 1994.
  - [28] P. C. Chang and Y. H. W. Lee, "Extracellular autoproducting of a metalloprotease from *Streptomyces cacaui*," *Journal of Biological Chemistry*, vol. 267, no. 6, pp. 3952–3958, 1992.
  - [29] B. Holmquist and B. L. Vallee, "Metal substitutions and inhibition of thermolysin: spectra of the cobalt enzyme," *Journal of Biological Chemistry*, vol. 249, no. 14, pp. 4601–4607, 1974.
  - [30] P. S. T. Tan, K. M. Pos, and W. N. Konings, "Purification and characterization of an endopeptidase from *Lactococcus lactis* subsp. *cremoris* Wg2," *Applied and Environmental Microbiology*, vol. 57, no. 12, pp. 3593–3599, 1991.
  - [31] R. Baankreis, S. Van Schalkwijk, A. C. Alting, and F. A. Exterkate, "The occurrence of two intracellular oligoendopeptidases in *Lactococcus lactis* and their significance for peptide conversion in cheese," *Applied Microbiology and Biotechnology*, vol. 44, no. 3–4, pp. 386–392, 1995.
  - [32] S. Ayora and F. Götz, "Genetic and biochemical properties of an extracellular neutral metalloprotease from *Staphylococcus hyicus* subsp. *hyicus*," *Molecular and General Genetics*, vol. 242, no. 4, pp. 421–430, 1994.
  - [33] L. A. Dreyfus and B. H. Iglewski, "Purification and characterization of an extracellular protease of *Legionella pneumophila*," *Infection and Immunity*, vol. 51, no. 3, pp. 736–743, 1986.
  - [34] P. Diermayr, S. Kroll, and H. Klostermeyer, "Influence of EDTA and metal ions on a metalloproteinase from *Pseudomonas fluorescens* biotype I," *Biological Chemistry Hoppe-Seyler*, vol. 368, no. 1, pp. 57–61, 1987.

- [35] T. Nonaka, Y. Hashimoto, and K. Takio, "Kinetic characterization of lysine-specific metalloendopeptidases from *Grifola frondosa* and *Pleurotus ostreatus* fruiting bodies," *Journal of Biochemistry*, vol. 124, no. 1, pp. 157–162, 1998.
- [36] Y. Nitanaï, Y. Satow, H. Adachi, and M. Tsujimoto, "Crystal structure of human renal dipeptidase involved in β -lactam hydrolysis," *Journal of Molecular Biology*, vol. 321, no. 2, pp. 177–184, 2002.
- [37] T. H. Tahirov, H. Oki, T. Tsukihara et al., "High-resolution crystals of methionine aminopeptidase from *Pyrococcus furiosus* obtained by water-mediated transformation," *Journal of Structural Biology*, vol. 121, no. 1, pp. 68–72, 1998.
- [38] M. C. J. Wilce, C. S. Bond, N. E. Dixon et al., "Structure and mechanism of a proline-specific aminopeptidase from *Escherichia coli*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 7, pp. 3472–3477, 1998.
- [39] E. Dobrovetsky, A. Dong, A. Seitova et al., "Crystal structure of human dipeptidyl peptidase III," in *Structural Genomics Consortium (SGC)*, 2009.
- [40] S. I. Miyoshi, K. Kawata, K. I. Tomochika, and S. Shinoda, "The hemagglutinating activity of *Vibrio vulnificus* metallo-protease," *Microbiology and Immunology*, vol. 43, no. 1, pp. 79–82, 1999.