Research Article

Finite Groups Whose Certain Subgroups of Prime Power Order Are S -Semipermutable

Mustafa Obaid

Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80111, Jeddah 21589, Saudi Arabia

Correspondence should be addressed to Mustafa Obaid, drmobaid@gmail.com

Received 17 July 2011; Accepted 3 August 2011

Academic Editor: A. Kiliçman

Copyright © 2011 Mustafa Obaid. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let G be a finite group. A subgroup H of G is said to be S -semipermutable in G if H permutes with every Sylow p -subgroup of G with $(p, |H|) = 1$. In this paper, we study the influence of S -permutability property of certain abelian subgroups of prime power order of a finite group on its structure.

1. Introduction

All groups considered in this paper will be finite. Two subgroups H and K of a group G are said to permute if $HK = KH$. It is easily seen that H and K permute if and only if HK is a subgroup of G . We say, following Kegel [1], that a subgroup of G is S -quasinormal in G if it permutes with every Sylow subgroup of G . Chen [2] introduced the following concept: a subgroup H of G is said to be S -semipermutable in G if H permutes with every Sylow p -subgroup of G with $(p, |H|) = 1$. Obviously, every S -quasinormal subgroup of G is an S -semipermutable subgroup of G . In contrast to the fact that every S -quasinormal subgroup of G is a subnormal subgroup of G (see [1]), it does not hold in general that every S -semipermutable subgroup of G is a subnormal subgroup of G . It suffices to consider the alternating group of degree 4.

Several authors have investigated the structure of a finite group when some information is known about some subgroups of prime power order in the group. Huppert [3] proved that a finite group G is solvable provided that all subgroups of prime order are normal in G . Buckley [4], proved that a group G of odd order is supersolvable provided that all subgroups of prime order are normal in G . Srinivasan [5], and proved that a finite group G is supersolvable if the maximal subgroups of every Sylow subgroup of G are normal in G .

Developing the result of Srinivasan, Ramadan [6] proved that if G is a solvable group and the maximal subgroups of every Sylow subgroup of the Fitting subgroup $F(G)$ of G are normal in G , then G is supersolvable.

For a finite p -group P , we denote

$$\Omega(P) = \Omega_1(P) \quad \text{if } p > 2, \quad \Omega(P) = \langle \Omega_1(P), \Omega_2(P) \rangle \quad \text{if } p = 2, \quad (1.1)$$

where $\Omega_i(P) = \langle x \in P : |x| = p^i \rangle$.

Of late there has been a considerable interest to investigate the influence of the abelian subgroups of largest possible exponent of prime power order (we call such subgroups ALPE-subgroups) on the structure of the group. Asaad et al. [7] proved that if G is a group such that for every prime p and every Sylow p -subgroup G_p of G , the ALPE-subgroups of G_p (resp., $\Omega(G_p)$) are normal in G , then G is supersolvable. Ramadan [8] proved the following two results. (1) Let G be a group such that for every prime p and every Sylow p -subgroup G_p of G , the ALPE-subgroups of G_p (resp., $\Omega(G_p)$) are S -quasinormal in G , then G is supersolvable. (2) Let K be a normal subgroup of G such that G/K is supersolvable. If for every prime p and every Sylow p -subgroup K_p of K , the ALPE-subgroups of K_p (resp., $\Omega(K_p)$) are S -quasinormal in G , then G is supersolvable.

In this paper, we study the structure of a finite group under the assumption that certain subgroups of prime power order are S -semipermutable in the group. We focus our attention on S -semipermutability property of the ALPE-subgroups of a fixed ALPE-subgroup having maximal order of the Sylow subgroups of a finite group. Furthermore, we improve and extend the above-mentioned results by using the concept of S -semipermutability.

2. Preliminaries

In this section, we give some results which will be useful in the sequel.

Lemma 2.1 (see [2, Lemmas 1 and 2]). *Let G be a group.*

- (i) *If H is a S -semipermutable subgroup of G and K is a subgroup of G such that $H \leq K \leq G$, then H is S -semipermutable in K .*
- (ii) *Let π be a set of primes, N a normal π' -subgroup of G , and H a π -subgroup of G . If H is S -semipermutable in G , then HN/N is S -semipermutable in G/N .*

Lemma 2.2 (see [9, Lemma A]). *Let H be a p -subgroup of G ; for some prime p . Then H is S -quasinormal in G if and only if $O^p(G) \leq N_G(H)$, where $O^p(G)$ is the normal subgroup of G generated by all p' -elements of G .*

Lemma 2.3. *Let H be a p -subgroup of G , p is a prime. Then the following statements are equivalent:*

- (i) *H is S -quasinormal in G ;*
- (ii) *$H \leq O_p(G)$ and H is S -semipermutable in G .*

Proof. (i) $\Rightarrow$ (ii): Suppose that H is S -quasinormal in G . So it follows by [1, Satz 1, page 209] that H is subnormal in G and then by [10, Lemma 8.6(a), page 28] that $H \leq O_p(G)$. Since H is S -quasinormal in G , obviously, it is S -semipermutable in G . Thus (ii) holds.

(ii) $\Rightarrow$ (i): Since H is S -semipermutable in G , then $HG_q = G_qH$ for every Sylow q -subgroup G_q of G with $(q, |H|) = 1$. Clearly, $H = O_p(G) \cap HG_q$ is normal in HG_q and so $G_q \leq N_G(H)$. Thus $O^p(G) \leq N_G(H)$. Applying Lemma 2.2, we have that H is S -quasinormal in G . Thus (i) holds. $\square$

Lemma 2.4 (see [7, Theorem 4, page 253]). *Let P be a normal p -subgroup of G . If the ALPE-subgroups of P are normal in G , then P is supersolvably embedded in G .*

Lemma 2.5 (see [11, Lemma 3.8, page 2245]). *Let p be the smallest prime dividing the order of a group G , and let G_p be a Sylow p -subgroup of G . If $\Omega(G_p) \leq \text{genz}_\infty(G)$, then G is p -nilpotent.*

Lemma 2.6 (see [12, Lemma 2.6]). *Let N be a nontrivial normal subgroup of a group G . If $N \cap \Phi(G) = 1$, then the Fitting subgroup $F(N)$ of N is the direct product of minimal normal subgroups of G which are contained in $F(N)$.*

Lemma 2.7 (see [13, Lemma 3.3.1, page 23]). *Suppose that G_p is a normal Sylow p -subgroup of G and that $\Omega(G_p)K$ is supersolvable, where K is a p' -Hall subgroup of G . Then G is supersolvable.*

3. Main Results

Theorem 3.1. *Let p be the smallest prime dividing the order of a group G , and let G_p be a Sylow p -subgroup of G . Fix an ALPE-subgroup P of G_p having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is p -nilpotent.*

Proof. Suppose that the theorem is false, and let G be a counterexample of minimal order. We prove the following steps.

(1) *If $P \leq M < G$, Then M Is p -Nilpotent*

It is clear to see by Lemma 2.1 that the ALPE-subgroups of P are S -semipermutable in M , so that M satisfies the hypothesis of the theorem. Thus, the minimality of G yields that M is p -nilpotent.

(2) *$N_G(P)$ Is p -Nilpotent*

Suppose that P is normal in G . Let H be an ALPE-subgroup of P (in particular, we may take $H = P$). By hypothesis, H is S -semipermutable in G and so by Lemma 2.3, we have that H is S -quasinormal in G . Hence HG_q is a subgroup of G , where G_q is a Sylow q -subgroup of G with $q \neq p$. Clearly, H is a subnormal Hall subgroup of HG_q . Thus H is normal in HG_q and hence H is normal in PG_q as P is abelian. Thus P is supersolvably embedded in PG_q by Lemma 2.4 and so $P \leq Q_\infty(PG_q)$. Since $Q_\infty(PG_q) \leq \text{genz}_\infty(PG_q)$ by [14, page 34], it follows by Lemma 2.5 that PG_q is p -nilpotent. Thus $PG_q = P \times G_q$. Hence $G_q \leq C_G(P)$, so that $O^p(G) \leq C_G(P)$. If $C_G(P) < G$, then $C_G(P)$ is p -nilpotent by (1). Thus $O^p(G)$ is p -nilpotent and so G is p -nilpotent: a contradiction. Thus we may assume that $C_G(P) = G$. Then $P \leq Z(G)$, in particular, $P \leq Z(G_p)$. So, $P = G_p$ by the maximality of P and we have by [15, Theorem 4.3, page 252] that G is p -nilpotent: a contradiction. Thus we may assume that $N_G(P) < G$. According to (1), we have that $N_G(P)$ is p -nilpotent.

(3) $O_{p'}(G) = 1$

If $O_{p'}(G) \neq 1$, we consider the quotient group $G/O_{p'}(G)$. Clearly, $G_p O_{p'}(G)/O_{p'}(G)$ is a Sylow p -subgroup of $G/O_{p'}(G)$ and $PO_{p'}(G)/O_{p'}(G)$ is an ALPE-Subgroup of $G_p O_{p'}(G)/O_{p'}(G)$ having maximal order. By Lemma 2.1, the hypotheses are inherited over $G/O_{p'}(G)$. Thus, the minimality of G implies that $G/O_{p'}(G)$ is p -nilpotent, hence G is p -nilpotent, which is a contradiction.

(4) $G = G_p G_q$, Where G_q Is a Sylow q -Subgroup of G with $q \neq p$

Since G is not p -nilpotent by [15, Theorem 4.5, page 253], there exists a subgroup H of G_p such that $N_G(H)$ is not p -nilpotent. But $N_G(G_p)$ is p -nilpotent by a similar argument of the proof of the step (2). Thus we may choose a subgroup H of G_p such that $N_G(H)$ is not p -nilpotent but $N_G(K)$ is p -nilpotent for every subgroup K of G_p with $H < K \leq G_p$. It is easy to see that $N_G(G_p) \leq N_G(H) \leq G$. If $N_G(H) < G$, it follows by (1) that $N_G(H)$ is p -nilpotent: a contradiction. Thus $N_G(H) = G$. This leads to $O_p(G) \neq 1$ and $N_G(K)$ is p -nilpotent for every subgroup K of G_p with $O_p(G) < K \leq G_p$. Now, by [15, Theorem 4.5, page 253] again, we see that $G/O_p(G)$ is p -nilpotent and therefore that G is p -solvable. Since G is p -solvable, for any $q \in \pi(G)$ with $q \neq p$, there exists a Sylow q -subgroup G_q of G such that $G_p G_q \leq G$ by [15, Theorem 3.5, page 229]. If $G_p G_q < G$, then $G_p G_q$ is p -nilpotent by (1) and hence $O_p(G)G_q$ is p -nilpotent. Thus $O_p(G)G_q = O_p(G) \times G_q$. This leads to $G_q \leq C_G(O_p(G)) \leq O_p(G)$ by [15, Theorem 3.2, page 228] as $O_{p'}(G) = 1$ by (3), which is a contradiction. Thus $G = G_p G_q$.

(5) *The Final Contradiction*

Let N be a minimal normal subgroup of G such that $N \leq O_p(G)$. Clearly, $N \cap Z(G_p) \neq 1$ and so $Z(G_p) \leq P$ by the maximality of P . Hence $1 \neq N \cap Z(G_p) \leq N \cap P$. By hypothesis, $PG_q \leq G$ for any Sylow q -subgroup G_q of G with $(q, |P|) = 1$. It is easy to see that $N \cap P = N \cap PG_q \triangleleft PG_q$. Thus $O^p(G) \leq N_G(N \cap P)$. If $N_G(N \cap P) < G$, then by (1), $N_G(N \cap P)$ is p -nilpotent. Hence $O^p(G)$ is p -nilpotent and so also does G : a contradiction. Thus we may assume that $N_G(N \cap P) = G$. By the minimality of N and since $N \cap P \neq 1$, we have that $N \cap P = N$ and so $N \leq P$. If $PG_q < G$, then PG_q is p -nilpotent by (1) and hence NG_q is p -nilpotent. Thus $NG_q = N \times G_q$ and so $G_q \leq C_G(N)$. Thus by (4), $G/C_G(N)$ is a p -group and so by [14, Theorem 6.3, page 221], $N \leq Z_\infty(G)$. Since $Z_\infty(G) \leq Q_\infty(G)$, we have that $N \leq Q_\infty(G)$ which implies that N is supersolvably embedded in G and so clearly that $|N| = p$. Thus, it is easy to see that the quotient group G/N satisfies the hypothesis of the theorem by Lemma 2.1. Now, by the minimality of G , we see that G/N is p -nilpotent. Since the class of all p -nilpotent groups is a saturated formation, we have that N is the unique minimal normal subgroup of G and $N \not\leq \Phi(G)$. Thus $\Phi(G) = 1$ and hence $N = O_p(G)$ by Lemma 2.6 and so $F(G) = O_p(G) = N$ by (3). Hence $G_q \leq C_G(F(G))$. Since G is solvable, it follows by [15, Theorem 2.6, page 216] that $C_G(F(G)) \leq F(G) = O_p(G)$: a contradiction. Thus we must have $G = PG_q$. Let G_q^* be a Sylow q -subgroup of $N_G(P)$. By (2), we have that $G_q^* \triangleleft N_G(P)$. Hence $N_G(P) = PG_q^* = P \times G_q^*$. Thus $P \leq Z(N_G(P))$, and, therefore, G is p -nilpotent by [15, Theorem 4.3, page 252]: a final contradiction. $\square$

We need the following result.

Theorem 3.2. *Let $\mathcal{F}$ be a saturated formation containing the class of supersolvable groups $\mathcal{U}$. Let G_p be a normal Sylow p -subgroup of a group G such that $G/G_p \in \mathcal{F}$. Fix an ALPE-subgroup P of G_p having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then $G \in \mathcal{F}$.*

Proof. We treat the following two cases.

Case 1. $O_{p'}(G) \neq 1$.

Clearly, $G_p O_{p'}(G)/O_{p'}(G)$ is a normal Sylow p -subgroup of $G/O_{p'}(G)$ and $PO_{p'}(G)/O_{p'}(G)$ is an ALPE-subgroup of $G_p O_{p'}(G)/O_{p'}(G)$ having maximal order. By hypothesis and Lemma 2.1, the ALPE-subgroups of $PO_{p'}(G)/O_{p'}(G)$ are S -semipermutable in $G/O_{p'}(G)$. Clearly,

$$\frac{(G/G_p)}{(G_p O_{p'}(G)/G_p)} \cong \frac{G}{G_p O_{p'}(G)} \cong \frac{(G/O_{p'}(G))}{(G_p O_{p'}(G)/O_{p'}(G))} \in \mathcal{F}. \quad (3.1)$$

Thus, our hypothesis carries over to $G/O_{p'}(G)$ and so $G/O_{p'}(G) \in \mathcal{F}$ by induction on the order of G . Therefore, $G/(O_{p'}(G) \cap G_p) \cong G \in \mathcal{F}$.

Case 2. $O_{p'}(G) = 1$.

Let H be an ALPE-subgroup of P . Then H is S -quasinormal in G by Lemma 2.3 and hence $O^p(G) \leq N_G(H)$ by Lemma 2.2. Let $T = PO^p(G)$. Then H is normal in T . Thus Lemma 2.4 implies that P is supersolvably embedded in T . Then, $T/C_T(P)$ is supersolvable by [14, Lemma 7.15, page 35]. Clearly, $T_p = G_p \cap T \triangleleft T$, where T_p is a Sylow p -subgroup of T . Let Q be a p' -subgroup of $C_T(P)$. Then $QP = Q \times P$ is a group of automorphisms of $T_p = O_p(T)$. But $C_{T_p}(P) = P$, and consequently, Q acts trivially on $C_{T_p}(P)$. Then Q acts trivially on T_p by [15, Theorem 3.4, page 179], that is, $Q \leq C_T(T_p)$. It is easy to see that T is subnormal in G and so $O_{p'}(T) \leq O_{p'}(G) = 1$. Hence $F(T) = T_p$. Since T is solvable, it follows by [15, Theorem 2.6, page 216] that $Q \leq C_T(F(T)) \leq F(T) = T_p$: a contradiction. Hence $C_T(P)$ must be a p -group and so $C_T(P) = P$. Thus, $T/C_T(P) = T/P$ is supersolvable which implies that T is supersolvable by [16, Theorem 4]. Thus $O^p(G)$ is supersolvable and therefore, $G = G_p O^p(G)$ is supersolvable by [17, Exercise 7.2.23, page 159]. Hence, $G \in \mathcal{U} \subseteq \mathcal{F}$. □

As an immediate consequence of Theorem 3.2, we have the following theorem.

Corollary 3.3. *Let G_p be a normal Sylow p -subgroup of a group G such that G/G_p is supersolvable. Fix an ALPE-subgroup P of G_p having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

We now prove the following theorem.

Theorem 3.4. *Let G be a group. For every prime p and every Sylow p -subgroup G_p of G , fix an ALPE-subgroup P of G_p having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

Proof. By repeated applications of Theorem 3.1, the group G has a Sylow tower of supersolvable type. Hence G has a normal Sylow p -subgroup G_p , where p is the largest prime dividing the order of G . By Lemma 2.1, our hypothesis carries over to G/G_p . Thus G/G_p is

supersolvable by induction on the order of G . Now, it follows from Corollary 3.3 that G is supersolvable. $\square$

As an immediate consequence of Theorem 3.4, we have the following corollary.

Corollary 3.5 (Asaad et al. [7]). *If G is a group such that the ALPE-subgroups of every Sylow subgroup of G are normal in G , then G is supersolvable.*

Corollary 3.6 (Ramadan [8]). *If G is a group such that the ALPE-subgroups of every Sylow subgroup of G are S -quasinormal in G , then G is supersolvable.*

We need the following Lemma.

Lemma 3.7. *Let K be a normal p -subgroup of a group G such that G/K is supersolvable. Fix an ALPE-subgroup P of K having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

Proof. Let G_p be a Sylow p -subgroup of G . We treat the following two cases.

Case 1. $K = G_p$.

Then by Corollary 3.3, G is supersolvable.

Case 2. $K < G_p$.

Put $\pi(G) = \{p_1, p_2, \dots, p_n\}$, where $p_1 > p_2 > \dots > p_n$. Since G/K is supersolvable, it follows by [18, Theorem 5, page 5] that G/K possesses supersolvable subgroups M/K and L/K such that $|G/K : M/K| = p_1$ and $|G/K : L/K| = p_n$. Since M/K and L/K are supersolvable, it follows that M and L are supersolvable by induction on the order of G . Since $|G : M| = |G/K : M/K| = p_1$ and $|G : L| = |G/K : L/K| = p_n$, it follows again by [18, Theorem 5, page 5] that G is supersolvable. $\square$

Now, we can prove the following theorem.

Theorem 3.8. *Let K be a normal subgroup of G such that G/K is supersolvable. For every prime p dividing the order of K and every Sylow p -subgroup K_p of K , fix an ALPE-subgroup P of K_p having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

Proof. By Lemma 2.1, the ALPE-subgroups of P are S -semipermutable in K . Hence K is supersolvable by Theorem 3.4. Thus K has a normal Sylow p -subgroup K_p , where p is the largest prime dividing the order of K . Since K_p is characteristic in K and $K \triangleleft G$, we have that $K_p \triangleleft G$. Clearly, $(G/K_p)/(K/K_p) \cong G/K$ is supersolvable. By Lemma 2.1, our hypothesis carries over to G/K_p and hence G/K_p is supersolvable by induction on the order of G . Now, it follows from Lemma 3.7 that G is supersolvable. $\square$

As an immediate consequence of Theorem 3.8, we have the following corollary.

Corollary 3.9 (Ramadan [8]). *Let K be a normal subgroup of a group G such that G/K is supersolvable. If the ALPE-subgroups of every Sylow subgroup of K are S -quasinormal in G , then G is supersolvable.*

4. Similar Results

Following similar arguments to those of Theorem 3.1, it is possible to prove the following result.

Theorem 4.1. *Let p be the smallest prime dividing the order of a group G and let G_p be a Sylow p -subgroup of G . Fix an ALPE-subgroup P of $\Omega(G_p)$ having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is p -nilpotent.*

We prove the following lemma.

Lemma 4.2. *Let K be a normal p -subgroup of a group G such that G/K is supersolvable. Fix an ALPE-subgroup P of $\Omega(K)$ having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

Proof. Let G_p be a Sylow p -subgroup of G . We treat the following two cases.

Case 1. By [15, Theorem 2.1, page 221], there exists a p' -Hall subgroup T , which is a complement to G_p in G . Hence $G/G_p \cong T$ is supersolvable. Since $\Omega(G_p)$ is characteristic in G_p and $G_p \triangleleft G$, we have that $\Omega(G_p) \triangleleft G$. Clearly, $\Omega(G_p)T/\Omega(G_p) \cong T$ is supersolvable. Thus, our hypothesis and Corollary 3.3 imply that $\Omega(G_p)T$ is supersolvable. Therefore, G is supersolvable by Lemma 2.7.

Case 2. Put $\pi(G) = \{p_1, p_2, \dots, p_n\}$, where $p_1 > p_2 > \dots > p_n$. Since G/K is supersolvable, it follows by [18, Theorem 5, page 5] that G/K possesses supersolvable subgroups M/K and L/K such that $|G/K : M/K| = p_1$ and $|G/K : L/K| = p_n$. Since M/K and L/K are supersolvable, it follows that M and L are supersolvable by induction on the order of G . Since $|G : M| = |G/K : M/K| = p_1$ and $|G : L| = |G/K : L/K| = p_n$, it follows again by [18, Theorem 5, page 5] that G is supersolvable. □

By a similar proof to the proof of Theorem 3.4, we can prove the following theorem.

Theorem 4.3. *Let G be a group. For every prime p and every Sylow p -subgroup G_p of G , fix an ALPE-subgroup P of $\Omega(G_p)$ having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

As an immediate consequence of Theorem 4.3, we have the following corollary.

Corollary 4.4 (Asaad et al. [7]). *If G is a group such that for every prime p and every Sylow p -subgroup G_p , the ALPE-subgroups of $\Omega(G_p)$ are normal in G , then G is supersolvable.*

Corollary 4.5 (Ramadan [8]). *If G is a group such that for every prime p and every Sylow p -subgroup G_p , the ALPE-subgroups of $\Omega(G_p)$ are S -quasinormal in G , then G is supersolvable.*

We can now prove the following corollary.

Corollary 4.6. *Let K be a normal subgroup of G such that G/K is supersolvable. For every prime p dividing the order of K and every Sylow p -subgroup K_p of K , fix an ALPE-subgroup P of $\Omega(K_p)$ having maximal order. If the ALPE-subgroups of P are S -semipermutable in G , then G is supersolvable.*

Proof. By Lemma 2.1, the ALPE-subgroups of P are S -semipermutable in K . Hence K is supersolvable by Theorem 4.3. Thus K has a normal Sylow p -subgroup K_p , where p is the largest prime dividing the order of K . Since K_p is characteristic in K and $K \triangleleft G$, we have that $K_p \triangleleft G$. Clearly, $(G/K_p)/(K/K_p) \cong G/K$ is supersolvable. By Lemma 2.1, the hypothesis of our theorem carries over to G/K_p . Thus G/K_p is supersolvable by induction on the order of G and it follows that G is supersolvable by Lemma 4.2. $\square$

Remarks 4.7. (a) The converse of Theorem 3.4 is not true. For example, set $G = S_3 \times Z_3$, where $S_3 = \langle x, y \mid x^3 = y^2 = 1, yx = x^2y \rangle$ and $Z_3 = \langle z \mid z^3 = 1 \rangle$. Clearly, G is supersolvable and G has an abelian Sylow 3-subgroup of exponent 3. It is easy to check that G contains a subgroup $\langle xz \rangle$ of order 3 which fails to be S -semipermutable in G .

(b) Theorem 4.3 is not true when the smallest prime dividing the order of G is even and $\Omega(G_p) = \Omega_1(G_p)$, where G_p is a Sylow p -subgroup of G . For example, if Q is the quaternion group $\langle a, b \mid a^4 = 1, b^2 = a^2, b^{-1}ab = a^{-1} \rangle$, C_9 is a cyclic group of order 9 with generator c , and the action of C_9 on Q is given by $a^c = b$, $b^c = ab$, then the semidirect product of Q by C_9 is a group of even order in which every subgroup of prime order is S -semipermutable. Clearly, the semidirect product of Q by C_9 is not supersolvable (see Buckley [4, Examples (ii)]).

References

- [1] O. H. Kegel, "Sylow-gruppen und subnormalteiler endlicher gruppen," *Mathematische Zeitschrift*, vol. 78, pp. 205–221, 1962.
- [2] Z. Chen, "Generalization of the Schur-Zassenhaus theorem," *Journal of Mathematics*, vol. 18, no. 3, pp. 290–294, 1998.
- [3] B. Huppert, *Endliche Gruppen I*, Springer, Berlin, Germany, 1967.
- [4] J. Buckley, "Finite groups whose minimal subgroups are normal," *Mathematische Zeitschrift*, vol. 116, pp. 15–17, 1970.
- [5] S. Srinivasan, "Two sufficient conditions for supersolvability of finite groups," *Israel Journal of Mathematics*, vol. 35, no. 3, pp. 210–214, 1980.
- [6] M. Ramadan, "Influence of normality on maximal subgroups of Sylow subgroups of a finite group," *Acta Mathematica Hungarica*, vol. 59, no. 1-2, pp. 107–110, 1992.
- [7] M. Asaad, M. Ramadan, and M. Ezzat, "Finite groups with normal subgroups of prime power orders," *Pure Mathematics and Applications*, vol. 5, no. 3, pp. 251–256, 1994.
- [8] M. Ramadan, "The influence of S -quasinormality of some subgroups of prime power order on the structure of finite groups," *Archiv der Mathematik*, vol. 77, no. 2, pp. 143–148, 2001.
- [9] P. Schmid, "Subgroups permutable with all Sylow subgroups," *Journal of Algebra*, vol. 207, no. 1, pp. 285–293, 1998.
- [10] K. Doerk and T. Hawkes, *Finite Soluble Groups*, vol. 4, Walter de Gruyter, Berlin, Germany, 1992.
- [11] M. Asaad and M. Ezzat Mohamed, "On generalized hypercenter of a finite group," *Communications in Algebra*, vol. 29, no. 5, pp. 2239–2248, 2001.
- [12] Y. Wang, "Finite groups with some subgroups of Sylow subgroups c -supplemented," *Journal of Algebra*, vol. 224, no. 2, pp. 467–478, 2000.
- [13] M. Ezzat, *Finite groups in which some subgroups of prime power order are normal*, M.S. thesis, Cairo University, 1995.
- [14] Weinstein M., Ed., *Between Nilpotent and Solvable*, Polygonal Publishing House, New Jersey, NJ, USA, 1982.
- [15] D. Gorenstien, *Finite Groups*, Harper & Row, New York, NY, USA, 1967.
- [16] A. Yokoyama, "Finite solvable groups whose F -hypercenter contains all minimal subgroups Π ," *Archiv der Mathematik*, vol. 27, pp. 572–575, 1976.
- [17] W. R. Scott, *Group Theory*, Englewood cliffs, New Jersey, NJ, USA, 1964.
- [18] M. Asaad, "On the supersolvability of finite groups. I," *Annales Universitatis Scientiarum Budapestinensis*, vol. 18, pp. 3–7, 1975.