Research Article

Mutation Analysis of the *APC* Gene in a Chinese FAP Pedigree with Unusual Phenotype

S. Chen,¹ J. Zhou,² X. Zhang,¹ X. Zhou,² M. Zhu,¹ Y. Zhang,¹ G. Ma,¹ and J. Li¹

¹Laboratory of Genetics and Molecular Biology, Jiangsu Institute of Cancer Research, Nanjing 210009, China

²Department of General Surgery, Jiangsu Cancer Hospital, Nanjing 210009, China

Correspondence should be addressed to S. Chen, chensenqing2008@126.com

Received 2 January 2011; Accepted 19 January 2011

Academic Editors: A. J. Karayiannakis and A. A. te Velde

Copyright © 2011 S. Chen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background and Aim. Germline mutations of the adenomatous polyposis coli (*APC*) gene cause familial adenomatous polyposis (FAP), an autosomal dominant inherited disease mainly characterized by colorectal adenomatous polyposis. Genetic studies of FAP have shown that somatic *APC* mutations are dependent on the position of the germline *APC* mutation. However, the molecular mechanism underlying these genotype-phenotype associations for *APC* in Chinese remain largely unknown. **Patients and Methods.** In this study, we investigated the *APC* gene mutation in a Chinese FAP family by systematic screening with multiplex ligation-dependent probe amplification (MLPA), denaturing high-performance liquid chromatography (dHPLC), and DNA sequencing. Promoter methylation was detected by methylation-specific PCR. **Results.** The identical germline mutation c.1999 C>T (Q667X) of *APC* was identified in 5 affected members, among which 2 members carried somatic mutations of *APC*, one with promoter hypermethylation and the other with loss of wild-type allele in their adenomas. The somatic mutations were shown connected with the disease severity, demonstrating a unique genotype-phenotype association in this FAP pedigree. **Conclusion.** The study revealed the existence of novel pathogenic mutations in Chinese patients with FAP. Somatic mutations are of particular interest because of the unusual phenotypic features shown by patients.

1. Introduction

Familial adenomatous polyposis (FAP; MIM #175100) is a common hereditary syndrome characterized by early onset form of colorectal cancer based on multiple adenomas in the colon and rectum [1]. The penetrance of the genetic defect is approximately 100%. The responsible gene for FAP, the adenomatous polyposis coli (*APC*) gene, was isolated and localized to chromosome 5q21-22 by positional cloning [2, 3]. More than 900 different mutations have been registered in the Human Gene Mutation Database (<http://www.hgmd.cf.ac.uk/ac/>). Most mutations in *APC* are nonsense or frameshift mutations that cause premature truncation of the *APC* protein, and more than 60% have been localized within exon 15 [4, 5]. In addition, several studies revealed that the exon-spanning fragment variations in *APC* gene may be a frequent cause of FAP [6, 7]. In several studies, an association between the location of the *APC* mutation and the phenotype in FAP patients has been described [8]. Identification of the *APC* mutation in one affected individual of

a family opens the possibility of testing the at-risk members for the same mutation, thus preventing the development of colorectal cancer by the prophylactic removal of the bowel. Although the *APC* gene has been extensively studied in the Caucasian population, it has not been widely described in the Chinese population. The molecular mechanisms underlying the genotype-phenotype associations for *APC* remain largely unknown [9, 10]. Furthermore, the possible role of *APC* inactivation by methylation need to be further assessed. In this study, we screened the whole coding region of *APC* gene by MLPA, dHPLC, and DNA sequencing to identify the germline and somatic mutations in all of the affected members of a suspicious FAP family and discussed the genotype-phenotype association.

2. Patients and Methods

The FAP family pedigree (Figure 1) consists of 7 members, with 5 affected and 2 unaffected members. The proband

FIGURE 1: Pedigree of the family with autosomal dominant inheritance of familial adenomatous coli. d = died; Age = current age; Blackened symbols indicate affected individuals. Proband is denoted with a filled arrowhead, and roman numbers indicate generations.

was a 40-year-old female presenting to our hospital in August, 2005, with increasing intermittent rectal bleeding and abdominal pain for several months. Her family history included her father and two younger sisters, who died of rectal cancer in the age of 49, colorectal cancer at age of 38, and endometrial cancer at age of 27, respectively. Colonoscopy performed in the proband revealed numerous polyps carpeting the entire colon and rectum. Histological examination of biopsied specimens revealed findings of poorly differentiated adenocarcinomas, and the other polyps were diagnosed as tubular adenomas with mainly heavy-grade dysplasia. On the basis of these findings, the patient was diagnosed to have FAP. With informed consent from the patient and her family members, we collected their peripheral blood. For the numbered family members, colonoscopy examination was performed, and the found adenomas, adenocarcinomas, and the paired normal tissues were obtained. Each biopsy was assessed by the analysis of hematoxylin and eosin-stained sections. Fifteen control subjects were selected from medical checkup people in our hospital free from cancer.

DNA was isolated from peripheral blood and obtained tissue from consenting family members with the Qiaamp DNA Blood and Tissue Extraction kit (Qiagen Corp, Germany), respectively. MLPA analysis to determine the large fragment duplication or deletion and allelic loss in *APC* gene was performed using the Salsa MLPA kit P043 *APC* (MRC-Holland Corp, Amsterdam, The Netherlands) according to the protocol supplied with the kit with a few modifications according to the condition in our laboratory. The WAVE System 3500 (Transgenomic, Inc., Omaha, NE) and associated WAVE-Maker 4.1 software were used to analyze the exons 1 to 15 of *APC* gene, each of them in

FIGURE 2: *APC* exon 15A by denaturing high-performance liquid chromatography (dHPLC) in the FAP pedigree family. f1 is the proband; f2 and f7 are the unaffected members; f3–f6 are the affected members.

the range of 200–500 bp. Preparation and analysis of the PCR products was carried out as previously described [11]. In brief, oligonucleotide primers used in PCR for dHPLC had been reported previously [1, 4, 12]. PCR products from analyzed samples were denatured and then cooled slowly to room temperature before dHPLC to maximize heteroduplex formation. The precise gradients for temperature and buffers giving the optimal *T_M* for each fragment were calculated by WAVE-Maker 4.1 software (Transgenomic Inc., USA). The sequence variations could be detected using automated sequencing of variant fragments observed by dHPLC. Analysis of DNA sequencing was carried out on the ABI Prism

FIGURE 3: Sequence analysis: sequence spanning c.1999 C>T of individual 3, an affected family member. Arrow indicates a double peak at the site of the single nucleotide mutation, which leads to Q667X.

3100 Avant Genetic Analyzer (Applied Biosystems, Foster City, CA).

Methylation-specific PCR was used for methylation analysis of the APC promoter. Briefly, 1 μ g DNA was treated with sodium bisulfite using the CpGenome kit (Cat#S7820, Chemicon, USA) according to manufacturer's instructions. DNA methylation in the CpG islands of APC was determined by methylation-specific PCR. Primer sequences of APC promoter for the unmethylated DNA were 5'-GGGCTAGGGCTAGGCAGG-3' (sense) and 5'-GCTGCACCAATACAGCCAC-3' (antisense) amplifying the 98-bp product, while those for the methylated DNA were 5'-GGGTTAGGGTTAGGTAGG-3' (sense) and 5'-AACTACACCAATACAACCAC-3' (antisense). The PCR reaction was carried out using a hot start with a denaturation cycle of 94°C for 2 min and 80°C for 3 min, followed by 30 cycles of 30 s at 95°C, 30 s at 55°C, 40 s at 72°C, and final extension for 5 min at 72°C. The specificity of the reactions for methylated DNA is confirmed using unmethylated human sperm DNA and CpGenome Universal Methylated DNA.

3. Results

In this study, systematic screening strategy including MLPA, dHPLC, and sequencing was used to enable optimal mutation detection of germline and somatic mutations of APC gene in this FAP family.

3.1. Germline Mutations. The whole coding region of APC, exon by exon, was subsequently screened for mutation by dHPLC in constitutional DNA from this family. Only one aberrant peak elution profiles in the fragment of exon 15A from constitutional DNA were shown for each of Number 1, 3, 4, 5, and 6 (Figure 2), which indicates they harbor the identical germline mutation. Sequence analysis revealed a c.1999 C>T substitution predicted to cause the substitution at codon 667 from Glutamine to STOP (Q667X) (Figure 3). While the 2 unaffected family members and 15 normal individuals as control were homozygous C at the same position. By MLPA, large deletion or duplication of APC gene was not found in constitutional DNA. All members and 15

controls showed consistently normal control peak pattern as described by the kit's manufacturer.

3.2. Somatic Mutations. Somatic mutations of APC gene were screening by dHPLC and MLPA in the collected tissues from this family. Out of 5 affected members' tissues, the adenomas from Number 3 were exclusively found somatic loss of wild-type APC allele (Figure 4). The APC promoter methylation was exclusively identified in both carcinomas and adenomas from Number 1 and in none of the other members (Figure 5).

3.3. Clinical Manifestation of the Family Members. Colonoscopic examination performed in the affected members (nos. 1, 3, 4, 5, and 6) found that the number of colonic polyps presented in the colorectum substantially increased along with their growth older. For Number 1 with additional APC promoter, hypermethylation has a severe form of FAP with colon carcinomas and hepatoblastoma, the typical extracolonic manifestations. For Number 3 with loss of wild-type allele, the severe phenotype was manifested with tubular adenomas with mainly heavy-grade dysplasia in colon and gastric fundal polyps. No polyps were found in the colonoscopy performed in Numbers 2 and 7, corresponding to their APC-negative mutation.

4. Discussion

Examination of APC germline mutations in FAP families is being used as an efficient tool for predictive testing of subjects at risk. However, its relevance for clinical management of individual patients is still open to question.

By dHPLC and sequence analysis, we found an APC Q667X in constitutional DNA, which was identical in 5 affected members and also presented in their tissues. The affected members have manifested an aggressive form of FAP with early onset of colonic adenoma and extracolonic manifestations, and the unaffected members have shown normal phenotype. We suggest the germline mutation of Q667X is the cause of clinical phenotype of the family with FAP.

FIGURE 4: Peak pattern of exons from *APC* genes in MLPA. The *x*-axis shows the size of PCR products, and the *y*-axis reflects the relative quantity of PCR products. The arrows indicate a deletion of promoter region 1-2, exon 1–15 in the *APC* gene discovered in the adenomas from Number 3 of the family (a). The amount of PCR products in the *APC* gene was only half of a normal control (b).

FIGURE 5: Methylation of the *APC* promoter was exclusively identified in the adenomas and carcinomas from Number 1. Primer sets for methylation-specific PCR are designated as methylated (M) and unmethylated (U) DNA. N (1): normal cancer-free tissue from Number 1. A (1): Adenoma from Number 1. C (1): Carcinoma from Number 1. +: Positive control. -: blank control.

Recent studies revealed that 5% of classic FAP was ascribed to the large deletions in the *APC*, which cannot easily be detected by conventional mutation-detection techniques [6, 13]. Therefore, we used MLPA test in our mutation analysis strategy to screen for exon-spanning deletions in this family though no large deletion was found in the constitutional DNA from this family. However, Number 3 was found somatic loss of wild-type *APC* allele in the adenomas.

The *APC* protein contains seven 20-amino acid repeats (20AARs) which are involved in degrading the transcriptional cofactor beta-catenin and hence negatively regulate

Wnt signaling [14, 15]. The germline mutation Q667X encoded a limited amount of functional *APC* protein (23.4% of normal *APC* protein) and led to the truncated protein, which leaves zero 20AARs, implying a potential unfolding of beta-catenin. Moreover, the somatic loss of wild-type *APC* allele, as the “second hit”, was found in the adenomas from Number 3. Interestingly, the member carrying this somatic mutation had a more severe form of FAP with extracolonic manifestation though severe dysplasia may accumulate over time. We suggested that the somatic mutation may not only reduce the residual activity of the mutant germline allele but also inactivate the *APC* wild-type allele, and thus probably initiate the tumorigenesis. Several studies [16, 17] have shown that usually germline mutations between codons 1285 and 1378 were strongly associated with somatic loss of the wild-type *APC* allele in colorectal polyps in Caucasian population, which was inconsistent with our study. We thought it was the unique somatic *APC* mutation pattern that presents the clinical manifestation in this Chinese family pedigree.

The multitude of genetic changes may originate with a single mutation, which evolves towards tumorigenesis in conjunction with other genetic and epigenetic abnormalities, in particular CpG island hypermethylation [18]. In this family, we suggested that the promoter hypermethylation observed in adenomas and carcinomas from the proband may inactivate the second allele of *APC* and manifest the severe phenotype. These changes sometimes appear to be accumulated in all of the stages of colorectal tumorigenesis.

In FAP, correlations between site of mutation in the *APC* gene and severity of colonic polyposis or extracolonic

manifestations are demonstrated [19–21]. However, several findings indicated that the geographic differences in APC mutation pattern may manifest population differences in the FAP phenotype [22–24]. Therefore, further studies with more Chinese family pedigrees analysis are necessary to give the overall view of the mutation pattern and to evaluate the genotype-phenotype association in Chinese FAP families compared to FAP families in other populations.

Acknowledgments

The authors are grateful to the patients and their families for participating in this research study. This work was supported in part by the grants from Jiangsu Department of Health (H200846), and the “Six Peaks Talents Project” (no. 2008-B21) from Jiangsu Personnel Ministry.

References

- [1] J. Groden, A. Thliveris, W. Samowitz et al., “Identification and characterization of the familial adenomatous polyposis coli gene,” *Cell*, vol. 66, no. 3, pp. 599–600, 1991.
- [2] K. W. Kinzler, M. C. Nilbert, L. K. Su et al., “Identification of FAP locus genes from chromosome 5q21,” *Science*, vol. 253, no. 5020, pp. 661–665, 1991.
- [3] I. Nishisho, Y. Nakamura, Y. Miyoshi et al., “Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients,” *Science*, vol. 253, no. 5020, pp. 665–669, 1991.
- [4] H. Nagase and Y. Nakamura, “Mutations of the APC (Adenomatous Polyposis Coli) gene,” *Human Mutation*, vol. 2, no. 6, pp. 425–434, 1993.
- [5] C. Beroud and T. Soussi, “APC gene: database of germline and somatic mutations in human tumors and cell lines,” *Nucleic Acids Research*, vol. 24, no. 1, pp. 121–124, 1996.
- [6] O. M. Sieber, H. Lamlum, M. D. Crabtree et al., “Whole-gene APC deletions cause classical familial adenomatous polyposis, but not attenuated polyposis or “multiple” colorectal adenomas,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 5, pp. 2954–2958, 2002.
- [7] J. Meuller, G. Kanter-Smoler, A. O. H. Nygren et al., “Identification of genomic deletions of the APC gene in familial adenomatous polyposis by two independent quantitative techniques,” *Genetic Testing*, vol. 8, no. 3, pp. 248–256, 2004.
- [8] M. H. Nieuwenhuis and H. F. A. Vasen, “Correlations between mutation site in APC and phenotype of familial adenomatous polyposis (FAP): a review of the literature,” *Critical Reviews in Oncology/Hematology*, vol. 61, no. 2, pp. 153–161, 2007.
- [9] C. Albuquerque, C. Breukel, R. van der Luijt et al., “The ‘just-right’ signaling model: APC somatic mutations are selected based on a specific level of activation of the β -catenin signaling cascade,” *Human Molecular Genetics*, vol. 11, no. 13, pp. 1549–1560, 2002.
- [10] M. Crabtree, O. M. Sieber, L. Lipton et al., “Refining the relation between ‘first hits’ and ‘second hits’ at the APC locus: The ‘loose fit’ model and evidence for differences in somatic mutation spectra among patients,” *Oncogene*, vol. 22, no. 27, pp. 4257–4265, 2003.
- [11] M. Mihalatos, A. Apepos, J. K. Triantafyllidis et al., “Evaluation of dHPLC in mutation screening of the APC gene in a Greek FAP cohort,” *Anticancer Research*, vol. 23, no. 3, pp. 2691–2695, 2003.
- [12] Y. Miyoshi, H. Ando, H. Nagase et al., “Germ-line mutations of the APC gene in 53 familial adenomatous polyposis patients,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 89, no. 10, pp. 4452–4456, 1992.
- [13] E. T. Renkonen, P. Nieminen, W. M. Abdel-Rahman et al., “Adenomatous polyposis families that screen APC mutation-negative by conventional methods are genetically heterogeneous,” *Journal of Clinical Oncology*, vol. 23, no. 24, pp. 5651–5659, 2005.
- [14] B. Rubinfeld, B. Souza, I. Albert, S. Munemitsu, and P. Polakis, “The APC protein and E-cadherin form similar but independent complexes with α -catenin, β -catenin, and plakoglobin,” *Journal of Biological Chemistry*, vol. 270, no. 10, pp. 5549–5555, 1995.
- [15] L. K. Su, B. Vogelstein, and K. W. Kinzler, “Association of the APC tumor suppressor protein with catenins,” *Science*, vol. 262, no. 5140, pp. 1734–1737, 1993.
- [16] O. M. Sieber, S. Segditsas, A. L. Knudsen et al., “Disease severity and genetic pathways in attenuated familial adenomatous polyposis vary greatly but depend on the site of the germline mutation,” *Gut*, vol. 55, no. 10, pp. 1440–1448, 2006.
- [17] O. M. Sieber, K. Heinimann, P. Gorman et al., “Analysis of chromosomal instability in human colorectal adenomas with two mutational hits at APC,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 99, no. 26, pp. 16910–16915, 2002.
- [18] J. F. Costello, M. C. Frühwald, D. J. Smiraglia et al., “Aberrant CpG-island methylation has non-random and tumour-type-specific patterns,” *Nature Genetics*, vol. 24, no. 2, pp. 132–138, 2000.
- [19] W. Friedl, R. Caspari, M. Sengteller et al., “Can APC mutation analysis contribute to therapeutic decisions in familial adenomatous polyposis? Experience from 680 FAP families,” *Gut*, vol. 48, no. 4, pp. 515–521, 2001.
- [20] S. Aretz, S. Uhlhaas, Y. Sun et al., “Familial adenomatous polyposis: aberrant splicing due to missense or silent mutations in the APC gene,” *Human Mutation*, vol. 24, no. 5, pp. 370–380, 2004.
- [21] L. Bertario, A. Russo, P. Sala et al., “Multiple approach to the exploration of genotype-phenotype correlations in familial adenomatous polyposis,” *Journal of Clinical Oncology*, vol. 21, no. 9, pp. 1698–1707, 2003.
- [22] T. M. Attard, R. J. Young, J. A. Stoner, and H. T. Lynch, “Population differences in familial adenomatous polyposis may be an expression of geographic differences in APC mutation pattern,” *Cancer Genetics and Cytogenetics*, vol. 172, no. 2, pp. 180–182, 2007.
- [23] S.-H. Han, J.-S. Ryu, Y.-J. Kim, H.-I. Cho, Y.-H. Yang, and K.-R. Lee, “Mutation analysis of the APC gene in unrelated Korean patients with FAP: four novel mutations with unusual phenotype,” *Familial Cancer*, vol. 10, no. 1, pp. 21–26, 2010.
- [24] F. Fostira, G. Thodi, R. Sandaltzopoulos, G. Fountzilias, and D. Yannoukakos, “Mutational spectrum of APC and genotype-phenotype correlations in Greek FAP patients,” *BMC Cancer*, vol. 10, article 389, 2010.