Clinical Study

Does Number of Ports Affect Outcomes in Patients Undergoing Laparoscopic Pyloromyotomy? Retrospective Chart-Review Study

Tariq O. Abbas and Adel Ismail

Pediatric Surgery Department, Hamad General Hospital, P.O. Box 3050, Doha, Qatar

Correspondence should be addressed to Adel Ismail, adel11252@yahoo.com

Received 20 September 2011; Accepted 3 November 2011

Academic Editor: A. Polydorou

Copyright © 2012 T. O. Abbas and A. Ismail. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Although open Ramstedt's pyloromyotomy is the gold standard for the surgical management of infantile hypertrophic pyloric stenosis, laparoscopic pyloromyotomy has been found highly successful. Various factors, however, can affect the outcomes of surgical interventions in these patients. We observed a relationship between the number of ports used and outcome in patients undergoing laparoscopic pyloromyotomies. **Methods.** We retrospectively assessed the medical records of selected group of patients who underwent laparoscopic pyloromyotomy in our institution. Factors analyzed included operation time, length of hospital stay, postoperative complications, and time to postoperative full feeding. **Results.** We observed failure of myotomy in both two patients who underwent laparoscopic pyloromyotomy using only two working ports compared to successful myotomies in the remaining patients. **Conclusion.** Laparoscopy provides good results in terms of intraoperative exposure and cosmesis. However, standardized surgical technique with two working ports is advisable, and this can trigger further research to be ascertained.

1. Introduction

Infantile hypertrophic pyloric stenosis (IHPS) is a common condition affecting young infants; despite its frequency, IHPS has been recognized as a condition for only a little over a century and its etiology remains unknown [1]. Among the approaches used for pyloromyotomy are umbilical skin-fold incision [2] and laparoscopy [3].

Several factors can affect the outcomes of surgical intervention in these patients. In this case series, we observed a relationship between the number of ports and the outcome in patients undergoing laparoscopic pyloromyotomy.

2. Methods

We assessed patients who had undergone laparoscopic pyloromyotomy at our institution. IHPS was diagnosed by pyloric muscle thickness >4 mm and length >14 mm on ultrasonography. The study protocol was approved by our institution's Internal Review Board. There was no external source of funding.

Patients were excluded if they had been born prematurely (before 37 weeks of gestation) or had recent respiratory infections, major developmental anomalies, or had undergone prior abdominal surgery.

All the procedures were performed by one surgeon. The case notes of these patients were retrospectively reviewed for age of the patients at operation, subsequent surgical interventions, and patient outcomes.

The primary end points of this study were rate of postoperative vomiting and ultimately the need for revision pyloromyotomy. Secondary outcome measures included operation time, length of hospital stay, postoperative complications, and follow-up findings.

3. Results

We assessed 7 children who fit the inclusion criteria mentioned above and underwent laparoscopic pyloromyotomy (Table 1). Only one of the patients has a positive family history of the disease.

TABLE 1

	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patient 6	Patient 7
Duration of symptoms (d)	2	15	3	7	7	10	4
Family history	−ve	−ve	−ve	−ve	−ve	−ve	+ve
Preoperative weight (kg)	3	4.5	4.6	4	4	3	5.4
Pylorus length (mm)	17 × 10	24 × 3	25 × 5	25 × 7	15 × 4	NA	18 × 19
Serum electrolytes (mmol/L)							
Before operation							
Chloride	132	104	87	84	85	87	104
Sodium	—	141	137	136	136	137	137
Bicarb.	28	30	29	32	38	22	—
Number of working ports	3	2	3	2	3	3	3
Conversion to open	−ve	−ve	−ve	−ve	−ve	−ve	−ve
Postoperative complications	−ve	−ve	−ve	−ve	−ve	−ve	−ve
Postoperative vomiting	−ve	+ve	−ve	+ve	−ve	+ve	−ve
Revision pyloromyotomy	−ve	+ve	−ve	+ve	−ve	−ve	−ve
Duration of surgery (min)	35	35	50	—	30	40	30
Duration of anesthesia (min)	65	55	85	80	50	50	65
Time to full feeding (h)	96	96	48	168	22	46	21
Postoperative length of hospital stay (d)	4	4.5	2.5	9	1	3	1
Wound infection	−ve	−ve	−ve	−ve	−ve	−ve	−ve
Wound dehiscence	−ve	−ve	−ve	−ve	−ve	−ve	−ve

However, there is no remarkable difference in the operative or anesthesia time between the two groups of patients who underwent laparoscopic pyloromyotomy using either two or three working ports. On the other hand, procedures performed using only two working ports resulted in failure of complete myotomy and both needed revision pyloromyotomy consequently. No postoperative wound infection or dehiscence was recorded.

4. Discussion

The surgical procedure for pyloromyotomy was first described in 1912. Attention to detail is necessary to minimize morbidity and mortality [4]. When compared with open pyloromyotomy, the laparoscopic approach appears equally safe and effective, with superior cosmetic results [5]. A systematic review in 2009 found that the rate of total complications was significantly lower with the laparoscopic than the open approach, due primarily to a lower wound complication rate [6]. Moreover, the laparoscopic procedure is as quick as the open procedure, has low morbidity, is devoid of major wound-related problems, and can be easily taught [7].

An adequate pyloromyotomy must balance between the risk of perforation and the risk of incomplete myotomy, although an inability to palpate the divided pylorus makes the evaluation of these risks particularly challenging [8].

In one of the largest series to date, the incidence of incomplete pyloromyotomy was 4%, suggesting the need for conversion whenever the quality of the myotomy was reduced. In contrast, about 15% of patients in that trial experienced postoperative vomiting, with this factor not

FIGURE 1

influenced by experience with laparoscopic pyloromyotomy [7].

However, we observed a relationship between the number of ports and postoperative feeding tolerance in patients undergoing laparoscopic pyloromyotomies. A representative diagram of the time to postoperative feeding tolerance is shown in Figure 1.

Procedures performed using only 2 working ports resulted in failure of complete myotomy, despite being able to visualize the bulging mucosa. This may have resulted from

an oblique angle to the myotomy line, leading to the easing of both sides of the muscle and refusion.

Similar to previous findings, none of our patients experienced postoperative wound infection or dehiscence. By contrast, the incidence of these complications after open pyloromyotomy has been reported to be 6.7% [5, 9, 10].

In conclusion, laparoscopy provides good results in terms of intraoperative exposure and cosmesis; however, standardized surgical technique with two working ports is advisable.

References

- [1] M. Hernanz-Schulman, "Infantile hypertrophic pyloric stenosis," *Radiology*, vol. 227, no. 2, pp. 319–331, 2003.
- [2] K. C. Tan and A. Bianchi, "Circumumbilical incision for pyloromyotomy," *British Journal of Surgery*, vol. 73, no. 5, p. 399, 1986.
- [3] R. D. Spicer, "Infantile hypertrophic pyloric stenosis: a review," *British Journal of Surgery*, vol. 69, no. 3, pp. 128–135, 1982.
- [4] J. L. Alain, D. Grousseau, and G. Terrier, "Extramucosal pylorotomy by laparoscopy," *Journal of Pediatric Surgery*, vol. 26, no. 10, pp. 1191–1192, 1991.
- [5] K. L. Greason, W. R. Thompson, E. C. Downey, and B. L. Sasso, "Laparoscopic pyloromyotomy for infantile hypertrophic pyloric stenosis: report of 11 cases," *Journal of Pediatric Surgery*, vol. 30, no. 11, pp. 1571–1574, 1995.
- [6] J. E. Sola and H. L. Neville, "Laparoscopic vs open pyloromyotomy: a systematic review and meta-analysis," *Journal of Pediatric Surgery*, vol. 44, no. 8, pp. 1631–1637, 2009.
- [7] J. D. W. Van Der Bilt, W. L. M. Kramer, D. C. Van Der Zee, and N. M. A. Bax, "Laparoscopic pyloromyotomy for hypertrophic pyloric stenosis: impact of experience on the results in 182 cases," *Surgical Endoscopy and Other Interventional Techniques*, vol. 18, no. 6, pp. 907–909, 2004.
- [8] M. D. Leclair, V. Plattner, E. Mirallie et al., "Laparoscopic pyloromyotomy for hypertrophic pyloric stenosis: a prospective, randomized controlled trial," *Journal of Pediatric Surgery*, vol. 42, no. 4, pp. 692–698, 2007.
- [9] P. F. Leahy, R. Farrell, and B. O'Donnell, "300 infants with hypertrophic pyloric stenosis: presentation and outcome," *Irish Medical Journal*, vol. 79, no. 5, pp. 114–116, 1986.
- [10] L. W. E. Van Heurn, P. Vos, and G. Sie, "Recurrent vomiting after successful pyloromyotomy," *Pediatric Surgery International*, vol. 15, no. 5-6, pp. 385–386, 1999.