Research Article

On the Jacobson Radical of an (m, n) -Semiring

Yongwen Zhu

School of Mathematics and Information Science, Yantai University, Yantai City 264005, China

Correspondence should be addressed to Yongwen Zhu; zyw@ytu.edu.cn

Received 28 March 2013; Accepted 9 July 2013

Academic Editor: Yiqiang Zhou

Copyright © 2013 Yongwen Zhu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The notion of n -ary semimodules is introduced so that the Jacobson radical of an (m, n) -semiring is studied and some well-known results concerning the Jacobson radical of a ring (a semiring or a ternary semiring) are generalized to an (m, n) -semiring.

1. Introduction

The concept of semigroups [1] was generalized to that of ternary semigroups [2], that of n -ary semigroups [3–6], and even to that of (n, m) -semigroups [7]. Similarly, it was natural to generalize the notion of rings to that of ternary semirings, that of n -ary semirings, and even that of (m, n) -semirings.

Indeed, there were some research articles on semirings, (see, for example, [8–14]), specially on the radical of a semiring; see [15–18]. Semigroups over semirings were studied in [19] and semimodules over semirings were studied in [14]. The notion of semirings can be generalized to ternary semirings [20] and Γ -semirings [21], even to (m, n) -semirings [22–24]. The radicals of ternary semirings and of Γ -semirings were studied in [20, 21], respectively. The concept of (m, n) -semirings was introduced and accordingly some simple properties were discussed in [22–24], where the concept of radicals was not mentioned.

The notion of the Jacobson radicals was first introduced by Jacobson in the ring theory in 1945. Jacobson [25] defined the radical of R , which we call the Jacobson radical, to be the join of all quasi-regular right ideals and verified that the radical is a two-sided ideal and can also be defined to be the join of the left quasi-regular ideals.

The concept of the Jacobson radical of a semiring has been introduced internally by Bourne [15], where it was proved that the right and left Jacobson radicals coincide; thus one could say the Jacobson radical briefly. These and some other results were generalizations of well-known results of Jacobson [25].

In 1958, by associating a suitable ring with the semiring, Bourne and Zassenhaus defined the semiradical of the semiring [16]. In [18] it was proved that the concepts of the Jacobson radical and the semiradical coincide.

Iizuka [17] considered the Jacobson radical of a semiring from the point of view of the representation theory [15] without reducing it to the ring theory. The external notion of the radical was proved to be related to internal one; at the same time, it was shown that the radical defined in [17] coincides with the Jacobson radical and with the semiradical of the semiring.

In the present paper, we investigate (n, m) -semirings by means of n -ary semimodules so that we can define externally the Jacobson radical of an (n, m) -semiring, and then we establish the radical properties of the Jacobson radical of an (n, m) -semiring. Some necessary notions such as irreducible n -ary semimodules over an (n, m) -semiring are adequately defined. All results in this paper generalize the corresponding ones concerning the radical of a ring [25], of a semiring [15–18], or of a ternary semiring [20].

2. Preliminaries

We used following convention as followed by [4]: The sequence $x_i, x_{i+1}, \dots, x_m$ is denoted by x_i^m . Thus the following expression

$$f(x_1, \dots, x_i, y_{i+1}, \dots, y_j, z_{j+1}, \dots, z_m) \quad (1)$$

is represented as

$$f(x_1^i, y_{i+1}^j, z_{j+1}^m). \quad (2)$$

In the case when $y_{i+1} = \dots = y_j = y$, then (2) is expressed as

$$f\left(x_1^i, \overset{(j-i)}{y}, z_{j+1}^m\right). \quad (3)$$

If $x_1 = \dots = x_i = y_{i+1} = \dots = y_j = z_{j+1} = \dots = z_m = f(a_1^m)$, then (2) can be written as $f(\overset{(m)}{f(a_1^m)})$.

Recall that an n -ary semigroup (S, f) is defined as a nonempty set S with an n -ary associative operation $f : S^n = \underbrace{S \times \dots \times S}_n \rightarrow S$; that is,

$$f(x_1^{i-1}, f(x_i^{n+i-1}, x_{n+i}^{2n-1}), x_{n+j}^{2n-1}) = f(x_1^{j-1}, f(x_j^{n+j-1}, x_{n+j}^{2n-1}), x_{n+j}^{2n-1}) \quad (4)$$

for all $x_1^{2n-1} \in S$ and all $1 \leq i < j \leq n$. Whence we may denote $f(f(x_1^n), x_{n+1}^{2n-1})$ by $f(x_1^{2n-1})$ briefly. Generally, we have the notation $f(x_1^{t(m-1)+1})$ for each positive t and all $x_j \in S$. Thus for positive integer k , $f(x_1^k)$ is well defined if and only if $k \equiv 1 \pmod{n-1}$; see [7, Lemma 1.1]. An n -ary semigroup (S, f) is called cancellative if

$$f(x_1^{i-1}, a, x_{i+1}^n) = f(x_1^{i-1}, b, x_{i+1}^n) \implies a = b \quad (5)$$

for all $a, b, x_j \in S$.

The next definition is a generalization of the concept of ternary semirings in [20] and similar to the notion of the (m, n) -semirings in [24].

Definition 1. A nonempty set R together with an m -ary operation f , called *addition*, and an n -ary operation g , called *multiplication*, is said to be an (m, n) -semiring if the following conditions are satisfied.

- (1) (R, f) is an m -ary semigroup and (R, g) is an n -ary semigroup.
- (2) g is distributive with respect to operation f ; that is, for every $a_1^{i-1}, b_1^m, a_{i+1}^n \in R$,

$$g(a_1^{i-1}, f(b_1^m), a_{i+1}^n) = f(g(a_1^{i-1}, b_1, a_{i+1}^n), \dots, g(a_1^{i-1}, b_m, a_{i+1}^n)). \quad (6)$$

- (3) (R, f) is commutative; that is, for every permutation p of $\{1, 2, \dots, m\}$ and all $x_1^m \in R$,

$$f(x_1, x_2, \dots, x_m) = f(x_{p(1)}, x_{p(2)}, \dots, x_{p(m)}). \quad (7)$$

- (4) There is an element 0, called the *zero* of (R, f, g) , satisfying the following two properties:

$$(4A) \quad 0 \text{ is an } f\text{-identity; that is, for every } x \in R, f(\overset{(m-1)}{0}, x) = x;$$

$$(4B) \quad 0 \text{ is a } g\text{-zero; that is, for all } x_1^m \in R, g(x_1^n) = 0 \text{ whenever there exists } i \text{ such that } x_i = 0.$$

It is clear that the zero of an (m, n) -semiring R is necessarily unique.

Definition 2. An (m, n) -semiring (R, f, g) is called *additively cancellative* if the m -ary semigroup (R, f) is cancellative and *multiplicatively cancellative* if the n -ary semigroup (R, g) is cancellative.

Recall that for an n -ary semigroup (S, f) , a nonempty subset T of S is called a subsemigroup of (S, f) if $f(a_1^n) \in T$ whenever all $a_1^n \in T$. For $i \leq n$, we call T an i -ideal of S if $f(a_1^n) \in T$ whenever $a_i \in T$. T is called an ideal of S if and only if it is an i -ideal of S . See, for example, [7, Definition 1.6].

Definition 3. A nonempty subset T of an (m, n) -semiring (R, f, g) is called an n -ary subsemiring of (R, f, g) if T is a subsemigroup of (R, f) as well as a subsemigroup of (R, g) and an (i) -ideal of (R, f, g) if T is a subsemigroup of (R, f) as well as an (i) -ideal of (R, g) (where $i \leq n$). An 1-ideal is also called a *right ideal* and an n -ideal is also called a *left ideal*. An ideal T of (R, f, g) is called a k -ideal if $f(x, y_2^m) \in T$; $x \in R$ and $y_2^m \in T$ imply that $x \in T$. An ideal T of (R, f, g) is called an h -ideal if $f(x, y_2^m, z_2^m) = f(y_1, z_2^m)$; $x, z_2^m \in R$ and $y_1^m \in T$ imply that $x \in T$.

Let A be an ideal of (R, f, g) . Then the k -closure of A , denoted by $\overline{A}$, is defined by $\overline{A} = \{x \in R : f(x, a_2^m) = a_1^m \text{ for some } a_1^m \in A\}$. Similarly, the h -closure of A , denoted by $\widehat{A}$, is defined by $\widehat{A} = \{x \in R : f(x, a_2^m, y_2^m) = f(a_1, y_2^m) \text{ for some } a_1^m \in A \text{ and some } y_2^m \in R\}$. One can show that $\overline{A}$ is a k -ideal and $\widehat{A}$ is an h -ideal. Furthermore, it is shown that an ideal A of (R, f, g) is a k -ideal if and only if $\overline{A} = A$ and that A is an h -ideal if and only if $\widehat{A} = A$.

Definition 4. An equivalence relation ρ on an (m, n) -semiring (R, f, g) is said to be a *congruence relation* or simply a *congruence* of (R, f, g) if the following conditions are satisfied:

- (1) $a_1^m \rho b_1^m \implies f(a_1^m) \rho f(b_1^m)$ for all $a_1^m, b_1^m \in R$,
- (2) $a_1^n \rho b_1^n \implies g(a_1^n) \rho g(b_1^n)$ for all $a_1^n, b_1^n \in R$.

Let I be a proper ideal of an (m, n) -semiring (R, f, g) . Then the congruence on (R, f, g) , denoted by k_I , and defined by setting $x k_I y$ if and only if $f(x, a_2^m) = f(y, b_2^m)$ for some $a_2^m, b_2^m \in I$, is called the *Bourne congruence* on (R, f, g) defined by the ideal I . We denote the Bourne congruence class of an element x by x/I and denote the set of all such congruence classes of (R, f, g) by R/I . If the Bourne congruence k_I is proper, that is, $0/I \neq R$, then we can define two operations, m -ary addition and n -ary multiplication on R/I by $\overline{f}(a_1^m/I) = f(a_1^m)/I$ and $\overline{g}(b_1^n/I) = g(b_1^n)/I$ for all $a_1^m, b_1^n \in R$. Then $(R/I, \overline{f}, \overline{g})$ is an (m, n) -semiring and is called the *Bourne factor* (m, n) -semiring.

Similarly, the congruence on (R, f, g) , denoted by h_I , and defined by setting $x h_I y$ if and only if $f(x, a_2^m, y_2^m) = f(y, b_2^m, y_2^m)$ for some $a_2^m, b_2^m \in I$ and some $y_2^m \in R$, is

called the *Iizuka congruence* on (R, f, g) defined by the ideal I . We denote the Iizuka congruence class of an element x by $x[/I]$ and denote the set of all such congruence classes of (R, f, g) by $R[/I]$. If the Iizuka congruence h_I is proper, that is, $0[/I] \neq R$, then we can define two operations, m -ary addition and n -ary multiplication on $R[/I]$ by $\hat{f}(a_1^m[/I]) = f(a_1^m)$ and $\hat{g}(b_1^n[/I]) = g(b_1^n)$ for all $a_1^m, b_1^n \in R$. Then $(R[/I], \hat{f}, \hat{g})$ is an (m, n) -semiring and we call it the *Iizuka factor (m, n) -semiring*.

The next definition is a generalization of [20, Definition 2.13].

Definition 5. A commutative m -ary semigroup (M, f_0) with an identity 0_M (operation f_0 to be called addition) is called a *right n -ary semimodule* over an (m, n) -semiring (R, f, g) or simply an n -ary R -semimodule if there exists a mapping $M \times \underbrace{R \times \cdots \times R}_{n-1} \rightarrow M$ (images to be denoted by $ar_2 \cdots r_n$ or briefly by ar_2^n for all $a \in M$ and $r_2^n \in R$) satisfying the following conditions:

- (1) $f_0(a_1^m)r_2 \cdots r_n = f_0(a_1^m r_2 \cdots r_n)$ for all $a_1^m \in M$ and all $r_2^n \in R$;
- (2) $ar_2 \cdots (f(s_1^m)) \cdots r_n = f_0(ar_2 \cdots s_1^m \cdots r_n)$ for all $a \in M$ and all $r_2^n, s_1^m \in R$;
- (3) $(ar_2 \cdots r_n)r_{n+1} \cdots r_{2n-1} = ar_2 \cdots g(r_{k+1}^{k+n}) \cdots r_{2n-1}$ for all $a \in M, k \in \{1, \dots, n-1\}$, and $r_{k+1}^{k+n} \in R$;
- (4) $0_M r_2 \cdots r_n = 0_M$ for all $r_2^n \in R$;
- (5) $ar_2 \cdots r_n = 0_M$ whenever $a \in M, r_2^n \in R$, and $r_i = 0$ for some i .

Definition 6. A nonempty subset N of a right n -ary semimodule (M, f_0) over an (m, n) -semiring (R, f, g) is called an *n -ary subsemimodule* of M if (i) $f_0(a_1^m) \in N$ and (ii) $a_1 x_2^n \in N$ for all $a_1^m \in N$ and $x_2^n \in R$.

An n -ary subsemimodule N of M is called an *n -ary k -subsemimodule* if $f_0(a, b_2^m) \in N$; $a \in M$ and $b_2^m \in N$ imply that $a \in N$. An n -ary subsemimodule N of M is called an *n -ary h -subsemimodule* if $f(a, b_2^m, c_2^m) = f(b_1, c_2^m)$; $a, c_2^m \in R$ and $b_1^m \in N$ imply that $a \in N$.

For example, an (m, n) -semiring (R, f, g) can be regarded as a right n -ary R -semimodule naturally. Then if I is a k -ideal (an h -ideal) of the (m, n) -semiring (R, f, g) , then I is also an n -ary k -(h -)subsemimodule of this right n -ary R -semimodule R .

Definition 7. A right n -ary R -semimodule (M, f_0) is said to be *cancellative* if (M, f_0) is a cancellative m -ary semigroup.

Definition 8. An equivalence relation ρ on right n -ary R -semimodule (M, f_0) is said to be a *congruence relation* or

simply a *congruence* of (M, f_0) if the following conditions are satisfied:

- (1) $a_1^m \rho b_1^m \Rightarrow f_0(a_1^m) \rho f_0(b_1^m)$ for all $a_1^m, b_1^m \in M$,
- (2) $a \rho b \Rightarrow ar_2^n \rho br_2^n$ for all $a, b \in M$ and all $r_2^n \in R$.

We say that a congruence ρ of (M, f_0) admits the cancellation law (of addition) if

- (3) $f_0(x, a_2^m) \rho f_0(y, b_2^m)$ and $a_2^m \rho b_2^m$ imply $x \rho y$.

Let N be an n -ary subsemimodule of an n -ary right semimodule (M, f_0) over an (m, n) -semiring R . Then the congruence on (M, f_0) , denoted by k_N , and defined by setting

$$xk_N y \text{ iff } f_0(x, a_2^m) = f_0(y, b_2^m) \text{ for some } a_2^m, b_2^m \in N, \quad (8)$$

is called the *Bourne congruence* on M defined by the n -ary subsemimodule N . We denote the Bourne congruence class of an element x by x/N and denote the set of all such congruence classes of M by M/N . Define two operations, m -ary addition and n -ary scalar multiplication on M/N , by $\overline{f_0(a_1^m/N)} = f_0(a_1^m)/N$ and $(a_1/N)r_2^n = (a_1 r_2^n)/N$ for all $a_1^m \in M$ and all $r_2^n \in R$. With these two operations, M/N is an n -ary right semimodule over R and we call it the *Bourne factor n -ary semimodule*.

Similarly, we can define the *Iizuka congruence* h_N and the *Iizuka factor n -ary semimodule* $M[/I]N$. It is easy to show that $M[/I]N$ is cancellative.

In what follows, we always assume that the n -ary right semimodule is cancellative.

3. Primitive (m, n) -Semirings

Definition 9. Let (R, f, g) be an (m, n) -semiring with zero 0. Then the *zeroid* of R , denoted by $Z(R)$, is defined as

$$Z(R) = \{x \in R : f(x, y_2^m) = f(0, y_2^m) \text{ for some } y_2^m \in R\}. \quad (9)$$

Clearly, the zero element 0 of R belongs to $Z(R)$. Furthermore, we have the following.

Lemma 10. The zeroid $Z(R)$ of an (m, n) -semiring (R, f, g) is the smallest h -ideal of (R, f, g) .

Proof. It is easily verified that $Z(R)$ is an ideal of R . To show $Z(R)$ is an h -ideal of R , we suppose $f(x, y_2^m, z_2^m) = f(y_1, z_2^m)$, where $x, z_2^m \in R$ and $y_1^m \in Z(R)$. For each $i \in \{1, \dots, m\}$ there exist $u(i)_2^m \in R$ such that $f(y_i, u(i)_2^m) = f(0, u(i)_2^m)$, so we have

$$f(f(x, y_2^m, z_2^m), u(1)_2^m, \dots, u(m)_2^m) = f(f(y_1, z_2^m), u(1)_2^m, \dots, u(m)_2^m), \quad (10)$$

that is,

$$f(x, u(1)_2^m, f(y_2, u(2)_2^m), \dots, f(y_m, u(m)_2^m), z_2^m) = f(f(y_1, u(1)_2^m), u(2)_2^m, \dots, u(m)_2^m, z_2^m). \quad (11)$$

It follows that

$$\begin{aligned} f(x, u(1)_2^m, f(0, u(2)_2^m), \dots, f(0, u(m)_2^m), z_2^m) \\ = f(f(0, u(1)_2^m), u(2)_2^m, \dots, u(m)_2^m, z_2^m). \end{aligned} \quad (12)$$

Hence we obtain

$$\begin{aligned} f(x, u(1)_2^m, u(2)_2^m, \dots, u(m)_2^m, z_2^m) \\ = f(0, u(1)_2^m, u(2)_2^m, \dots, u(m)_2^m, z_2^m), \end{aligned} \quad (13)$$

which shows that $x \in Z(R)$, so that $Z(R)$ is an h -ideal of R .

At last, suppose that I is an arbitrary h -ideal of R . We aim to show $Z(R) \subseteq I$. For this, let $x \in Z(R)$. Then there exist $y_2^m \in R$ such that $f(x, y_2^m) = f(0, y_2^m)$, so $f(x, 0^{m-1}, y_2^m) = f(0, y_2^m)$. It follows that $x \in I$ since I is an h -ideal and $0 \in I$. Thus $Z(R) \subseteq I$. $\square$

Definition 11. Let M be a right n -ary R -semimodule. The annihilator of M in R , denoted by $(0 : M)$ or $A_R(M)$, is defined as the subset

$$\begin{aligned} \{r \in R : ar_2^m = 0_M \text{ whenever } a \in M, \\ r_2^m \in R \text{ and } r_i = r \text{ for some } i\}. \end{aligned} \quad (14)$$

Lemma 12. $A_R(M)$ is an h -ideal of R .

Proof. It is obvious that $A_R(M)$ is an ideal of R . To show that it is an h -ideal, suppose $f(x, y_2^m, z_2^m) = f(y_1, z_2^m)$, where $x, z_2^m \in R$ and $y_1^m \in A_R(M)$. Then for all $r_i \in R$,

$$ar_2 \cdots f(x, y_2^m, z_2^m) \cdots r_n = ar_2 \cdots f(y_1, z_2^m) \cdots r_n, \quad (15)$$

that is,

$$\begin{aligned} f_0(ar_2 \cdots x \cdots r_n, ar_2 \cdots y_2^m \cdots r_n, ar_2 \cdots z_2^m \cdots r_n) \\ = f_0(ar_2 \cdots y_1 \cdots r_n, ar_2 \cdots z_2^m \cdots r_n), \end{aligned} \quad (16)$$

which deduces that

$$\begin{aligned} f_0(ar_2 \cdots x \cdots r_n, 0^{m-1}, ar_2 \cdots z_2^m \cdots r_n) \\ = f_0(0, ar_2 \cdots z_2^m \cdots r_n) \end{aligned} \quad (17)$$

since $ar_2 \cdots y_j \cdots r_n = 0$ for each $j \in \{1, \dots, m\}$. Thus we have

$$f_0(ar_2 \cdots x \cdots r_n, ar_2 \cdots z_2^m \cdots r_n) = f_0(0, ar_2 \cdots z_2^m \cdots r_n). \quad (18)$$

By cancellation law of M , $ar_2 \cdots x \cdots r_n = 0$. Hence $x \in A_R(M)$, as required. $\square$

Definition 13. A right n -ary R -semimodule M is said to be faithful if $Z(R) = A_R(M)$.

One of difficulties when studying the radical of an (m, n) -semiring R is how to give an appropriate definition of the irreducibility of a right n -ary R -semimodule. The next definition is a generalization of [20, Definition 3.9].

Definition 14. A right n -ary R -semimodule M is said to be irreducible if for every arbitrary fixed pair $u_2^m, v_2^m \in M$ with $u_j \neq v_j$ for some j and for any $x \in M$, there exist $a(i)_2^m, b(i)_2^m \in R$ with $i = 2, \dots, m$ such that

$$\begin{aligned} f_0(x, u_2a(2)_2^n, \dots, u_ma(m)_2^n, v_2b(2)_2^n, \dots, v_mb(m)_2^n) \\ = f_0(0_M, u_2b(2)_2^n, \dots, u_mb(m)_2^n, \\ v_2a(2)_2^n, \dots, v_ma(m)_2^n). \end{aligned} \quad (19)$$

Remark 15. Since M is cancellative, it is easily seen that a right n -ary R -semimodule M is irreducible if and only if for every arbitrary fixed pair $u_2^m, v_2^m \in M$ with $u_j \neq v_j$ for all j and for any $x \in M$, there exist $a(i)_2^m, b(i)_2^m \in R$ with $i = 2, \dots, m$ such that equality (2) holds.

Lemma 16. Let I be an h -ideal of an (m, n) -semiring R . If M is an irreducible right n -ary R/I -semimodule, then M is an irreducible right n -ary R -semimodule.

Proof. Let M be an irreducible right n -ary R -semimodule. Then we can define an n -ary action on M by $ar_2 \cdots r_n = a(r_2/I) \cdots (r_n/I)$ for all $a \in M$ and for all $r_2^n \in R$, and this makes M into an irreducible right n -ary R -semimodule. $\square$

The converse of Lemma 16 is not necessarily true. But in particular we have the following lemma.

Lemma 17. If M is a right n -ary R -semimodule then M is a right n -ary $R/A_R(M)$ -semimodule, where $R/A_R(M)$ is the Bourne factor semiring. Moreover, if M is an irreducible right n -ary R -semimodule, then M is also an irreducible right n -ary $R/A_R(M)$ -semimodule.

Proof. Suppose M is a right n -ary R -semimodule. We define an n -ary action on M as follows: $ar_2/I \cdots r_n/I = ar_2 \cdots r_n$ where $I = A_R(M)$, for all $a \in M$ and for all $r_2^n \in R$. We now show that this definition is well-defined. If for each $i = 2, \dots, n$, $r_i/I = s_i/I$, then $r_i k_I s_i$, that is, there exist $x(i)_2^m, y(i)_2^m \in I$ such that $f(r_i, x(i)_2^m) = f(s_i, y(i)_2^m)$. It follows that $af(r_2, x(2)_2^m) \cdots f(r_n, x(n)_2^m) = af(s_2, y(2)_2^m) \cdots f(s_n, y(n)_2^m)$, which implies that $ar_2 \cdots r_n = as_2 \cdots s_n$ since $x(i)_2^m, y(i)_2^m \in A_R(M)$. Thus $ar_2/I \cdots r_n/I = as_2/I \cdots s_n/I$, as required. It is easy to see that the above definition makes M into a right n -ary R -semimodule.

Moreover, if M is an irreducible right n -ary R -semimodule then it is routine to verify that M is also an irreducible right n -ary $R/A_R(M)$ -semimodule by (2). $\square$

Lemma 18. Let M be a right n -ary R -semimodule. Then $A_{R/A_R(M)}(M) = \{0/A_R(M)\}$.

Proof. Let $x/I \in A_{R/I}(M)$, where $I = A_R(M)$. Then for any $a \in M$ $ar_2/I \cdots r_n/I = 0_M$ whenever $r_2^n \in R$ and $r_i = x$ for some $i \in \{2, \dots, n\}$. It follows that $ar_2 \cdots r_n = 0_M$ where $r_i = x$

for some $i \in \{2, \dots, n\}$. This shows that $x \in I$ and so that $x/I = 0/I$. Consequently, $A_{R/A_R(M)}(M) = \{0/A_R(M)\}$. $\square$

Lemma 19. Any right n -ary R -semimodule M is a faithful $R/A_R(M)$ -semimodule.

Proof. Let M be a right n -ary R -semimodule. Then in view of Lemma 17, M is an $R/A_R(M)$ -semimodule. On the one hand, by Lemma 12, $A_{R/A_R(M)}(M)$ is an h -ideal of $R/A_R(M)$. On the other hand, by Lemma 10, $Z(R/A_R(M))$ is the smallest h -ideal of $R/A_R(M)$. Thus $Z(R/A_R(M)) \subseteq A_{R/A_R(M)}(M)$. According to Lemma 18, $A_{R/A_R(M)}(M) = \{0/A_R(M)\}$. So $Z(R/A_R(M)) = \{0/A_R(M)\} = A_{R/A_R(M)}(M)$, which means that M is a faithful $R/A_R(M)$ -semimodule. $\square$

Lemma 20. If I is an h -ideal of an (m, n) -semiring R then $Z(R/I) = \{0/I\}$ where R/I is the Bourne factor semiring.

Proof. Suppose $x/I \in Z(R/I)$. Then we have $f(x/I, y_2^m/I) = f(0/I, y_2^m/I)$ for some $y_2^m/I \in R/I$. Thus we have $f(x, y_2^m)/I = f(0, y_2^m)/I$ which implies that $f(x, y_2^m, a_2^m) = f(0, y_2^m, b_2^m)$ for some $a_2^m, b_2^m \in I$. Hence $f(x, a_2^m, y_2^m) = f(0, b_2^m, y_2^m)$. This shows that $x \in I$ since I is an h -ideal of R . Consequently, $x/I = 0/I$. Thus $Z(R/I) = \{0/I\}$. $\square$

Definition 21. An (m, n) -semiring R is said to be *primitive* if it has a faithful irreducible cancellative n -ary R -semimodule. An ideal P is said to be *primitive* if the Bourne factor semiring R/P is primitive.

Evidently, an (m, n) -semiring R is primitive if and only if $\{0\}$ is a primitive ideal of R . The following theorem characterizes primitive ideals of an (m, n) -semiring.

Theorem 22. An h -ideal P of (m, n) -semiring R is primitive if and only if $P = A_R(M)$ for some irreducible right n -ary R -semimodule M .

Proof. Let P be an h -ideal of R such that $P = A_R(M)$ for some irreducible right n -ary R -semimodule M . Then by Lemmas 17 and 19 M is a faithful irreducible n -ary R/P -semimodule. This shows that R/P is primitive and hence P is a primitive h -ideal of R .

Conversely, let P be a primitive h -ideal of R . Then R/P is a primitive (m, n) -semiring. So there exists a faithful irreducible n -ary R/P -semimodule M . Now by Lemma 16 M is an irreducible n -ary R -semimodule. It remains to show that $P = A_R(M)$. Now $x \in A_R(M) \Leftrightarrow$ for all $a \in M$ and $r_2^n \in R$, $ar_2 \cdots r_n = 0_M$ whenever $x = r_i$ for some $i \in \{2, \dots, n\} \Leftrightarrow ar_2/P \cdots r_n/P = 0_M/P$ whenever $x/P = r_i/P$ for some $i \in \{2, \dots, n\} \Leftrightarrow x/P \in A_{R/P}(M) = Z(R/P)$ since M is a faithful n -ary R/P -semimodule $\Leftrightarrow x/P \in A_{R/P}(M) = \{0/P\}$, by Lemma 20 $\Leftrightarrow x/P = 0/P \Leftrightarrow x \in P$. Thus $P = A_R(M)$ as desired. $\square$

4. Jacobson Radical of an (m, n) -Semiring

Let us begin this section by defining the semi-irreducibility of a right n -ary R -semimodule.

Definition 23. A right n -ary R -semimodule M is said to be *semi-irreducible* if $MR^{n-1} \neq \{0_M\}$; that is, $ar_2 \cdots r_n \neq 0_M$ for some $a \in M$ and some $r_2^n \in R$, and M does not contain any n -ary k -subsemimodule other than $\{0_M\}$ and M .

Lemma 24. Let I be a subset of an (m, n) -semiring R and M a right n -ary R -semimodule with $MR^{i-2}IR^{n-i} \neq \{0_M\}$ for some $i \in \{2, \dots, n\}$. In the case where $i = 2$, we assume further that I is a left ideal of R . Then the following statements are true:

- (1) If M is semi-irreducible and $a \in M$, then $a = 0$ if and only if $aR^{i-2}IR^{n-i} = \{0_M\}$;
- (2) If M is irreducible and $a, b \in M$, then $a = b$ if and only if $ar_2^{i-1}sr_{i+1}^n = br_2^{i-1}sr_{i+1}^n$ for all $r_j \in R$ and all $s \in I$.

Proof. Suppose that (M, f_0) is a semi-irreducible right n -ary semimodule over an (m, n) -semiring (R, f, g) , and that I is a subset of R such that $MR^{i-2}IR^{n-i} \neq \{0_M\}$ for some $i \in \{2, \dots, n\}$. In the case where $i = 2$, we further assume that I is a left ideal of R .

(1) Assume that M is semi-irreducible. Let $a \in M$ be such that

$$aR^{i-2}IR^{n-i} = \{0_M\}. \quad (20)$$

Set

$$M_0 = \{x \in M : xR^{i-2}IR^{n-i} = \{0_M\}\}. \quad (21)$$

It is clear that $a \in M_0$, and it is easy to show that M_0 is a subsemimodule of M . Let $f_0(x, y_2^m) \in M_0$ and $y_2^m \in M_0$. Then $f_0(x, y_2^m)R^{i-2}IR^{n-i} = \{0_M\}$ and $y_2^mR^{i-2}IR^{n-i} = \{0_M\}$. Thus $xR^{i-2}IR^{n-i} = \{0_M\}$; that is, $x \in M_0$. This shows that M_0 is a k -subsemimodule of M . Since $MR^{i-2}IR^{n-i} \neq \{0_M\}$, $M_0 \neq M$. Since M is semi-irreducible, $M_0 = \{0_M\}$ and therefore $a = 0$.

The converse part is obvious.

(2) Assume that M is irreducible. Let $a, b \in M$ be such that $a \neq b$. Set $u_j = a$, $v_j = b$ for $j = 2, \dots, m$. Since $MR^{i-2}IR^{n-i} \neq \{0_M\}$, we have $xr_2^{i-1}sr_{i+1}^{n-i} \neq 0_M$ for some $x \in M$, $s \in I$ and $r_j \in R$. Since M is irreducible, according to Definition 14, there exist $a(i)_2^m, b(i)_2^m \in R$ with $i = 2, \dots, m$ such that

$$\begin{aligned} & f_0(x, u_2a(2)_2^n, \dots, u_ma(m)_2^n, \\ & \quad v_2b(2)_2^n, \dots, v_mb(m)_2^n) \\ & = f_0(0_M, u_2b(2)_2^n, \dots, u_mb(m)_2^n, \\ & \quad v_2a(2)_2^n, \dots, v_ma(m)_2^n). \end{aligned} \quad (22)$$

Hence

$$\begin{aligned}
& f_0 \left(x r_2^{i-1} s r_{i+1}^{n-i}, u_2 a(2)_2^{n-i-1} s r_{i+1}^{n-i}, \dots, u_m a(m)_2^{n-i-1} s r_{i+1}^{n-i}, \right. \\
& \quad \left. v_2 b(2)_2^{n-i-1} s r_{i+1}^{n-i}, \dots, v_m b(m)_2^{n-i-1} s r_{i+1}^{n-i} \right) \\
&= f_0 \left(0_M r_2^{i-1} s r_{i+1}^{n-i}, u_2 b(2)_2^{n-i-1} s r_{i+1}^{n-i}, \dots, \right. \\
& \quad \left. u_m b(m)_2^{n-i-1} s r_{i+1}^{n-i}, \right. \\
& \quad \left. v_2 a(2)_2^{n-i-1} s r_{i+1}^{n-i}, \dots, v_m a(m)_2^{n-i-1} s r_{i+1}^{n-i} \right) \\
&= f_0 \left(0_M, u_2 b(2)_2^{n-i-1} s r_{i+1}^{n-i}, \dots, u_m b(m)_2^{n-i-1} s r_{i+1}^{n-i}, \right. \\
& \quad \left. v_2 a(2)_2^{n-i-1} s r_{i+1}^{n-i}, \dots, v_m a(m)_2^{n-i-1} s r_{i+1}^{n-i} \right). \tag{23}
\end{aligned}$$

Since M is cancellative and $x r_2^{i-1} s r_{i+1}^{n-i} \neq 0_M$, at least one of the following $2(m-1)$ equalities does not hold:

$$\begin{aligned}
& u_j a(j)_2^{n-i-1} s r_{i+1}^{n-i} = v_j a(2)_2^{n-i-1} s r_{i+1}^{n-i}, \\
& \quad \text{where } j = 2, \dots, m; \\
& u_j b(j)_2^{n-i-1} s r_{i+1}^{n-i} = v_j b(2)_2^{n-i-1} s r_{i+1}^{n-i}, \\
& \quad \text{where } j = 2, \dots, m. \tag{24}
\end{aligned}$$

So we conclude that if $a r_2^{i-1} s r_{i+1}^n = b r_2^{i-1} s r_{i+1}^n$ for all $r_j \in R$ and all $s \in I$, then $a = b$.

The converse part follows easily. $\square$

Lemma 25. *Let M be a right n -ary R -semimodule and $M \neq \{0_M\}$. Then M is semi-irreducible if and only if for every nonzero $a \in M$, $\overline{aR^{n-1}} = M$.*

Proof. Assume that M is a semi-irreducible right n -ary R -semimodule and $M \neq \{0_M\}$. Let $a \in M$ be such that $a \neq 0_M$. Then by Lemma 24 $\overline{aR^{n-1}} \neq \{0_M\}$. Since $\overline{aR^{n-1}}$ is an n -ary k -subsemimodule of M , $\overline{aR^{n-1}} = M$.

Conversely, suppose that for any nonzero $a \in M$, $\overline{aR^{n-1}} = M$. Let $N \neq \{0_M\}$ be an n -ary k -subsemimodule of M . Then there exists $b \in N$ such that $b \neq 0_M$. So by hypothesis, $\overline{bR^{n-1}} = M$. Hence for any $x \in M$, there exist $a_1^m \in bR^{n-1}$ such that $f_0(x, a_2^m) = a_1$. Since $bR^{n-1} \subseteq N$, we have $a_1^m \in N$. Since N is an n -ary k -subsemimodule, $f_0(x, a_2^m) = a_1$ implies that $x \in N$. This shows that $N = M$. Now if $\overline{MR^{n-1}} = \{0_M\}$ then $\overline{aR^{n-1}} = \{0_M\}$ for all $a \in M$. Hence $\overline{aR^{n-1}} = \{0_M\}$. So we have $M = \{0_M\}$, a contradiction. Therefore, $\overline{MR^{n-1}} \neq \{0_M\}$. Thus M is semi-irreducible. $\square$

Corollary 26. *If a right n -ary R -semimodule M is irreducible, then it is semi-irreducible and $\overline{MR^{n-1}} = M$.*

Proof. Assume that M is an irreducible right n -ary R -semimodule. Then $M \neq \{0_M\}$ and, consequently, there exists a nonzero $y \in M$. In view of (2) with $u_j = y$ and $v_j = 0_M$

for $j = 2, \dots, m$, we obtain that for any $x \in M$ there exist $a(j)_2^n \in R$ with $j = 2, \dots, m$ such that

$$\begin{aligned}
& f_0 \left(x, ya(2)_2^n, \dots, ya(m)_2^n, 0_M^{(m-1)} \right) \\
&= f_0 \left(0_M, yb(2)_2^n, \dots, yb(m)_2^n, 0_M^{(m-1)} \right), \tag{25}
\end{aligned}$$

so that

$$\begin{aligned}
& f_0(x, ya(2)_2^n, \dots, ya(m)_2^n) \\
&= f_0(0_M, yb(2)_2^n, \dots, yb(m)_2^n) \in yR^{n-1}. \tag{26}
\end{aligned}$$

It follows that $x \in \overline{yR^{n-1}}$. Thus $\overline{yR^{n-1}} = M$. By Lemma 25, M is semi-irreducible.

Furthermore, $\overline{MR^{n-1}} \neq \{0_M\}$, which implies that $\overline{MR^{n-1}} \neq \{0_M\}$. Since $\overline{MR^{n-1}}$ is an n -ary k -subsemimodule of M , $\overline{MR^{n-1}} = M$ as required. $\square$

Now we can define the Jacobson radical of an (m, n) -semiring in an external way.

Definition 27. Let R be an (m, n) -semiring and Δ be the set of all irreducible right n -ary R -semimodules. Then $J(R) = \bigcap_{M \in \Delta} A_R(M)$ is called the *Jacobson radical* of R . If Δ is empty then R itself is considered as $J(R)$; that is, $J(R) = R$, and in this case, we say that R is a *radical (m, n) -semiring*. An (m, n) -semiring R is said to be *Jacobson semisimple* or *J-semisimple* if $J(R) = \{0\}$.

By Lemma 12, $A_R(M)$ is an h -ideal of R . Note that the intersection of any family of h -ideals is again an h -ideal. Consequently, we obtain the following.

Lemma 28. *$J(R)$ is an h -ideal of R .*

Lemma 29. *If M is a right n -ary R -semimodule then M is a right n -ary $R/J(R)$ -semimodule, where $R/J(R)$ is the Bourne factor semiring. Moreover, if M is an irreducible right n -ary R -semimodule, then M is also an irreducible right n -ary $R/J(R)$ -semimodule.*

Proof. This lemma can be proved by the same method as in proving Lemma 17. $\square$

Theorem 30. *If R is an (m, n) -semiring, then the Bourne factor semiring $R/J(R)$ is Jacobson semisimple.*

Proof. By Δ and Λ , we denote the set of all irreducible right n -ary R -semimodules and the set of all irreducible right n -ary $R/J(R)$ -semimodules, respectively. Then according to Lemmas 28, 16, and, 29, we obtain that $\Delta = \Lambda$. For any $x \in J(R/J(R))$ and any $M \in \Delta$, we have $x \in A_{R/J(R)}(M)$, which means that for any $a \in M$, $ar_2/J(R) \cdots r_n/J(R) = 0_M$ whenever $r_2^n \in R$ and $r_i = x$ for some $i \in \{2, \dots, n\}$. Thus $ar_2 \cdots r_n = 0_M$ whenever $r_2^n \in R$ and $r_i = x$ for some $i \in \{2, \dots, n\}$, so $x \in A_R(M)$ for all $M \in \Lambda$. That is, $x \in J(R)$. Hence $x/J(R) = 0/J(R)$. We have shown that

$J(R/J(R)) = \{0/J(R)\}$. By Definition 27, $R/J(R)$ is Jacobson semisimple. $\square$

The next theorem is a direct corollary of Theorem 22, giving an internal characterization of the Jacobson radical of an (m, n) -semiring.

Theorem 31. $J(R)$ is the intersection of all primitive h -ideals of R .

Definition 32. Let P be an i -ideal of an (m, n) -semiring R for some $i \in \{1, \dots, n\}$. Then P is said to be *strongly seminilpotent* if there exists a positive integer t such that

$(PR^{n-2})^{t-1}P \subseteq Z(R)$, where $R^{n-2} = \overbrace{R \cdots R}^{n-2}$, $(PR^{n-2})^{t-1} = (PR^{n-2})(PR^{n-2}) \cdots t-1$ times, $(PR^{n-1})^0P = P$. P is said to be *strongly nilpotent* if there exists a positive integer t such that $(PR^{n-2})^{t-1}P = \{0\}$.

Theorem 33. If P is a strongly semi-nilpotent left ideal of R , then $P \subseteq J(R)$.

Proof. Suppose on the contrary that

$$P \not\subseteq J(R) = \bigcap_{M \in \Delta} A_R(M), \quad (27)$$

where R is an (m, n) -semiring and Δ is the set of all irreducible right n -ary R -semimodules. Then there exists an $M \in \Delta$ such that $P \not\subseteq A_R(M)$. Thus there exists $i \in \{2, \dots, n\}$ such that

$$MR^{i-2}PR^{n-i} \neq \{0_M\}. \quad (28)$$

Since P is strongly semi-nilpotent, there exists a positive integer t such that $(PR^{n-2})^{t-1}P \subseteq Z(R)$. By Lemmas 10 and 12, $Z(R) \subseteq A_R(M)$. It follows that $(PR^{n-2})^{t-1}P \subseteq A_R(M)$, which implies that

$$MR^{i-2}(PR^{n-2})^{t-1}PR^{n-i} = \{0_M\}. \quad (29)$$

If (29) holds for all positive integers t 's, then in particular it is true for $t = 1$ and in this case we have $MR^{i-2}PR^{n-i} = \{0_M\}$, a contradiction to (28). If (29) does not hold for all t , then there exist $x \in M$ and positive s such that

$$\begin{aligned} xR^{i-2}(PR^{n-2})^{s-1}PR^{n-i} &\neq \{0_M\}, \\ xR^{i-2}(PR^{n-2})^sPR^{n-i} &= \{0_M\}. \end{aligned} \quad (30)$$

Thus $x \neq 0_M$ and there exists $a \in xR^{i-2}(PR^{n-2})^{s-1}PR^{n-i}$ such that $a \neq 0_M$. It follows that

$$\begin{aligned} aR^{i-2}PR^{n-i} &\subseteq xR^{i-2}(PR^{n-2})^{s-1}PR^{n-i}R^{i-2}PR^{n-i} \\ &= xR^{i-2}(PR^{n-2})^sPR^{n-i} = \{0_M\}, \end{aligned} \quad (31)$$

so we have

$$aR^{i-2}PR^{n-i} = \{0_M\}. \quad (32)$$

By Lemma 24, we obtain $a = 0_M$, again a contradiction. This completes the proof. $\square$

The next result is a direct corollary of Theorem 33.

Corollary 34. If an (m, n) -semiring R is Jacobson semisimple then R does not contain any non-zero strongly semi-nilpotent left ideal and hence R does not contain any nontrivial strongly nilpotent left ideal.

Lemma 35. If M is a (semi-)irreducible right n -ary R -semimodule and $N \neq \{0_M\}$ is an arbitrary R -subsemimodule (and $NR^{n-1} \neq \{0_M\}$), then N is (semi-)irreducible, and for any $x_2^n, y_2^n \in R$ the following statement is true: the equality $ux_2^n = uy_2^n$ holds for all $u \in M$ if and only if the same equality holds for all $u \in N$. Furthermore, $A_R(M) = A_R(N)$.

Proof. Assume M is an irreducible right n -ary R -semimodule. Then from (2), it follows that N is irreducible. If M is a semi-irreducible and $NR^{n-1} \neq \{0_M\}$, then N is semi-irreducible by Definition 23 since any subsemimodule of N is clearly a subsemimodule of M .

Let $x_2^n, y_2^n \in R$ be such that the equality $ux_2^n = uy_2^n$ holds for all $u \in M$. Since M is semi-irreducible, for any $a(\neq 0_M) \in M$ and any $b(\neq 0_M) \in N$, there exist positive p, q and $x(i)_2^p, y(j)_2^q \in R$ such that $p \equiv q \equiv 0 \pmod{m-1}$ and

$$f_0(a, bx(2)_2^n, \dots, bx(p)_2^n) = f_0(0_M, by(2)_2^n, \dots, by(q)_2^n). \quad (33)$$

Thus we have the following two equalities:

$$\begin{aligned} f_0(ax_2^n, bx(2)_2^n x_2^n, \dots, bx(p)_2^n x_2^n) \\ = f_0(0_M, by(2)_2^n x_2^n, \dots, by(q)_2^n x_2^n), \\ f_0(ay_2^n, bx(2)_2^n y_2^n, \dots, bx(p)_2^n y_2^n) \\ = f_0(0_M, by(2)_2^n y_2^n, \dots, by(q)_2^n y_2^n). \end{aligned} \quad (34)$$

It follows that

$$\begin{aligned} f_0(0_M, ax_2^n, bx(2)_2^n x_2^n, \dots, bx(p)_2^n x_2^n, \\ by(2)_2^n y_2^n, \dots, by(q)_2^n y_2^n) \\ = f_0(0_M, ay_2^n, bx(2)_2^n y_2^n, \dots, bx(p)_2^n y_2^n, \\ by(2)_2^n x_2^n, \dots, by(q)_2^n x_2^n). \end{aligned} \quad (35)$$

Observing that $bx(i)_2^n, by(j)_2^n \in N$, since N is a submodule, we have $bx(i)_2^n x_2^n = bx(i)_2^n y_2^n$ and $by(2)_2^n x_2^n = by(2)_2^n y_2^n$ for all i, j by the assumption. Hence by cancellation law, (35) deduces that $ax_2^n = ay_2^n$. The converse implication is clear.

Furthermore, letting $y_i = 0$ for some i , we get that the equality $ux_2^n = 0_M$ holds for all $u \in M$ if and only if the same equality holds for all $u \in N$. Thus $A_R(M) = A_R(N)$. $\square$

Lemma 36. Let I be an ideal of an (m, n) -semiring R .

- (1) If M is an (semi-)irreducible right n -ary R -semimodule (and $MR^{n-1} \neq \{0_M\}$), then M is an (semi-)irreducible right n -ary I -semimodule.

(2) If M is an irreducible right n -ary I -semimodule, then there exists an irreducible right n -ary R -semimodule M^* , which can be regarded as an I -subsemimodule N of M .

Proof. (1) Let M be an irreducible R -semimodule and $u_2^m, v_2^m \in M$ be such that $u_j \neq v_j$ for some j . Without loss of generality, we suppose that $M \neq \{0_M\}$. From (2) we deduce that $MR^{n-1} \neq \{0_M\}$. By Lemma 24, $u_j c_2^n \neq v_j c_2^n$ for some $c_2^n \in I$. Since M is irreducible, by (19) there exist $a(i)_2^m, b(i)_2^m \in R$ with $i = 2, \dots, m$ such that

$$\begin{aligned} & f_0(x, (u_2 c_2^n) a(2)_2^n, \dots, (u_m c_2^n) a(m)_2^n, \\ & (v_2 c_2^n) b(2)_2^n, \dots, (v_m c_2^n) b(m)_2^n) \\ &= f_0(0_M, (u_2 c_2^n) b(2)_2^n, \dots, (u_m c_2^n) b(m)_2^n, \\ & (v_2 c_2^n) a(2)_2^n, \dots, (v_m c_2^n) a(m)_2^n), \end{aligned} \quad (36)$$

that is,

$$\begin{aligned} & f_0(x, u_2 (c_2^n a(2)_2^n), \dots, u_m (c_2^n a(m)_2^n), \\ & v_2 (c_2^n b(2)_2^n), \dots, v_m (c_2^n b(m)_2^n)) \\ &= f_0(0_M, u_2 (c_2^n b(2)_2^n), \dots, u_m (c_2^n b(m)_2^n), \\ & v_2 (c_2^n a(2)_2^n), \dots, v_m (c_2^n a(m)_2^n)), \end{aligned} \quad (37)$$

which means that M is an irreducible I -semimodule by (19) again since for all $j \in \{2, \dots, n\}$ $c_2^n a(j)_2^n, c_2^n b(j)_2^n \in I^{n-1}$.

Assume that M is a semi-irreducible R -semimodule and $MI^{n-1} \neq \{0_M\}$. According to Lemma 24, for any $u (\neq 0_M) \in M$ there exist $b_2^n \in R$ such that $ub_2^n \neq 0_M$. By Lemma 25, $(ub_2^n)R^{n-1} = M$, so for any $x \in M$ there exist positive integers p, q and $x(i)_2^n, y(j)_2^n \in R$ such that $p \equiv q \equiv 0 \pmod{m-1}$ and

$$\begin{aligned} & f_0(x, (ub_2^n) x(2)_2^n, \dots, (ub_2^n) x(p)_2^n) \\ &= f_0(0_M, (ub_2^n) y(2)_2^n, \dots, (ub_2^n) y(q)_2^n), \end{aligned} \quad (38)$$

which shows that

$$\begin{aligned} & f_0(x, (ub_2^n x(2)_2^n), \dots, u(b_2^n x(p)_2^n)) \\ &= f_0(0_M, u(b_2^n x(2)_2^n), \dots, u(b_2^n x(q)_2^n)). \end{aligned} \quad (39)$$

Note that for all i, j , $(b_2^n x(i)_2^n), (b_2^n x(j)_2^n) \in I^{n-1}$. Thus we obtain $\overline{uI^{n-1}} = M$. By Lemma 25 again, M is a semi-irreducible right n -ary I -semimodule.

(2) Let M be an irreducible right n -ary I -semimodule, and let $N = MI^{n-1}$. Then $N \neq \{0_M\}$ and N is an I -subsemimodule of M . Thus by Lemma 35, N is irreducible and for any $x_2^n, y_2^n \in R$ the following conclusion is true: the equality $ux_2^n = uy_2^n$ holds for all $u \in M$ if and only if the same equality holds for all $u \in N$. If $f_0(a(1)x(1)_2^n, \dots, a(p)x(p)_2^n) =$

$f_0(b(1)y(1)_2^n, \dots, b(q)y(q)_2^n)$ for some $x(i)_2^n, y(j)_2^n \in I$ and $a(i), b(j) \in N$, then for any $r_2^n \in R$ and any $z_2^n \in I$,

$$\begin{aligned} & f_0(a(1)(x(1)_2^n r_2^n), \dots, a(p)(x(p)_2^n r_2^n)) z_2^n \\ &= f_0(a(1)(x(1)_2^n r_2^n z_2^n), \dots, a(p)(x(p)_2^n r_2^n z_2^n)) \\ &= f_0(a(1)x(1)_2^n, \dots, a(p)x(p)_2^n)(r_2^n z_2^n) \\ &= f_0(b(1)y(1)_2^n, \dots, b(q)y(q)_2^n)(r_2^n z_2^n) \\ &= f_0(b(1)(y(1)_2^n r_2^n z_2^n), \dots, b(q)(y(q)_2^n r_2^n z_2^n)) \\ &= f_0(b(1)(y(1)_2^n r_2^n), \dots, b(q)(y(q)_2^n r_2^n)) z_2^n, \end{aligned} \quad (40)$$

which implies that

$$\begin{aligned} & f_0(a(1)(x(1)_2^n r_2^n), \dots, a(p)(x(p)_2^n r_2^n)) \\ &= f_0(b(1)(y(1)_2^n r_2^n), \dots, b(q)(y(q)_2^n r_2^n)) \end{aligned} \quad (41)$$

by Lemma 24 since M is an irreducible right n -ary I -semimodule. Thus we can define an operation on NR^{n-1} into N by setting

$$\begin{aligned} & f_0(a(1)x(1)_2^n, \dots, a(p)x(p)_2^n) r_2^n \\ &= f_0(a(1)(x(1)_2^n r_2^n), \dots, a(p)(x(p)_2^n r_2^n)), \end{aligned} \quad (42)$$

where $x(i)_2^n, y(j)_2^n \in I$ and $a(i), b(j) \in N$. Thus N with the addition and the above operation becomes a right n -ary R -semimodule M^* which, as a right n -ary I -semimodule, is isomorphic to the right n -ary I -semimodule N . It is clear that M^* is an irreducible right n -ary R -semimodule. $\square$

Now we are ready to generalize [17, Theorem 2].

Theorem 37. If I is an ideal of an (m, n) -semiring R , then $J(I) = J(R) \cap I$.

Proof. Let R be an (m, n) -semiring and let Δ be the set of all irreducible right n -ary R -semimodules. Then by Definition 27 $J(R) = \bigcap_{M \in \Delta} A_R(M)$. If I is an ideal of an (m, n) -semiring R , then $J(I) = \bigcap_{M \in \Lambda} A_I(M)$, where Λ is the set of all irreducible right n -ary I -semimodules.

For any $M \in \Delta$, according to Lemma 36, we have $M \in \Lambda$. It is evident that $A_I(M) = A_R(M) \cap I$. This shows that $J(I) \subseteq J(R) \cap I$.

For any $M \in \Lambda$, according to Lemma 36, we have that $M^* \in \Delta$ and that M^* can be regarded as an I -subsemimodule N of M . By Lemma 35, we have that $A_I(M) = A_I(N) = A_I(M^*) = A_R(M^*) \cap I$. This shows that $J(I) \supseteq J(R) \cap I$.

Summarizing the above, we obtain that $J(I) = J(R) \cap I$. $\square$

Consequently, we have.

Theorem 38. For an (m, n) -semiring R , $J(R)$ is a radical (m, n) -semiring; that is, $J(J(R)) = J(R)$.

Acknowledgments

This work was supported by the National Natural Foundation of China (no. 10571005) and the Shandong Province Natural Science Foundation ZR2011AM005.

References

- [1] J. M. Howie, *Fundamentals of Semigroup Theory*, vol. 12 of *London Mathematical Society Monographs. New Series*, The Clarendon Press; Oxford University Press; Oxford Science Publications, New York, NY, USA, 1995.
- [2] M. L. Santiago and S. Sri Bala, "Ternary semigroups," *Semigroup Forum*, vol. 81, no. 2, pp. 380–388, 2010.
- [3] W. A. Dudek and I. Grodzinska, "On ideals in regular n -semigroups," *Matematichki Bilten*, vol. 29-30, no. 3-4, pp. 35–44, 1980.
- [4] W. A. Dudek and V. V. Mukhin, "On topological n -ary semigroups," *Quasigroups and Related Systems*, vol. 3, pp. 73–88, 1996.
- [5] W. A. Dudek, "Idempotents in n -ary semigroups," *Southeast Asian Bulletin of Mathematics*, vol. 25, no. 1, pp. 97–104, 2001.
- [6] W. A. Dudek and V. V. Mukhin, "On n -ary semigroups with adjoint neutral element," *Quasigroups and Related Systems*, vol. 14, no. 2, pp. 163–168, 2006.
- [7] Y. Zhu, "On (n, m) -semigroups," *Semigroup Forum*, vol. 84, no. 2, pp. 342–364, 2012.
- [8] R. El Bashir and T. Kepka, "Congruence-simple semirings," *Semigroup Forum*, vol. 75, no. 3, pp. 589–609, 2007.
- [9] A. K. Bhuniya and T. K. Mondal, "Distributive lattice decompositions of semirings with a semilattice additive reduct," *Semigroup Forum*, vol. 80, no. 2, pp. 293–301, 2010.
- [10] Y. Katsov and T. G. Nam, "Morita equivalence and homological characterization of semirings," *Journal of Algebra and its Applications*, vol. 10, no. 3, pp. 445–473, 2011.
- [11] Y. Shitov, "Inequalities for Gondran-Minoux rank and idempotent semirings," *Linear Algebra and its Applications*, vol. 435, no. 7, pp. 1769–1777, 2011.
- [12] R. Pereira, "Bijective linear rank preservers for spaces of matrices over antinegative semirings," *Linear Algebra and its Applications*, vol. 435, no. 7, pp. 1666–1671, 2011.
- [13] M. K. Sen and A. K. Bhuniya, "On semirings whose additive reduct is a semilattice," *Semigroup Forum*, vol. 82, no. 1, pp. 131–140, 2011.
- [14] S. Ebrahimi Atani and M. Shajari Kohan, "A note on finitely generated multiplication semimodules over commutative semirings," *International Journal of Algebra*, vol. 4, no. 5–8, pp. 389–396, 2010.
- [15] S. Bourne, "The Jacobson radical of a semiring," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 37, pp. 163–170, 1951.
- [16] S. Bourne and H. Zassenhaus, "On the semiradical of a semiring," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 44, pp. 907–914, 1958.
- [17] K. Iizuka, "On the Jacobson radical of a semiring," *The Tohoku Mathematical Journal*, vol. 11, pp. 409–421, 1959.
- [18] K. Iizuka and I. Nakahara, "A note on the semiradical of a semiring," *Kumamoto Journal of Science A*, vol. 4, pp. 1–3, 1959.
- [19] Z. Izhakian, J. Rhodes, and B. Steinberg, "Representation theory of finite semigroups over semirings," *Journal of Algebra*, vol. 336, pp. 139–157, 2011.
- [20] T. K. Dutta and S. Kar, "On the Jacobson radical of a ternary semiring," *Southeast Asian Bulletin of Mathematics*, vol. 28, no. 1, pp. 1–13, 2004.
- [21] S. K. Sardar, "On Jacobson radical of a Γ -semiring," *Novi Sad Journal of Mathematics*, vol. 35, no. 1, pp. 1–9, 2005.
- [22] A. Pop, "Remarks on embedding theorems of (m, n) -semirings, Dedicated to Maria Sanziana Pop on her 60th birthday," *Buletinul Stiintific al Universitatii Baia Mare B*, vol. 16, no. 2, pp. 297–302, 2000.
- [23] A. Pop and M. S. Pop, "On partially ordered n -semigroups and (n, m) -semirings," *Creative Mathematics and Informatics*, vol. 18, no. 2, pp. 194–199, 2009.
- [24] S. E. Alam, S. Rao, and B. Davvaz, " (m, n) -semirings and a generalized fault-tolerance algebra of systems," *Journal of Applied Mathematics*, vol. 2013, Article ID 482391, 10 pages, 2013.
- [25] N. Jacobson, "The radical and semi-simplicity for arbitrary rings," *American Journal of Mathematics*, vol. 67, pp. 300–320, 1945.