Conference Paper

Modulation Effect in Oncothermia

Oliver Szasz,¹ Gabor Andocs,² and Nora Meggyeshazi³

¹ Department of Biotechnics, St. Istvan University, Pater K. u. 1., Godollo 2100, Hungary

² Department of Veterinary Clinical Medicine, Faculty of Veterinary Science, Tottori University, 4-101 Minami, Koyama-Cho, Tottori Prefecture 680-8553, Japan

³ 1st Department of Pathology and Experimental Cancer Research, Semmelweis University, Üllői út 26, Budapest 1085, Hungary

Correspondence should be addressed to Oliver Szasz; droliverszasz@gmail.com

Received 15 January 2013; Accepted 9 May 2013

Academic Editors: G. F. Baronzio, M. Jackson, and A. Szasz

This Conference Paper is based on a presentation given by Oliver Szasz at “Conference of the International Clinical Hyperthermia Society 2012” held from 12 October 2012 to 14 October 2012 in Budapest, Hungary.

Copyright © 2013 Oliver Szasz et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Conventional hyperthermia is based on the local or systemic heating, which is measured by the realized temperature in the process. Oncothermia applies nanoheating, which means high energy absorption in the nanoscopic range of the malignant cell membrane selectively. This high temperature and its consequent stress create special effects: it evolves the possibility for chaperone proteins to be expressed on the outer membrane by softening the membrane and starts various excitations for programmed cell death of the targeted malignant cell. The process needs special delivery of the energy which selects as desired. A strict 13.56 MHz sinusoidal carrier frequency is amplitude modulated by time-fractal signals. The modulation is far from any sinus or other periodic patterns; it is a 1/f spectrum having definite templates for its construction. In some personalized cases, a definite template is used for the fractal pattern, which is copied from the actual character of the tumor pathology or any other specialty of the target.

1. Introduction

To understand the principle of modulation, let us start with a simple everyday task: to listen to our favorite radiobroadcasts, like 107.1 MHz Cologne, 91.8 MHz Frankfurt, Radio Energy (Munich) 93.3 MHz, and so forth. We should choose the frequency (tune the radio to select it), and we can enjoy the broadcast. The carrier frequency which was the basis of the tuning never meets the ear; it is too high to sense, and anyway it would be a too monotonic sound as it is only a single frequency. Instead of monotony, we hear the music or other information carried by this chosen frequency. The carrier frequency delivers the real information coded in its modulation (see Figure 1).

The carrier frequency carries two important information characters:

- (i) its modulation finds the target on cellular level;
- (ii) its energy heats up the selected cells from the outside by its neighboring extracellular matrix.

The modulation method is similar to the process when light goes through the window's glass. When the glass is transparent to a specific set of colors (visible light, definite interval of frequencies), its absorption is almost zero; all energy goes through it. However, when it has any bubbles, grains, precipitations, and so forth, those irregularities will absorb a bigger part of the energy, their transparency will be locally low, their energy absorption will be high, and they will be heated up locally. It is a self-selection depending on the material and the frequency (color) which we apply in the given example. The carrier frequency delivers the information (modulation frequencies), for which the cancer cells are much less “transparent” than their healthy counterpart is. Malignant cells are heated up by the selectively absorbed energy.

The applied signal has synergy with the heating of the extracellular matrix, constructing zero-mode noise component which surpasses the thermal noise [1]. The “demodulation” of the noise [2] uses the ratchet mechanism [3] and the stochastic resonance phenomena [4] combined with the membrane rectification [5] and nonlinearity [6].

FIGURE 1: The amplitude modulation. Carrier frequency (a), modulation signal (b), modulated signal showing the frame of the modulator (c), and modulated signal alone (d).

The applied time-fractal modulation is one of the specialties, which only oncoterminia has in hyperthermia applications in oncology. Recently, the research of amplitude-modulated RF in human medicine is intensified [7]. Clinical trials show its progress [8, 9]. Research had shown how an AC electric field inhibits the metastatic spread of a solid lung tumor [10]. The time-fractal modulation follows the natural (healthy) dynamic fluctuations of the tissue, adding a selection facility to distinguish it from the malignant counterpart. The cell junctions and other cellular connections ensure that the proper communications between the cells are dominantly broken [11], making isolation of the malignant cells from each other [11, 12].

2. Method

The living material is not an ordered solid. Contrary to the crystals, it is hard to introduce the cooperativity. The living matter is in an aqueous solution, which is mostly well ordered [13] in the living state. This relative order formed the

“dilute salted water” into the system having entirely different mechanical, chemical, physical, and other behaviors from the normal aqueous solutions. Indeed, the important role of the living systems of the so-called ordered water was pointed out in the middle of the sixties, and later, it was proven [14]. At first, the ordered water was suggested to be as much as 50% of the total amount of the water in the living bodies [15]. The systematic investigations showed more ordered water [16, 17] than it was expected before. Probably, the ordered water bound to the membrane is oriented (ordered) by the membrane potential, which probably decreases the order of the connected water, so it increases the electric permeability of the water [18] and so decreases the cell-cell adhesion and could be the cause of the cell division even for the proliferation [18].

According to Warburg’s effect, the metabolism gradually favors the fermentation in malignancy [19]. The end products of both the metabolic processes are ions in the aqua-based electrolyte. The oxidative cycle products dissociate like $6\text{CO}_2 + 6\text{H}_2\text{O} \leftrightarrow 12\text{H}^+ + 6\text{CO}_3^{2-}$, while the lactate produced by fermentation dissociates: $2\text{CH}_3\text{CHOHCOOH} \leftrightarrow 2\text{CH}_3\text{CHOHCOO}^- + 2\text{H}^+$. Assuming the equal proton production (by more intensive fermentation energy flux), the main difference is in the negative ions. The complex lactate ion concentration grows rapidly and increases its osmotic pressure. To keep the pressure normal, the dissolvent (the monomer water) has to be increased as well, seeking to solvent by nonordered water. Indeed, it is measured in various malignancies that the water changes to be disordered [20–22], so in these cases, the ordered water concentration in cancerous cells is smaller than in their healthy counterpart. Consequently, the hydrogen ionic transmitter becomes weak, and the removal of the hydrogen ions becomes less active. This decreases the intracellular pH and the proton gradient in mitochondria, which directly worsens the efficacy of ATP production. To compensate the lowered proton gradient, the membrane potential of mitochondria grows. This lowers the permeability of the membrane and decreases the mitochondrial permeability transition, which have crucial roles in apoptosis [23, 24]. The high mitochondrial membrane potential and low K-channel expression were observed in cancerous processes [25]. These processes lead to apoptosis resistance, and for the cell energizing the ATP production of the host cell (fermentation) becomes supported. The free-ion concentration increases in the cytoplasm, and so the HSP chaperone stress proteins start to be produced. This process needs more ATP, and it is an antiapoptotic agent, so the process could lead to the complete block of apoptosis. Rearranging (disordering) the water structure needs energy [26]. It is similar to the way the ice is melted with latent heat from zero centigrade solid to liquid with unchanged temperature conditions. This drastic change (phase transition) modifies the physical properties (like the dielectric constant) of the material without changing the composition (only the microscopic ordering) of the medium itself.

The decisional role of the two metabolic pathways (the oxidative and the fermentative) was studied by Szent-Györgyi [18], having an etiology approach and using additional

FIGURE 2: The modulation reestablishes the apoptosis, the natural cell-killing process, and after 48 h, the effect is obviously acting. Immediately after the treatment, there is no difference between the samples, but 24 hours later, the difference is obvious (HT29 xenograft model on single-tumor-bearing mice, heating single shot, 30 min to 42°C. Animals were sacrificed 24 h after the treatment).

formulation. His interpretation describes the cellular states by two different stages. The alpha state of the cell is the fermentative status.

What makes the difference in the absorption? It is the missing collective order in malignancy. The healthy cells live collectively. They have special “social” signals [27] commonly regulating and controlling their life. They are specialized for work division in the organism, and their life cycle is determined by the collective “decisions.” The cancerous cells behave noncollectively; they are autonomic. They are “individual fighters,” having no common control over them, only the available nutrients regulate their life. The order, which characterizes the healthy tissue, is lost in their malignant version, and the cellular communications disappear [11].

The problem of the autonomy of the malignant cells makes the treatment very much complicated, because cancer has its own fractal structure [28]. The analysis of the fractal structures of malignancies could even indicate the stage of the disease [29]. Careful fractal analysis can make predictive information for the prognosis as well [30].

3. Results

The effect of modulation was measured on immunodeficient nude mice xenograft model made by HT29 human colorectal carcinoma cell line, like what was described elsewhere [31]. The single shot oncothermia was used for 30 min keeping 42°C in the tumor. A day after oncothermia, a definite difference could be detected between the modulated and unmodulated effects, which became very emphasized after two days (see Figure 2 [32]). This is one of the reasons why we propose the treatment frequency in the protocols of oncothermia every other day.

The multiple fractal physiological proofs are extended by the oncothermia specialized experimental results too. We used the same xenograft model on a high number of nude mice (30 tumors were examined, 5-5 mice having double

FIGURE 3: The modulation makes a definitely and significantly higher tumor destruction compared to the nontreated side. The high error bars are consequences of the biovariability, making the results trivially different but mathematically not significant. Research is in progress to improve the accuracy. (HT29 cell line in nude mice, xenograft model, single shot for 30 min keeping at 40°C. 5-5 mice were used in both arms; sampling was taken 48 h after the treatment).

tumors in two arms, modulated (active) arm, and nonmodulated arm (passive arm)). The single shot experiment was also for 30 min, but the tumors were treated only on 40°C. We know from other experiments that this temperature is generally not enough to make a hyperthermia effect in a classical heating approach. The animals were sacrificed after 48 h, and the results (see Figure 3) show the modulation effect well: the treated arm in modulated cases had 45.8% higher cell-distortion than that of the nontreated part, while the effect in the nonmodulated mice was only 3.9%.

More detailed explanation and background of the modulation applications in oncothermia could be obtained from

the oncothermia book [33]. The modulation method has patent applications [34–36].

4. Conclusion

Oncothermia modulation is one of the three specialties of this treatment. Its efficacy and its role in the personalization process have introduced an effective tool for the apoptotic cancer cell destruction.

Acknowledgment

The authors acknowledge the experimental work and fruitful discussions of Professor Andras Szasz.

References

- [1] G. Vincze, N. Szasz, and A. Szasz, "On the thermal noise limit of cellular membranes," *Bioelectromagnetics*, vol. 26, no. 1, pp. 28–35, 2005.
- [2] J. Silny, "Demodulation in tissue, the relevant parameters and the implications for limiting exposure," *Health Physics*, vol. 92, no. 6, pp. 604–608, 2007.
- [3] T. Y. Tsong and C. H. Chang, "Ion pump as Brownian motor: theory of electroconformational coupling and proof of ratchet mechanism for Na,K-ATPase action," *Physica A*, vol. 321, no. 1–2, pp. 124–138, 2003.
- [4] L. Gammaitoni, P. Hänggi, P. Jung, and F. Marchesoni, "Stochastic resonance," *Reviews of Modern Physics*, vol. 70, no. 1, pp. 223–287, 1998.
- [5] R. D. Astumian, J. C. Weaver, and R. K. Adair, "Rectification and signal averaging of weak electric fields by biological cells," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 92, no. 9, pp. 3740–3743, 1995.
- [6] A. Beuter, L. Glass, M. C. Mackey et al., Eds., *Nonlinear Dynamics in Physiology and Medicine*, vol. 25 of *Interdisciplinary Applied Mathematics*, Springer, New York, NY, USA, 2003.
- [7] A. Barbault, F. P. Costa, B. Bottger et al., "Amplitude-modulated electromagnetic fields for the treatment of cancer: discovery of tumor-specific frequencies and assessment of a novel therapeutic approach," *Journal of Experimental and Clinical Cancer Research*, vol. 28, no. 1, article 51, 2009.
- [8] E. D. Kirson, V. Dbal, C. Rochlitz et al., "Treatment of locally advanced solid tumors using alternating electric fields (TTFields)—a translational study. Clinical research 17: phase II and III adult clinical trials 1," *Proceedings of American Association Cancer Research*, vol. 47, abstract #5259, 2006.
- [9] National Institutes of Health US, Low Levels of Electromagnetic Fields to Treat Advanced Cancer (ADLG3), <http://clinicaltrials.gov/ct2/show/NCT00805337>.
- [10] E. D. Kirson, M. Giladi, Z. Gurvich et al., "Alternating electric fields (TTFields) inhibit metastatic spread of solid tumors to the lungs," *Clinical and Experimental Metastasis*, vol. 26, no. 7, pp. 633–640, 2009.
- [11] W. R. Loewenstein and Y. Kanno, "Intercellular communication and tissue growth. I. Cancerous growth," *Journal of Cell Biology*, vol. 33, no. 2, pp. 225–234, 1967.
- [12] W. R. Loewenstein, *The Touchstone of Life, Molecular Information, Cell Communication and the Foundations of the Life*, Oxford University Press, Oxford, UK, 1999.
- [13] F. W. Cope, "Nuclear magnetic resonance evidence using D₂O for structured water in muscle and brain," *Biophysical Journal*, vol. 9, no. 3, pp. 303–319, 1969.
- [14] R. Damadian, "Tumor detection by nuclear magnetic resonance," *Science*, vol. 171, no. 3976, pp. 1151–1153, 1971.
- [15] F. W. Cope, "A review of the applications of solid state physics concepts to biological systems," *Journal of Biological Physics*, vol. 3, no. 1, pp. 1–41, 1975.
- [16] C. F. Hazlewood, B. L. Nichols, and N. F. Chamberlain, "Evidence for the existence of a minimum of two phases of ordered water in skeletal muscle," *Nature*, vol. 222, no. 5195, pp. 747–750, 1969.
- [17] C. F. Hazlewood, D. C. Chang, D. Medina, G. Cleveland, and B. L. Nichols, "Distinction between the preneoplastic and neoplastic state of murine mammary glands," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 69, no. 6, pp. 1478–1480, 1972.
- [18] A. Szent-Györgyi, "The living state and cancer," *Physiological Chemistry and Physics*, vol. 12, no. 2, pp. 99–110, 1980.
- [19] O. Warburg, *Oxygen, the Creator of Differentiation*, *Biochemical Energetics*, Academic Press, New York, NY, USA, 1966.
- [20] M. Gniadecka, O. F. Nielsen, and H. C. Wulf, "Water content and structure in malignant and benign skin tumours," *Journal of Molecular Structure*, vol. 661–662, no. 1–3, pp. 405–410, 2003.
- [21] P. T. Beall, B. B. Asch, D. C. Chang, D. Medina, and C. F. Hazlewood, "Water-relaxation times of normal, preneoplastic, and malignant primary cell cultures of mouse mammary gland," in *Proceedings of the 23rd Annual Meeting of the Biophysical Society*, Atlanta, Ga, USA, February 1979.
- [22] S. H. Chung, A. E. Cerussi, C. Klifa et al., "In vivo water state measurements in breast cancer using broadband diffuse optical spectroscopy," *Physics in Medicine and Biology*, vol. 53, no. 23, pp. 6713–6727, 2008.
- [23] G. Fiskum, "Mitochondrial participation in ischemic and traumatic neural cell death," *Journal of Neurotrauma*, vol. 17, no. 10, pp. 843–855, 2000.
- [24] F. Ichas and J. P. Mazat, "From calcium signaling to cell death: two conformations for the mitochondrial permeability transition pore. Switching from low- to high-conductance state," *Biochimica et Biophysica Acta*, vol. 1366, no. 1–2, pp. 33–50, 1998.
- [25] S. Bonnet, S. L. Archer, J. Allalunis-Turner et al., "A mitochondria-K⁺ channel axis is suppressed in cancer and its normalization promotes apoptosis and inhibits cancer growth," *Cancer Cell*, vol. 11, no. 1, pp. 37–51, 2007.
- [26] R. Chidambaram and M. Ramanadham, "Hydrogen bonding in biological molecules—an update," *Physica B*, vol. 174, no. 1–4, pp. 300–305, 1991.
- [27] M. C. Raff, "Social controls on cell survival and cell death," *Nature*, vol. 356, no. 6368, pp. 397–400, 1992.
- [28] L. Ballerini Franzen, "Fractal analysis of microscopic images of breast tissue," 2012, <http://www.wseas.us/e-library/conferences/digest2003/papers/466-198.pdf>.
- [29] M. Tambasco and A. M. Magliocco, "Relationship between tumor grade and computed architectural complexity in breast cancer specimens," *Human Pathology*, vol. 39, no. 5, pp. 740–746, 2008.
- [30] A. Delides, I. Panayiotides, A. Alegakis et al., "Fractal dimension as a prognostic factor for laryngeal carcinoma," *Anticancer Research*, vol. 25, no. 3 B, pp. 2141–2144, 2005.
- [31] G. Andocs, H. Renner, L. Balogh, L. Fonyad, C. Jakab, and A. Szasz, "Strong synergy of heat and modulated electromagnetic

- field in tumor cell killing,” *Strahlentherapie und Onkologie*, vol. 185, no. 2, pp. 120–126, 2009.
- [32] G. Andocs, “Unpublished experiments for oncothermia know-how,” 2008-2009.
- [33] A. Szasz, N. Szasz, and O. Szasz, *Oncothermia—Principles and Practices*, Springer, Heidelberg, Germany, 2010.
- [34] A. Szasz, N. Szasz, and O. Szasz, “Radiofrequency hyperthermia device with target feedback signal modulation,” European Patent Application No. EP, 08075820.4, 2009.
- [35] A. Szasz, N. Szasz, and O. Szasz, “Device and procedure for measuring and examining the signal of systems releasing measurable signals during operation or in response to external excitation,” European Patent Application No. EP, 05798498.1, 2011.
- [36] A. Szasz, N. Szasz, and O. Szasz, “Fractal templates and fractal feedback in homeostatic control,” European Patent Application, 2012.