

Retraction

Retracted: Naturally Occurring Xanthoness: Chemistry and Biology

Journal of Applied Chemistry

Received 3 February 2021; Accepted 3 February 2021; Published 15 March 2021

Copyright © 2021 Journal of Applied Chemistry. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Journal of Applied Chemistry has retracted the article titled “Naturally Occurring Xanthoness: Chemistry and Biology” [1]. The article was found to contain a substantial amount of overlapping material from previously published articles, including the following sources cited as [2–5]:

- (i) Kurt Hostettmann, Hildebert Wagner, “Xanthone glycosides,” *Phytochemistry*, Volume 16, Issue 7, 1977, Pages 821–829, ISSN 0031-9422, doi: 10.1016/S0031-9422(00)86673-X [2].
- (ii) L. M.M. Vieira, A. Kijjoo, Naturally-occurring xanthoness: recent developments, *Current Medicinal Chemistry*, Volume 12, Issue 21, 2005, doi: 10.2174/092986705774370682 [3].
- (iii) S. R. Jensen, J. Schripsema, Chemotaxonomy and pharmacology of *Gentianaceae*. In: Lena Struwe, Victor A. Albert (eds.), *Gentianaceae*, 573-632, January 2001. doi: 10.1017/cbo9780511541865.007 [4].
- (iv) Ozlem Demirkiran, (2007) Xanthoness in *Hypericum*: synthesis and biological activities. In: Khan M. T. H. (eds) *Bioactive Heterocycles III. Topics in Heterocyclic Chemistry*, vol. 9. Springer, Berlin, Heidelberg, Germany, doi: 10.1007/7081_2007_079 [5].

Additionally, 507 words were reproduced from the authors’ earlier review article [6], cited as reference 108 in the article.

The article is being retracted due to this overlap, with the agreement of the editorial board. The authors did not respond to these concerns.

References

- [1] S. Negi, V. K. Bisht, P. Singh, M. S. M. Rawat, and G. P. Joshi, “Naturally occurring xanthoness: chemistry and biology,” *Journal of Applied Chemistry*, vol. 2013, Article ID 621459, 9 pages, 2013.
- [2] H. Kurt and H. Wagner, “Xanthone glycosides,” *Phytochemistry*, vol. 16, no. 7, pp. 821–829, 1977.
- [3] L. M. M. Vieira and A. Kijjoo, “Naturally-occurring xanthoness: recent developments,” *Current Medicinal Chemistry*, vol. 12, no. 21, 2005.
- [4] S. R. Jensen and J. Schripsema, “Chemotaxonomy and pharmacology of *Gentianaceae*,” in *Gentianaceae-Systematics and Natural History*, V. A. Albert, Ed., Cambridge University Press, Cambridge, UK, 2001.
- [5] O. Demirkiran, “Xanthoness in *Hypericum*: synthesis and biological activities,” in *Bioactive Heterocycles III. Topics in Heterocyclic Chemistry*, M. T. H. Khan, Ed., Springer, Berlin, Heidelberg, Germany, 2007.
- [6] J. S. Negi, P. Singh, and B. Rawat, “Chemical constituents and biological importance of *Swertia*: a review,” *Current Research in Chemistry*, vol. 3, pp. 1–15, 2011.

Review Article

Naturally Occurring Xanthenes: Chemistry and Biology

J. S. Negi,¹ V. K. Bisht,¹ P. Singh,² M. S. M. Rawat,² and G. P. Joshi²

¹ Herbal Analytical Laboratory, Herbal Research and Development Institute, Mandal, Gopeshwar, Uttarakhand 246401, India

² Department of Chemistry, HNB Garhwal University, Srinagar, Garhwal, Uttarakhand 246174, India

Correspondence should be addressed to J. S. Negi; negijs@yahoo.com

Received 4 April 2013; Accepted 24 September 2013

Academic Editor: Ming-Jer Lee

Copyright © 2013 J. S. Negi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Xanthenes are one of the biggest classes of compounds in natural product chemistry. A number of xanthenes have been isolated from natural sources of higher plants, fungi, ferns, and lichens. They have gradually risen to great importance because of their medicinal properties. This review focuses on the types, isolation, characterization, biological applications, and biosynthesis of naturally occurring xanthenes isolated so far. Different physicochemical and instrumental methods such as liquid-solid and liquid-liquid extraction, TLC, flash chromatography, column chromatography, IR, ¹H NMR and ¹³C NMR spectroscopy, GLC, HPLC, GC, and LCMS have been widely used for isolation and structural elucidation of xanthenes. Hepatoprotective, anticarcinogenic, antileprosy, antimalarial, antioxidant, anticholinergic, mutagenicity, radioprotective, immunomodulatory, antibone resorption, antiparasitic, neuraminidase inhibitory, anticomplement, antibacterial, antifungal, algicidal, anti-HIV, cardioprotective, antitumoral, antidiabetes, antihyperlipidemic, antiatherogenic, anti-inflammatory, antiulcer, antidiabetic, hypolipidemic, analgesic, antiasthmatic, antihistaminic, antiamebic, diuretic, antidiarrheal, larvicidal, and ovicidal activities have been reported for natural occurring xanthenes. To a certain extent, this review provides necessary foundation for further research and development of new medicines.

1. Introduction

Xanthenes are secondary metabolites commonly occurring in higher plant families, fungi, and lichen [1]. Their pharmacological properties have raised great interest. Structures of xanthenes are related to that of flavonoids and their chromatographic behaviours are also similar. Flavonoids are frequently encountered in nature, whereas xanthenes are found in limited number of families. Xanthenes always occur in the families Gentianaceae, Guttiferae, Moraceae, Clusiaceae, and Polygalaceae. Xanthenes are sometimes found as the parent polyhydroxylated compounds but most are mono- or polymethyl ethers or are found as glycosides [2]. Unlike iridoids, xanthenes are apparently not present in all plant species investigated in the family Gentianaceae. This is documented by the systematic work of Hostettmann et al. [3]. Natural occurrence of 12 xanthenes in higher plants and 4 in fungi has been reviewed by Roberts in 1961 and by Dean in 1963 [4, 5]. Gottlieb [6] mentioned the isolation of 60 xanthenes from higher plants and 7 from fungi, whereas Carpenter et al. [7] listed 82 xanthenes from higher plants.

Gunasekera [8] recorded 183 xanthenes from 5 families of tracheophyta. According to Vieira and Kijjoa [9], out of total 515 xanthenes, 278 were reported from natural sources. These xanthenes have been isolated from 20 families of higher plants (122 species in 44 genera), fungi (19 species), and lichens (3 species). In this period, the xanthenes from higher plants appear to be associated mainly with the families Clusiaceae (55 species in 12 genera) and Gentianaceae (28 species in 8 genera). Bo and Liu [10] have reviewed separation methods used for pharmacologically active xanthenes. Jose Pedraza-Chaverri et al. [11] reviewed the isolated chemical constituents and medicinal properties of *C. Garcinia* (mangostana). Some of the plants, ferns, and fungus species which contain xanthenes are *Artocarpus*, *Anthocleista*, *Allanblackia*, *Andrographis*, *Aspergillus*, *Bersama*, *Blackstonia*, *Calophyllum*, *Canscora*, *Centaurium*, *Chironia*, *Cratoxylum*, *Comastoma*, *Garcinia*, *Cudrania*, *Eustoma*, *Emericella*, *Frasera*, *Garcinia*, *Gentiana*, *Gentianella*, *Gentianopsis*, *Halenia*, *Hoppea*, *Hypericum*, *Ixanthus*, *Lomatogonium*, *Mesua*, *Morinda*, *Macrocarpaea*, *Mangrove* fungi, *Orphium*, *Peperomia*, *Pentadesma*, *Polygala*, *Penicillium*, *Phoma*, *Phomopsis*, *Rheedia*,

Rhus, *Securidaca*, *Symphonia*, *Schultesia*, *Swertia*, *Tripterospermum*, *Vismia*, *Veratrilla*, and *Xylaria*.

2. Classification

Xanthenes isolated from natural sources are classified into six main groups, namely, simple xanthenes, xanthone glycosides, prenylated xanthenes, xanthonolignoids, bisxanthenes, and miscellaneous xanthenes.

2.1. Simple Oxygenated Xanthenes. Simple oxygenated xanthenes are subdivided according to the degree of oxygenation into non-, mono-, di-, tri-, tetra-, penta-, and hexaoxygenated substances [9, 12, 13]. In these xanthenes the substituents are simple hydroxy, methoxy, or methyl groups. About 150 simple oxygenated xanthenes have been reported.

2.1.1. Nonoxygenated Simple Xanthenes. The nonoxygenated xanthenes, namely, methylxanthenes (1-,2-,3-,4-methyl-xanthone), were reported in crude oils from off-shore Norway [14]. This was the first description of xanthenes in fossil organic matter. These xanthenes might have been generated as diagenetic products, formed by oxidation of xanthenes in the reservoir, or might have originated by biosynthesis from aromatic precursors.

2.1.2. Monooxygenated Xanthenes. Besides, six monooxygenated xanthenes from *Swertia*, 2-hydroxyxanthone, 4-hydroxyxanthone, and 2-methoxyxanthone have been isolated from four genera, namely, *Calophyllum*, *Kielmeyera*, *Mesua*, and *Ochrocarpus*.

2.1.3. Dioxxygenated Xanthenes. More than fifteen dioxxygenated xanthenes were reported from plants of the families Clusiaceae and Euphorbiaceae. 1,5-Dihydroxyxanthone, 1,7-dihydroxyxanthone, and 2,6-dihydroxyxanthone are found fairly extensively. Other deoxygenated xanthenes such as 1-hydroxy-5-methoxyxanthone, 1-hydroxy-7-methoxyxanthone, 2-hydroxy-1-methoxy-xanthone, 3-hydroxy-2-methoxyxanthone, 3-hydroxy-4-methoxyxanthone, 5-hydroxy-1-methoxyxanthone, and 1,2-methylenedioxyxanthone have been reported from eleven plants genera.

2.1.4. Trioxxygenated Xanthenes. Forty-five trioxxygenated xanthenes have been reported; out of these fifteen were described as new. Among these, only two natural sulfonated xanthenes, namely, 1,3-dihydroxy-5-methoxyxanthone-4-sulfonate and 5-O- β -D-glucopyranosyl-1,3-dihydroxyxanthone-4-sulfonate, are reported from *Hypericum sampsonii*. These sulfonated xanthenes were found to exhibit significant cytotoxicity against cancer cell line [15, 16]. 1,3,5-, 1,5,6-, 1,6,7-, and 2,3,4-trihydroxyxanthone, seventeen methyl ethers, and two methylenedioxy derivatives from nine genera have been reported.

2.1.5. Tetraoxxygenated Xanthenes. Among the 53 tetraoxxygenated xanthenes identified so far, 21 were found to be new

natural products. These xanthenes were mainly reported from plants of the families Gentianaceae, Clusiaceae, and Polygalaceae. Interestingly, 7-chloro-1,2,3-trihydroxy-6-methoxyxanthone isolated from *Polygala vulgaris* [17] appeared to be the first chloroxanthone of the family Polygalaceae. This compound exhibited antiproliferative activity against the human intestinal adenocarcinoma cell line. The free hydroxyxanthenes are 1,3,5,6-, 1,3,5,7-, and 1,3,6,7-tetrahydroxyxanthone [18].

2.1.6. Pentaoxxygenated Xanthenes. Twenty-seven pentaoxxygenated xanthenes have been identified. Four partially methylated pentaoxxygenated xanthenes, namely, 1,8-dihydroxy-2,3,7-trimethoxyxanthone, 5,6-dihydroxy-1,3,7-trimethoxyxanthone, 1,7-dihydroxy-2,3,8-trimethoxyxanthone, 3,8-dihydroxy-1,2,6-trimethoxyxanthone [19], and 3,7-dihydroxy-1,5,6-trimethoxyxanthone, have been isolated from three plants genera.

2.1.7. Hexaoxygenated Xanthenes. Two hexaoxygenated xanthenes, 8-hydroxy-1,2,3,4,6-pentamethoxyxanthone [15, 20] and 1,8-dihydroxy-2,3,4,6-tetramethoxyxanthone [21], are isolated from two *Centaurium* species and 3-hydroxy-1,2,5,6,7-pentamethoxyxanthone was isolated from the roots of *Polygala japonica*. The natural occurrence of pentaoxxygenated, hexaoxygenated, and dimeric xanthenes has been reviewed by Peres and Nagem [22].

2.2. Xanthone Glycosides. Sixty-one naturally occurring glycosylated xanthenes, thirty-nine of which are new compounds, have been reported predominantly in the families Gentianaceae and Polygalaceae as C- or O-glycosides. The details of naturally occurring xanthone glycosides have been reviewed [2] and distinction between C-glycosides and O-glycosides has also been made. In C-glycosides, C-C bond links the sugar moiety to the xanthone nucleus and they are resistant to acidic and enzymatic hydrolysis whereas the O-glycosides have typical glycosidic linkage.

2.2.1. C-Glycosides. C-glycosides are rare; thus, only seven C-glycosides were mentioned in Sultanbawa's review [13] and 17 in Al-Hazimi's review [23]. Mangiferin and isomangiferin are the most common C-glycosides. Mangiferin (2-C- β -D-glucopyranosyl-1,3,6,7-tetrahydroxyxanthone) is of widespread occurrence in angiosperms and ferns and was first isolated from *Mangifera indica* [24–26]. An isomer, isomangiferin (4-C- β -D-glucopyranosyl-1,3,6,7-tetrahydroxyxanthone), has been isolated from the aerial parts of *Anemarrhena asphodeloides* [27]. Homomangiferin (2-C- β -D-glucopyranosyl-3-methoxy-1,6,7-trihydroxyxanthone) has also been isolated from the bark of *Mangifera indica* [28]. In 1973, another glycoxanthone (2-C- β -D-glucopyranosyl-1,3,5,6-tetrahydroxyxanthone) with an oxidation pattern other than that of mangiferin was found in *Canscora decussate* [29]. Arisawa and Morita [30] have isolated tetraoxxygenated xanthone glycoside 2-C- β -D-glucopyranosyl-5-methoxy-1,3,6-trihydroxyxanthone from *Iris florentina*.

2.2.2. O-Glycosides. More than 20 xanthone O-glycosides are known. A few are from natural sources, namely, gentiocaloside from *Gentiana acaulis*, gentioside from *G. lutea*, and swertianolin from *Swertia japonica* [31]. Their natural occurrence is restricted to the family Gentianaceae. The first xanthone O-glycoside, norswertianin-1-O-glucosyl-3-O-glucoside, was isolated from *S. perennis* [2]. A tetraoxygenated xanthone O-glycoside (3,7,8-trihydroxyxanthone-1-O- β -laminaribioside) was isolated from the fern species [32]. 1-Hydroxy-7-methoxy-3-O-primeverosylxanthone [33] and 1-methoxy-5-hydroxyxanthone-3-O-rutinoside [34] have been isolated from *Gentiana* species and *Canscora decussata*.

2.3. Prenylated and Related Xanthenes. Among 285 prenylated xanthenes, 173 were described as new compounds. The occurrence of prenylated xanthenes is restricted to the plant species of the family Guttiferae. The major C₅ unit of the substituents included the commonly found 3-methylbut-2-enyl or isoprenyl group as in isoemericellin and the less frequent 3-hydroxy-3-methylbutyl as in nigrolineaxanthone P and 1,1-dimethylprop-2-enyl as in globuxanthone, respectively [35–37]. Prenylated xanthenes, caloxanthone O and caloxanthone P, were isolated from *Calophyllum inophyllum* [38] and polyprenylated xanthenes and benzophenones from *Garcinia oblongifolia* [39].

2.4. Xanthonolignoids. Naturally occurring xanthonolignoids are rare, so only five compounds are known. The first xanthonolignoid was isolated from *Kielmeyera* species by Castelh o Jr. et al. [40]. They also isolated two other xanthonolignoids named Cadensins A and B from *Caraipa densiflora*. A xanthonolignoid Kielcorin was obtained from *Hypericum* species [41]. Recently, kielcorin was also isolated from *Vismia guaramirangae* [42], *Kielmeyera variabilis* [43], and *Hypericum canariensis* [44], whereas cadensin C and cadensin D from *Vismia guaramirangae* and *Hypericum canariensis* have been reported [45].

2.5. Bisxanthenes. A total of twelve bisxanthenes, five from higher plants, one from lichen, and six from fungi, have been reported to date. These include jacarellhyperols A and B [46], from the aerial parts of *Hypericum japonicum* and dimeric xanthone, and globulixanthone E, from the roots of *Symphonia globulifera* [47]. Three C₂-C₂' dimeric tetrahydroxyxanthenes dicerandrols A, B, and C, are also isolated from the fungus *Phomopsis longicolla* [48].

2.6. Miscellaneous. Xanthenes with substituents other than those mentioned above are included in this group. Xanthofulvin and vinaxanthone were isolated from *Penicillium* species [49]. A polycyclic substance (xanthopterin) with the ability to inhibit the HSP47 (heat shock protein) gene expression was isolated from the culture broth of a *Streptomyces* species [50]. Xantholiptin is a potent inhibitor of collagen production induced by treatment with TGF- β in human dermal fibroblasts. Xanthenes have been synthesized by different methods. The elements of synthetic methods such as building blocks,

Diels-Alder reaction, and heterogeneous catalysts have also been reviewed [51].

3. Methods for Isolation and Characterization of Xanthenes

Plants xanthenes are commonly isolated by column chromatography on silica gel using different solvent mixtures with increasing polarity [52–55]. Xanthone glycosides are usually crystallized from MeOH. They may also be separated and identified using TLC [56] and HPLC [57–61] by comparison with authentic samples. The structure of xanthenes has been established on the basis of UV, IR, MS, and NMR data [13, 62–72]. Preparative TLC on silica gel using AcOEt, MeOH, and H₂O (21:4:3) as mobile phase has been used in instances of difficult separation. Frequently used solvents in TLC are on polyamide, MeOH-H₂O (9:1) and MeOH-H₂O-AcOH (90:5:5); on cellulose, HOAc (5–30%); on silica gel, Py-H₂O-AcOEt-MeOH (12:10:80:5) and AcOEt-MeOH-H₂O (21:4:3) and chromatoplates are viewed in UV light. In certain cases, spraying with 5% KOH in MeOH or 5% aqueous H₂SO₄ has been advantageous [33]. Polyamide columns are frequently applied for the separation of xanthone glycosides. Purification of xanthenes on Sephadex LH20 column has also been carried out [2]. Xanthenes are also isolated from resin of *Garcinia hanburyi* [73] and from the fermentation products of an endophytic fungus *Phomopsis* [74].

HPLC has been proved as the best technique for separation, identification, and quantification of xanthenes. Several HPLC methods have been developed for naturally occurring xanthenes using microporous chemically bonded silica gel (Micropak CN column), solvent hexane-chloroform (13:7, v/v), isooctane-CHCl₃ (3:17, v/v), or dioxane-dichloromethane (1:9) detected at 254 nm by UV detector [60]. Polar aglycones as well as glycosides of xanthenes are also resolved on reversed phase column (C₈ and C₁₈) using acetonitrile-water as mobile phase [75, 76]. High-speed counter current chromatography (HSCCC) and high performance centrifugal partition chromatography (HPCPC) were also used for the separation and isolation of mangiferin and neomangiferin from an extract of *Anemarrhena asphodeloides* [77] and α -mangostins and γ -mangostins from mangosteen pericarp, respectively [78].

3.1. Ultraviolet Visible Spectroscopy (UV). Ultraviolet visible spectroscopy technique is useful for locating free hydroxyl groups in xanthenes. In particular, the OH group at position 3 is easily detected by addition of NaOAc which results in a bathochromic shift of the 300–330 nm bands with increased intensity. Three or four bands of maximum absorption are always found in the region 220–410 nm and it is noteworthy that all bands show high intensity. Most of the substances show a marked absorption in the 400 nm regions, which accounts for their yellow colour [79].

3.2. Infrared Spectroscopy (IR). The carbonyl group in xanthenes is always easily detectable in IR spectra as a strong band (stretching frequency) in the region of 1657 cm⁻¹ [53].

The presence of a hydroxyl group in the 1 or 8 position lowers the frequency to about 1650 cm^{-1} by hydrogen bonding. Substituents in the 3 or 6 position of the xanthone nucleus may have a marked effect upon the carbonyl stretching frequency [80].

3.3. Proton Nuclear Magnetic Resonance Spectroscopy (^1H NMR). 1D and 2D-NMR spectra (^1H , ^{13}C , DEPT, COSY, TOCSY, HROESY, HSQC, HMBC, and NOESY) have been used for characterization of the xanthenes. The ^1H NMR spectrum appears predominantly in the range of 0–12 ppm downfield from the reference signal of TMS. The integral of the signals is proportional to the number of protons present. ^1H NMR gives information about the substitution pattern on each ring. Acetylated derivatives have been utilised in the structure determination of glycosides [81]. The number and relative position of acetyl and methoxy groups can be determined by observing the shift for the position of absorption for the aromatic protons which occurs upon replacing methoxy group by an acetyl group. Signals between δ 2.40–2.50 are indicative of acetylation at peri-position to the carbonyl group (1 or 8 position) since for other positions the acetyl signals fall between δ 2.30 and 2.35. In nonacetylated xanthenes the presence of hydrogen bonded OH at δ 12–13 also confirms hydroxyl substitution at 1 or 8. But when these positions are unsubstituted, then absorption for the aromatic protons appears at δ 7.70–8.05 [82]. Tetraoxygenated xanthenes, namely, 1,3,7,8- and 1,3,5,8-, showed two meta- and two ortho-coupled protons in the ^1H NMR spectrum. They can also be distinguished by the fact that the presence for the ortho-coupled proton in the 1,3,7,8- system appears at lower field [83] than that for 1,3,5,8- (bellidifolin) system [84]. The signals of 2''-O-acetyl methyl protons of 8-C-glucosyl flavone acetate are found at higher field than those of corresponding 6-C-glucosyl flavone acetate [85]. In a similar manner, 2-C and 4-C isomeric glucosyl xanthenes can be distinguished.

3.4. Carbon Nuclear Magnetic Resonance Spectroscopy (^{13}C NMR). The number of signals in the ^{13}C NMR spectrum indicates the number of different types of C atoms. It gives the information about the total number of the C atoms present in the molecule. It is particularly diagnostic for determining the sugar linkage in di- or polysaccharides; the signal of the carbon carrying the primary alcohols appears at δ 62 in glucose. This signal is shifted to δ 67 in disaccharides possessing a 1–6 linkage [60, 61]. The chemical shift for carbonyl carbon is δ 184.5 when positions 1 and 8 are substituted by hydroxyl groups. But when one of these positions is occupied either by a methoxy or a sugar moiety, the carbonyl signal is shifted upfield by about 4 ppm. If both positions are occupied by a methoxy group or sugar moieties, the upfield shift is about 10 ppm. When methoxy groups are located in position 1 or 8, the corresponding absorption appears at δ 60–61, whereas they appear at about δ 56 when the methoxy group is located in the remaining positions on xanthone nucleus [53].

3.5. Mass Spectrometry (MS). Mass spectrometry is also a useful tool in the structure elucidation of xanthone glycosides. Prox [86] established the fragmentation pattern of

mangiferin and related C-glycosides. Aritomi and Kawasaki [27, 28] obtained satisfactory results using peracetylated derivatives of the same and analogous compounds. In mass spectrum of O-glycosides, no discernible molecular ion peak can be observed, but an important fragment ion peak due to the aglycone moiety appears, followed by further fragmentation. Significant fragment ions from the loss of OH, H_2O , and CHO are typical for xanthenes and related compounds with a methoxy substituent peri to the carbonyl group [34, 53, 87].

4. Biological Activities of Xanthenes

Plants belonging to the family Gentianaceae are best known for their bitter taste due to the presence of xanthenes and are used in traditional remedies against loss of appetite and fever and are still included in many “tonic” formulations [88]. Some specific activities have been reported for xanthenes and iridoids from Gentianaceae. Xanthenes (especially mangiferin) are reported to give CNS stimulation [89, 90] and have anti-inflammatory activity [12]. For bellidifolin and swerchirin, a strong hypoglycemic activity has been reported [91–93]. A crude extract of *Swertia* has been reported to display insect repellent activity [94]. The extracts of most of the *Swertia* species show mutagenic activity [95]. An extract from *S. paniculata* is used in the Indian System of Medicine as a bitter tonic and in the treatment of some mental disorders [96]. *S. hookeri* extract is used in the treatment of microbial infections and as a mood elevator [97]. Swertifranchaside isolated from *S. franchetiana* was found to be potent inhibitor of the DNA polymerase activity of human immunodeficiency virus-1 reverse transcriptase [98]. Naturally occurring xanthenes have emerged as an important class of organic compounds in view of their remarkable pharmacological and other biological activities. It has now been observed that a number of plant products which are in regular use as chemotherapeutic agents contain xanthenes as active constituents. Mangiferin was the first xanthone to be investigated pharmacologically and has been found to exhibit a broad spectrum of biological activities. It shows monoamine oxidase inhibition, cardiostonic, convulsant, and choleric activities [29, 89]. Pronounced anti-inflammatory activity has also been observed for mangiferin [99]. Oral and topical compounds containing mangiferin are useful for the treatment of diseases caused by herpes virus. Mangiferin has been found to protect the liver of the rats from high altitude hypoxia. On the other hand, Ghosal and Chaudhuri [100] have observed the opposite CNS depressant effect for xanthone-O-glycosides in mice and rats. The antimalarial drug AYUSH-64 contains *S. chirata* as one of the ingredients. Xanthenes from *S. chirata* are reported to produce CNS depression [29]. The total extract of *S. chirata* showed significant antifeedant activity against *Jute semilooper* [101]. Nor-swertianolin, an O-glycoside, has been reported to produce antitubercular activity. The O-glycosides of *S. purpurascens* are known to produce CNS depression in albino rats and mice [102]. Xanthenes of *Mammea americana* exhibited inhibitory




FIGURE 1: Biosynthetic pathways leading to the xanthenes (1) and (2).

activity against sarcoma 180 tumor cell [103]. 1,8-Dihydroxy-3,5-dimethoxyxanthone (swerchirin), isolated from the hexane fraction from *Swertia chirayita*, has a very significant blood sugar lowering effect in fasted, fed, glucose loaded, and tolbutamide pretreated albino rats. The ED_{50} for 40% glycaemia lowering in CF male albino rats was 23.1 mg/kg when orally administered [104]. *Swertia* species have also been investigated for the presence of essential elements [105–107]. Xanthenes have been reported to display hepatoprotective, antimicrobial, anticarcinogenic, antileprosy, antioxidant, anticholinergic, mutagenicity [108, 109], and radioprotective effect [110], immunomodulatory effect [111], antibone resorption [112], and antiparasitic effects [113], neuraminidase inhibitory [114], antimalarial [115], anticomplement [116], antifungal and algicidal [117], and anti-HIV activity [118], and cardioprotective, antitumoral, antibacterial, antidiabetes, antihyperlipidemic, antiatherogenic, immunomodulator, anti-inflammatory, antiulcer, antiviral, antifungal [119], antidiabetic, hypolipidemic [120], analgesic, antiasthmatic, antihistaminic, antiamoebic, diuretic, antidiarrheal, larvicidal, ovicidal, antiprotozoal, antileptospirosis, anti-TMV, and anticancer activities [121–124]. Xanthenes from *S. mussotii* were evaluated for their anti-hepatitis B virus activity on HepG 2.2.15 cells line; they exhibited significant activity inhibiting hepatitis B virus DNA replication with IC_{50} values from 0.01 mM to 0.13 mM [125].

5. Biosynthesis of Xanthenes

Biosynthetically xanthenes are of mixed shikimate and acetate origin (Figure 1). Thus, phenylalanine, which is formed from shikimate, loses two carbon atoms from the side-chain and is oxidized to form *m*-hydroxybenzoic acid. This combines with three units of acetate (via malonate) to give the intermediate. The shikimate-acetate intermediate undergoes ring-closure to give substituted benzophenone, which by an oxidative phenol coupling generates the central ring of the xanthone moiety. This oxidative coupling can take place in two ways depending on the folding of the benzophenone either in the *ortho* or in the *para* position to the hydroxyl substituent in the potential B-ring to give 1,3,5-trihydroxyxanthone (1) or the 1,3,7-substituted analogue gentisin (2), respectively. Thus, depending on the orientation of the intermediate, two different hydroxylation patterns can be found. Experimental proof for the overall pathway has been obtained from experiments performed using *Gentiana lutea* [126, 127].

When plants were fed ^{14}C -labeled phenylalanine, the label was recovered solely in the B-ring (Figure 1). Conversely, feeding of ^{14}C -labeled acetate gave incorporation of the main part in the A-ring. The alternative ring closure to (1) has recently been shown to take place in cultured cells of *Centaureum erythraea*, where 2,3',4,6-tetrahydroxybenzophenone

is the precursor for 1,3,5-trihydroxyxanthone [128]. Furthermore, in these cell cultures, compound (1) is selectively oxidized by a xanthone 6-hydroxylase to 1,3,5,6-tetrahydroxyxanthone [129]. Explored methods for synthesis of simple oxygenated xanthenes have been documented by Sousa and Pinto [130].

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors are thankful to Director HRDI and Professor M. S. M. Rawat, Dean School of Sciences, HNB Garhwal University, Srinagar, Uttarakhand, India.

References

- [1] M. L. Cardona, I. Fernández, J. R. Pedro, and A. Serrano, "Xanthenes from *Hypericum reflexum*," *Phytochemistry*, vol. 29, no. 9, pp. 3003–3006, 1990.
- [2] K. Hostettmann and I. Miura, "A new Xanthone diglucoside from *Swertia perennis* L.," *Helvetica Chimica Acta*, vol. 60, pp. 262–264, 1977.
- [3] K. M. Hostettmann, K. Hostettmann, and O. Sticher, "Xanthenes, flavones and secoiridoids of american *Gentiana* species," *Phytochemistry*, vol. 20, no. 3, pp. 443–446, 1981.
- [4] J. C. Roberts, "Naturally occurring xanthenes," *Chemical Reviews*, vol. 61, no. 6, pp. 591–605, 1961.
- [5] F. M. Dean, *Naturally Occurring Oxygen Ring Compounds*, Butterworth, London, UK, 1963.
- [6] O. R. Gottlieb, "Biogenetic proposals regarding aucuparins and xanthenes," *Phytochemistry*, vol. 7, no. 3, pp. 411–421, 1968.
- [7] I. Carpenter, H. D. Locksley, and F. Scheinmann, "Xanthenes in higher plants: biogenetic proposals and a chemotaxonomic survey," *Phytochemistry*, vol. 8, no. 10, pp. 2013–2025, 1969.
- [8] S. P. Gunasekera, University of Sri Lanka, Peradeniya Campus, Peradeniya, Sri Lanka, 1976.
- [9] L. M. M. Vieira and A. Kijjoa, "Naturally-occurring xanthenes: recent developments," *Current Medicinal Chemistry*, vol. 12, no. 21, pp. 2413–2446, 2005.
- [10] T. Bo and H. Liu, "Separation methods for pharmacologically active xanthenes," *Journal of Chromatography B*, vol. 812, no. 1–2, pp. 165–174, 2004.
- [11] J. Pedraza-Chaverri, N. Cárdenas-Rodríguez, M. Orozco-Ibarra, and J. M. Pérez-Rojas, "Medicinal properties of mangosteen (*Garcinia mangostana*)," *Food and Chemical Toxicology*, vol. 46, no. 10, pp. 3227–3239, 2008.
- [12] S. Mandal, P. C. Das, and P. C. Joshi, "Naturally occurring xanthenes from terrestrial flora," *Journal of Indian Chemical Society*, vol. 69, pp. 611–636, 1992.
- [13] M. U. S. Sultanbawa, "Xanthonoids of tropical plants," *Tetrahedron*, vol. 36, no. 11, pp. 1465–1506, 1980.
- [14] T. B. P. Oldenburg, H. Wilkes, B. Horsfield, A. C. T. Van Duin, D. Stoddart, and A. Wilhelms, "Xanthenes—novel aromatic oxygen-containing compounds in crude oils," *Organic Geochemistry*, vol. 33, no. 5, pp. 595–609, 2002.
- [15] T. Janković, D. Krstić, K. Šavikin-Fodulović, N. Menković, and D. Grubišić, "Xanthenes and secoiridoids from hairy root cultures of *Centaurium erythraea* and *C. pulchellum*," *Planta Medica*, vol. 68, no. 10, pp. 944–946, 2002.
- [16] M.-J. Don, Y.-J. Huang, R.-L. Huang, and Y.-L. Lin, "New phenolic principles from *Hypericum sampsonii*," *Chemical and Pharmaceutical Bulletin*, vol. 52, no. 7, pp. 866–869, 2004.
- [17] S. Dall'Acqua, G. Innocenti, G. Viola, A. Piovan, R. Caniato, and E. M. Cappelletti, "Cytotoxic compounds from *Polygala vulgaris*," *Chemical and Pharmaceutical Bulletin*, vol. 50, no. 11, pp. 1499–1501, 2002.
- [18] Y. Chen, G. K. Wang, C. Wu, and M. J. Qin, "Chemical constituents of *Gentiana rhodantha*," *Zhongguo Zhong Yao Za Zhi*, vol. 38, no. 3, pp. 362–365, 2013.
- [19] Q.-C. Xue, C.-J. Li, L. Zuo, J.-Z. Yang, and D.-M. Zhang, "Three new xanthenes from the roots of *Polygala japonica* houtt," *Journal of Asian Natural Products Research*, vol. 11, no. 5, pp. 465–469, 2009.
- [20] P. Valentão, P. B. Andrade, E. Silva et al., "Methoxylated xanthenes in the quality control of small centaury (*Centaurium erythraea*) flowering tops," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 3, pp. 460–463, 2002.
- [21] D. Krstic, T. Jankovic, K. Savikin-Fodulovic, N. Menkovic, and A. Grubisic, "Secoiridoids and xanthenes in the shoots and roots of *Centaurium pulchellum* cultured in-vitro," *In Vitro Cellular and Developmental Biology*, vol. 39, pp. 203–207, 2003.
- [22] V. Peres and T. J. Nagem, "Naturally occurring penta-oxygenated, hexaoxygenated and dimeric xanthenes: a literature survey," *Quimica Nova*, vol. 20, no. 4, pp. 388–397, 1997.
- [23] H. M. G. Al-Hazimi and G. A. Miana, "Naturally occurring xanthenes in higher plants and ferns," *Journal of the Chemical Society of Pakistan*, vol. 12, no. 2, pp. 174–188, 1990.
- [24] S. Iseda, "Isolation of 1,3,6,7-tetrahydroxyxanthone and the skeletal structure of Mangiferin," *Bulletin of the Chemical Society of Japan*, vol. 30, pp. 625–629, 1957.
- [25] L. J. Haynes and D. R. Taylor, "C-glycosyl compounds. Part V. Mangiferin; the nuclear magnetic resonance spectra of xanthenes," *Journal of the Chemical Society C*, pp. 1685–1687, 1966.
- [26] V. K. Bhatia, J. D. Ramanathan, and T. R. Seshadri, "Constitution of mangiferin," *Tetrahedron*, vol. 23, no. 3, pp. 1363–1368, 1967.
- [27] M. Aritomi and T. Kawasaki, "A New xanthone C-glucoside, position isomer of Mangiferin, from *Anemarrhena asphodeloides* Bunge," *Chemical and Pharmaceutical Bulletin*, vol. 18, pp. 2327–2333, 1970.
- [28] M. Aritomi and T. Kawasaki, "A Mangiferin monomethyl ether from *Mangifera indica* L.," *Chemical and Pharmaceutical Bulletin*, vol. 18, pp. 2224–2234, 1970.
- [29] S. Ghosal and R. K. Chaudhuri, "New tetraoxygenated xanthenes of *Canscora decussata*," *Phytochemistry*, vol. 12, no. 8, pp. 2035–2038, 1973.
- [30] M. Arisawa and N. Morita, "Studies on constituents of genus *Iris*. VII. The constituents of *Iris unguicularis* POIR. (I)," *Chemical and Pharmaceutical Bulletin*, vol. 24, no. 4, pp. 815–817, 1976.
- [31] V. Plouvier, J. Massicot, and P. Rivaille, "On gentiacauleine, a new tetra-substituted xanthone, aglycone of gentiacauleside of *Gentiana acaulis* L.," *Comptes Rendus Hebdomadaires des Seances de l'Academie des Sciences D*, vol. 264, no. 9, pp. 1219–1222, 1967.

- [32] F. Imperato, "A xanthone-O-glycoside from *Asplenium adiantum-nigrum*," *Phytochemistry*, vol. 19, no. 9, pp. 2030–2031, 1980.
- [33] A. M. Verney and A. M. Debelmas, "Xanthonnes of *Gentiana lutea*, *G. purpurea*, *G. punctata*, *G. pannonica*," *Annales Pharmaceutiques Francaises*, vol. 31, pp. 415–420, 1973.
- [34] S. Ghosal, R. B. P. S. Chauhan, K. Biswas, and R. K. Chaudhuri, "New 1,3,5-trioxygenated xanthonnes in *Canscora decussata*," *Phytochemistry*, vol. 15, no. 6, pp. 1041–1043, 1976.
- [35] G. Bringmann, G. Lang, S. Steffens, E. Günther, and K. Schumann, "Evariquinone, isoemicellin, and stromemycin from a sponge derived strain of the fungus *Emericella varicolor*," *Phytochemistry*, vol. 63, no. 4, pp. 437–443, 2003.
- [36] L. H. D. Nguyen and L. J. Harrison, "Xanthonnes and triterpenoids from the bark of *Garcinia vilsianiana*," *Phytochemistry*, vol. 53, no. 1, pp. 111–114, 2000.
- [37] V. Rukachaisirikul, M. Kamkaew, D. Sukavisit, S. Phongpaichit, P. Sawangchote, and W. C. Taylor, "Antibacterial Xanthonnes from the Leaves of *Garcinia nigrolineata*," *Journal of Natural Products*, vol. 66, no. 12, pp. 1531–1535, 2003.
- [38] H.-F. Dai, Y.-B. Zeng, Q. Xiao, Z. Han, Y.-X. Zhao, and W.-L. Mei, "Caloxanthonnes O and P: two new prenylated xanthonnes from *Calophyllum inophyllum*," *Molecules*, vol. 15, no. 2, pp. 606–612, 2010.
- [39] W. G. Shan, T. S. Lin, H. N. Yu, Y. Chen, and Z. J. Zhan, "Polyprenylated Xanthonnes and Benzophenones from the Bark of *Garcinia oblongifolia*," *Helvetica Chimica Acta*, vol. 95, no. 8, pp. 1442–1448, 2012.
- [40] J. F. Castela Jr., O. R. Gottlieb, R. A. De Lima, A. A. L. Mesquita, H. E. Gottlieb, and E. Wenkert, "Xanthonolignoids from *Kielmeyera* and *Caraipa* species-13C NMR spectroscopy of xanthonnes," *Phytochemistry*, vol. 16, no. 6, pp. 735–740, 1977.
- [41] H. Nielsen and P. Arends, "Structure of the xanthonolignoid kielcorin," *Phytochemistry*, vol. 17, no. 11, pp. 2040–2041, 1978.
- [42] F. D. Monache, M. M. Mac-Quhae, G. D. Monache, G. B. M. Bettolo, and R. A. De Lima, "Xanthonnes, xanthonolignoids and other constituents of the roots of *Vismia guaramirangae*," *Phytochemistry*, vol. 22, no. 1, pp. 227–232, 1983.
- [43] L. Pinheiro, C. V. Nakamura, B. P. Dias Filho, A. G. Ferreira, M. C. M. Young, and D. A. Garcia Cortez, "Antibacterial Xanthonnes from *Kielmeyera variabilis* Mart. (Clusiaceae)," *Memorias do Instituto Oswaldo Cruz*, vol. 98, no. 4, pp. 549–552, 2003.
- [44] M. L. Cardona, M. I. Fernández, J. R. Pedro, E. Seoane, and R. Vidal, "Additional new xanthonnes and xanthonolignoids from *Hypericum canariensis*," *Journal of Natural Products*, vol. 49, no. 1, pp. 95–100, 1986.
- [45] G. G. Nikolaeva, V. I. Glyzin, M. S. Mladentseva, V. I. Sheichenko, and A. V. Patudin, "Xanthonnes of *Gentiana lutea*," *Chemistry of Natural Compounds*, vol. 19, no. 1, pp. 106–107, 1983.
- [46] K. Ishiguro, S. Nagata, H. Oku, and Y. Masae, "Bisxanthonnes from *Hypericum japonicum*: inhibitors of PAF-induced hypotension," *Planta Medica*, vol. 68, no. 3, pp. 258–261, 2002.
- [47] A. E. Nkengfack, P. Mkounga, M. Meyer, Z. T. Fomum, and B. Bodo, "Globulixanthonnes C, D and E: three prenylated xanthonnes with antimicrobial properties from the root bark of *Symphonia globulifera*," *Phytochemistry*, vol. 61, no. 2, pp. 181–187, 2002.
- [48] M. M. Wagenaar and J. Clardy, "Dicerandrols, new antibiotic and cytotoxic dimers produced by the fungus *Phomopsis longicolla* isolated from an endangered mint," *Journal of Natural Products*, vol. 64, no. 8, pp. 1006–1009, 2001.
- [49] K. Kumagai, N. Hosotani, K. Kikuchi, T. Kimura, and I. Saji, "Xanthofulvin, a novel semaphorin inhibitor produced by a strain of *Penicillium*," *The Journal of Antibiotics*, vol. 56, no. 7, pp. 610–616, 2003.
- [50] Y. Terui, C. Yiwen, L. Jun-Ying et al., "Xantholipin, a novel inhibitor of HSP47 gene expression produced by *Streptomyces* sp.," *Tetrahedron Letters*, vol. 44, no. 29, pp. 5427–5430, 2003.
- [51] C. Yang, M. A. Li, W. E. I. Zhen-ping, H. A. N. Feng, and G. A. O. Jing, "Advances in isolation and synthesis of xanthone derivatives," *Chinese Herbal Medicines*, vol. 4, no. 2, pp. 87–102, 2012.
- [52] J. B. Harbirbe, *Phytochemical Methods*, Chapman and Hall, London, UK, 1973.
- [53] O. Purev, K. Oyun, G. Odontuya et al., "Isolation and structure elucidation of two new xanthonnes from *Gentiana azureum* Bunge (Fam. Gentianaceae)," *Zeitschrift für Naturforschung B*, vol. 57, no. 3, pp. 331–334, 2002.
- [54] A. Khan, M. Rahman, and S. Islam, "Isolation and bioactivity of a xanthone glycoside from *Peperomia pellucida*," *Life Sciences and Medicine Research*, pp. 1–15, 2010.
- [55] V. K. Dua, V. P. Ojha, R. Roy et al., "Anti-malarial activity of some xanthonnes isolated from the roots of *Andrographis paniculata*," *Journal of Ethnopharmacology*, vol. 95, no. 2–3, pp. 247–251, 2004.
- [56] H. M. Chawla, S. S. Chibber, and U. Khera, "Separation and identification of some hydroxyxanthonnes by thin-layer chromatography," *Journal of Chromatography A*, vol. 111, no. 1, pp. 246–247, 1975.
- [57] S. Suryawanshi, N. Mehrotra, R. K. Asthana, and R. C. Gupta, "Liquid chromatography/tandem mass spectrometric study and analysis of xanthone and secoiridoid glycoside composition of *Swertia chirata*, a potent antidiabetic," *Rapid Communications in Mass Spectrometry*, vol. 20, no. 24, pp. 3761–3768, 2006.
- [58] Y. Song, F. Hu, P. Lin, Y. Lu, and Z. Shi, "Determination of xanthonnes in *Swertia mussotii* and *Swertia franchetiana* by high performance liquid chromatography," *Chinese Journal of Chromatography*, vol. 22, no. 1, pp. 51–53, 2004.
- [59] J.-S. Zhang, X.-M. Wang, X.-H. Dong, H.-Y. Yang, and G.-P. Li, "Studies on chemical constituents of *Swertia mussotii*," *Zhong Yao Cai*, vol. 32, no. 4, pp. 511–514, 2009.
- [60] J. S. Negi and P. Singh, "HPLC fingerprinting of highly demanding medicinal plant *Swertia*: an overview," *International Journal of Medicinal and Aromatic Plants*, vol. 1, no. 3, pp. 333–337, 2011.
- [61] J. S. Negi, P. Singh, G. J. Pant, and M. S. M. Rawat, "RP-HPLC analysis and antidiabetic activity of *Swertia paniculata*," *Natural Product Communications*, vol. 5, no. 6, pp. 907–910, 2010.
- [62] R. R. Sun, F. P. Miao, J. Zhang, G. Wang, X. L. Yin, and N. Y. Ji, "Three new xanthone derivatives from an algicolous isolate of *Aspergillus wentii*," *Magnetic Resonance in Chemistry*, vol. 51, no. 1, pp. 65–68, 2013.
- [63] H. H. F. Koolen, L. S. Menezes, M. P. Souza et al., "Talaroxanthone, a novel xanthone dimer from the endophytic fungus *Talaromyces* sp. associated with *Duguetia stelechanthan* (Diels) R. E. Fries," *Journal of the Brazilian Chemical Society*, vol. 24, no. 5, pp. 880–883, 2013.
- [64] A. K. Chakravarty, T. Sarkar, K. Masuda et al., "Structure and synthesis of glycoborine, a new carbazole alkaloid from the roots of *Glycosmis arborea*: a note on the structure of glycozolicine," *Indian Journal of Chemistry B*, vol. 40, no. 6, pp. 484–489, 2001.
- [65] S. Mandal and A. Chatterjee, *Seminar on Research in Ayurveda and Siddha*, CCRAS, New Delhi, India, 1994.

- [66] A. K. Chakravarty, S. Mukhopadhyay, S. K. Moitra, and B. Das, "(-)-Syringaresinol, a hepatoprotective agent and other constituents from *Swertia chirata*," *Indian Journal of Chemistry*, vol. 33, pp. 405–408, 1994.
- [67] K. S. Khetwal and R. S. Bisht, "A xanthone glycoside from *Swertia speciosa*," *Phytochemistry*, vol. 27, no. 6, pp. 1910–1911, 1988.
- [68] D. L. Verma and K. S. Khetwal, "Phenolics in the roots of *Swertia paniculata* Wall.," *Science and Culture*, vol. 51, pp. 305–306, 1985.
- [69] M. Kikuchi and M. Kikuchi, "Studies on the constituents of *Swertia japonica* MAKINO I. On the structures of new secoiridoid diglycosides," *Chemical and Pharmaceutical Bulletin*, vol. 52, no. 10, pp. 1210–1214, 2004.
- [70] H. Yang, C. Ding, Y. Duan, and J. Liu, "Variation of active constituents of an important Tibet folk medicine *Swertia mussootii* Franch. (Gentianaceae) between artificially cultivated and naturally distributed," *Journal of Ethnopharmacology*, vol. 98, no. 1–2, pp. 31–35, 2005.
- [71] S.-S. Wang, W.-J. Zhao, X.-W. Han, and X.-M. Liang, "Two new iridoid glycosides from the Tibetan folk medicine *Swertia franchetiana*," *Chemical and Pharmaceutical Bulletin*, vol. 53, no. 6, pp. 674–676, 2005.
- [72] H. Hajimehdipour, Y. Amanzadeh, S. E. Sadat Ebrahimi, and V. Mozaffarian, "Three tetraoxygenated xanthenes from *Swertia longifolia*," *Pharmaceutical Biology*, vol. 41, no. 7, pp. 497–499, 2003.
- [73] Y. X. Deng, T. Guo, Z. Y. Shao, H. Xie, and S. L. Pan, "Three new xanthenes from the resin of *Garcinia hanburyi*," *Planta Medica*, vol. 79, no. 9, pp. 792–796, 2013.
- [74] H. Y. Yang, Y. H. Gao, D. Y. Niu et al., "Xanthone derivatives from the fermentation products of an endophytic fungus *Phomopsis* sp.," *Fitoterapia*, vol. S0367-326X, no. 13, pp. 00239–00246, 2013.
- [75] A. Aberham, S. Schwaiger, H. Stuppner, and M. Ganzera, "Quantitative analysis of iridoids, secoiridoids, xanthenes and xanthone glycosides in *Gentiana lutea* L. roots by RP-HPLC and LC-MS," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 45, no. 3, pp. 437–442, 2007.
- [76] R. Dai, J. Gao, and K. Bi, "High-performance liquid chromatographic method for the determination and pharmacokinetic study of mangiferin in plasma of rats having taken the traditional Chinese medicinal preparation Zi-Shen Pill," *Journal of Chromatographic Science*, vol. 42, no. 2, pp. 88–90, 2004.
- [77] T. T. Zhou, Z. Y. Zhu, C. Wang et al., "On-line purity monitoring in high-speed counter-current chromatography: application of HSCCC-HPLC-DAD for the preparation of 5-HMF, neomangiferin and mangiferin from *Anemarrhena asphodeloides* Bunge," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 44, no. 1, pp. 96–100, 2007.
- [78] Y. Shan and W. Zhang, "Preparative separation of major xanthenes from mangosteen pericarp using highperformance centrifugal partition chromatography," *Journal of Separation Science*, vol. 33, no. 9, pp. 1274–1278, 2010.
- [79] K. S. Van and B. Nessler, "Ueber einige oxyxanthone," *Chemische Berichte*, vol. 24, pp. 3980–3984, 1891.
- [80] W. E. Whitman and L. A. Wiles, "The polarographic reduction of xanthone and methoxyxanthenes," *Journal of Chemical Society*, pp. 3016–3019, 1956.
- [81] D. Barraclough, H. D. Locksley, F. Scheinmann, M. Taveira Magalhaes, and O. R. Gottlieb, "Applications of proton magnetic resonance spectroscopy in the structural investigation of xanthenes," *Journal of the Chemical Society B*, pp. 603–612, 1970.
- [82] M. Kaldas, K. Hostettmann, and A. Jacot-Guillarmod, "Contribution à la phytochimie du genre *Gentiana* IX. Etude de composés flavoniques et xanthoniques dans les feuilles de *Gentiana campestris* L. 1ère communication," *Helvetica Chimica Acta*, vol. 57, pp. 2557–2561, 1974.
- [83] K. Hostettmann, R. Tabacchi, and A. Jacot-Guillarmod, "Contribution à la phytochimie du genre *Gentiana*, VI. Etude des xanthenes dans les feuilles de *Gentiana bavarica* L.," *Helvetica Chimica Acta*, vol. 57, pp. 294–301, 1974.
- [84] B. Gentili and R. M. Horowitz, "Flavonoids of *Citrus*. IX. Some new c-glycosylflavones and a nuclear magnetic resonance method for differentiating 6- and 8-c-glycosyl isomers," *Journal of Organic Chemistry*, vol. 33, no. 4, pp. 1571–1577, 1968.
- [85] D. K. Holdsworth, "A method to differentiate isomeric C-glucosyl chromones, isoflavones and xanthenes," *Phytochemistry*, vol. 12, no. 8, pp. 2011–2015, 1973.
- [86] A. Prox, "Massenspektrometrische untersuchung einiger natürlicher C-glukosyl-verbindungen," *Tetrahedron*, vol. 24, no. 9, pp. 3697–3715, 1968.
- [87] P. Arends, P. Helboe, and Moller, "Mass spectrometry of xanthenes-I: the electron-impact-induced fragmentation of xanthone, monohydroxy and monomethoxyxanthenes," *Journal of Organic Mass Spectrometry*, vol. 7, pp. 667–681, 1998.
- [88] B. Martindale, *The Extra Pharmacopoeia*, The Pharmaceutical Press, London, UK, 28th edition, 1982.
- [89] S. K. Bhattacharya, S. Ghosal, R. K. Chaudhuri, and A. K. Sanyal, "*Canscora decussata* (Gentianaceae) xanthenes. 3. Pharmacological studies," *Journal of Pharmaceutical Sciences*, vol. 61, no. 11, pp. 1838–1840, 1972.
- [90] C. N. Lin, M. I. Chung, M. Arisawa, M. Shimizu, and N. Morita, "The constituents of *Tripterospermum taiwanense* Satake var. *alpinum* Satake and pharmacological activity of some xanthone derivatives," *Shoyakugaku Zasshi*, vol. 38, pp. 80–82, 1984.
- [91] A. M. Saxena, M. B. Bajpai, P. S. R. Murthy, and S. K. Mukherjee, "Mechanism of blood sugar lowering by a swerchirin-containing hexane fraction (SWI) of *Swertia chirayita*," *Indian Journal of Experimental Biology*, vol. 31, no. 2, pp. 178–181, 1993.
- [92] P. Basnet, S. Kadota, T. Namba, and M. Shimizu, "The hypoglycaemic activity of *Swertia japonica* extract in streptozotocin induced hyperglycaemic rats," *Phytotherapy Research*, vol. 8, no. 1, pp. 55–57, 1994.
- [93] P. Basnet, S. Kadota, M. Shimizu, Y. Takata, M. Kobayashi, and T. Namba, "Bellidifolin stimulates glucose uptake in rat 1 fibroblasts and ameliorates hyperglycemia in streptozotocin (STZ)-induced diabetic rats," *Planta Medica*, vol. 61, no. 5, pp. 402–405, 1995.
- [94] T. Okada, "Insect repellent," Japan. Patent 7875, 327, (Chemical Abstract no. 89: 192516, (1978)), 1977.
- [95] I. Morimoto, F. Watanabe, and T. Osawa, "Mutagenicity screening of crude drugs with *Bacillus subtilis* rec-assay and Salmonella/microsome reversion assay," *Mutation Research*, vol. 97, no. 2, pp. 81–102, 1982.
- [96] A. Prakash, P. C. Basumatary, S. Ghosal, and S. S. Handa, "Chemical constituents of *Swertia paniculata*," *Planta Medica*, vol. 45, no. 1, pp. 61–62, 1982.
- [97] S. Ghosal, K. Biswas, and D. K. Jaiswal, "Xanthone and flavonol constituents of *Swertia hookeri*," *Phytochemistry*, vol. 19, no. 1, pp. 123–126, 1980.
- [98] T. Pengsuparp, L. Cai, H. Constant et al., "Mechanistic evaluation of new plant-derived compounds that inhibit HIV-1 reverse transcriptase," *Journal of Natural Products*, vol. 58, no. 7, pp. 1024–1031, 1995.

- [99] P. C. Das, S. Sarkar, A. Das et al., *Proceeding of the 5th ISHS International Symposium on Medicinal, Aromatic and Spice Plants, Darjeeling, India*, 1985.
- [100] S. Ghosal and R. K. Chaudhuri, "Chemical constituents of gentianaceae XVI: antitubercular activity of xanthenes of *Canscora decussata* Schult," *Journal of Pharmaceutical Sciences*, vol. 64, no. 5, pp. 888–889, 1975.
- [101] R. N. Mullick, P. C. Das, and S. M. Chatterji, "Antifeeding properties of *Swertia chirata* against jute semilooper (*Anomis sabulifera* Guen)," *Current Science*, vol. 54, pp. 1110–1112, 1985.
- [102] S. Ghosal, P. V. Sharma, and R. K. Chaudhuri, "Chemical constituents of gentianaceae X: xanthone O glucosides of *Swertia purpurascens* Wall," *Journal of Pharmaceutical Sciences*, vol. 63, no. 8, pp. 1286–1290, 1974.
- [103] R. A. Finnegan, K. E. Merkel, and J. K. Patel, "Constituents of *Mammea americana* L. XII. Biological data for xanthenes and benzophenones," *Journal of Pharmaceutical Sciences*, vol. 62, no. 3, pp. 483–485, 1973.
- [104] M. B. Bajpai, R. K. Asthana, N. K. Sharma, S. K. Chatterjee, and S. K. Mukherjee, "Hypoglycemic effect of swerchirin from the hexane fraction of *Swertia chirayita*," *Planta Medica*, vol. 57, no. 2, pp. 102–104, 1991.
- [105] J. S. Negi, P. Singh, G. J. Nee Pant, and M. S. M. Rawat, "Study on the variations of mineral elements in *Swertia speciosa* (G. Don)," *Biological Trace Element Research*, vol. 138, no. 1–3, pp. 300–306, 2010.
- [106] J. S. Negi, "Evaluation of trace element contents in *Swertia paniculata* Wall," *Natural Product Research*, vol. 26, no. 1, pp. 72–76, 2012.
- [107] J. S. Negi, P. Singh, M. S. M. Rawat, and G. J. Pant, "Study on the trace elements in *Swertia chirayita* (Roxb.) H. Karsten," *Biological Trace Element Research*, vol. 133, no. 3, pp. 350–356, 2010.
- [108] J. S. Negi, P. Singh, and B. Rawat, "Chemical constituents and biological importance of *Swertia*: a review," *Current Research in Chemistry*, vol. 3, no. 1, pp. 1–15, 2011.
- [109] C. T. Luo, S. S. Mao, F. L. Liu et al., "Antioxidant xanthenes from *Swertia mussotii*, a high altitude plant," *Fitoterapia*, vol. S0367-326X, no. 13, pp. 00229–00233, 2013.
- [110] G. C. Jagetia and V. A. Venkatesha, "Mangiferin, a glucosylxanthone, protects against the radiation-induced micronuclei formation in the cultured human peripheral blood lymphocytes," *International Congress Series*, vol. 1276, pp. 195–196, 2005.
- [111] J. Leiro, J. A. Arranz, M. Yáñez, F. M. Ubeira, M. L. Sanmartín, and F. Orallo, "Expression profiles of genes involved in the mouse nuclear factor-kappa B signal transduction pathway are modulated by mangiferin," *International Immunopharmacology*, vol. 4, no. 6, pp. 763–778, 2004.
- [112] H. Li, T. Miyahara, Y. Tezuka et al., "The effect of *Kampo formulae* on bone resorption in vitro and in vivo. I. Active constituents of *Tsu-kan-gan*," *Biological and Pharmaceutical Bulletin*, vol. 21, no. 12, pp. 1322–1326, 1998.
- [113] S. Perrucci, G. Fichi, C. Buggiani, G. Rossi, and G. Flamini, "Efficacy of mangiferin against *Cryptosporidium parvum* in a neonatal mouse model," *Parasitology Research*, vol. 99, no. 2, pp. 184–188, 2006.
- [114] H. W. Ryu, M. J. Curtis-Long, S. Jung et al., "Xanthenes with neuraminidase inhibitory activity from the seedcases of *Garcinia mangostana*," *Bioorganic and Medicinal Chemistry*, vol. 18, no. 17, pp. 6258–6264, 2010.
- [115] M. V. Ignatushchenko, R. W. Winter, and M. Riscoe, "Xanthenes as antimalarial agents: sage specificity," *The American Journal of Tropical Medicine and Hygiene*, vol. 62, no. 1, pp. 77–81, 2000.
- [116] G.-H. Quan, S.-R. Oh, J.-H. Kim, H.-K. Lee, A. D. Kinghorn, and Y.-W. Chin, "Xanthone constituents of the fruits of *Garcinia mangostana* with anticomplement activity," *Phytotherapy Research*, vol. 24, no. 10, pp. 1575–1577, 2010.
- [117] F. M. Tala, K. Krohn, H. Hussain et al., "Laurentixanthone C: a new antifungal and algicidal xanthone from stem bark of *Vismia laurentii*," *Zeitschrift für Naturforschung B*, vol. 62, no. 4, pp. 565–568, 2007.
- [118] A. Groweiss, J. H. Cardellina II, and M. R. Boyd, "HIV-inhibitory prenylated xanthenes and flavones from *Maclura tinctoria*," *Journal of Natural Products*, vol. 63, no. 11, pp. 1537–1539, 2000.
- [119] M. M. M. Pinto, M. E. Sousa, and M. S. J. Nascimento, "Xanthone derivatives: new insights in biological activities," *Current Medicinal Chemistry*, vol. 12, no. 21, pp. 2517–2538, 2005.
- [120] B. Dineshkumar, A. Mitra, and M. Manjunatha, "Studies on the anti-diabetic and hypolipidemic potentials of mangiferin (xanthone glucoside) in streptozotocin-induced type 1 and type 2 diabetic model rats," *International Journal of Advances in Pharmaceutical Sciences*, vol. 1, no. 1, pp. 75–85, 2010.
- [121] T. Shan, Q. Ma, K. Guo et al., "Xanthenes from mangosteen extracts as natural chemopreventive agents: potential anticancer drugs," *Current Molecular Medicine*, vol. 11, no. 8, pp. 666–677, 2011.
- [122] W. Seesom, A. Jaratrungratawee, S. Suksumran et al., "Antileptospiral activity of xanthenes from *Garcinia mangostana* and synergy of gamma-mangostin with penicillin G," *BMC Complementary and Alternative Medicine*, vol. 13, no. 1, p. 182, 2013.
- [123] S. M. Al-Massarani, A. A. El Gamal, N. M. Al-Musayeib et al., "Phytochemical, antimicrobial and antiprotozoal evaluation of *Garcinia mangostana* pericarp and a-mangostin, its major xanthone derivative," *Molecules*, vol. 18, no. 9, pp. 10599–10608, 2013.
- [124] Y. P. Wu, W. Zhao, Z. Y. Xia et al., "Three novel xanthenes from *Garcinia paucinervis* and their anti-TMV activity," *Molecules*, vol. 18, no. 8, pp. 9663–9669, 2013.
- [125] T. W. Cao, C. A. Geng, Y. B. Ma et al., "Xanthenes with anti-hepatitis B virus activity from *Swertia mussotii*," *Planta Medica*, vol. 79, no. 8, pp. 697–700, 2013.
- [126] J. E. Atkinson, P. Gupta, and J. R. Lewis, "Benzophenone participation in xanthone biosynthesis (Gentianaceae)," *Chemical Communications*, no. 22, pp. 1386–1387, 1968.
- [127] P. Gupta and J. R. Lewis, "Biogenesis of xanthenes in *Gentiana lutea*," *Journal of the Chemical Society C*, pp. 629–631, 1971.
- [128] S. Peters, W. Schmidt, and L. Beerhues, "Regioselective oxidative phenol couplings of 2,3',4,6-tetrahydroxybenzophenone in cell cultures of *Centaurium erythraea* RAFN and *Hypericum androsaemum* L," *Planta*, vol. 204, no. 1, pp. 64–69, 1998.
- [129] W. Schmidt, S. Peters, and L. Beerhues, "Xanthone 6-hydroxylase from cell cultures of *Centaurium erythraea* RAFN and *Hypericum androsaemum* L," *Phytochemistry*, vol. 53, no. 4, pp. 427–431, 2000.
- [130] M. E. Sousa and M. M. M. Pinto, "Synthesis of xanthenes: an overview," *Current Medicinal Chemistry*, vol. 12, no. 21, pp. 2447–2479, 2005.

