

Research Article

Oxygen Consumption at 30 W of Exercise Is Surrogate for Peak Oxygen Consumption in Evaluation of Cardiorespiratory Fitness in Young-Adult African-American Females

Richard M. Millis,¹ Vernon Bond Jr.,² M. Sadegh Asadi,³
Georges E. Haddad,¹ and Richard G. Adams⁴

¹ Department of Physiology & Biophysics, The Howard University College of Medicine, Washington, DC 20059, USA

² Department of Health, Human Performance & Leisure Studies, The Howard University College of Medicine, Washington, DC 20059, USA

³ Department of Neurology, The Howard University College of Medicine, Washington, DC 20059, USA

⁴ Department of Medicine, The Howard University College of Medicine, Washington, DC 20059, USA

Correspondence should be addressed to Richard M. Millis; rickmillis@aol.com

Received 5 July 2013; Accepted 1 September 2013

Academic Editors: S. Perrey, M. Spencer, and A. A. Steiner

Copyright © 2013 Richard M. Millis et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Body mass index (BMI) is negatively correlated with cardiorespiratory fitness, measured by maximal or peak oxygen consumption (VO_{2peak}). VO_{2peak} measurements require heavy aerobic exercise to near exhaustion which increases the potential for adverse cardiovascular events. This study tests the hypothesis that VO_2 measured at a fixed submaximal workload of 30 W is a surrogate for VO_{2peak} . We studied 42 normotensive African-American female university students, 18–25 years of age. We measured VO_{2peak} , blood pressure, and VO_2 at a 30 W exercise workload and computed BMI. We found significant negative correlations between BMI and VO_{2peak} ($r = -0.41$, $P < 0.01$) and between BMI and VO_2 at 30 W ($r = -0.53$, $P < 0.001$). Compared to VO_{2peak} , VO_2 at 30 W increased the significance of the negative correlation with BMI. The heart rate-systolic pressure product at 30 W was positively correlated with BMI ($r = 0.36$, $P < 0.01$) and negatively correlated with VO_{2peak} ($r = -0.38$, $P < 0.001$). The positive correlation between BMI and the heart rate-systolic pressure product and the greater negative correlation between VO_2 and BMI at 30 W of exercise than that at exercise to fatigue suggest that normalized measurements of VO_2 at the fixed exercise workload of 30 W could be useful surrogates for measurements of VO_{2peak} .

1. Introduction

Aerobic exercise testing provides valuable data for measuring a person's cardiorespiratory fitness and overall health. Such testing is also a basis for developing individualized, safe exercise prescriptions. Maximal and peak oxygen consumption (VO_{2max} , VO_{2peak}) are gold standard measuring cardiorespiratory fitness [1]. However, low cardiorespiratory fitness makes it difficult for sedentary, overweight, and/or obese individuals to complete the high-intensity protocols required for VO_{2max} or VO_{2peak} determinations [2]. Moreover, such exercise may put individuals with low cardiorespiratory fitness at risk for adverse cardiovascular events because

determinations of VO_{2max} and VO_{2peak} require substantial exertion to near exhaustion or fatigue [3, 4]. These limitations are consistent with the report that positive electrocardiographic indicators of cardiovascular disease are only 75% sensitive in women, compared to 90% sensitive in men and that African-American women appear to exhibit lower VO_{2peak} than a matched population of Caucasian women [3]. These findings suggest that lack of reliable measures of cardiorespiratory fitness at submaximal workloads may limit our ability to evaluate the health status and prescribe appropriate exercise regimens for women. These impediments have been addressed by the usage of submaximal aerobic exercise tests that are shown to be equally as reliable as VO_{2max} and

VO_{2peak} for measuring cardiorespiratory fitness in sedentary populations [4–9]. Although body mass index (BMI) is a reliable inverse correlate of VO_{2max} and VO_{2peak} in most populations, no studies have been performed to determine the robustness of the correlation between BMI and VO_{2peak} , compared to that at a fixed submaximal workload. Therefore, the present study tests the hypothesis that VO_2 measured during submaximal exercise at a 30 W exercise workload is a surrogate for the VO_{2peak} measurement, as a correlate of BMI and, therefore, of cardiorespiratory fitness.

2. Methods

2.1. Subjects. Forty-two healthy young-adult women volunteered to participate in the study. Their anthropomorphic and physiological characteristics are summarized in Table 1. Participants were normotensive, free of any medication, nonsmoking, and nondrinkers. No participant engaged in regular physical activity and was informed of the study risks. The institutional review board at Howard University granted ethical approval, and informed consent was obtained from all subjects prior to study participation.

2.2. Study Protocol. Subjects participated in three separate sessions in the laboratory. The first session was used to familiarize the participant with the study monitors and devices. In the second laboratory visit, participants were instructed to abstain from exercise and caffeine or any energy drinks for 6 h and food for 3 h prior to entering the laboratory. Body height and weight were measured using standard laboratory procedures. The participant then performed a progressive test of VO_{2peak} . Approximately 1-2 weeks after the second laboratory visit, participants performed the third laboratory visit. Prior to entering the laboratory, the participants were reminded of the prior physical activity and fasting instructions upon entering the laboratory. The participant then performed a submaximal steady-state exercise test using a work output of 30 W.

2.3. Peak Oxygen Consumption Test. VO_{2peak} was measured during a standardized incremental cycle task with a SensorMedics Ergoline-800 ergometer (SensorMedics Corp., Yorba Linda, CA). Participants were instructed to cycle continuously at 70–75 rpm, at a starting work intensity of 25 W. The work rate was increased by 25 W every 3 min until volitional fatigue. During the incremental exercise test, expired gas fractions of VO_2 , carbon dioxide, and minute ventilation (expired) were measured using the method of open-circuit indirect calorimetry (Physio-Dyne Max II Metabolic System, Quogue, NY). The gas analyzers were calibrated using known medical grade gas concentrations. The pneumatic gas volume was calibrated using a 3-L syringe. The VO_2 value achieved during the last minute of the incremental exercise test was defined as VO_{2peak} .

2.4. Submaximal Exercise Test. Participants cycled on the ergometer at an absolute work output of 30 W for a duration of 10 min. This low intensity work load of 30 W was selected

TABLE 1: Descriptive characteristics of the study subjects.

| Variables | Subjects ($n = 42$) |
|--|-----------------------|
| Age (yr) | 20.7 ± 2.2 |
| Height (cm) | 165.3 ± 7.9 |
| Weight (kg) | 68.4 ± 11.7 |
| VO_{2peak} (mL·kg ⁻¹ ·min ⁻¹) | 26.4 ± 3.9 |
| HR _{peak} (beats·min ⁻¹) | 182.3 ± 12.3 |
| $VO_{2@30W}$ (mL·kg ⁻¹ ·min ⁻¹) | 11.2 ± 2.2 |
| HR _{@30W} (beats·min ⁻¹) | 80.9 ± 12.3 |
| Systolic pressure at 30 W (mm Hg) | 119.0 ± 2.8 |
| Diastolic pressure at 30 W (mm Hg) | 76.8 ± 2.2 |
| Rate-pressure product at 30 W (bpm·mm Hg) | 9,445 ± 1,589 |

HR_{peak}: peak heart rate; VO_{2peak} : peak oxygen consumption. HR_{@30W}: heart rate during exercise at the submaximal workload of 30 W; $VO_{2@30W}$: oxygen consumption during exercise at the submaximal workload of aerobic exercise. Systolic pressure, diastolic pressure, and rate-pressure product at 30 W = systolic blood pressure, diastolic blood pressure, and heart rate × systolic pressure product during exercise at the fixed workload of 30 W. Data are means ± standard deviations.

because of the sedentary lifestyle of the study participants. Prior to the study, the electric brake ergometer was calibrated. Prior to the submaximal steady-state workload, the participants were instrumented with the SunTech Tango (Sun-Tech Medical Inc., Raleigh, NC) automated blood pressure monitor that gates the R-wave with the Korotkoff sound to determine blood pressure. Heart rate was determined by electrocardiograph recordings of three electrodes positioned at the RA, LA, and V₅ anatomical positions using the automated blood pressure device. Baseline blood pressure and heart rate measures were collected during the last 5 min of a 10 min sitting rest position. After baseline recordings, the participants performed 10 min of submaximal exercise on the cycle ergometer at a work intensity of 30 W. Heart rate, systolic and diastolic blood pressure were recorded during the last minute of the exercise.

2.5. Statistical Analysis. Pearson's product-moment coefficient (r) and parametric linear regression analysis were used to compare the one-sided significance of correlations between BMI and VO_{2peak} , between BMI and VO_2 at 30 W of exercise, and between BMI and the heart rate-systolic blood pressure product at 30 W of exercise (Microsoft Excel, 2007).

3. Results

Table 1 presented the anthropomorphic and physiological characteristics of the study population. The subjects were mainly normotensive young-adult women with sedentary life style and hence low levels of cardiovascular fitness and low correlating VO_{2peak} levels. Figure 1 depicts the results of linear regression analysis demonstrating a significant negative correlation between both BMI and VO_{2peak} ($r = -0.41, P < 0.01$). The negative correlation between body mass and VO_{2peak} is not shown ($r = -0.45, P < 0.001$). Figure 2 presents the linear regression analysis and significant negative correlation between BMI and VO_2 at 30 W of exercise




FIGURE 1: Linear regression analysis of the relationship between body mass index (BMI) and peak oxygen consumption (VO_{2peak}). Subjects were 42 disease-free, normotensive, sedentary young-adult African-American females. VO_{2peak} was found to be significantly correlated with BMI ($r = -0.41$, $P < 0.01$).




FIGURE 2: Linear regression analysis of the relationship between body mass index (BMI) and oxygen consumption (VO_2) at the fixed submaximal workload of 30 W. Subjects were 42 disease-free, normotensive, sedentary young-adult African-American females. VO_2 was found to be significantly correlated with BMI ($r = -0.53$, $P < 0.001$).

($r = -0.53$, $P < 0.001$). The negative correlation between body mass and VO_2 at 30 W of exercise is also not shown ($r = -0.55$, $P < 0.001$). Correlations between BMI or body mass and heart rates and systolic and diastolic blood pressure at 30 W of exercise were not significant ($P > 0.1$). Both BMI and body weight were positively correlated with the heart rate-systolic pressure product at 30 W of exercise ($r = 0.36$, $P < 0.01$ and $r = 0.39$, $P < 0.001$, resp.). The heart rate-systolic pressure product was negatively correlated with VO_{2peak} ($r = -0.38$, $P < 0.001$).

4. Discussion

This study is the first to compare significance of the correlation between BMI and VO_{2peak} to that between BMI and VO_2 at a fixed, submaximal exercise workload of 30 W in a disease-free population. The participants of this study were normotensive African-American female university students, 18–25 years of age, nonsmokers, nondrinkers, and free of any medication. The main finding of this study is a more significant correlation between BMI or body weight and VO_2

at the fixed workload of 30 W than that between BMI or body weight and VO_{2peak} . Overweight or obese subjects often experience difficulty and adverse cardiovascular events while performing cardiorespiratory fitness tests requiring maximal or fatiguing exertion. A similar study has not been performed in another population. Thus, the correlation coefficients reported herein cannot be compared to those reported in previous studies.

Heretofore, the popular wisdom was that VO_{2max} or VO_{2peak} are the most reliable measures of aerobic capacity and, therefore, cardiorespiratory fitness [3, 4]. However, it is reported that several submaximal exercise protocols such as perceptually regulated, graded exercise with computation of an aerobic power index, step tests, and dance tests provide reliable alternatives to VO_{2max} or VO_{2peak} for measuring cardiorespiratory fitness [4–8]. Measuring cardiorespiratory fitness by submaximal exercise testing and estimating workload at a fixed heart rate are also a promising approach, yielding highly linear, significant correlation coefficients between heart rates and workloads >0.9 [10].

Measurements of cardiorespiratory fitness using exercise tests at submaximal workloads can help determine occupational fitness and evaluate work-related disabilities associated with jobs requiring large physical workloads [11]. Another important use of submaximal cardiorespiratory fitness tests is to evaluate the advertised safety, cost-benefit, and health outcome claims of exercise and dietary regimens [12]. The association of low VO_{2peak} and decreased motor strength in a population of 60-year-old healthy men [11] suggests that submaximal cardiorespiratory fitness testing, physical therapy counseling, and interventions in such populations might decrease their high rate of daily activity- and work-related injuries, as well as the associated health care costs. The further importance of screening such a population for cardiorespiratory fitness is underscored by a report that low cardiorespiratory fitness is associated with high risk for sudden cardiac death in a population of middle-aged men [13]. Thus, results of the present study imply that cardiorespiratory fitness can be reliably measured in populations of overweight, elderly, or otherwise frail subjects by cycle ergometer exercise at a workload of 30 W, thereby reducing the potential for adverse cardiovascular events.

This is also the first study to show a significant negative correlation between VO_{2peak} and the heart rate-systolic pressure product, as well as positive correlations between BMI or body weight and the heart rate-pressure product during aerobic exercise at a fixed submaximal workload. The finding of significant correlation between body mass and the heart rate-pressure product, a measure of myocardial oxygen demand [14], indicates a significant association of an increased requirement for coronary blood flow during exercise in overweight or obese compared to normal-weight persons. The coronary is the circulation with the lowest venous oxygen content, oxygen extraction ratio, and, therefore, oxygen demand at rest. Increases in myocardial oxygen demand must be met, mainly, by increases in coronary blood flow which, when compromised, can result in adverse cardiac events and sudden death [15]. The fact that coronary blood flow is limited by arterial narrowing in atherosclerosis is

well known [16], but other causes such as smoking, nicotine, and cocaine use are less well appreciated. Coronary arterial luminal diameters and areas are shown to be significantly smaller in females than in males, as well as in overweight than in normal-weight individuals [17]. Normotensive African-Americans, especially women, have also been shown to have limitations of endothelial function known to affect the coronary circulation [18, 19]. Therefore, our finding of a significant association between large body mass and large heart rate-pressure product during exercise at 30 W in disease-free normotensive, sedentary young-adult females may be indicative of the potential for limitations of coronary blood flow linked to adverse cardiac events associated with aerobic exercise, smoking, and cocaine use in this population [20]. This finding also supports the hypothesis that experiencing adverse cardiac events during exercise could explain lack of participation of persons with low aerobic capacity in exercise programs [21], thereby creating a vicious cycle of exercise avoidance, omitting effective strategies for weight loss, and improving cardiorespiratory fitness.

5. Limitations and Conclusions

Limitations of this study were (1) inclusion of only sedentary subjects with a relatively low level of cardiorespiratory fitness, thereby limiting our ability to extrapolate the results to a wider range of aerobic capacity; (2) exclusion of obese subjects; (3) not randomizing the exercise procedures and, therefore, not varying the order of presentation and measurement of the $\text{VO}_{2\text{peak}}$ test before and after measuring $\text{VO}_{2\text{peak}}$ at the submaximal, 30 W workload; and (4) performing the cardiovascular measurements such as heart rate and blood pressure only during the submaximal exercise trial. These cardiovascular measurements were made to determine whether the study subjects exhibited physiological responses at this submaximal, absolute workload and whether the subjects exhibiting the highest cardiorespiratory fitness during the $\text{VO}_{2\text{peak}}$ test would exhibit the lowest cardiac oxygen demand at the submaximal 30 W workload, as expected.

In summary, this study demonstrates significant associations between large body mass, low oxygen consumption, and high myocardial oxygen demand during aerobic exercise at a fixed workload of 30 W in a population of normotensive, sedentary, young-adult African-American females. The greater correlations between BMI or body weight and oxygen consumption found at 30 W of submaximal exercise than those between BMI or body weight and $\text{VO}_{2\text{peak}}$ in this population suggest that normalized measurements of VO_2 during exercise at submaximal workloads may be useful surrogates for measurements of $\text{VO}_{2\text{peak}}$ to limit adverse cardiac events without loss of reliability in evaluations of cardiorespiratory fitness.

Conflict of Interests

The authors have no conflict of interests, financial or otherwise, to disclose.




Acknowledgments

This work was supported by NIH-MBRS Program and in part by Grant 2 G12 RR003048 from the RCMI Program Division of Research Infrastructure, National Center for Research, National Institutes of Health.

References

- [1] D. L. Huggett, D. M. Connelly, and T. J. Overend, "Maximal aerobic capacity testing of older adults: a critical review," *The Journals of Gerontology Series A*, vol. 60, no. 1, pp. 57–66, 2005.
- [2] R. M. Ross, J. N. Murthy, I. D. Wollak, and A. S. Jackson, "The six minute walk test accurately estimates mean peak oxygen uptake," *BMC Pulmonary Medicine*, vol. 10, article 31, 2010.
- [3] C. L. Melby, R. C. Ho, K. Jeckel, L. Beal, M. Goran, and W. T. Donahoo, "Comparison of risk factors for obesity in young, nonobese African-American and Caucasian women," *International Journal of Obesity*, vol. 24, no. 11, pp. 1514–1522, 2000.
- [4] K. E. Wallman and L. Campbell, "Test-retest reliability of the Aerobic Power Index submaximal exercise test in an obese population," *Journal of Science and Medicine in Sport*, vol. 10, no. 3, pp. 141–146, 2007.
- [5] K. Wallman, C. Goodman, A. Morton, R. Grove, and B. Dawson, "Test-retest reliability of the aerobic power index component of the tri-level fitness profile in a sedentary population," *Journal of Science and Medicine in Sport*, vol. 6, no. 4, pp. 443–454, 2003.
- [6] J. P. Buckley, J. Sim, R. G. Eston, R. Hession, and R. Fox, "Reliability and validity of measures taken during the Chester step test to predict aerobic power and to prescribe aerobic exercise," *British Journal of Sports Medicine*, vol. 38, no. 2, pp. 197–205, 2004.
- [7] R. G. Eston, J. A. Faulkner, E. A. Mason, and G. Parfitt, "The validity of predicting maximal oxygen uptake from perceptually regulated graded exercise tests of different durations," *European Journal of Applied Physiology*, vol. 97, no. 5, pp. 535–541, 2006.
- [8] M. S. Olson, H. N. Williford, D. L. Blessing, G. D. Wilson, and G. Halpin, "A test to estimate $\text{VO}_{2\text{max}}$ in females using aerobic dance, heart rate, BMI, and age," *Journal of Sports Medicine and Physical Fitness*, vol. 35, no. 3, pp. 159–168, 1995.
- [9] S. McMahon and H. A. Wenger, "The relationship between aerobic fitness and both power output and subsequent recovery during maximal intermittent exercise," *Journal of Science and Medicine in Sport*, vol. 1, no. 4, pp. 219–227, 1998.
- [10] T. G. Lohman, K. Ring, K. Pfeiffer et al., "Relationships among fitness, body composition, and physical activity," *Medicine and Science in Sports and Exercise*, vol. 40, no. 6, pp. 1163–1170, 2008.
- [11] E. L. Cadore, R. S. Pinto, C. L. Alberton et al., "Neuromuscular economy, strength, and endurance in healthy elderly men," *Journal of Strength and Conditioning Research*, vol. 25, no. 4, pp. 997–1003, 2011.
- [12] J. L. Graef, A. E. Smith, K. L. Kendall et al., "The effects of four weeks of creatine supplementation and high-intensity interval training on cardiorespiratory fitness: a randomized controlled trial," *Journal of the International Society of Sports Nutrition*, vol. 6, article 18, 2009.
- [13] J. A. Laukkanen, T. H. Mikkilä, R. Rauramaa, V. Kiviniemi, K. Ronkainen, and S. Kurl, "Cardiorespiratory fitness is related to the risk of sudden cardiac death: a population-based follow-up

- study,” *Journal of the American College of Cardiology*, vol. 56, no. 18, pp. 1476–1483, 2010.
- [14] M. D. Muller, J. L. Mast, H. Patel, and L. I. Sinoway, “Cardiac mechanics are impaired during fatiguing exercise and cold pressor test in healthy older adults,” *Journal of Applied Physiology*, vol. 114, no. 2, pp. 186–194, 2013.
- [15] J. D. Tune, M. W. Gorman, and E. O. Feigl, “Matching coronary blood flow to myocardial oxygen consumption,” *Journal of Applied Physiology*, vol. 97, no. 1, pp. 404–415, 2004.
- [16] C. Berry, S. Noble, R. Ibrahim et al., “Remodeling is a more important determinant of lumen size than atheroma burden in left main coronary artery disease,” *American Heart Journal*, vol. 160, no. 1, pp. 188–e1, 2010.
- [17] S.-G. Kim, S. Apple, G. S. Mintz et al., “The importance of gender on coronary artery size: in-vivo assessment by intravascular ultrasound,” *Clinical Cardiology*, vol. 27, no. 5, pp. 291–294, 2004.
- [18] J. L. Houghton, A. A. Carr, D. S. Strogatz et al., “Coronary vasomotor reactivity among normotensive African and white American subjects with chest pain,” *American Journal of Medicine*, vol. 102, no. 3, pp. 245–251, 1997.
- [19] J. L. Houghton, E. F. Philbin, D. S. Strogatz et al., “The presence of African American race predicts improvement in coronary endothelial function after supplementary L-arginine,” *Journal of the American College of Cardiology*, vol. 39, no. 8, pp. 1314–1322, 2002.
- [20] J. Lucena, M. Blanco, C. Jurado et al., “Cocaine-related sudden death: a prospective investigation in south-west Spain,” *European Heart Journal*, vol. 31, no. 3, pp. 318–329, 2010.
- [21] J. Goodman, S. Thomas, and J. F. Burr, “Physical activity series: cardiovascular risks of physical activity in apparently healthy individuals: risk evaluation for exercise clearance and prescription,” *Canadian Family Physician*, vol. 59, no. 1, pp. 46–49, 2013.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

