Review Article

Mycotoxin Analysis: New Proposals for Sample Treatment

**Natalia Arroyo-Manzanares, José F. Huertas-Pérez,
Ana M. García-Campaña, and Laura Gámiz-Gracia**

Department of Analytical Chemistry, Faculty of Sciences, University of Granada, Campus Fuentenueva s/n, E-18071 Granada, Spain

Correspondence should be addressed to Laura Gámiz-Gracia; lgamiz@ugr.es

Received 10 February 2014; Revised 22 July 2014; Accepted 22 July 2014; Published 7 September 2014

Academic Editor: Brijesh Tiwari

Copyright © 2014 Natalia Arroyo-Manzanares et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Mycotoxins are toxic secondary metabolites produced by different fungi, with different chemical structures. Mycotoxins contaminate food, feed, or raw materials used in their production and cause diseases and disorders in humans and livestock. Because of their great variety of toxic effects and their extreme heat resistance, the presence of mycotoxins in food and feed is considered a high risk to human and animal health. In order to ensure food quality and health consumers, European legislation has set maximum contents of some mycotoxins in different matrices. However, there are still some food commodities susceptible to fungal contamination, which were not contemplated in this legislation. In this context, we have developed new analytical techniques for the multiclass determination of mycotoxins in a great variety of food commodities (some of them scarcely studied), such as cereals, pseudocereals, cereal syrups, nuts, edible seeds, and botanicals. Considering the latest technical developments, ultrahigh performance liquid chromatography coupled to tandem mass spectrometry has been chosen as an efficient, fast, and selective powerful analytical technique. In addition, alternative sample treatments based on emerging methodologies, such as dispersive liquid-liquid microextraction and QuEChERS, have been developed, which allow an increased efficiency and sample throughput, as well as reducing contaminant waste.

1. Introduction

Mycotoxins are toxic natural secondary metabolites produced by several species of fungi (as *Fusarium*, *Aspergillus*, and *Penicillium* genera) on agricultural commodities. The presence of mycotoxins in food and feed may affect human and animal health, as they may cause many different adverse effects such as estrogenic, gastrointestinal, and kidney disorders, induction of cancer, and mutagenicity. Furthermore, some mycotoxins are also immunosuppressive and reduce resistance to infectious diseases [1, 2]. Mycotoxins grow under a wide range of climatic conditions and the Food and Agriculture Organization (FAO) has estimated that they affect 25% of the world crops. On the other hand, mycotoxins are the hazard category with the highest number of border rejections reported by the Rapid Alert System for Food and Feed (RASFF) [3]; therefore their impact on economy is evident.

Hundreds of mycotoxins have been recognized with diverse chemical structures, different toxicity, and biological effects. The most relevant groups of mycotoxins found in

food are aflatoxins (aflatoxin B1 (AFB₁), included in group 1 of carcinogenic to humans by the International Agency for Research on Cancer (IARC) [4]), aflatoxin B2 (AFB₂), aflatoxin G1 (AFG₁), aflatoxin G2 (AFG₂), and aflatoxin M1 (AFM₁, metabolite of AFB₁, excreted in the milk of mammals); ochratoxin A (OTA); trichothecenes (HT-2 and T-2 toxin and deoxynivalenol (DON)); zearalenone (ZEN); fumonisins B1 and B2 (FB₁ and FB₂); citrinin (CIT); patulin (PAT) and ergot alkaloids [2, 5]. Figure 1 shows the structures of some common mycotoxins and Table 1 includes the most important toxins, their main producing fungi, and typical food commodities that may be contaminated by them.

In order to protect consumer health, international institutions and organizations have proposed regulatory limits for some mycotoxins. Thus, the European Commission (EC) establishes maximum permitted levels for most mycotoxins in foods by means of the Commission Regulation (EC) number 1881/2006 [6] (or recommended levels for other mycotoxins [7]), as well as methods of sampling and analysis for

FIGURE 1: Structures of some common mycotoxins: (a) citrinin; (b) aflatoxin B₁; (c) fumonisin B₁; (d) deoxynivalenol; (e) patulin; (f) ochratoxin A.

their control by Commission Regulation (EC) number 401/2006 [8], which have been subsequently amended. Regulations are also established by the US Food and Drug Administration (FDA). FDA mycotoxin compliance programs provide introductory information about mycotoxins, products prone to contamination, and analytical methods [9].

In this context, the use of robust analytical methodologies for sampling, sample treatment, and identification/quantification of mycotoxins in food and feed is mandatory, in order to protect the consumer health [10–12].

2. Current Analytical Methods for Determination of Mycotoxins in Food

Different analytical methods have been proposed for mycotoxin determination in food, such as thin layer chromatography (TLC) [13], ELISA [14], gas chromatography (GC) [15] or capillary electrophoresis (CE) [16]. However, the most popular technique is high performance liquid chromatography (HPLC) with UV/Vis, fluorescence (FL) [17–19], or mass spectrometry (MS) detection [20–22]. Recently, ultrahigh performance liquid chromatography (UHPLC) coupled with tandem mass spectrometry (MS/MS) has become very popular, especially for multiclass determination of mycotoxins and

for multiresidue determination with other contaminants [23–26].

Because of the complexity of food matrices, an extraction and clean-up purification step is usually required before analysis. Different approaches have been proposed. The most common methodology implies solid-liquid extraction (SLE) followed by solid phase extraction (SPE) with immunoaffinity columns (IACs), which contain specific antibodies to the analyte of interest [10]. Several reviews present an overview of the different methodologies proposed for the determination of mycotoxins in food including the most frequent sample treatments [12, 27–30].

However, IACs are expensive and complex purification systems which suffer from low recoveries for some mycotoxins and their use in multiclass analysis is limited because of their high selectivity. As a consequence, simpler, more efficient, multiclass, and environmentally friendly extraction systems are demanded. Among the different proposals, the so-called QuEChERS (quick, easy, cheap, effective, rugged, and safe) and dispersive liquid-liquid microextraction (DLLME) are becoming increasingly popular treatments.

DLLME is based on the use of a ternary component solvent system; an appropriate mixture of a few microliters of an organic extraction solvent and a small volume of

TABLE 1: Most common mycotoxins, main producing fungal species, and food commodities frequently contaminated.

Fungal species	Mycotoxin	Food commodity	Maximum permitted levels ^a ($\mu\text{g Kg}^{-1}$)
<i>Aspergillus parasiticus</i>	Aflatoxins B1, B2, G1, G2	Maize, wheat, rice, sorghum, ground nuts, tree nuts, and figs	B1: 0.10–12 B1 + B2 + G1 + G2: 4–15
<i>Aspergillus flavus</i>	Aflatoxins B1 y B2	Idem	
Metabolite of aflatoxin B1 in mammals	Aflatoxin M1	Milk and milk products	0.025–0.050
<i>Fusarium sporotrichioides</i>	Toxins T-2 and HT-2	Cereals and cereal products	T-2 + HT-2: 15–2000 ^b
<i>Fusarium graminearum</i>	Deoxynivalenol zearalenone	Cereals and cereal products	DON: 200–1750 ZEN: 20–400
<i>Fusarium moniliforme</i> (<i>F. verticillioides</i>)	Fumonisin B1, B2	Maize, maize products, sorghum, and asparagus	B1 + B2: 200–4000
<i>Penicillium verrucosum</i> <i>Aspergillus ochraceus</i>	Ochratoxin A	Cereals, wine, fruits, coffee, and spices	0.50–80
<i>Penicillium expansum</i>	Patulin	Apples, apple juice, and apple products	25–50
<i>Aspergillus</i> , <i>Penicillium</i> , and <i>Monascus</i>	Citrinin	Cereals, red rice, fruits, and cheese	nr
<i>Aspergillus versicolor</i>	Sterigmatocystin	Cereals, coffee, ham, pepper, and cheese	nr
Hypocreales (<i>Claviceps purpurea</i>), Eurotiales	Ergot alkaloids	Cereals	nr

^aRange of maximum permitted levels in the EU, depending on the food commodity [6].

^bRecommended level [7].

nr: not regulated.

a disperser solvent (miscible with the extraction solvent and with water) is rapidly injected into an aqueous medium, with the result of a stable emulsion. The organic analytes present in the aqueous medium rapidly migrate to the extraction solvent, because of the large contact surface between the organic and the aqueous phases. After phases separation, the organic phase with the analytes of interest is collected and analysed by an appropriate technique [31–33]. DLLME has been applied for the determination of OTA in wine by HPLC-MS [34], cereals by HPLC-FL [35], and patulin in apple juices by CE-UV [36]. Also, we have developed two DLLME methods (one of them using an ionic liquid as extraction solvent) for the determination of OTA in wine by capillary-HPLC with laser induced fluorescence detection (LIF) with excellent results [18, 19].

On the other hand, QuEChERS is a fast and inexpensive method widely used in the last years, mainly for the extraction of pesticides and presents some advantages such as its simplicity, minimum steps, and effectiveness for cleaning-up complex samples [37, 38]. It comprises two steps: (i) an extraction based on partitioning via salting-out, involving the equilibrium between an aqueous and an organic layer; (ii) a dispersive SPE (dSPE) for further clean-up using combinations of MgSO_4 and different sorbents, such as C_{18} or primary and secondary amine (PSA). QuEChERS-based methods have been recently reported for the extraction of different mycotoxins in cereal products [39–41], bread [42], eggs [43], or spices [44] and in the multiresidue extraction of different contaminants (including mycotoxins) in foods [45, 46]. Also, we proposed this methodology for the determination of OTA in wine samples by capillary HPLC-LIF [18].

3. New Proposals for Determination of Mycotoxins in Different Foods

Considering the above described advances in sample treatments, we have proposed a multiclass method for the determination of mycotoxins in different food commodities. Taking advantage of UHPLC-MS/MS characteristics, we optimised a separation method that allows the determination of 15 mycotoxins in only four minutes. The studied mycotoxins are included in Regulation (EC) number 1881/2006 or considered as dangerous by the IARC [6, 47].

Moreover, in order to propose alternative methods for multiclass mycotoxins determination in scarcely investigated matrices and considering previous results obtained for the determination of OTA, we have tried to explore the advantages of the above mentioned sample treatments (DLLME and QuEChERS) as green, easy, and simple alternatives to other well-established methodologies, as IAC or SPE.

Below, we will explain the UHPLC-MS/MS conditions (common to all the methods) and then we will focus on the studied samples: cereals and pseudocereals, cereal syrups, edible nuts and seeds, and milk thistle. In all cases, a validation was performed in order to assess the compliance with the current requirements for mycotoxin determination in foods [8]. The validation included matrix effect study, establishment of matrix-matched calibrations, limits of detection (LODs) and quantification (LOQs), and intraday and intermediate precision. Moreover, recovery studies at three different concentration levels were carried out by comparison of the signal obtained for a sample spiked with a known concentration of mycotoxins before the sample treatment with the signal of

FIGURE 2: Sample treatment for the determination of mycotoxins in cereals and pseudocereals (MeCN: acetonitrile; MeOH: methanol).

an spiked extract obtained after the sample treatment. As a summary, the most significant analytical characteristics of the developed methods are shown in Table 2.

3.1. Chromatographic and MS Conditions. UHPLC separations were performed on an Agilent 1290 Infinity LC under the conditions summarised in Table 2.

The triple quadrupole mass spectrometer API 3200 (AB Sciex) worked with electrospray ionization in positive mode (ESI+), under multiple reaction monitoring (MRM) conditions shown in Table 3. The ionization source parameters were source temperature 500°C; curtain gas (nitrogen) 30 psi; ion spray voltage 5000 V; GAS 1 and GAS 2 (both of them nitrogen) 50 psi. Moreover, a precursor ion and two product ions (the most abundant for quantification and the other one for confirmation) were selected, obtaining four identification points, fulfilling the requirements established by European Union (EU) for confirmation of contaminants in foodstuff [48].

3.2. Analysis of Cereals and Pseudocereals. Cereals are a commodity of great interest, highly prone to microbial contamination because of their chemical composition. Rice is one of the most consumed cereals in the world. Moreover, brown rice and red rice (obtained by the fermentation of rice with *Monascus* fungi [49], that can produce CIT [50]) are increasingly chosen by customers because of their health benefits. Other cereal of interest is spelt. Its nutritional properties, high resistance in unfavourable environments, and low fertilization requirements made it increasingly valuable for food product manufacturers and consumers [51]. Matrices of concern are also pseudocereals, such as amaranth, quinoa, and buckwheat. Though botanically they are not true cereal grains, they produce starch-rich seeds consumed like cereals. Pseudocereals are also susceptible to fungal growth and

therefore to mycotoxin contamination. However, this issue has received little attention in literature.

Taking into account the interest and the scarce data about the determination of mycotoxins in some of the above mentioned matrices, we developed and validated an analytical method for the simultaneous identification and quantification of 15 mycotoxins (AFB_1 , AFB_2 , AFG_1 , AFG_2 , OTA, FB_1 , FB_2 , T-2, HT-2, CIT, STE, F-X, NIV, DON, and ZEN) in pseudocereals, spelt, and white, red, and brown rice. As a sample treatment we proposed a simple salting out assisted solid-liquid extraction (i.e., a QuEChERS-based extraction, see Figure 2). No further clean-up was required, although matrix effect was higher than [20%] for some mycotoxins (aflatoxins, DON, and NIV). Thus matrix-matched calibration was applied. A typical chromatogram corresponding to a spiked white rice sample submitted to the proposed method is shown in Figure 3(a). This methodology has proved to be a suitable and efficient choice for multiclass mycotoxin determination in these matrices, with LOQs below the contents currently regulated. It provides good recoveries (between 60.0% and 103.5%) and precision (RSD lower than 12% in all cases), allows extraction time reduction, and is environmentally friendly. Among all the samples analysed, a red rice sample was positive for AFB_1 (Figure 4(a)). The result was confirmed by comparison with a standard method [52].

3.3. Analysis of Cereal Syrups. Cereal syrups are obtained by isolation of starch after wet milling of grains, hydrolysis, and further purification and are widely used in food and pharmaceutical industry. Mycotoxins may also be found in cereal syrups as result of using contaminated raw material, or because of contamination of the final manufactured product by microorganisms during storage [53]. However, there are very few methods for the determination of mycotoxins in these matrices.

TABLE 2: Analytical characteristics of proposed methods.

Analytes	Matrix	Sample treatment	UHPLC chromatographic conditions	LOQs ($\mu\text{g Kg}^{-1}$)	Recovery (%)	Intraday precision (%RSD, $n = 9$)	Intermediate precision (%RSD, $n = 15$)
AFB ₁ , AFB ₂ , AFG ₁ , AFG ₂ , OTA, FB ₁ , FB ₂ , T-2, HT-2, STE, CIT, DON, NIV, F-X, ZEN	Cereals and pseudocereals	QuEChERS-based extraction	Mobile phase: (A) H ₂ O with 0.3% formic acid and 5 mM ammonium formate (B) MeOH with 0.3% formic acid and 5 mM ammonium formate Gradient profile: 0 min: 5% B; 1 min: 50% B; 2 min: 72% B; 4 min: 80% B and 6 min: 90% B Column: Zorbax Eclipse Plus RRHD C ₁₈ (50 mm $\times$ 2.1 mm, 1.8 μm) Flow-rate: 0.4 mL min ⁻¹ Temperature: 35°C Injection volume: 5 μL	From 0.23 (AFG ₁) to 233 (NIV)	From 60.0 (NIV in red rice) to 103.5 (AFG ₂ in white rice)	From 1.3 (FB ₂) to 8.8 (ZEN)	From 6.4 (HT-2) to 11.9 (FB ₁ , FX)
OTA, FB ₁ , FB ₂ , T-2, HT-2, STE, CIT, DON, F-X, ZEN	Cereal syrups	QuEChERS-based extraction		From 0.45 (STE) to 75.2 (DON)	From 62.8 (STE in wheat syrup) to 100.6 (CIT in rice syrup)	From 1.3 (T-2, CIT) to 8.4 (DON)	From 2.6 (CIT) to 11.5 (OTA)
AFB ₁ , AFB ₂ , AFG ₁ , AFG ₂ , OTA, FB ₁ , FB ₂ , T-2, HT-2, STE, CIT, DON, F-X, ZEN	Edible nuts and seeds	QuEChERS-based extraction + DLLME		From 0.57 (OTA) to 150 (F-X)	From 61.7 (STE in almond) to 104.3 (T-2 in hazelnut)	From 0.6 (ZEN) to 8.9 (CIT)	From 6.0 (T-2) to 10.6 (DON)
AFB ₁ , AFB ₂ , AFG ₁ , AFG ₂ , OTA, FB ₁ , FB ₂ , T-2, HT-2, STE, CIT, DON, NIV, F-X, ZEN	Milk thistle	QuEChERS-based extraction + DLLME		From 1.50 (AFG ₁) to 1530 (NIV)	From 60.0 (CIT in extract) to 98.9 (AFB ₂ in extract)	From 4.4 (OTA) to 9.3 (AFG ₂)	From 5.3 (FB ₁) to 9.9 (T2)

TABLE 3: Monitored ions of target analytes and MS/MS parameters.

Analyte	Retention time (min)	Precursor ion (m/z)	Molecular ion	DP ^a	EP ^a	CEP ^a	Product ions ^b	CEn ^a	CXP ^a
NIV	1.36	313.1	$[M + H]^+$	41.0	5.0	14.0	174.9 (Q) 128.0 (I)	17.0 75.0	4.0 6.0
DON	1.70	297.1	$[M + H]^+$	36.0	5.5	16.0	249.2 (Q) 161.0 (I)	17.0 29.0	4.0 4.0
F-X	2.02	355.1	$[M + H]^+$	26.0	12.0	18.0	174.7 (Q) 137.1 (I)	23.0 31.0	4.0 4.0
AFG ₂	2.52	331.1	$[M + H]^+$	61.0	6.0	42.0	245.1 (Q) 313.1 (I)	39.0 27.0	4.0 6.0
AFG ₁	2.61	329.0	$[M + H]^+$	76.0	9.5	16.0	243.1 (Q) 311.1 (I)	39.0 29.0	6.0 6.0
AFB ₂	2.72	315.1	$[M + H]^+$	81.0	4.0	34.0	286.9 (Q) 259.0 (I)	33.0 39.0	6.0 8.0
AFB ₁	2.79	313.1	$[M + H]^+$	46.0	12.0	26.0	241.0 (Q) 284.9 (I)	41.0 39.0	4.0 4.0
CIT	2.90	251.2	$[M + H]^+$	26.0	11.0	18.0	233.0 (Q) 204.8 (I)	23.0 73.0	23.0 10.0
HT-2	3.18	442.0	$[M + NH_4]^+$	21.0	5.5	21.0	262.8 (Q) 215.4 (I)	22.0 19.0	8.0 4.0
FB ₁	3.24	722.2	$[M + H]^+$	71.0	10.0	30.0	334.2 (Q) 352.2 (I)	51.0 47.0	6.0 6.0
T-2	3.44	484.0	$[M + NH_4]^+$	21.0	10.0	22.0	215.0 (Q) 185.0 (I)	22.0 29.0	4.0 4.0
ZEN	3.71	319.0	$[M + H]^+$	26.0	8.0	20.0	282.9 (Q) 301.0 (I)	19.0 15.0	4.0 10.0
OTA	3.76	404.0	$[M + H]^+$	41.0	7.5	16.0	238.9 (Q) 102.1 (I)	31.0 91.0	6.0 6.0
FB ₂	3.86	706.2	$[M + H]^+$	71.0	10.5	20.0	336.3 (Q) 318.3 (I)	43.0 45.0	14.0 12.0
STE	3.88	325.1	$[M + H]^+$	66.0	3.5	26.0	281.0 (Q) 310.0 (I)	43.0 37.0	4.0 4.0

^a(DP) Declustering potential, (EP) entrance potential, (CEP) collision cell entrance potential, (CXP) collision cell exit potential, and (CEn) collision energy. All expressed in voltage.

^bProduct ions: (Q) transition used for quantification and (I) transition employed to confirm the identification.

FIGURE 3: (a) Chromatogram of a spiked white rice sample; (b) chromatogram of a spiked barley syrup sample applying the proposed QuEChERS-UHPLC-MS/MS methodology. (Aflatoxins, OTA and STE: $25 \mu\text{g kg}^{-1}$; CIT: $50 \mu\text{g kg}^{-1}$; FB₁, FB₂, T-2, HT-2, and ZEN: $250 \mu\text{g kg}^{-1}$; DON: $1000 \mu\text{g kg}^{-1}$; F-X, NIV: $2500 \mu\text{g kg}^{-1}$). 1: NIV; 2: DON; 3: F-X; 4: AFG₂; 5: AFG₁; 6: AFB₂; 7: AFB₁; 8: CIT; 9: HT-2; 10: FB₁; 11: T-2; 12: ZEN; 13: OTA; 14: FB₂; 15: STE.

FIGURE 4: Extracted ion chromatogram of positive samples: (a) red rice ($8.3 \mu\text{g kg}^{-1}$ of AFB₁); (b) sunflower seed sample ($3.7 \mu\text{g kg}^{-1}$ of STE); (c) walnut sample ($222 \mu\text{g kg}^{-1}$ of ZEN and $346 \mu\text{g kg}^{-1}$ of DON); (d) macadamia nut sample ($2466 \mu\text{g kg}^{-1}$ of F-X); (e) milk thistle ($827 \mu\text{g kg}^{-1}$ of T-2 and $944 \mu\text{g kg}^{-1}$ of HT-2).

Trying to fill this gap, we modified the sample treatment previously described for cereals and pseudocereals (Figure 2) to make it suitable for the determination of 10 mycotoxins (OTA, FB₁, FB₂, T-2, HT-2, CIT, STE, F-X, ZEN, and DON) in wheat, barley, and rice syrup. Unfortunately, very low recoveries were obtained for aflatoxins and NIV and their quantification was impossible in these matrices. Good recoveries were obtained (between 62.8% and 100.6%), with RSD lower than 11.5% [54]. A chromatogram corresponding to a spiked barley syrup sample submitted to the proposed method is shown in Figure 3(b). It must be pointed out

that there is no specific legislation for this kind of matrices, although the low LOQs obtained allowed the determination of these mycotoxins at concentrations lower than the maximum contents usually established by current legislation in different foodstuffs.

3.4. Analysis of Edible Nuts and Seeds. It is well known that nuts and seeds are susceptible to mould growth and consequently to mycotoxin contamination [55–57]. Insect feeding-damage is the principal factor leading to preharvest fungal infection and subsequent mycotoxin contamination, but

FIGURE 5: Sample treatment for the determination of mycotoxins in nuts and seeds and milk thistle (MeCN: acetonitrile; MeOH: methanol).

infection may also occur after harvesting and storage. Current EU food safety legislation only regulates the content of aflatoxins in these matrices, with maximum permitted levels which depend on the kind of nut or seed for direct human consumption [6, 58].

Considering the good results obtained with the previously described matrices, the same approach was attempted. In this case, the QuEChERS-based extraction allowed the determination of OTA, T-2, HT-2, STE, CIT, ZEN, FB₁, FB₂, DON, and F-X. However, a further purification of the extracts was required for subsequent determination of aflatoxins. Thus, an additional clean-up step based on DLLME was proposed in order to reduce matrix effect, allowing the determination of aflatoxins. A flow chart of the whole procedure is shown in Figure 5. This methodology was applied for the determination of 14 mycotoxins in different nuts and seeds (almonds, peanuts, sunflower seeds, pumpkin seeds, walnuts, macadamia nuts, pistachios, hazelnuts, and pine nuts). The low LOQs obtained for aflatoxins, the only mycotoxins regulated in nut and seed matrices, allowed their quantification at concentrations lower than the maximum level established by current legislation. RSD was lower than 11% and recoveries ranged from 60.7% to 104.3%. Among the samples analysed, a sunflower seed sample showed a high content of STE (Figure 4(b)), a walnut sample of ZEN and

DON (Figure 4(c)), and a macadamia nut sample of F-X (Figure 4(d)) [59].

3.5. Analysis of Herbal Products: Milk Thistle. The consumption of products with specific nutritional and/or functional characteristics has significantly increased during the last decade. Among them, there are many food supplements containing herbal products and/or their derivatives as ingredients. Previous studies have demonstrated that these materials can suffer from fungi and mycotoxin contamination [60, 61]. However, maximum levels for mycotoxins in food supplements are not established; only, the European Pharmacopoeia sets a maximum level for AFB₁ ($2 \mu\text{g kg}^{-1}$) and for the sum of AFB₁, AFB₂, AFG₁, and AFG₂ ($4 \mu\text{g kg}^{-1}$) for herbal products used as drug ingredients [62].

Most of the scarce methods proposed for the determination of mycotoxins in herbal products use SLE and IAC for clean-up, including the method recommended by the Pharmacopoeia for the determination of AFB₁ [62].

In this context, we proposed a method for the multiclass determination of mycotoxins in milk thistle (*Silybum marianum*), a botanical consumed as food supplement because of its protective effects on the liver. Moreover, although analytical methods for studying the occurrence of mycotoxins in other herbal products (as tea, ginseng, or ginger) have been

previously reported, milk thistle has been scarcely studied and only for aflatoxin content [63].

The sample treatment optimized for this matrix is similar to that proposed for nuts and seeds, with some modifications (Figure 5). After QuEChERS-based extraction, FB₁, FB₂, NIV, DON, and F-X were quantified. However, a second clean-up step based on DLLME was needed for the determination of AFB₁, AFB₂, AFG₁, AFG₂, OTA, T-2, HT-2, STE, CIT, and ZEN. The method allowed the quantification of aflatoxins at concentrations lower than their maximum level established in botanicals by Pharmacopoeia. The rest of mycotoxins were determined at concentrations lower than their usual established limits in different foodstuff. Good recoveries were obtained (between 62.3% and 98.9%, except for ZEN in seed samples and CIT in extract). Among the different commercial samples of milk thistle (seeds and natural extract), two were contaminated with T-2 and HT-2 (Figure 4(e)) and ZEN was detected in one of them [64].

4. Future Trends

Currently, mycotoxins analysis presents several challenges that still need to be addressed and overcome. On the first place, carryover of mycotoxins from contaminated feeds to animal tissues and biological fluids and eventually to products intended for human consumption (meat, milk, and eggs) is a matter of concern. In some animals, mycotoxins undergo metabolic processes and are transformed into other compounds with different toxicity. For instance, most mammals metabolize AFB₁ into AFM₁ that is transferred to the milk, while poultries metabolize AFB₁ into toxic hydroxylated metabolites that may pass to eggs [65]. Animal feed is the first link in food chain and, therefore, the production of safe food depends not only on the manufacturers compliance with current legislation, but also on the use of safe feed by the farmers. As a solution, the use of detoxifying agents (a new group of feed additives) has been proposed in order to decrease the effect of mycotoxin contamination in feeds. These products are mainly adsorbent agents (i.e., activated charcoal, clays, silicates, and some synthetic polymers) that reduce the absorption of mycotoxins in the gastrointestinal tract of animal. Also, some enzymes and microorganisms are used as agents capable of transforming the mycotoxins by modification of their structure, although their use is more limited [66]. However, because of the different properties of mycotoxins, an adsorbent may be effective against a mycotoxin while ineffective against others. Recently, some published studies deal with various natural extracts, such as derivatives of honey [67] and organosulfur compounds derived from allium, like garlic [68], which could be natural alternatives to the usual binders, opening a very interesting field of research.

Another issue of great importance is the evaluation of occurrence of the so-called “masked mycotoxins” in food and feed. The metabolism of some plants (which have natural detoxification mechanisms) can generate conjugated compounds (masked mycotoxins), with different chemical behaviors than the mycotoxins of origin. Thus, some plants are able to transform the relatively nonpolar trichothecenes and

ZEN into more polar derivatives by conjugation with sugars, amino acids, or sulfate groups, which are then isolated into the vacuoles. However, these forms can be hydrolyzed to their precursors in the animal digestive tract, thus showing similar toxicity than free mycotoxins. Although this phenomenon has been studied mainly on *Fusarium* toxins (trichothecenes, zearalenone, and fumonisins), it has also been described for other mycotoxins. In addition, some technological food processings play an important role in the mechanisms of masking, mainly in cereal products, as they can induce reactions with macromolecules such as sugars, proteins, or lipids and inversely release the native forms of mycotoxins by decomposition of masked derivatives. Nowadays, toxicological data on masked mycotoxins are scarce, although several studies highlight the potential threat of these compounds for consumer safety. In particular, the possible hydrolysis of masked mycotoxin (generating the initial mycotoxin) during mammalian digestion would be a risk factor to be considered. For instance, products with an apparent low mycotoxin contamination have induced toxic effects due to the presence of “occult” fumonisins liberated upon hydrolysis and not detected in routine analysis. In this way, masked mycotoxins can quantitatively contribute to the total amount of mycotoxins, especially in cereals [69, 70]. Consequently, the development of analytical methods for multiclass analysis of mycotoxins including their transformation products is a challenge, in order to assess their real risk to the health of consumers.

Finally, the study of the so-called emerging mycotoxins derived from *Fusarium* fungi (as fusaproliferin, moniliformin, beauvericin, and enniatins), present prevalently in foods from northern Europe and Mediterranean countries, must also be highlighted. Unlike other better studied mycotoxins, permitted maximum levels have not yet been established for these mycotoxins. This is mainly because of the scarce data related to their presence in food, level of contamination, and toxicity. Although no cases of mycotoxicosis by the intake of these mycotoxins have been described, some studies (most *in vitro*) revealed the possible toxicity of these compounds, which could be increased as a result of the interaction of several mycotoxins present in food [71, 72]. For this reason, more studies are mandatory in order to evaluate the risk of these emerging toxins. Once again, reliable analytical methods are urgently required.

5. Conclusions

Alternative UHPLC-MS/MS analytical methods for multiclass determination of mycotoxins based on QuEChERS and DLLME for sample treatment have been developed. The proposed methods have been evaluated in diverse food commodities, most of them scarcely investigated. They showed as general advantages their efficacy, simplicity, versatility, and accuracy, as well as their low impact on the environment, shorter analysis time, and the relatively low-cost, compared with conventional IAC. Thus they fulfill the current requirements of analytical methods for the determination of contaminants. However, there are some aspects concerning mycotoxin determination that still are a challenge for

the scientific community, as the development of new analytical methods including the determination of masked or emerging mycotoxins. The proposed methodologies could be applied also to these analytes, opening new perspectives that, combined with powerful analytical techniques, such as UHPLC-MS/MS, offer interesting perspectives in this field.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] European Food Safety Authority (EFSA), 2012, <http://www.efsa.europa.eu/en/topics/topic/mycotoxins.htm>.
- [2] S. Marin, A. J. Ramos, G. Cano-Sancho, and V. Sanchis, "Mycotoxins: occurrence, toxicology, and exposure assessment," *Food and Chemical Toxicology*, vol. 60, pp. 218–237, 2013.
- [3] http://ec.europa.eu/food/safety/rasff/index_en.htm.
- [4] IARC, *Monographs on the Evaluation of Carcinogenic Risks to Humans*, vol. 56, IARC, Lyon, France, 1993.
- [5] A. L. Capriotti, G. Caruso, C. Cavaliere, P. Foglia, R. Samperi, and A. Laganà, "Multiclass mycotoxin analysis in food, environmental and biological matrices with chromatography/mass spectrometry," *Mass Spectrometry Reviews*, vol. 31, no. 4, pp. 466–503, 2012.
- [6] "Commission regulation (EC) No. 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs," *Official Journal of the European Union*, pp. L364/5–L364/24, 2006.
- [7] "Commission recommendation 2013/165/EU on the presence of T-2 and HT-2 toxin in cereals and cereal products," *Official Journal of the European Union L*, vol. 91, p. 12, 2013.
- [8] "Commission Regulation (EC) No. 401/2006 laying down the methods of sampling and analysis for the official control of the levels of mycotoxins in foodstuffs," *Official Journal of the European Union*, vol. L70, pp. 12–34, 2006.
- [9] FDA, "Molecular biology and natural toxins. Mycotoxins in domestic and imported foods," in *Compliance Program Guidance Manual*, chapter 7, Food and Drug Administration, Washington, DC, USA, 2008, <http://www.fda.gov/Food/ComplianceEnforcement/FoodCompliancePrograms/ucm071496.htm>.
- [10] G. S. Shephard, "Determination of mycotoxins in human foods," *Chemical Society Reviews*, vol. 37, pp. 2468–2477, 2008.
- [11] S. de Saeger, Ed., *Determining Mycotoxins and Mycotoxigenic Fungi in Food and Feed*, Woodhead, Cambridge, UK, 2011.
- [12] G. S. Shephard, F. Berthiller, P. A. Burdaspal et al., "Developments in mycotoxin analysis: an update for 2011–2012," *World Mycotoxin Journal*, vol. 6, no. 1, pp. 3–30, 2013.
- [13] D. Heperkan, F. K. Güler, and H. I. Oktay, "Mycoflora and natural occurrence of aflatoxin, cyclopiazonic acid, fumonisin and ochratoxin A in dried figs," *Food Additives and Contaminants*, vol. 29, pp. 277–286, 2012.
- [14] J. S. Dos Santos, C. R. Takabayashi, E. Y. S. Ono et al., "Immunoassay based on monoclonal antibodies versus LC-MS: deoxynivalenol in wheat and flour in Southern Brazil," *Food Additives & Contaminants A*, vol. 28, no. 8, pp. 1083–1090, 2011.
- [15] S. C. Cunha and J. O. Fernandes, "Development and validation of a method based on a QuEChERS procedure and heart-cutting GC-MS for determination of five mycotoxins in cereal products," *Journal of Separation Science*, vol. 33, no. 4–5, pp. 600–609, 2010.
- [16] N. Arroyo-Manzanares, L. Gámiz-Gracia, A. M. Gámiz-Gracia, J. J. Soto-Chinchilla, A. M. García-Campaña, and L. E. García-Ayuso, "On-line preconcentration for the determination of aflatoxins in rice samples by micellar electrokinetic capillary chromatography with laser-induced fluorescence detection," *Electrophoresis*, vol. 31, no. 13, pp. 2180–2185, 2010.
- [17] L. Campone, A. L. Piccinelli, R. Celano, and L. Rastrelli, "Application of dispersive liquid-liquid microextraction for the determination of aflatoxins B1, B2, G1 and G2 in cereal products," *Journal of Chromatography A*, vol. 1218, no. 42, pp. 7648–7654, 2011.
- [18] N. Arroyo-Manzanares, A. M. García-Campaña, and L. Gámiz-Gracia, "Comparison of different sample treatments for the analysis of ochratoxin A in wine by capillary HPLC with laser-induced fluorescence detection," *Analytical and Bioanalytical Chemistry*, vol. 401, p. 2987, 2011.
- [19] N. Arroyo-Manzanares, L. Gámiz-Gracia, and A. M. García-Campaña, "Determination of ochratoxin A in wines by capillary liquid chromatography with laser induced fluorescence detection using dispersive liquid-liquid microextraction," *Food Chemistry*, vol. 135, no. 2, pp. 368–372, 2012.
- [20] J. Rubert, C. Soler, and J. Mañes, "Application of an HPLC-MS/MS method for mycotoxin analysis in commercial baby foods," *Food Chemistry*, vol. 133, pp. 176–183, 2012.
- [21] I. Sospedra, J. Blesa, J. M. Soriano, and J. Mañes, "Use of the modified quick easy cheap effective rugged and safe sample preparation approach for the simultaneous analysis of type A- and B-trichothecenes in wheat flour," *Journal of Chromatography A*, vol. 1217, no. 9, pp. 1437–1440, 2010.
- [22] P. Li, Z. Zhang, X. Hu, and Q. Zhang, "Advanced hyphenated chromatographic-mass spectrometry in mycotoxin determination: Current status and prospects," *Mass Spectrometry Reviews*, vol. 32, no. 6, pp. 420–452, 2013.
- [23] M. Zachariasova, O. Lacina, A. Malachova et al., "Novel approaches in analysis of Fusarium mycotoxins in cereals employing ultra performance liquid chromatography coupled with high resolution mass spectrometry," *Analytica Chimica Acta*, vol. 662, no. 1, pp. 51–61, 2010.
- [24] M. M. Aguilera-Luiz, P. Plaza-Bolaños, R. Romero-González, J. L. Martínez-Vidal, and A. Garrido-Frenich, "Comparison of the efficiency of different extraction methods for the simultaneous determination of mycotoxins and pesticides in milk samples by ultra high-performance liquid chromatography-tandem mass spectrometry," *Analytical and Bioanalytical Chemistry*, vol. 399, no. 8, pp. 2863–2875, 2011.
- [25] P. Pérez-Ortega, B. Gilbert-López, J. F. García-Reyes, and A. Molina-Díaz, "Generic sample treatment method for simultaneous determination of multiclass pesticides and mycotoxins in wines by liquid chromatography-mass spectrometry," *Journal of Chromatography A*, vol. 1249, pp. 32–40, 2012.
- [26] J. O'Mahony, L. Clarke, M. Whelan et al., "The use of ultra-high pressure liquid chromatography with tandem mass spectrometric detection in the analysis of agrochemical residues and mycotoxins in food—challenges and applications," *Journal of Chromatography A*, vol. 1292, pp. 83–95, 2013.
- [27] J. P. Meneely, F. Ricci, H. P. van Egmond, and C. T. Elliott, "Current methods of analysis for the determination of trichothecene

- mycotoxins in food," *TrAC—Trends in Analytical Chemistry*, vol. 30, no. 2, pp. 192–203, 2011.
- [28] A. Veršilovskis and S. de Saeger, "Sterigmatocystin: occurrence in foodstuffs and analytical methods—an overview," *Molecular Nutrition & Food Research*, vol. 54, no. 1, pp. 136–147, 2010.
- [29] R. Köppen, M. Koch, D. Siegel, S. Merkel, R. Maul, and I. Nehls, "Determination of mycotoxins in foods: current state of analytical methods and limitations," *Applied Microbiology and Biotechnology*, vol. 86, no. 6, pp. 1595–1612, 2010.
- [30] E. Reiter, J. Zentek, and E. Razzazi, "Review on sample preparation strategies and methods used for the analysis of aflatoxins in food and feed," *Molecular Nutrition and Food Research*, vol. 53, no. 4, pp. 508–524, 2009.
- [31] A. Zgoła-Grzeskowiak and T. Grzeskowiak, "Dispersive liquid-liquid microextraction," *TrAC Trends in Analytical Chemistry*, vol. 30, no. 9, pp. 1382–1399, 2011.
- [32] V. Andruch, I. S. Balogh, L. Kocúrová, and J. Sandrejová, "Five years of dispersive liquid-liquid microextraction," *Applied Spectroscopy Reviews*, vol. 48, no. 3, pp. 161–259, 2013.
- [33] H. Yan, H. Wang, and J. Chromatogr, "Recent development and applications of dispersive liquid-liquid microextraction," *Journal of Chromatography A*, vol. 1295, pp. 1–15, 2013.
- [34] L. Campone, A. L. Piccinelli, and L. Rastrelli, "Dispersive liquid-liquid microextraction combined with high-performance liquid chromatography-tandem mass spectrometry for the identification and the accurate quantification by isotope dilution assay of Ochratoxin A in wine samples," *Analytical and Bioanalytical Chemistry*, vol. 399, no. 3, pp. 1279–1286, 2011.
- [35] L. Campone, A. L. Piccinelli, R. Celano, and L. Rastrelli, "PH-controlled dispersive liquid-liquid microextraction for the analysis of ionisable compounds in complex matrices: case study of ochratoxin A in cereals," *Analytica Chimica Acta*, vol. 754, pp. 61–66, 2012.
- [36] M. D. Víctor-Ortega, F. J. Lara, A. M. García-Campaña, M. del Olmo-Iruela, and A. M. García-Campaña, "Evaluation of dispersive liquid-liquid microextraction for the determination of patulin in apple juices using micellar electrokinetic capillary chromatography," *Food Control*, vol. 31, no. 2, pp. 353–358, 2013.
- [37] S. J. Lehotay, M. Anastassiades, and R. E. Majors, "QuEChERS, a sample preparation technique that is "catching on": an up-to-date interview with the inventors," *LC-GC North America*, vol. 28, no. 7, pp. 504–516, 2010.
- [38] M. Anastassiades, S. J. Lehotay, D. Stajnbaher, and F. J. Schenck, "Fast and easy multiresidue method employing acetonitrile extraction/partitioning and "dispersive solid-phase extraction" for the determination of pesticide residues in produce," *Journal of AOAC International*, vol. 86, no. 2, pp. 412–431, 2003.
- [39] A. Desmarchelier, J. M. Oberson, P. Tella, E. Gremaud, W. Seefelder, and P. Mottier, "Development and comparison of two multiresidue methods for the analysis of 17 mycotoxins in cereals by liquid chromatography electrospray ionization tandem mass spectrometry," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 13, pp. 7510–7519, 2010.
- [40] L. Vaclavik, M. Zachariasova, V. Hrbek, and J. Hajslova, "Analysis of multiple mycotoxins in cereals under ambient conditions using direct analysis in real time (DART) ionization coupled to high resolution mass spectrometry," *Talanta*, vol. 82, no. 5, pp. 1950–1957, 2010.
- [41] U. Koesukwiat, K. Sanguankaew, and N. Leepipatpiboon, "Evaluation of a modified QuEChERS method for analysis of mycotoxins in rice," *Food Chemistry*, vol. 153, pp. 44–51, 2014.
- [42] P. Paíga, S. Morais, T. Oliva-Teles et al., "Extraction of ochratoxin A in bread samples by the QuEChERS methodology," *Food Chemistry*, vol. 135, no. 4, pp. 2522–2528, 2012.
- [43] A. Garrido-Frenich, R. Romero-González, M. L. Gómez-Pérez, J. L. Martínez-Vidal, and J. Chromatogr, "Multi-mycotoxin analysis in eggs using a QuEChERS-based extraction procedure and ultra-high-pressure liquid chromatography coupled to triple quadrupole mass spectrometry," *Journal of Chromatography A*, vol. 1218, no. 28, pp. 4349–4356, 2011.
- [44] P. Yogendrarajah, C. van Poucke, B. de Meulenaer, and S. de Saeger, "Development and validation of a QuEChERS based liquid chromatography tandem mass spectrometry method for the determination of multiple mycotoxins in spices," *Journal of Chromatography A*, vol. 1297, pp. 1–11, 2013.
- [45] R. Romero-González, A. Garrido-Frenich, J. L. Martínez-Vidal, O. D. Prestes, and S. L. Grió, "Simultaneous determination of pesticides, biopesticides and mycotoxins in organic products applying a quick, easy, cheap, effective, rugged and safe extraction procedure and ultra-high performance liquid chromatography-tandem mass spectrometry," *Journal of Chromatography A*, vol. 1218, no. 11, pp. 1477–1485, 2011.
- [46] J. M. Zhang, Y. L. Wu, and Y. B. Lu, "Simultaneous determination of carbamate insecticides and mycotoxins in cereals by reversed phase liquid chromatography tandem mass spectrometry using a quick, easy, cheap, effective, rugged and safe extraction procedure," *Journal of Chromatography B*, vol. 915–916, pp. 13–20, 2013.
- [47] International Agency for Research on Cancer (IARC), <http://www.iarc.fr>.
- [48] "Commission Decision of 12 August 2002 implementing Council Directive 96/23/EC concerning the performance of analytical methods and the interpretation of results (2002/657/EC)," *Official Journal of the European Communities L*, vol. 221, p. 8, 2002.
- [49] T. Wang and T. Lin, "Monascus rice products," *Advances in Food and Nutrition Research*, vol. 53, pp. 123–159, 2007.
- [50] N. I. Samsudin and N. Abdullah, "A preliminary survey on the occurrence of mycotoxigenic fungi and mycotoxins contaminating red rice at consumer level in Selangor, Malaysia," *Mycotoxin Research*, vol. 29, pp. 89–96, 2013.
- [51] G. Bonafaccia, V. Galli, R. Francisci, V. Mair, V. Skrabanja, and I. Kreft, "Characteristics of spelt wheat products and nutritional value of spelt wheat-based bread," *Food Chemistry*, vol. 68, no. 4, pp. 437–441, 2000.
- [52] N. Arroyo-Manzanares, J. F. Huertas-Pérez, A. M. García-Campaña, and L. Gámiz-Gracia, "Simple methodology for the determination of mycotoxins in pseudocereals, spelt and rice," *Food Control*, vol. 36, no. 1, pp. 94–101, 2014.
- [53] C. M. Hazel and S. Patel, "Influence of processing on trichothecene levels," *Toxicology Letters*, vol. 153, no. 1, pp. 51–59, 2004.
- [54] N. Arroyo-Manzanares, J. F. Huertas-Pérez, L. Gámiz-Gracia, and A. M. García-Campaña, *Food Chemistry*. In press.
- [55] A. C. Baquião, P. Zorzeta, T. A. Reisa, E. Assunção, S. Vergueiro, and B. Correa, "Mycoflora and mycotoxins in field samples of Brazil nuts," *Food Control*, vol. 28, no. 2, pp. 224–229, 2012.
- [56] C. N. Ezekiel, M. Sulyok, B. Warth, A. C. Odebo, and R. Krska, "Natural occurrence of mycotoxins in peanut cake from Nigeria," *Food Control*, vol. 27, no. 2, pp. 338–342, 2012.
- [57] J. Rubert, C. Soler, and J. Mañes, "Occurrence of fourteen mycotoxins in tiger-nuts," *Food Control*, vol. 25, no. 1, pp. 374–379, 2012.

- [58] “Commission Regulation (EU) No. 165/2010 amending Regulation (EC) No 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards aflatoxins,” *Official Journal of the European Union L*, vol. 50, p. 8, 2010.
- [59] N. Arroyo-Manzanares, J. F. Huertas-Pérez, L. Gámiz-Gracia, and A. M. García-Campaña, “A new approach in sample treatment combined with UHPLC-MS/MS for the determination of multiclass mycotoxins in edible nuts and seeds,” *Talanta*, vol. 115, pp. 61–67, 2013.
- [60] L. Santos, S. Marín, V. Sanchis, and A. J. Ramos, “Screening of mycotoxin multicontamination in medicinal and aromatic herbs sampled in Spain,” *Journal of the Science of Food and Agriculture*, vol. 89, pp. 1802–1807, 2009.
- [61] J. D. di Mavungu, S. Monbaliu, M.-L. Scippo et al., “LC-MS/MS multi-analyte method for mycotoxin determination in food supplements,” *Food Additives and Contaminants A*, vol. 26, no. 6, pp. 885–895, 2009.
- [62] European Pharmacopoeia 6.0, “01/2008:20818. Determination of aflatoxin B1 in herbal drugs,” pp. 256–257.
- [63] V. H. Tournasa, C. Sapp, and M. W. Trucksess, “Occurrence of aflatoxins in milk thistle herbal supplements,” *Food Additives & Contaminants: Part A*, vol. 29, no. 6, p. 994, 2012.
- [64] N. Arroyo-Manzanares, A. M. García-Campaña, and L. Gámiz-Gracia, “Multiclass mycotoxin analysis in *Silybum marianum* by ultra high performance liquid chromatography–tandem mass spectrometry using a procedure based on QuEChERS and dispersive liquid–liquid microextraction,” *Journal of Chromatography A*, vol. 1282, pp. 11–19, 2013.
- [65] L. Afsah-Hejri, S. Jinap, P. Hajeb, S. Radu, and S. Shakibazadeh, “A review on mycotoxins in food and feed: Malaysia case study,” *Comprehensive Reviews in Food Science and Food Safety*, vol. 12, no. 6, pp. 629–651, 2013.
- [66] G. Jard, T. Liboz, F. Mathieu, A. Guyonvarch, and A. Lebrihi, “Review of mycotoxin reduction in food and feed: from prevention in the field to detoxification by adsorption or transformation,” *Food Additives & Contaminants A*, vol. 28, no. 11, pp. 1590–1609, 2011.
- [67] C. Siddoo-Atwal and A. S. Atwal, “A possible role for honey bee products in the detoxification of mycotoxins,” *Acta Horticulturae*, vol. 963, pp. 237–245, 2012.
- [68] K. Mylona, *Fusarium species in grains: dry matter losses, mycotoxin contamination and control strategies using Ozone and chemical compounds [Ph.D. thesis]*, Cranfield University, 2012.
- [69] G. Galaverna, C. Dallsta, M. A. Mangia, A. Dossena, and R. Marchelli, “Masked mycotoxins: an emerging issue for food safety,” *Czech Journal of Food Sciences*, vol. 27, pp. S89–S92, 2009.
- [70] F. Berthiller, C. Crews, C. Dall’Asta et al., “Masked mycotoxins: a review,” *Molecular Nutrition and Food Research*, vol. 57, no. 1, pp. 165–186, 2013.
- [71] M. Jestoi, “Emerging *Fusarium*-mycotoxins fusaproliferin, beauvericin, enniatins, and moniliformin—a review,” *Critical Reviews in Food Science and Nutrition*, vol. 48, no. 1, pp. 21–49, 2008.
- [72] A. Santini, G. Meca, S. Uhlig, and A. Ritieni, “Fusaproliferin, beauvericin and enniatins: occurrence in food: a review,” *World Mycotoxin Journal*, vol. 5, no. 1, pp. 71–81, 2012.