SINGULAR NONLINEAR ELLIPTIC EQUATIONS IN $\mathbf{R}^N$

C. O. ALVES, J. V. GONCALVES* AND L. A. MAIA

ABSTRACT. This paper deals with existence, uniqueness and regularity of positive generalized solutions of singular nonlinear equations of the form $-\Delta u + a(x)u = h(x)u^{-\gamma}$ in $\mathbf{R}^N$ where a, h are given, not necessarily continuous functions, and γ is a positive number. We explore both situations where a, h are radial functions, with a being eventually identically zero, and cases where no symmetry is required from either a or h . Schauder's fixed point theorem, combined with penalty arguments, is exploited.

1. INTRODUCTION

This paper addresses existence, uniqueness and regularity questions on generalized solutions of the singular nonlinear elliptic problem

$$(*) \quad \begin{cases} -\Delta u + a(x)u = h(x)u^{-\gamma} & \text{in } \mathbf{R}^N \\ u > 0 & \text{in } \mathbf{R}^N \end{cases}$$

where a, h are nonnegative L_{loc}^∞ functions, $h \not\equiv 0$, (eventually we consider $a \equiv 0$), $\gamma > 0$ and $N \geq 3$. We point out that the search of positive solutions of the Dirichlet problem for the equation

$$-\Delta u + a(x)u = h(x)u^{-\gamma} \quad \text{in } \Omega$$

where Ω is a smooth bounded domain has deserved the attention of many authors. Nowosad [1] studied a related Hammerstein equation, namely

$$u(x) = \int_0^1 K(x, y)(u(x))^{-\gamma} dy,$$

1991 *Mathematics Subject Classification.* 35J60.

Key words and phrases. singular nonlinear elliptic equations, Schauder's fixed point theorem, existence, uniqueness, regularity, positive solutions.

* Partially supported by CNPq/Brasil.

Received: April 8, 1998.

where $\gamma = 1$, $\int_0^1 K(x, y) dy \geq \delta > 0$ and $K(x, y)$ is positive semidefinite. Nowosad's work was extended by Karlin and Nirenberg [2] where more general Hammerstein equations were considered including the case $\gamma > 0$ in the equation above. Crandall-Rabinowitz and Tartar [3] studied the Dirichlet problem

$$Lu = f(x, u) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega$$

where L is a linear second order elliptic operator and $f : \Omega \times (0, +\infty) \rightarrow \mathbf{R}$ is singular in the sense that $f(x, r) \rightarrow \infty$ as $r \rightarrow 0^+$. Examples such as $f(x, r) = r^{-\gamma}$ with $\gamma > 1$, $\gamma < 1$ or $\gamma = 1$ were covered.

There is by now an extensive literature on singular elliptic problems. With respect to the case of bounded domains $\Omega \subset \mathbf{R}^N$ we would like to further mention Gomes [4], Lazer and McKenna [5], Cac and Hernandez [8], Chen [9], Lair and Shaker [10], Shangbin [13] while for the case $\Omega = \mathbf{R}^N$ we recall Kuzano and Swanson [11], Lair and Shaker [12, 14]. This reference list is far from complete. In the earlier papers concerning $\Omega = \mathbf{R}^N$, $h(x)$ is assumed at least continuous and several techniques are developed such as the method of lower and upper solutions. In this paper we assume $h(x)$ only integrable and use the Schauder fixed point theorem and elliptic estimates. Singular equations appear in the theory of heat conduction in electrically conducting materials, (Fulks and Maybee [6]), in binary communications by signals (Nowosad [1]) and in the theory of pseudoplastic fluids (Nachman and Callegari [7]).

The following condition on a will be required in the first one of our main results stated below:

$$(a)_R \quad a(x) \geq a_0 \text{ for } |x| \geq R \text{ for some } a_0, R > 0.$$

In what follows we take $\gamma, \alpha \in (0, 1)$ and $h \in L^\theta \cap L^2$ where $\theta \equiv \frac{2}{2-(1-\gamma)}$.

Theorem 1. *Assume $(a)_R$. Then $(*)$ has a unique solution $u \in \mathcal{D}^{1,2} \cap W_{loc}^{2,p}$ where $1 < p < \infty$ with $\int a(x)u^2 < \infty$. If a, h are radial functions the solution is radial, as well, and in fact, $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Moreover if $a, h \in C_{loc}^{\alpha}$ then $u \in C_{loc}^{2,\alpha}$.*

In our second result we take $a \equiv 0$ and h radially symmetric that is, we study the problem

$$(*)_o \quad \begin{cases} -\Delta u = h(|x|)u^{-\gamma} & \text{in } \mathbf{R}^N \\ u > 0 & \text{in } \mathbf{R}^N. \end{cases}$$

This problem shall be treated by first perturbing the equation by a radially symmetric term, then using the earlier result in the case a, h are radial functions and finally taking limits.

Theorem 2. *Let $a \equiv 0$ and let h be radially symmetric. Then $(*)_o$ has a unique radially symmetric solution $u \in \mathcal{D}^{1,2} \cap W_{loc}^{2,p}$, $1 < p < \infty$ and $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Moreover, if $h \in C_{loc}^{\alpha}$ then $u \in C_{loc}^{2,\alpha}$.*

2. PRELIMINARIES

The main goal in this section is to prove theorem 1. For that purpose let $\epsilon > 0$ and consider the problem

$$(2.1) \quad \begin{cases} -\Delta u + a(x)u = \frac{h(x)}{(u+\epsilon)^\gamma} & \text{in } \mathbf{R}^N \\ u > 0 & \text{in } \mathbf{R}^N. \end{cases}$$

We are going to show by applying the Schauder fixed point theorem that (2.1) has a solution $u_\epsilon \in W_{loc}^{2,p}$, $1 < p < \infty$, and then by passing to the limit as $\epsilon \rightarrow 0$ we arrive at a solution of (*).

In order to deal with a first step namely, existence of a solution of (2.1), let $f \in L^2$ and consider the linear equation

$$(2.2) \quad -\Delta u + a(x)u = f(x) \text{ in } \mathbf{R}^N.$$

Recalling that the Hilbert space $\mathcal{D}^{1,2}$ is defined as the closure of C_0^∞ with respect to the gradient norm $\|\varphi\|_1^2 = \int |\nabla \varphi|^2$ we introduce the space

$$E \equiv \left\{ u \in \mathcal{D}^{1,2} \mid \int a u^2 < \infty \right\}$$

which endowed with the inner product and norm given respectively by

$$\langle u, v \rangle = \int (\nabla u \cdot \nabla v + a u v) \text{ and } \|u\|^2 = \langle u, u \rangle$$

is itself a Hilbert space. Under condition $(a)_R$ it follows that $u \in E$ iff $u \in W^{1,2}(\mathbf{R}^N)$.

Yet if $f \in L^2$ it follows by minimizing over E the energy functional associated with (2.2),

$$I(u) = \frac{1}{2} \|u\|^2 - \int f u$$

that (2.2) has a weak solution $u \in E$, that is,

$$\int (\nabla u \cdot \nabla \varphi + a u \varphi) = \int f(x) \varphi, \quad \varphi \in E.$$

The solution u is, in fact, unique. Letting $S : L^2 \rightarrow E$ be the solution operator associated to (2.2) that is $Sf = u$ for $f \in L^2$ it follows that S is linear and moreover

$$\|Sf\| \leq C \|f\|_2, \quad f \in L^2$$

for some $C > 0$. In addition, splitting u into $u^+ - u^-$ where $u^\pm$ are respectively the positive and negative parts of u , taking $\varphi = -u^-$ above and noticing that $u^- \in E$ we infer that

$$Sf \geq 0 \text{ whenever } f \geq 0.$$

Now let $u \in L^2$ with $u \geq 0$. Since

$$(2.3) \quad 0 \leq \frac{h(x)}{(u+\epsilon)^\gamma} \leq \frac{h(x)}{\epsilon^\gamma}$$

and $\frac{h(x)}{\epsilon^\gamma} \in L^2$ the operator

$$Tu \equiv S \left[\frac{h(x)}{(u + \epsilon)^\gamma} \right]$$

is continuous in L^2 , and as a matter of fact, letting $w \equiv T(0)$ we have

$$w = S \left[\frac{h(x)}{\epsilon^\gamma} \right].$$

Considering

$$K \equiv \left\{ v \in L^2 \mid 0 \leq v \leq w \text{ a.e. in } \mathbf{R}^N \right\}$$

we shall prove that the following result holds true.

Lemma 3. *The set $K \subset L^2$ is closed, convex and bounded and moreover $T(K) \subset K$ and $\overline{T(K)}$ is a compact subset of L^2 .*

Using lemma 3 and the Schauder fixed point theorem there is some $u_\epsilon \in K$ satisfying

$$u_\epsilon = S \left[\frac{h(x)}{(u_\epsilon + \epsilon)^\gamma} \right]$$

that is

$$\begin{cases} \int (\nabla u_\epsilon \nabla \varphi + a u_\epsilon \varphi) = \int \frac{h(x) \varphi}{(u_\epsilon + \epsilon)^\gamma}, & \varphi \in E \\ u_\epsilon \geq 0 \text{ a.e. in } \mathbf{R}^N, & u_\epsilon \in E. \end{cases}$$

Now since by (2.3)

$$\frac{h(x)}{(u_\epsilon + \epsilon)^\gamma} \in L_{loc}^\infty$$

it follows by the elliptic regularity theory that $u_\epsilon \in W_{loc}^{2,p}$, $1 < p < \infty$, and further if $B \subset \mathbf{R}^N$ is a ball, then

$$-\Delta u_\epsilon + a(x)u_\epsilon = \frac{h(x)}{(u_\epsilon + \epsilon)^\gamma} \text{ a.e. in } B.$$

In fact, it follows by the maximum principle that $u_\epsilon > 0$ in B and so

$$\begin{cases} -\Delta u_\epsilon + a(x)u_\epsilon = \frac{h(x)}{(u_\epsilon + \epsilon)^\gamma} \text{ a.e. in } \mathbf{R}^N \\ u_\epsilon > 0 \text{ in } \mathbf{R}^N. \end{cases}$$

On the other hand, if $f \in L_{rad}^2$ we get by minimizing the functional I above over the space

$$E_{rad} \equiv \left\{ u \in W_{rad}^{1,2} \mid \int a(r)u^2 < \infty \right\}$$

which endowed with the inner product and norm given above is also a Hilbert space, a weak solution $u \in E_{rad}$ of (2.2) that is

$$\int (\nabla u \nabla \varphi + a u \varphi) = \int f(x) \varphi, \quad \varphi \in E_{rad}.$$

The solution is also unique and as before the solution operator associated to (2.2), namely $S : L_{rad}^2 \rightarrow E_{rad}$ satisfies

$$\|Sf\| \leq C\|f\|_2$$

for $f \in L^2_{rad}$ and further

$$Sf \geq 0 \text{ whenever } f \geq 0.$$

Letting

$$K \equiv \left\{ v \in L^2_{rad} \mid 0 \leq v \leq w \text{ a.e. in } \mathbf{R}^N \right\}$$

we have a corresponding symmetric variant of lemma 3 and so there is some $u_\epsilon \in E_{rad}$ with

$$\int (\nabla u_\epsilon \nabla \varphi + a(r) u_\epsilon \varphi) = \int \frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} \varphi, \quad \varphi \in E_{rad}.$$

Proof of Lemma 3.

It is easy to show that K is convex, closed and bounded. So we will only show that $T(K) \subset K$ and $\overline{T(K)}$ is compact in L^2 . If $v \in K$ then

$$T(0) - T(v) = S \left[h \left(\frac{1}{\epsilon^\gamma} - \frac{1}{(v + \epsilon)^\gamma} \right) \right] \geq 0$$

that is $T(v) \leq w$ and hence $T(K) \subset K$.

In order to show that $\overline{T(K)} \subset L^2$ is compact let v_n be a sequence in $T(K)$ say $v_n = T(u_n)$ for some $u_n \in K$. By (2.3)

$$\frac{h(x)}{(u_n + \epsilon)^\gamma} \text{ is bounded in } L^2$$

so that

$$T(u_n) = S \left[\frac{h(x)}{(u_n + \epsilon)^\gamma} \right] \text{ is bounded in } E.$$

Thus, passing to subsequences,

$$T(u_n) \rightarrow v \text{ for some } v \in E$$

and

$$T(u_n) \rightarrow v \text{ a.e. in } \mathbf{R}^N.$$

On the other hand, since $0 \leq T(u_n) \leq w$ it follows by Lebesgue's theorem that

$$T(u_n) \rightarrow v \text{ in } L^2.$$

showing that $\overline{T(K)}$ is compact in L^2 , ending the proof of lemma 3. The radial case is handled similarly. ■

The next result states that the family u_ϵ increases when ϵ decreases.

Lemma 4. *If $0 < \epsilon < \epsilon'$ then $u_{\epsilon'} \leq u_\epsilon$ in $\mathbf{R}^N$.*

Proof of Lemma 4.

Letting $\omega \equiv u_{\epsilon'} - u_\epsilon$ we get

$$-\Delta \omega + a(x)\omega = h(x) \left[\frac{1}{(u_{\epsilon'} + \epsilon')^\gamma} - \frac{1}{(u_\epsilon + \epsilon)^\gamma} \right] \text{ a.e. in } \mathbf{R}^N$$

which gives

$$\int |\nabla \omega^+|^2 + a(x)\omega^{+2} = \int h(x) \left[\frac{1}{(u_{\epsilon'} + \epsilon')^\gamma} - \frac{1}{(u_\epsilon + \epsilon)^\gamma} \right] \omega^+ \leq 0$$

showing that $\omega^+ = 0$ and thus $u_{\epsilon'} \leq u_\epsilon$ a.e. in $\mathbf{R}^N$, finishing the proof of lemma 4. ■

3. PROOFS OF MAIN RESULTS

Proof of Theorem 1.

Step 1 (the non-symmetric case).

Let $\epsilon_n > 0$ be a decreasing sequence converging to 0 and set $u_n = u_{\epsilon_n}$. We claim that

$$\|u_n\| \text{ is bounded.}$$

Indeed,

$$(3.1) \quad \int (|\nabla u_n|^2 + a|u_n|^2) = \int \frac{h(x)u_n}{(u_n + \epsilon_n)^\gamma} \leq \int h(x)u_n^{1-\gamma} \leq C|h|_\theta \|u_n\|^{1-\gamma}$$

for some $C > 0$, showing that u_n is bounded in E . Hence, passing to subsequences, we have

$$u_n \rightharpoonup u \text{ in } E, \text{ and } u_n \rightarrow u \text{ a.e. in } \mathbf{R}^N.$$

Moreover since by lemma 4 $0 < u_1 \leq u_n$ in $\mathbf{R}^N$ we infer that if $\varphi \in E$ has compact support then $\text{supp}(\varphi) \subset B$ for some ball $B \subset \mathbf{R}^N$ and

$$\frac{|h(x)\varphi|}{(u_n + \epsilon_n)^\gamma} \leq H(x) \text{ for some } H \in L^1$$

which gives, by applying Lebesgue's theorem to

$$\int (\nabla u_n \nabla \varphi + a u_n \varphi) = \int \frac{h(x)\varphi}{(u_n + \epsilon_n)^\gamma}$$

that

$$\begin{cases} \int (\nabla u \nabla \varphi + a u \varphi) = \int \frac{h(x)\varphi}{u^\gamma} \\ u \geq u_1 > 0 \text{ in } \mathbf{R}^N. \end{cases}$$

Using the regularity theory again we arrive at

$$\begin{cases} -\Delta u + a(x)u = h(x)u^{-\gamma} \text{ a.e. in } \mathbf{R}^N \\ u \in W_{loc}^{2,p}, \quad 1 < p < \infty \\ u > 0 \text{ in } \mathbf{R}^N. \end{cases}$$

In order to prove uniqueness let $M \in C_0^\infty$ such that

$$M(x) = 1 \text{ if } |x| \leq 1, \quad M(x) = 0 \text{ if } |x| \geq 2 \text{ and } 0 \leq M \leq 1.$$

Given $\varphi \in E$, an integer $j \geq 1$ and letting

$$\varphi^j(x) \equiv M\left(\frac{x}{j}\right)\varphi(x), \quad x \in \mathbf{R}^N$$

it follows that $\varphi^j \in E$ and $\text{supp}(\varphi^j)$ is compact. Moreover as we will show in the Appendix

$$(3.2) \quad \varphi^j \rightarrow \varphi \quad \text{in } E.$$

Now assume u, v are two solutions of $(*)$ and let $w_j \equiv u^j - v^j$. Then

$$\begin{aligned} \langle u - v, u^j - v^j \rangle &= \int (\nabla(u - v) \nabla w_j + a(x)(u - v)w_j) \\ &= \int h(x) \left(\frac{1}{u^\gamma} - \frac{1}{v^\gamma} \right) w_j. \end{aligned}$$

Assuming, by contradiction, that $u \neq v$ and once

$$\langle u - v, u^j - v^j \rangle \rightarrow \|u - v\|^2$$

we have

$$\int h(x) \left(\frac{1}{u^\gamma} - \frac{1}{v^\gamma} \right) w_j > 0$$

for large values of j . On the other hand,

$$\int h(x) \left(\frac{1}{u^\gamma} - \frac{1}{v^\gamma} \right) w_j \leq \int_{\Omega_j} h(x) u^{1-\gamma} + \int_{\Omega_j} h(x) v^{1-\gamma}$$

where $\Omega_j \equiv B_{2j} \setminus B_j$. Therefore, passing to the limit as $j \rightarrow \infty$ and noticing that the two integrals in the right hand side tend to zero we get a contradiction, that is $u = v$.

Assume now, $h \in C_{loc}^\alpha$. Then by the elliptic regularity theory more precisely, interior elliptic estimates, we get $u \in C_{loc}^{2,\alpha}$. This proves theorem 1 (in the case of Step 1).

Step 2 (the symmetric case: a, h are radial).

From section 2 we have found by Schauder's Theorem some radial function $u_\epsilon \in K$, $u_\epsilon \neq 0$ satisfying $u_\epsilon = Tu_\epsilon$, which means

$$(3.3) \quad \int (\nabla u_\epsilon \nabla v + a(r)u_\epsilon v) = \int \frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} v, \quad v \in E_{rad}.$$

We will show next that $u_\epsilon \in W_{loc}^{2,p}(\mathbf{R}^N \setminus \{0\})$ for $1 < p < \infty$, and

$$-\Delta u_\epsilon + a(r)u_\epsilon = \frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} \quad \text{a.e. in } \mathbf{R}^N \setminus \{0\}.$$

Indeed, changing variables we get from (3.3)

$$\int_S \int_0^\infty (u'_\epsilon v' + a(r)u_\epsilon v) r^{N-1} dr dS = \int_S \int_0^\infty \frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} v r^{N-1} dr dS$$

where $S \subset \mathbf{R}^N$ is the unit sphere. Making

$$v \equiv r^{-(N-1)} \psi, \quad r > 0, \quad \psi \in C_0^\infty(0, \infty)$$

we have

$$\int_0^\infty \left[\left(r^{(N-1)} u'_\epsilon \right) \left(r^{-(N-1)} \psi \right)' + a u_\epsilon \psi \right] dr = \int_0^\infty \frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} \psi(r) dr,$$

for $\psi \in C_0^\infty(0, \infty)$, and labelling

$$\frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} - a(r)u_\epsilon \equiv \widehat{H}(r), \quad r > 0$$

we get

$$-\frac{1}{r^{N-1}} (r^{(N-1)} u'_\epsilon)' = \widehat{H}(r) \quad \text{in } (0, \infty)$$

in the distribution sense. But since $a, h, u_\epsilon \in L_{loc}^p(0, \infty)$, $1 < p < \infty$ it follows that $\widehat{H} \in L_{loc}^p(0, \infty)$ and using the regularity theory we infer that $u_\epsilon \in W_{loc}^{2,p}(0, \infty)$ and

$$-\frac{1}{r^{N-1}} (r^{(N-1)} u'_\epsilon)' = \widehat{H}(r) \quad \text{a.e. in } (0, \infty).$$

By the maximum principle,

$$u_\epsilon > 0 \quad \text{in } (0, \infty).$$

Since $u_\epsilon \in W_{loc}^{2,p}(\mathbf{R}^N \setminus \{0\})$ and

$$-\Delta u_\epsilon = -\frac{1}{r^{N-1}} (r^{(N-1)} u'_\epsilon)'$$

we also have

$$-\Delta u_\epsilon + a(r)u_\epsilon = \frac{h(r)}{(u_\epsilon + \epsilon)^\gamma} \quad \text{a.e. in } \mathbf{R}^N \setminus \{0\}.$$

Now, let $\epsilon_n > 0$ such that $\epsilon_n \rightarrow 0$ and label $u_{\epsilon_n} \equiv u_n$. Following the proof of lemma 4 we have $u_n \geq u_1 > 0$. On the other hand we claim that

$$\|u_n\| \text{ is bounded.}$$

Indeed, as in (3.1) we have

$$\int \left(|\nabla u_n|^2 + a|u_n|^2 \right) \leq C \|h\|_\theta \|u_n\|^{1-\gamma}$$

so that u_n is bounded in E_{rad} . Passing to subsequences we have

$$u_n \rightharpoonup u \text{ in } E_{rad}, \text{ and } u_n \rightarrow u \text{ a.e. in } \mathbf{R}^N.$$

On the other hand, if $v \in E_{rad}$ has compact support then, as in section 1, applying Lebesgue's Theorem to

$$\int (\nabla u_n \nabla v + a(r)u_n v) = \int \frac{h(r)}{(u_n + \epsilon_n)^\gamma} v,$$

gives

$$\int (\nabla u \nabla v + a(r)uv) = \int \frac{h(r)}{u^\gamma} v.$$

Now changing variables, making again $v \equiv r^{-(N-1)}\psi$ where $r > 0$ and $\psi \in C_0^\infty(0, \infty)$ and arguing as above we obtain $u \in W_{loc}^{2,p}(\mathbf{R}^N \setminus \{0\})$ and

$$-\frac{1}{r^{N-1}}(r^{(N-1)}u')' + a(r)u = \frac{h(r)}{u^\gamma} \text{ a.e. in } (0, \infty)$$

and in addition,

$$-\Delta u + a(r)u = \frac{h(r)}{u^\gamma} \text{ a.e. in } \mathbf{R}^N \setminus \{0\}.$$

So, if $\varphi \in C_0^\infty(\mathbf{R}^N \setminus \{0\})$ then

$$\int (\nabla u \nabla \varphi + a(r)u\varphi) = \int \frac{h(r)}{u^\gamma} \varphi$$

that is

$$-\Delta u + a(r)u = \frac{h(r)}{u^\gamma} \text{ in } \mathbf{R}^N \setminus \{0\}$$

in the distribution sense. Next we show that $u \in W_{loc}^{2,p}(\mathbf{R}^N)$ and

$$\int (\nabla u \nabla \varphi + a(r)u\varphi) = \int \frac{h(r)}{u^\gamma} \varphi, \quad \varphi \in C_0^\infty(\mathbf{R}^N).$$

Indeed, let $\eta \in C^\infty(\mathbf{R}^N)$ such that

$$\eta(x) = 0 \text{ for } |x| \leq 1, \text{ and } \eta(x) = 1 \text{ for } |x| \geq 2$$

and let

$$\psi_\epsilon(x) \equiv \eta\left(\frac{x}{\epsilon}\right), \quad \epsilon > 0.$$

If $\varphi \in C_0^\infty(\mathbf{R}^N)$ then $\psi_\epsilon \varphi \in C_0^\infty(\mathbf{R}^N \setminus \{0\})$ and from above

$$\int (\nabla u \nabla (\psi_\epsilon \varphi) + a(r)u(\psi_\epsilon \varphi)) = \int \frac{h(r)}{u^\gamma} (\psi_\epsilon \varphi)$$

so that

$$\int (\psi_\epsilon \nabla u \nabla \varphi + \varphi \nabla u \nabla \psi_\epsilon + a(r)u\psi_\epsilon \varphi) = \int \frac{h(r)}{u^\gamma} \psi_\epsilon \varphi.$$

Making $\epsilon \rightarrow 0$ and using Lebesgues's Theorem we infer that

$$\int \psi_\epsilon \nabla u \nabla \varphi \rightarrow \int \nabla u \nabla \varphi,$$

$$\int a(r)u\psi_\epsilon \varphi \rightarrow \int a(r)u\varphi$$

and

$$\int \frac{h(r)}{u^\gamma} \psi_\epsilon \varphi \rightarrow \int \frac{h(r)}{u^\gamma} \varphi.$$

Claim.

$$\int \varphi \nabla u \nabla \psi_\epsilon \rightarrow 0.$$

Assuming the Claim has been proved we have

$$\int (\nabla u \nabla \varphi + a(r)u\varphi) = \int \frac{h(r)}{u^\gamma} \varphi$$

and since $a, h \in L_{loc}^\infty$ we get by the regularity theory that $u \in W_{loc}^{2,p}(\mathbf{R}^N)$ for $1 < p < \infty$ and

$$-\Delta u + a(r)u = \frac{h(r)}{u^\gamma} \text{ a.e. in } \mathbf{R}^N$$

and if in addition $a, h \in C_{loc}^\alpha$ then $u \in C_{loc}^{2,\alpha}$ by the interior Schauder estimates.

Verification of the Claim.

Using Schwarz inequality we have

$$\begin{aligned} \left| \int \varphi \nabla u \nabla \psi_\epsilon \right| &\leq |\varphi|_\infty \left(\int_{|x| \leq 2\epsilon} |\nabla u|^2 \right)^{\frac{1}{2}} \left(\int_{|x| \leq 2\epsilon} |\nabla \psi_\epsilon|^2 \right)^{\frac{1}{2}} \\ &\leq |\varphi|_\infty |\nabla \eta|_2 \left(\int_{|x| \leq 2\epsilon} |\nabla u|^2 \right)^{\frac{1}{2}} \epsilon^{\frac{N-2}{2}} \end{aligned}$$

where $N \geq 3$. Letting $\epsilon \rightarrow 0$ shows the Claim.

As for the uniqueness the argument in the proof of theorem 1 (Step 1) applies ending the proof of theorem 1 (in case of Step 2). The proof of theorem 1 is finished. ■

Proof of Theorem 2.

In order to solve $(*)_0$ we consider the family of problems

$$(3.4) \quad \begin{cases} -\Delta u + \frac{1}{k}u = h(|x|)u^{-\gamma} & \text{in } \mathbf{R}^N \\ u > 0 & \text{in } \mathbf{R}^N. \end{cases}$$

where $k \geq 1$ is an integer. Making $a(x) \equiv \frac{1}{k}$ in theorem 1 (radial case), it follows that (3.4) has a solution $u_k \in H_{rad}^1 \cap W_{loc}^{2,p}$, $1 < p < \infty$ satisfying

$$\int |\nabla u_k|^2 + \frac{1}{k}u_k^2 = \int h(r)u_k^{1-\gamma}.$$

Using both Hölder's inequality and the continuous embedding $D^{1,2} \rightarrow L^{2^*}$ in the equality above we infer that

$$(3.5) \quad \int |\nabla u_k|^2 \leq C_1 \text{ for some } C_1 > 0.$$

By a well known property of radial functions $u \in D^{1,2}$, namely

$$|u(x)| \leq \frac{C_2}{|x|^{\frac{N-2}{2}}} \|u\|_{D^{1,2}}, \quad x \neq 0 \text{ for some } C_2 > 0$$

we get

$$(3.6) \quad 0 \leq u_k(x) \leq \frac{C}{|x|^{\frac{N-2}{2}}}, \quad x \neq 0 \text{ for some } C > 0.$$

We shall need the following result which says that the sequence u_k increases with k .

Lemma 5. *If $k < k'$ then $u_k \leq u_{k'}$, a.e. in $\mathbf{R}^N$.*

By the boundedness of u_k in $D^{1,2}$ and lemma 5 there is some radial function $u \in D^{1,2}$ such that

$$u_k \rightharpoonup u \text{ in } D^{1,2}, \quad u_k \rightarrow u \text{ a.e. in } \mathbf{R}^N$$

and

$$u_1 \leq u_2 \leq \dots \leq u_k \leq \dots \leq u \text{ a.e. in } \mathbf{R}^N.$$

Now if $\varphi \in C_0^\infty(\mathbf{R}^N)$ then

$$(3.7) \quad \int \left(\nabla u_k \nabla \varphi + \frac{1}{k} u_k \varphi \right) = \int h u_k^{-\gamma} \varphi.$$

Let $\Omega \subset \mathbf{R}^N$ be a bounded domain such that $\text{supp}(\varphi) \subset \Omega$. Then

$$|h u_k^{-\gamma} \varphi| \leq h u_1^{-\gamma} |\varphi| \in L^p(\Omega), \quad 1 \leq p < \infty$$

and

$$\int h u_k^{-\gamma} \varphi \rightarrow \int h u^{-\gamma} \varphi.$$

On the other hand, using (3.6) we get

$$\frac{1}{k} \int u_k \varphi \rightarrow 0.$$

Passing to the limit in (3.7) gives

$$\int \nabla u \nabla \varphi = \int h u^{-\gamma} \varphi.$$

Since $0 < u_1 \leq u$ and $u_1 \in W_{loc}^{2,p}(\mathbf{R}^N)$ it follows that $h u^{-\gamma} \in L_{loc}^p(\mathbf{R}^N)$ and by the regularity theory $u \in W_{loc}^{2,p}(\mathbf{R}^N)$. In addition $u \in C_{loc}^{2,\alpha}$ when $h \in C_{loc}^\alpha$. This proves Theorem 2. ■

Proof of Lemma 5.

Letting $\omega = u_k - u_{k'}$ we have

$$\begin{aligned} \int |\nabla \omega^+|^2 + \frac{1}{k'} (\omega^+)^2 &\leq \int \nabla \omega \nabla \omega^+ + \frac{1}{k'} \omega \omega^+ \\ &\leq \int h \left(\frac{1}{u_k} - \frac{1}{u_{k'}} \right) \omega^+ \end{aligned}$$

showing that $\omega^+ = 0$ and so $\omega \leq 0$, ending the proof of lemma 5. ■

4. APPENDIX

Verification of (3.2).

Indeed,

$$a|\varphi^j - \varphi|^2 \leq 4a\varphi^2 \in L^1 \text{ and } a|\varphi^j - \varphi|^2 \rightarrow 0 \text{ a.e. in } \mathbf{R}^N$$

so that by Lebesgue's theorem

$$\int a|\varphi^j - \varphi|^2 \rightarrow 0.$$

Now

$$\frac{\partial \varphi^j}{\partial x_i} = \frac{1}{j} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi + M\left(\frac{x}{j}\right) \frac{\partial \varphi}{\partial x_i}.$$

Hence

$$\begin{aligned} \int \left| \frac{\partial \varphi^j}{\partial x_i} - \frac{\partial \varphi}{\partial x_i} \right|^2 &= \int \left| \frac{1}{j} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi + M\left(\frac{x}{j}\right) \frac{\partial \varphi}{\partial x_i} - \frac{\partial \varphi}{\partial x_i} \right|^2 \\ &\leq C \int \left| \frac{1}{j^2} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi \right|^2 + \int \left| M\left(\frac{x}{j}\right) \frac{\partial \varphi}{\partial x_i} - \frac{\partial \varphi}{\partial x_i} \right|^2. \end{aligned}$$

Arguing as above we infer that

$$M\left(\frac{x}{j}\right) \frac{\partial \varphi}{\partial x_i} \rightarrow \frac{\partial \varphi}{\partial x_i} \text{ in } L^2.$$

It remains to show that

$$\int \left| \frac{1}{j^2} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi \right|^2 \rightarrow 0.$$

At first we remark that

$$\begin{aligned} \int \left| \frac{1}{j^2} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi \right|^2 &= \int_{B_{2j} \setminus B_j} \left| \frac{1}{j^2} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi \right|^2 \\ &\leq \frac{C}{j^2} \int_{B_{2j} \setminus B_j} \varphi^2. \end{aligned}$$

Now using Hölder inequality with exponents $\frac{N}{N-2}$ and $\frac{N}{2}$ in the last integral we obtain

$$\begin{aligned} \int \left| \frac{1}{j^2} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi \right|^2 &\leq \frac{C}{j^2} \left(\int_{B_{2j} \setminus B_j} 1 dx \right)^{\frac{2}{N}} \left(\int_{B_{2j} \setminus B_j} |\varphi|^{2^*} \right)^{\frac{1}{2^*}} \\ &\leq \frac{C}{j^2} \left(\int_{B_{2j}} 1 dx \right)^{\frac{2}{N}} \left(\int_{B_{2j}^c} |\varphi|^{2^*} \right)^{\frac{1}{2^*}} \\ &\leq \frac{C \omega_N^{\frac{2}{N}} (2j)^2}{j^2} \left(\int_{B_{2j}^c} |\varphi|^{2^*} \right)^{\frac{1}{2^*}} \end{aligned}$$

where ω_N denotes the volume of the unit sphere of $\mathbf{R}^N$.

Next passing to the limit we get

$$\int \left| \frac{1}{j^2} \frac{\partial}{\partial x_i} M\left(\frac{x}{j}\right) \varphi \right|^2 \rightarrow 0.$$

This shows that $\varphi^j \rightarrow \varphi$ in E proving (3.2).

REFERENCES

- [1] P. Nowosad, *On the integral equation $Kf = \frac{1}{f}$ arising in a problem in communications*, J. Math. Anal. Appl. **14** (1966), 484–492.
- [2] S. Karlin and L. Nirenberg, *On a theorem of P. Nowosad*, J. Math. Anal. Appl. **17** (1967), 61–67.
- [3] M. Crandall, P. Rabinowitz and L. Tartar, *On a Dirichlet problem with singular nonlinearity*, Comm. Partial Differential Equations, **2** (1977), 193–222.
- [4] S. M. Gomes, *On a singular nonlinear elliptic problem*, SIAM J. Math. Anal. **17** (1986), 1359–1369.
- [5] A.C. Lazer and P. McKenna, *On a singular nonlinear elliptic boundary value problem*, Proc. Amer. Math. Soc. **111** (1991), 721–730.
- [6] W. Fulks and J. S. Maybee, *A singular nonlinear equation*, Osaka Math J. **12** (1960), 1–19.

- [7] A. Nachman and A. Callegari, *A nonlinear singular boundary value problem in the theory of pseudoplastic fluids*, SIAM J. Appl. Math. **28** (1986), 271–281.
- [8] N. P. Cac and G. Hernandez, *On a singular elliptic boundary value problem*, preprint.
- [9] H. Chen, *On a singular nonlinear elliptic equation*, Nonlinear Anal. **29** (1997), 337–345.
- [10] A. Lair and A. Shaker, *Uniqueness of solution to a singular quasilinear elliptic problem*, Nonlinear Anal. **28** (1997), 489–493.
- [11] T. Kusano and C. Swanson, *Entire positive solutions of singular semilinear elliptic equations*, Japan J. Math. **11** (1985), 145–156.
- [12] A. Lair and A. Shaker, *Classical and weak solutions of a singular semilinear elliptic problem*, J. Math. Anal. App. **211** (1997), 371–385.
- [13] C. Shangbin, *Positive solutions for Dirichlet problems associated to semilinear elliptic equations with singular nonlinearity*, Nonlinear Anal. **21** (1993), 181–190.
- [14] A. Shaker, *On singular semilinear elliptic equations*, J. Math. Anal. App. **173** (1993), 222–228.

C. O. ALVES

DEPARTAMENTO DE MATEMÁTICA E ESTATÍSTICA

UNIVERSIDADE FEDERAL DA PARAIBA

58109-970 CAMPINA GRANDE, PB BRAZIL

E-mail address: coalves@dme.ufpb.br

J. V. GONCALVES AND L. A. MAIA

DEPARTAMENTO DE MATEMÁTICA

UNIVERSIDADE DE BRASÍLIA

70910-900 BRASÍLIA DF, BRAZIL

E-mail addresses: jv@mat.unb.br, liliane@mat.unb.br