ON THE CONVERGENCE OF MIN/SUP POINTS IN OPTIMAL CONTROL PROBLEMS

ADIB BAGH

Received 26 March 2001

We modify the definition of lopsided convergence of bivariate functionals to obtain stability results for the min/sup points of some control problems. In particular, we develop a scheme of finite dimensional approximations to a large class of non-convex control problems.

1. Introduction

Let X be a topological space and consider the following problem:

Find $\bar{x}$ such that

$$f_0(\bar{x}) = \inf_{x \in X} f_0(x), \quad \text{subject to } x \in C \subset X. \quad (1.1)$$

In case f_0 and C are convex, a variety of primal/dual numerical methods can be used to find the saddle points of a Lagrangian L associated with this problem [4, 7]. These methods take advantage of the fact that the search for the saddle points of L is unconstrained, or conducted over sets much simpler than C . Furthermore, we can introduce penalties on f_0 in a way that regularizes L and makes it smoother. This in turn leads to a more convenient optimality condition of the form $(0, 0) \in \partial L$. In the case of a non-convex problem, similar methods can be used to find the saddle points of an augmented Lagrangian, and these points can be used to obtain a solution to the original problem (see [7, 8], and [11, Chapter 10, Sections I and K^*]).

The primal/dual methods are often combined with approximating L . The notion of epi/hypo convergence introduced by Attouch and Wets in [2] provides a setting for constructing a sequence L_n in such way that the saddle points $(\bar{x}_n, \bar{y}_n)$ of L_n converge to the saddle points of L . In this paper, however, we are interested in problems where L has a min/sup point rather than a saddle point,

and we are also interested in approximating L with a sequence of Lagrangians of simpler forms.

The notion of lopsided convergence of bivariate functionals defined on $X \times Y$, where X and Y are topological vector spaces, was introduced by Attouch and Wets in [1] in order to study the stability of min/sup points. A very similar notion was also introduced and studied in details by Lignola, Loridan, and Morgan in [5, 6]. In this paper, we will use the concept of lopsided convergence to approximate control problems. However, the original definition of lopsided convergence in [1] cannot be used directly due to the lack of compactness of the perturbation space for most control problems. Therefore, we modify the definition of lopsided convergence to better suit our applications.

In Section 3, we review different notions of convergence for bivariate functions. In Section 4, we show how the modified lopsided convergence can provide a simple way to recover a number of stability results that already exist in the literature. Section 5, contains the main application; we develop a scheme to approximate non-convex optimal control problems with a sequence of finite dimensional problems. The modified definition of lopsided convergence we introduce in Section 3 has further applications in two level programming and problems of existence of Stackelberg equilibria in a non compact setting. These applications, however, will be left for a subsequent paper.

2. Preliminaries

In this section, we review some basic definitions of set convergence that we will use in the following sections. Most of these definitions can be found in [4, 11]. Let (X, τ) be a topological vector space. Let C_n be subsets of X . We define the limit inferior and the limit superior of the collection C_n

$$\begin{aligned} \tau - \text{Li } C_n &= \{x \in X \mid \exists \text{ sequence } x_n \xrightarrow{\tau} x, x_n \in C_n\}, \\ \tau - \text{Ls } C_n &= \{x \in X \mid \exists \text{ subsequence } x_{n_k} \xrightarrow{\tau} x, x_{n_k} \in C_{n_k}\}. \end{aligned} \quad (2.1)$$

We define the following notions of set convergence

$$\begin{aligned} C_n &\xrightarrow{K^-} C_0, \quad \text{if } C_0 \subset \tau - \text{Li } C_n, \\ C_n &\xrightarrow{K^+} C_0, \quad \text{if } \tau - \text{Ls } C_n \subset C_0. \end{aligned} \quad (2.2)$$

We say C_n converge to C_0 in the *Painlevé-Kuratowski* sense, and we write $C_n \xrightarrow{p.k} C$, if

$$C_n \xrightarrow{K^+} C, \quad C_n \xrightarrow{K^-} C_0. \quad (2.3)$$

Let $f : X \rightarrow \bar{\mathbb{R}}$, where $\bar{\mathbb{R}}$ is the set of extended real numbers. The function f is *sequentially lower semi-continuous* (lsc), if for all $x \in X$ and $x_n \xrightarrow{\tau} x$,

$$\liminf_n f(x_n) \geq f(x). \quad (2.4)$$

Similarly, f is *sequentially upper semi-continuous* (usc), if for all $x \in X$ and $x_n \xrightarrow{\tau} x$,

$$\limsup_n f(x_n) \leq f(x). \quad (2.5)$$

The *domain* of f is

$$\text{dom } f = \{x \in X \mid f(x) < \infty\}, \quad (2.6)$$

and f is *proper*, if $\text{dom } f \neq \emptyset$ and f never assumes the value $-\infty$. The *epigraph* of f is

$$\text{epi } f = \{(x, \alpha) \in X \times \mathbb{R} \mid \alpha \geq f(x)\}. \quad (2.7)$$

The function f is *coercive*, if for any α , $\{x \mid |f(x)| \leq \alpha\}$ is bounded.

We say f_n τ -epi-converge to f on the topological space (X, τ) , and we write $f_n \xrightarrow{e} f$, if

$$(i) \quad \forall x \in X, \forall x_n \xrightarrow{\tau} x,$$

$$\liminf_n f(x_n) \geq f(x), \quad (2.8)$$

$$(ii) \quad \forall x \in X, \exists x_n \xrightarrow{\tau} x \text{ such that}$$

$$\limsup_n f(x_n) \leq f(x). \quad (2.9)$$

The set of minimizers of f is denoted by $\text{argmin } f = \{x \in X \mid f(x) = \inf_X f(x)\}$.

Theorem 2.1 is the main theorem regarding epi-convergence.

THEOREM 2.1 (see [11, Theorem 7.31]). *Let (X, τ) be a topological space and let $f_n : X \rightarrow \bar{\mathbb{R}}$ be a collection of functions that τ -epi-converge to a proper function f_0 . Then,*

$$\tau - \text{Lsargmin } f_n \subset \text{argmin } f_0. \quad (2.10)$$

Furthermore, if $\tau - \text{Lsargmin } f_n$ is not empty, then $\lim_n \inf f_n = \inf f_0$.

Let X and Y be topological spaces and suppose that $\langle \cdot, \cdot \rangle : X \times Y \rightarrow \mathbb{R}$ is a bilinear continuous map. Let $f : X \rightarrow \bar{\mathbb{R}}$ be a convex function. We define $f^* : Y \rightarrow \bar{\mathbb{R}}$, the conjugate of f , by

$$f^*(y) = \sup_{x \in X} \{ \langle x, y \rangle - f(x) \}. \quad (2.11)$$

38 On the convergence of min/sup points in optimal control problems

Consider the functions $F_0 : X \rightarrow \bar{\mathbb{R}}$ and $G : X \rightarrow \bar{\mathbb{R}}$, where G is lsc and convex. Let $F(x) = F_0(x) + G(\Lambda(x))$, where $\Lambda : X \rightarrow X$ is a continuous map. Then, a Lagrangian of F , with respect to a dual space Y , can be defined by

$$L(x, y) = F_0(x) + \langle \Lambda(x), y \rangle - G^*(y), \quad (2.12)$$

where

$$G^*(y) = \sup_{x \in X} \{ \langle x, y \rangle - G(x) \}. \quad (2.13)$$

The bivariate function L is a Lagrangian in the sense that

$$F(x) = \sup_{y \in Y} L(x, y). \quad (2.14)$$

We now consider a certain class of integral functionals on L^p spaces. Consider first the function $f : [0, T] \times \mathbb{R}^n \rightarrow \bar{\mathbb{R}}$. We say f is a normal integrand, if f is measurable in the first variable and lsc in the second. We define the integral functional I_f over the space $L^r([0, T], \mathbb{R}^n)$ for $r \in [1, +\infty]$,

$$I_f(u) = \int_0^T f(t, u(t)) dt. \quad (2.15)$$

Suppose f is a normal integrand that is also convex in the second argument, then I_f is a proper, convex, lsc (and weakly lsc) function on L^r . Furthermore, if $r \in [1, +\infty)$, then the conjugate of I_f is defined over $(L^r([0, T], \mathbb{R}^n))^*$ by the following equation

$$I_f^*(v) = \int_0^T f^*(t, v(t)) dt, \quad (2.16)$$

where for every t , $f^*(t, \cdot)$ is conjugate of $f(t, \cdot)$ over $\mathbb{R}^n$ ([4, Proposition 2.1, Chapter IX]). Note that for $r = +\infty$, and a conjugate space $(L^\infty)^*$, the above formula will not be valid.

We now list the definitions that we need regarding bivariate functions. Let $K : X \times Y \rightarrow \bar{\mathbb{R}}$ where X and Y are topological spaces. We say K is *proper*, if the function $V : X \rightarrow \bar{\mathbb{R}}$ defined by $V(x) = \sup_{y \in Y} K(x, y)$ is proper. A *saddle point* for K is a pair $(\bar{x}, \bar{y})$ such that

$$K(\bar{x}, y) \leq K(\bar{x}, \bar{y}) \leq K(x, \bar{y}), \quad \forall x \in X, \forall y \in Y. \quad (2.17)$$

A min/sup point for K is a point $\bar{x} \in X$ such that

$$\sup_{y \in Y} K(\bar{x}, y) \leq \sup_{y \in Y} K(x, y), \quad \forall x \in X. \quad (2.18)$$

We write $\arg \text{sp } K$ to denote the set of saddle points of K , and we write $\arg \text{minsup } K$ to denote its set of min/sup points.

Finally, in all what follows, we will use the standard convention of $\infty - \infty = \infty$. This will allow us to deal with constraints on the variables x and y in a consistent manner.

3. Lopsided convergence

We start by reviewing some notions of convergence for bivariate functions. Let (X, τ) , (Y, σ) be topological spaces and consider a sequence of bivariate functionals $K_n : X \times Y \rightarrow \bar{\mathbb{R}}$. The sequence $\{K_n\}$ epi/hypo converges to K_0 [2], if

- (i) For all $(x, y) \in X \times Y$, for all $x_n \xrightarrow{\tau} x$, there exists $y_n \xrightarrow{\sigma} y$ such that

$$\liminf K_n(x_n, y_n) \geq K_0(x, y). \quad (3.1)$$

- (ii) For all $(x, y) \in X \times Y$, for all $y_n \xrightarrow{\sigma} y$, there exists $x_n \xrightarrow{\tau} x$ such that

$$\limsup K_n(x_n, y_n) \leq K_0(x, y). \quad (3.2)$$

Epi/hypo convergence implies the convergence of saddle points.

THEOREM 3.1 (see [2]). *If K_n epi/hypo converge to K_0 , then*

$$\tau - \text{Ls arg sp } K_n \subset \text{arg sp } K_0. \quad (3.3)$$

Definition 3.2 is the original definition of the lopsided convergence [3].

Definition 3.2. The sequence $\{K_n\}$ lopsided converges to K_0 , denoted by $K_n \xrightarrow{\text{lo}} K_0$, if

- (i) For all $(x, y) \in X \times Y$, for all $x_n \xrightarrow{\tau} x$, there exists $y_n \xrightarrow{\sigma} y$ such that

$$\liminf K_n(x_n, y_n) \geq K_0(x, y). \quad (3.4)$$

- (ii) For all $x \in X$, there exists $x_n \xrightarrow{\tau} x$ such that for all $y \in Y$ and for all $y_n \xrightarrow{\sigma} y$,

$$\limsup K_n(x_n, y_n) \leq K_0(x, y). \quad (3.5)$$

Note that in the definition of epi/hypo convergence there is a symmetry between parts (i) and (ii). The term “lopsided” is meant to emphasize the lack of such symmetry in **Definition 3.2**. **Theorem 3.3** is the main theorem regarding lopsided convergence that concerns us.

THEOREM 3.3 (see [1]). *If $K_n \xrightarrow{\text{lo}} K_0$, Y is a compact space, then*

$$\tau - \text{Ls argminsup } K_n \subset \text{argminsup } K_0. \quad (3.6)$$

It is clear from the definitions that lopsided convergence implies epi/hypo convergence but the converse is not true. The compactness of Y is essential in the proof of **Theorem 3.3** even though the compactness of Y is not required for

the existence of a min/sup point [1]. In this paper, however, we will investigate control problems where the space Y is a space of perturbations for the state variables which is not compact in most cases (or not compact in a topology that is compatible with the continuity properties the problem has). Therefore, we introduce the following modification of [Definition 3.2](#).

Definition 3.4. Consider two topologies τ_1 and τ_2 on the space X and assume that τ_1 is stronger than τ_2 . Similarly, consider two topologies σ_1 and σ_2 on Y and assume that σ_1 is stronger than σ_2 . The sequence $\{K_n\}$ converges to K_0 , $K_n \xrightarrow{\text{lo2}} K_0$, if

(i) For all $(x, y) \in X \times Y$, for all $x_n \xrightarrow{\tau_2} x$, there exists $y_n \xrightarrow{\sigma_1} y$ such that

$$\liminf K_n(x_n, y_n) \geq K_0(x, y). \quad (3.7)$$

(ii) For all $x \in X$, there exists $x_n \xrightarrow{\tau_1} x$ such that for any sequence $\{y_n\}$ such that $K_n(x_n, y_n)$ is bounded below, there exists a subsequence n_k , there exists $\epsilon_{n_k} \searrow 0$, there exists $w_{n_k} \in Y$ such that $w_{n_k} - y_{n_k} \xrightarrow{\sigma_2} 0$, and that eventually

$$K_0(x, w_{n_k}) \geq K_{n_k}(x_{n_k}, y_{n_k}) - \epsilon_{n_k}. \quad (3.8)$$

Note that when $\tau_1 = \tau_2$, $\sigma_1 = \sigma_2$, and Y is compact, the two definitions of lopsided convergence are equivalent. Now we can prove the following theorem.

THEOREM 3.5. *If $K_n \xrightarrow{\text{lo2}} K_0$ and K_0 is proper, then*

$$\tau_2 - \text{Ls argminsup } K_n \subset \text{argminsup } K_0. \quad (3.9)$$

The proof of [Theorem 3.5](#) follows immediately from the following lemma and [Theorem 2.1](#).

LEMMA 3.6. *Let X, Y be topological spaces as in [Definition 3.4](#). Let $K_n : X \times Y \rightarrow \bar{\mathbb{R}}$, where K_0 is proper. Let*

$$V_n(x) = \sup_{y \in Y} K_n(x, y), \quad n = 0, 1, 2, \dots \quad (3.10)$$

Assume

(i) *For every $(x, y) \in X \times Y$, for all $x_n \xrightarrow{\tau_2} x$, there exists $y_n \xrightarrow{\sigma_1} y$ such that*

$$\liminf K_n(x_n, y_n) \geq K_0(x, y). \quad (3.11)$$

(ii) *For every $x \in X$, there exists $x_n \xrightarrow{\tau_1} x$ such that for any sequence $\{y_n\}$, where $K_n(x_n, y_n)$ is bounded below, there exists a subsequence n_k and there exist $\epsilon_{n_k} \searrow 0$ and $w_{n_k} \in Y$ such that eventually*

$$K_0(x, w_{n_k}) \geq K_{n_k}(x_{n_k}, y_{n_k}) - \epsilon_{n_k}. \quad (3.12)$$

Then,

$$\tau_2 - \text{Ls argmin } V_n \subset \text{argmin } V_0. \quad (3.13)$$

Furthermore, if $\tau_2 - \text{Ls argmin } V_n$ is not empty, then

$$\liminf_n V_n = \liminf V_0. \quad (3.14)$$

Proof. Condition (i) implies that for all $(x, y) \in X \times Y$, for all $x_n \xrightarrow{\tau_2} x$, there exists $y_n \xrightarrow{\sigma_1} y$ such that

$$\liminf_n \sup_{y \in Y} K_n(x_n, y) \geq \liminf_n K_n(x_n, y_n) \geq K_0(x, y). \quad (3.15)$$

Hence,

$$\liminf_n V_n(x_n) \geq V_0(x). \quad (3.16)$$

We also claim that for all $x \in X$, there exists $x_n \xrightarrow{\tau_1} x$ such that

$$\limsup_n V_n(x_n) \leq V_0(x). \quad (3.17)$$

Assume not, and assume that $V_0(x) < +\infty$. Then, for every sequence $x_n \xrightarrow{\tau_1} x$, we have $\limsup_n V_n(x_n) = \beta > V_0(x)$. Thus, there exists a subsequence of $V_n(x_n)$ that converges to β . To simplify the notation, we will also use n to index this subsequence. Hence, there exist $\epsilon > 0$, $y \in Y$, a sequence $\{y_n\}$ in Y , and there exists n_0 such that for all $n \geq n_0$,

$$K_n(x_n, y_n) > K_0(x, y) + \epsilon. \quad (3.18)$$

Assumption (ii) implies that there exist w_{n_k} and $\epsilon_{n_k} \searrow 0$ such that eventually

$$K_0(x, w_{n_k}) > K_{n_k}(x_{n_k}, y_{n_k}) - \epsilon_{n_k}. \quad (3.19)$$

Hence, for some $n_k > n_0$, and for some $\epsilon' > 0$, we have

$$K_0(x, w_{n_k}) > K_0(x, y) + \epsilon'. \quad (3.20)$$

Thus,

$$V_0(x) \geq K_0(x, y) + \epsilon', \quad (3.21)$$

which contradicts the definition of V_0 .

Finally (3.16) and (3.17) yield the required conclusion via [Theorem 2.1](#). $\square$

Remark 3.7. In [Definition 3.4](#) of lopsided convergence, the requirements that $y_n \xrightarrow{\sigma_1} y$ in part (i) and that $w_{n_k} - y_{n_k} \xrightarrow{\sigma_2} 0$ in part (ii) were not used in the proof of [Lemma 3.6](#). In fact, these requirements are not needed for the conclusion of [Theorem 3.5](#). However, they are needed to make the notion of lopsided convergence stable with respect to perturbation of some classes of functions (e.g., the class of $\tau_2 \times \sigma_2$ continuous biaffine functions).

Remark 3.8. There is a number of variations of the definition of epi/hypo convergence. These variations consider more than one topology on the spaces X and Y (see [3, 12]) in a way that is very similar to what we have in [Definition 3.4](#).

Remark 3.9. At this point, it may seem that the original definition of lopsided convergence can be directly used in problems defined over infinite dimensional spaces when the Lagrangian is coercive in the y variable. It may seem that all what is needed in this case is to restrict the Lagrangian to a bounded set of Y and to choose a weak enough topology that will make this bounded set compact. This, however, is not true, as the last remark of [Section 5](#) will illustrate. In fact, in some of our applications, a topology that is weak enough to make the modified space Y compact will make it very difficult for us to verify condition (ii) of [Definition 3.2](#).

4. Applications I

In our first application, we will consider problems with Lagrangians of the following form (for examples and details see [4, Chapter VII]):

$$L(u, v) = J(u) + \langle v, \Phi(u) \rangle + \delta_A(u) - \delta_B(v), \quad (4.1)$$

where $L : U \times V \rightarrow \bar{\mathbb{R}}$ and U and V are Hilbert spaces, $J : U \rightarrow \mathbb{R}$ is a convex, lsc, Gâteaux differentiable function. A and B are nonempty closed convex subsets of U and V , respectively. δ_A and δ_B are the indicator functions of these sets (0 on the set and $+\infty$ outside it).

The map $\Phi : A \rightarrow V$ is Lipschitzian in the sense that

$$\|\Phi(u_2) - \Phi(u_1)\| \leq c \|u_1 - u_2\|, \quad \forall u_1, u_2 \in A. \quad (4.2)$$

And for every $v \in V$, the function $u \rightarrow \langle v, \phi(u) \rangle$ is convex and lsc on A , where $\langle \cdot, \cdot \rangle$ is the inner product of V . Note that our assumptions imply that for all v , for all $u_n \xrightarrow{w} u$, we have

$$\liminf_n \langle v, \phi(u_n) \rangle \geq \phi(v, u). \quad (4.3)$$

A number of primal/dual methods that take advantage of the special form of the Lagrangian can be used to attack this problem (cf. [4, Chapter VII]). Most of these methods, however, require the following additional conditions on J , A , and B

- (a) B is bounded
- (b) J is coercive in the sense that for all $u_1, u_2 \in A$,

$$\langle J'(u_2) - J'(u_1), u_1 - u_2 \rangle \geq \alpha \|u_2 - u_1\|, \quad \alpha > 0. \quad (4.4)$$

In this section, we are interested in problems that do not satisfy the above conditions. Therefore, we perturb the original Lagrangian in such a way that the

resulting Lagrangians will be

$$L_n(u, v) = J(u) + \langle v, \Phi(u) \rangle + \frac{1}{n} \|u\|^2 + \delta_A(u) - \delta_{B_n}(v), \quad (4.5)$$

where $B_n = B \cap r_n \mathcal{B}$, where $\mathcal{B}$ is the unit ball in V , and r_n increases to infinity.

Clearly, L_n satisfies conditions (a) and (b). We now show that L_n lopsided converges to L .

THEOREM 4.1. *Let τ_1 and τ_2 be, respectively, the norm and the weak topology on U . Similarly, let σ_1 and σ_2 be, respectively, the norm and weak topology on V . Let L_0 and L_n be defined by (4.1) and (4.5), then*

$$L_n \xrightarrow{\text{lo2}} L_0. \quad (4.6)$$

Proof. Let $(u, v) \in U \times V$ such that $v \in B$. Let $u_n \xrightarrow{w} u$, $v_n \xrightarrow{s} v$, and $v_n \in B_n$. Due to (4.3), and since every convex lsc function on a Banach space is also weakly lsc, we immediately obtain

$$\begin{aligned} \liminf_n \left\{ J(u_n) + \delta_A(u_n) + \langle v_n, \phi(u_n) \rangle + \frac{1}{n} \|u_n\|^2 - \delta_{B_n}(v_n) \right\} \\ \geq J(u) + \delta_A(u) + \langle v, \phi(u) \rangle - \delta_B(v), \end{aligned} \quad (4.7)$$

which is condition (i) of Definition 3.4.

To verify condition (ii), we let $u \in U$. For any sequence v_n , there exists ϵ_n such that

$$\begin{aligned} J(u) + \delta_A(u) + \langle v_n, \phi(u) \rangle - \delta_B(v_n) \\ \geq J(u) + \delta_A(u) + \langle v_n, \phi(u) \rangle + \frac{1}{n} \|u\|^2 - \delta_{B_n}(v_n) - \epsilon_n. \end{aligned} \quad (4.8)$$

□

Remark 4.2. The above approximation method can be used when B is unbounded. Moreover, in some control problems, the set B has the following form:

$$B = \{u \in L^p(\Omega) \text{ such that } \|\text{grad } u\| \leq M\}, \quad (4.9)$$

where Ω is a bounded open set in $\mathbb{R}^n$ and M is some constant. The set B is bounded in L^p due to the Poincaré inequality. However, the bound of B (the Poincaré constant for the region Ω) may not be available. In this case, it is convenient numerically to use the above approximation and replace B with an increasing sequence of balls in L^p even though B is bounded.

Finally, we show how the modified lopsided convergence can be used to recover some stability results regarding some classical control problems.

44 On the convergence of min/sup points in optimal control problems

Consider the following problem:

Minimize $J : L^2(Q) \rightarrow \mathbb{R}$

$$J(u) = \int_Q |y(t, x; u) - y_d|^2 dx dt \quad (4.10)$$

subject to

$$|\text{grad } y(t, x, ; u)| \leq 1 \quad \text{a.e., } (t, x) \in Q, \quad (4.11)$$

where $y_d(t, x) \in L^2([0, T]; H_o^1(\Omega))$ is a given function, and $y(t, x, u) \in L^2([0, T]; H_o^1(\Omega))$ is the unique weak solution of the following dynamics:

$$\begin{aligned} \frac{\partial y}{\partial t} - \Delta y &= f + u \quad \text{in } Q, \\ y &= 0 \quad \text{on } \partial\Omega \times]0, T[, \\ y(0, x; u) &= y_0(x), \quad \forall x \in \Omega, \end{aligned} \quad (4.12)$$

where Ω is a subset in $\mathbb{R}^n$, $T > 0$, $Q =]0, T[\times \Omega$, $y_0 \in L^2(\Omega)$, and $f \in L^2(Q)$.

The Lagrangian of the above problem can be expressed as (see [4, Section 3.1] for details)

$$L(u, v) = \int_Q \left[\frac{1}{2} |\text{div}(u)|^2 - \text{div}(u) \cdot z_d + u \cdot \text{grad } \phi + u \cdot v \right] dx dt, \quad (4.13)$$

where $\phi(t, x) = y(t, x; 0)$, $z(t, x; u) = y(t, x; u) - y(t, x; 0)$, $z_d(t, x) = y_d(t, x) - y(t, x; 0)$, and $\phi = y(0)$ (see [4, Section 3.1, Chapter VII]).

Solving the original control problem corresponds to finding u , where the following infimum is attained:

$$\inf_{u \in L^2(Q)^n} \sup_{v \in B} L(u, v), \quad (4.14)$$

where $A = \{u \mid u \in L^2(Q)^n, \text{div } v \in L^2(Q)\}$ and $B = \{v \mid v \in L^2(Q)^n \text{ and } |v(x)| \leq 1 \text{ a.e.}\}$. Note that, despite the fact that new formulation of the problem involves a supremum and an infimum, the format of L and the simplicity of the set B make the new formulation of the problem easier to solve numerically. The Lagrangian in (4.13), however, does not satisfy requirement (b) for the direct application of numerical methods. Furthermore, (4.5) may not have a saddle point under our current assumptions, and the standard methods of approximating saddle points (via epi/hypo convergence) cannot be used. Therefore, for $\epsilon > 0$, we introduce

$$L_\epsilon(u, v) = \int_Q \left[\frac{1}{2} |\text{div } u|^2 - \text{div } u \cdot z_d + \frac{\epsilon}{2} |u|^2 + u \cdot \text{grad } \phi + u \cdot v \right] dx dt. \quad (4.15)$$

Now for every $\epsilon > 0$, the Lagrangian $L_\epsilon(u, v)$ satisfies conditions (a) and (b), and a standard numerical method, such as the Uzawa method [4], can be used

to find a saddle point (u_ϵ, v_ϵ) for L_ϵ . If we show that, as $\epsilon \rightarrow 0$, L_ϵ lopsided converges to L , then we will know that every cluster point of $\{u_\epsilon\}$ is a min/sup point of L . This is precisely the claim of the following theorem.

THEOREM 4.3 (also [4, Proposition 3.4, Chapter VII]). *Let $L_\epsilon(u, v)$ and $L(u, v)$ be defined by (4.13) and (4.15). As $\epsilon \rightarrow 0$, we have*

$$\begin{aligned} L_\epsilon(u, v) &\xrightarrow{\text{lo2}} L(u, v), \\ w - \text{Ls argminsup } L_\epsilon(u, v) &\subset \text{argminsup } L(u, v). \end{aligned} \quad (4.16)$$

Furthermore, if $w - \text{Ls argminsup } L_\epsilon(u, v) \neq \emptyset$, then

$$\inf_V \sup_U L_\epsilon(u, v) \longrightarrow \inf_V \sup_U L(u, v). \quad (4.17)$$

Proof. The L^2 norm in the space $A = \{u \mid u \in L^2(Q)^n, \text{div } v \in L^2(Q)\}$ is weakly lsc, and the conditions of Definition 3.4 can be easily verified with the appropriate choices for the topologies τ_1 , τ_2 , σ_1 , and σ_2 . $\square$

5. Application II

In this section, we consider the following control problem, minimize, over $L^p = L^p([0, T]; \mathbb{R}^n)$, with $p \in [2, +\infty)$, the functional

$$I_1(u) = \int_0^T \phi(t, u(t), x(t)) dt, \quad (5.1)$$

where $x(\cdot) : [0, T] \rightarrow \mathbb{R}^k$ is a solution to

$$Lx = u, \quad (5.2)$$

and L is an operator from $C([0, T], \mathbb{R}^k)$ to L^1 . The states and controls are subject to the constraint

$$(t, f(t, u(t), x(t))) \in \text{gph } E \quad \text{a.e.}, \quad (5.3)$$

where E is a set-valued map from $[0, T]$ to $\mathbb{R}^j$ and $\text{gph } E$ is the graph of the map E .

Assumptions on the operator L . The operator L has an inverse Λ such that u_n converges weakly to u in L^p which implies that Λu_n converges pointwise to Λu .

The following are examples of operators that satisfy our assumptions.

Example 5.1. L is the linear differential operator given by

$$x'(t) = A(t)x(t) + B(t)u(t), \quad x(0) = \alpha, \quad (5.4)$$

with the standard assumptions on A and B .

Example 5.2. L is a differential operator representing an evolution equation,

$$x'(t) = g(t, u(t), x(t)), \quad x(0) = \alpha, \quad (5.5)$$

under the usual assumptions of growth and Lipschitz continuity on g .

Example 5.3. Any differential operator L whose inverse can be expressed by an integral equation of the form

$$x(t) = \int_0^T g(t, s)u(s)ds, \quad (5.6)$$

where $g(t, \cdot)$ is in $L^{+\infty}$.

Assumptions on the constraint (5.3). The map $f : [0, T] \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}^j$ is measurable in the first argument, continuous in the rest. The map E is nonempty and convex-valued with a closed graph. We also posit a growth condition, there exists a function $s \in L^2$ such that

$$\sup_{z \in E(t)} \|z\| \leq s(t) \quad \text{a.e.} \quad (5.7)$$

The constraint (5.3) can model a combination of constraints such as

$$u(t) \in U(t), \quad x(t) \in V(t), \quad M(t, x(t), u(t)) \in C(t), \quad (5.8)$$

where U , V , and C are set-valued maps. In this case, we only need to define

$$\begin{aligned} E(t) &= U(t) \times V(t) \times C(t), \\ f(t, u(t), x(t)) &= (u(t), x(t), M(t, x(t), u(t))). \end{aligned} \quad (5.9)$$

Assumptions on the cost function I_1 . We will assume that I_1 is finite-valued, weakly lower semi-continuous, and strongly upper semi-continuous over L^2 . These assumptions can be formulated in terms of standard conditions on ϕ

(i) The map $\phi : [0, T] \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}$ is measurable in the first argument, convex in the second, and continuous in the third.

(ii) There exists a function $\Phi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that for all t , y , and z such that

$$\Phi(|z|) \leq \phi(t, z, y). \quad (5.10)$$

(iii) For all $u \in L^2$, there exists a neighborhood V of u and a function $h \in L^1$ such that for every $u' \in V$

$$\phi(t, u'(t), x'(t)) \leq h(t), \quad (5.11)$$

where $x' = \Lambda u'$.

The above conditions imply that I_1 is finite-valued. Note that assuming that I_1 is finite-valued does not limit the scope of this model, since we will deal with the constraints on the controls when we deal with constraint (5.3). Furthermore, conditions (i) and (ii) imply the weak lower semi-continuity of I_1 (see [4, Theorem 2.1, Chapter VIII]). The (strong) upper semi-continuity of I_1 follows from (iii), Fatou's lemma, and Lebesgue dominated convergence theorem.

Our goal in this section is to construct finite dimensional approximations for the above non-convex (partially convex) control problem. In [9, 10], Rockafellar developed the full duality theory for a similar type of control problems. However, he was mainly interested in cases where the existence of saddle points for the Lagrangians is guaranteed, and therefore he required the cost function to be convex (in the control and the state) and the dynamics to be linear. We will be able to relax these conditions, since we are only interested in the partial duality of the problem and since we only assume the existence of min/sup points.

To simplify the notation, we will assume that $f : [0, T] \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}^n$. Hence, for all u and for all x such that $\Lambda x = u$, we have $f(t, u(t), x(t)) \in L^2$ because of the growth condition on E . We introduce an exact but finite penalty function.

Let $\theta : [0, T] \times \mathbb{R}^n$ be a normal, convex (in the second argument) integrand

$$\theta(t, w) = \begin{cases} 0, & \text{if } (t, w) \in \text{gph } E, \\ > 0, & \text{otherwise.} \end{cases} \quad (5.12)$$

More specifically, we define

$$\theta(t, z) = d(z, E(t)), \quad (5.13)$$

where $d(z, C) = \inf_{c \in C} \|c - z\|$ for any closed subset C . Note that $\theta(t, \cdot)$ is finite over $\mathbb{R}^j$.

Now the problem can be expressed as minimizing, over L^p , the functional

$$F(u) = I_1(u) + I_2(w), \quad (5.14)$$

where $I_2 : L^p \rightarrow \mathbb{R}$ is defined by

$$I_2(w) = \int_0^T \theta(t, w(t)) dt, \quad (5.15)$$

$$I_1(u) = \int_0^T \phi(t, u(t), x(t)) dt, \quad (5.16)$$

where $w(t) = f(t, u(t), x(t))$, and $\Lambda u = x$.

A Lagrangian associated with the above problem is given by $L : L^p \times L^2 \rightarrow \mathbb{R}$ with

$$L(u, v) = I_1(u) - I_2^*(v) + \Gamma(v, w), \quad (5.17)$$

where

$$I_2^* = \int_0^T \theta^*(t, v(t)) dt, \quad (5.18)$$

$$\theta^*(t, z) = \sup_{y \in \mathbb{R}^J} \{ \langle y, z \rangle - \theta(t, y) \}, \quad (5.19)$$

where θ is still given by (5.13), and

$$\Gamma(u, v) = \int_0^T v(t)w(t) dt, \quad (5.20)$$

where again $w(t) = f(t, u(t), x(t))$, and $\Lambda u = x$. A direct calculation of $\theta^*(t, z)$, when θ is given by (5.13), gives us

$$\theta^*(t, z) = \sigma_{E(t)}(z) + \delta_B(z), \quad (5.21)$$

where $\sigma_{E(t)}(z) = \sup_{y \in E(t)} \langle y, z \rangle$ is the support function of the set $E(t)$, and B is the unit ball in $\mathbb{R}^n$ (see [11, Example 11.26, Chapter 11] for details). Note that $\theta^*(t, \cdot)$ is convex, proper, and lsc. Moreover, it is coercive over $\mathbb{R}^n$ since $\theta(t, \cdot)$ is finite everywhere.

The fact that $F(u) = \sup_{v \in L^2} L(u, v)$ follows from the definition of the conjugate function and from the fact that the conjugate of I_2 is actually given by (5.18) (see also (2.12) in the Preliminaries).

Using exact penalties for the joint state and control constraints causes serious computational complications (see [10] for details). Therefore, we introduce a sequence of finite non exact penalties $\theta_n : [0, T] \times \mathbb{R}^n \rightarrow \mathbb{R}$ such that $\theta_n(t, \cdot)$ increase continuously to $\theta(t, \cdot)$. More specifically, we will take Moreau envelopes of θ ,

$$\theta_n(t, z) = \inf_{y \in \mathbb{R}^J} \left\{ \theta(t, y) + \frac{n}{2} \|y - z\|^2 \right\}. \quad (5.22)$$

These approximating functions are strictly convex and differentiable. Moreover, the conjugates of $\theta_n^*(t, w)$ are given by (see [11, Chapter 11])

$$\theta_n^*(t, z) = \sigma_{E(t)}(z) + \delta_B(z) + \frac{1}{2n} \|z\|^2. \quad (5.23)$$

Since $\theta_n(t, \cdot)$ is strictly convex, $\theta_n^*(t, \cdot)$ is differentiable on the interior of its domain (see [11, Theorem 11.13]).

We now use a method developed by Wright [12] to discretize the control problem in the primal and dual variables at the same time. Let Π_n be an increasing sequence of partitions of $[0, T]$

$$\Pi_n = (0 = a_0^n < a_1^n < \dots < a_{s-1}^n < a_s^n < \dots < a_T^n = T), \quad (5.24)$$

such that $|a_s^n - a_{s-1}^n| \rightarrow 0$, uniformly in s , as $n \rightarrow +\infty$.

We define the following sets:

$$\begin{aligned} U_n &= \{u \in L^p \mid u \text{ is constant on } (a_{s-1}^n, a_s^n), s = 0, 1, \dots, T\}, \\ V_n &= \{v \in L^2 \mid v \text{ is constant on } (a_{s-1}^n, a_s^n), s = 0, 1, \dots, T\}. \end{aligned} \quad (5.25)$$

The sequence of approximating Lagrangians that we will consider is

$$L_n(u, v) = I_{1,n}(u) - I_{2,n}^*(v) + \Gamma(u, v), \quad (5.26)$$

where δ_{U_n} and δ_{V_n} are the indicator functions of U_n and V_n , respectively, and

$$I_{1,n}(u) = I_1(u) + \delta_{U_n}(u), \quad (5.27)$$

$$I_{2,n}^*(v) = \int_0^T \theta_n^*(t, v(t)) dt + \delta_{V_n}(v), \quad (5.28)$$

where $I_{2,n}^* : L^2 \rightarrow \mathbb{R}$ and θ_n^* is given by (5.19).

The standard convention of $\in - \infty = \infty$ implies that $L(u, v) = +\infty$, if $u \notin U_n$. Moreover, $L(u, v) = -\infty$, if $u \in U_n$ and $v \notin V_n$. Now, the problem of finding the min/sup points of L_n is a finite dimensional problem. Also note that for a step function v , calculating $\theta_n^*(t, v(t))$ reduces to solving a number of simple problems of convex minimization (concave maximization) in $\mathbb{R}^n$. Furthermore, the discretized problem will be governed by a difference equation (see [12] for details).

Since L_n is not convex in u (ϕ is not convex in x and Γ is not convex in u), L and L_n may not possess saddle points. However, we can use the method of augmented Lagrangians developed by Rockafellar in [8] to find the min/sup points of L_n . The idea of this method is to construct an augmented Lagrangian $\tilde{L}_n(u, v, r)$, for some $r \in (0, +\infty)$, and then use some standard primal/dual numerical method to find $(\bar{u}_n, \bar{v}_n)$, the saddle points of $\tilde{L}_n(u, v, r)$. For every n , the point $\bar{u}_n$ will then be a min/sup point for L_n (see [8] and [11, Section K^* , Chapter 11], also see [7, Chapter 17] for an actual algorithm using augmented Lagrangians). This approach will allow us to approximate, discretize, and numerically solve the original problem despite its lack of convexity.

Finally, we state the main theorem of the section which will show that our approximation scheme will actually work.

THEOREM 5.4. *Let L_n and L be defined by (5.17) and (5.26). Then*

$$w - Ls \operatorname{argminsup} L_n \subset \operatorname{argminsup} L. \quad (5.29)$$

Before we prove Theorem 5.4, we list a number of basic lemmas that we will need in our proof.

LEMMA 5.5. Consider the functions I_1 and $I_{1,n}$ defined by (5.16) and (5.27). Consider also the sets U_n (5.25). Let $u \in L^p$, then there exists a sequence $\hat{u}_n \in U_n$ such that

$$\lim I_{1,n}(\hat{u}_n) = I_1(u). \quad (5.30)$$

Proof. The proof is an immediate result of the fact that step functions are dense in L^p for $p \in [2, +\infty)$ and of our assumption that I_1 is norm continuous. $\square$

LEMMA 5.6. Let $v \in L^2$ such that $v(t) \in B$ a.e., where B is the unit ball in $\mathbb{R}^n$. Then there exists a sequence $\hat{v}_n \rightarrow v$ such that $v_n \in B \cap V_n$ and

$$\limsup_n \int_0^T \sigma_{E(t)}(\hat{v}_n(t)) dt \leq \int_0^T \sigma_{E(t)}(v(t)) dt. \quad (5.31)$$

Proof. Using some elementary facts from measure theory (see [12, Lemmas 1, 2] for details) and using the definition of the support function, we can find a sequence $\hat{v}_n \rightarrow v$ such that $v_n(t) \in B$, $\hat{v}_n(t) \rightarrow v(t)$, and $v_n(t) \leq v(t)$ a.e. in $[0, T]$. Hence, for all n , we have

$$\int_0^T \sigma_{E(t)}(\hat{v}_n(t)) dt \leq \int_0^T \sigma_{E(t)}(v(t)) dt, \quad (5.32)$$

and the assertion of the lemma is immediate. $\square$

LEMMA 5.7. Let $v \in L^2$ such that $v(t) \in B$ a.e. Then, there exists a sequence $\hat{v}_n \rightarrow v$ such that $v_n(t) \in B \cap V_n$ and

$$\limsup_n I_{2,n}^*(\hat{v}_n) \leq I_2^*(v). \quad (5.33)$$

Proof. Lemma 5.6 and the definition of $I_{2,n}^*$ given by (5.19) and (5.28). $\square$

Proof of Theorem 5.4. In light of Theorem 3.5, we only need to show that L_n lopsided converges to L when τ_1 and τ_2 are, respectively, the norm and weak topologies on L^p , and σ_1 and σ_2 are also, respectively, the norm and weak topologies on L^2 .

Verifying part (i) of Definition 3.4.

For all $(u, v) \in L^p \times L^2$ such that $v \in \text{dom } I_2^*$, take $\hat{v}_n$ as in Lemma 5.7. Note that, for all $u_n \xrightarrow{w} u$, we have

$$\liminf_n I_1(u_n) \geq I_1(u) \quad (5.34)$$

due to our assumption that I_1 is weak lsc. Moreover,

$$\lim_n \|\Gamma(\hat{u}_n, v_n) - \Gamma(u, v)\| = 0. \quad (5.35)$$

It is clear now from our choice of $\hat{v}_n$ and from (5.34) and (5.35) that part (i) of Definition 3.4 is satisfied.

Verifying part (ii) of Definition 3.4.

For all $(u, v) \in L^p \times L^2$ such that $\sup_v L(u, v) < +\infty$, take $\hat{u}_n \rightarrow u$ as in Lemma 5.5. Now, for any $v_n \in L^2$ such that $-\infty < L_n(\hat{u}_n, v_n) = I_{1,n}(\hat{u}_n) - I_{2,n}(v_n) + \Gamma(\hat{u}_n, v_n)$, we must have $\{v_n(t)\} \in B$ a.e. Hence $\{v_n\}$ must be bounded in L^2 . Hence,

$$\lim_n \|\Gamma(\hat{u}_n, v_n) - \Gamma(u, v_n)\| = 0. \quad (5.36)$$

Therefore, there exists $\epsilon_n \rightarrow 0$ such that eventually

$$I_1(u) - I_2(v_n) + \Gamma(u, v_n) \geq I_{1,n}(\hat{u}_n) - I_2(v_n) + \Gamma(\hat{u}_n, v_n) - \epsilon_n \quad (5.37)$$

and part (ii) of the definition is verified. $\square$

Remark 5.8. Theorem 5.4 is only a stability result, in the sense that $\text{argminsup } L$ might be empty. However, standard conditions on ϕ can be added to assure that w -Ls $\text{argminsup } L_n$ and $\text{argminsup } L$ will not be empty. In this case, the numerical method we suggested will produce a sequence of pairs $(u_n, v_n) \in L^p \times L^2$ which are the saddle points of the augmented approximating Lagrangians. The sequence v_n will be bounded in L^2 , which is important for numerical reasons, but its weak cluster point may fail to be a solution to the dual problem. More specifically, a weak $\times$ weak cluster point of (u_n, v_n) , may fail to be a saddle point for the Lagrangian of the original control problem. However, as Theorem 5.4 shows, any weak cluster point of u_n will be a solution of the original control problem. Finally, if we know a priori that the original control problem has saddle points, then Theorem 5.4 can be easily modified to show stability of the saddle points with respect to the discretization and approximation scheme of this section.

Remark 5.9. The convexity of ϕ in the second variable was needed only to obtain the weak lower semi-continuity of I_1 . In case the cost function is known to be weakly lsc on L^p , no such assumption is needed. A cost function that depends on the state only would be an example of such a case.

Remark 5.10. We finally elaborate further on Remark 3.9 of Section 3; under the appropriate coercivity conditions, we may attempt to use the original definition of lopsided convergence, with Y taken as a large enough ball in L^2 and σ as the (relative) weak topology. In this case, Y is compact in σ . However, verifying condition (ii) of Definition 3.2 is, in essence, equivalent to verifying

that for all $u_n \xrightarrow{w} u$ in Y ,

$$\liminf_n \int \sigma_{E(t)} v_n(t) dt \geq \int \sigma_{E(t)} v_n(t) dt. \quad (5.38)$$

Such statement cannot be verified without further conditions on the set-valued map E .

References

- [1] H. Attouch and R. J.-B. Wets, *Convergence de points min/sup et de points fixes* [Convergence of min/sup points and fixed points], C. R. Acad. Sci. Paris Sér. I Math. **296** (1983), no. 21, 865–867 (French). [MR 84i:49028](#).
- [2] ———, *A convergence theory for saddle functions*, Trans. Amer. Math. Soc. **280** (1983), no. 1, 1–41. [MR 85f:49024](#). [Zbl 525.49009](#).
- [3] A. Bagh, *Epi/hypo-convergence: the slice topology and saddle points approximation*, J. Appl. Anal. **2** (1996), no. 1, 13–39. [MR 98b:49013](#). [Zbl 880.49011](#).
- [4] I. Ekeland and R. Temam, *Convex Analysis and Variational Problems*, North-Holland Publishing and American Elsevier Publishing, Amsterdam and New York, 1976. [MR 57#3931b](#). [Zbl 322.90046](#).
- [5] M. B. Lignola and J. Morgan, *Topological existence and stability for Min Sup problems*, J. Math. Anal. Appl. **151** (1990), no. 1, 164–180. [MR 91f:49011](#). [Zbl 714.49012](#).
- [6] P. Loridan and J. Morgan, *New results on approximate solutions in two-level optimization*, Optimization **20** (1989), no. 6, 819–836. [MR 91a:90177](#). [Zbl 684.90089](#).
- [7] J. Nocedal and S. J. Wright, *Numerical Optimization*, Springer Series in Operations Research, Springer-Verlag, New York, 1999. [MR 2001b:90002](#). [Zbl 930.65067](#).
- [8] R. T. Rockafellar, *Augmented Lagrange multiplier functions and duality in nonconvex programming*, SIAM J. Control **12** (1974), no. 2, 268–285. [MR 52#5040](#). [Zbl 285.90063](#).
- [9] ———, *Linear-quadratic programming and optimal control*, SIAM J. Control Optim. **25** (1987), no. 3, 781–814. [MR 88e:49045](#). [Zbl 617.49010](#).
- [10] ———, *Hamiltonian trajectories and duality in the optimal control of linear systems with convex costs*, SIAM J. Control Optim. **27** (1989), no. 5, 1007–1025. [MR 90i:49017](#). [Zbl 682.49019](#).
- [11] R. T. Rockafellar and R. J.-B. Wets, *Variational Analysis*, Grundlehren der Mathematischen Wissenschaften, vol. 317, Springer-Verlag, Berlin, 1998. [MR 98m:49001](#). [Zbl 888.49001](#).
- [12] S. E. Wright, *Consistency of primal-dual approximations for convex optimal control problems*, SIAM J. Control Optim. **33** (1995), no. 5, 1489–1509. [MR 96h:49057](#). [Zbl 876.49026](#).

ADIB BAGH: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA-DAVIS, CA 95616, USA

E-mail address: bagh@math.ucdavis.edu