PERTURBATIONS NEAR RESONANCE FOR THE p -LAPLACIAN IN $\mathbb{R}^N$

TO FU MA AND MAURÍCIO LUCIANO PELICER

Received 10 September 2001

We study a multiplicity result for the perturbed p -Laplacian equation $-\Delta_p u - \lambda g(x)|u|^{p-2}u = f(x, u) + h(x)$ in $\mathbb{R}^N$, where $1 < p < N$ and λ is near λ_1 , the principal eigenvalue of the weighted eigenvalue problem $-\Delta_p u = \lambda g|u|^{p-2}u$ in $\mathbb{R}^N$. Depending on which side λ is from λ_1 , we prove the existence of one or three solutions. This kind of results was firstly obtained by Mawhin and Schmitt (1990) for a semilinear two-point boundary value problem.

1. Introduction

In this paper, we study a class of p -Laplacian equations of the form

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u + f(x, u) + h(x) \quad \text{in } D^{1,p}(\mathbb{R}^N), \quad (1.1)$$

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2}\nabla u)$, $1 < p < N$, and $g \geq 0$ is a weight function. Here, $D^{1,p}(\mathbb{R}^N)$ is the closure of $C_0^\infty(\mathbb{R}^N)$ with respect to the norm

$$\|u\|_{D^{1,p}} = \left(\int_{\mathbb{R}^N} |\nabla u|^p dx \right)^{1/p}. \quad (1.2)$$

This space, which is motivated by the embedding $W^{1,p}(\mathbb{R}^N) \hookrightarrow L^{p^*}(\mathbb{R}^N)$, where $p^* = Np/(N-p)$, is in fact a reflexive Banach space characterized by

$$D^{1,p}(\mathbb{R}^N) = \left\{ u \in L^{p^*}(\mathbb{R}^N); \frac{\partial u}{\partial x_i} \in L^p(\mathbb{R}^N), 1 \leq i \leq N \right\}. \quad (1.3)$$

We refer the reader to Ben-Naoum et al. [5] for a quite complete discussion on the space $D^{1,p}(\mathbb{R}^N)$.

Our study is based on a bifurcation result by Mawhin and Schmitt [15], related to the two-point boundary value problem,

$$-u'' - \lambda u = f(x, u) + h, \quad u(0) = u(\pi) = 0. \quad (1.4)$$

By assuming that f is bounded and satisfying a sign condition, they obtained the following result. If λ is sufficiently near to λ_1 from left, where $\lambda_1 = 1$ is the first eigenvalue of the corresponding linear problem, then (1.4) has at least three solutions. If $1 \leq \lambda < 4$, then problem (1.4) has at least one solution. Some extensions and variations of their result were considered by other authors (cf. Badiale and Lupo [3], Chiappinelli et al. [7], Sanchez [18], and Ma et al. [13]). In [14], the multiplicity part of that result was extended to the p -Laplacian operator in bounded domains, using critical point theory. Our objective is to extend this problem to the p -Laplacian in $\mathbb{R}^N$, with λ approaching to λ_1 from left and from right.

In order to state the Mawhin-Schmitt problem in the context of $D^{1,p}(\mathbb{R}^N)$, we recall some facts about the eigenvalue problem for the weighted p -Laplacian in $\mathbb{R}^N$

$$-\Delta_p u = \lambda g |u|^{p-2} u \quad \text{in } D^{1,p}(\mathbb{R}^N), \quad (1.5)$$

where $g \in L^\infty(\mathbb{R}^N) \cap L^{N/p}(\mathbb{R}^N)$ is a locally Hölder continuous weight function. It is known that for $g \geq 0$, there exists a first eigenvalue $\lambda_1 = \lambda_1(g)$, characterized by

$$\lambda_1 = \inf \left\{ \|u\|_{D^{1,p}}^p; u \in D^{1,p}(\mathbb{R}^N), \int_{\mathbb{R}^N} g |u|^p dx = 1 \right\}, \quad (1.6)$$

which is simple and positive. This implies that

$$\int_{\mathbb{R}^N} |\nabla u|^p dx \geq \lambda_1 \int_{\mathbb{R}^N} g |u|^p dx \quad \forall u \in D^{1,p}(\mathbb{R}^N). \quad (1.7)$$

Besides, the corresponding eigenfunction φ_1 belongs to $D^{1,p}(\mathbb{R}^N) \cap L^\infty(\mathbb{R}^N)$ and may be taken positive (see a complete proof in [12]). Putting

$$W = \left\{ w \in D^{1,p}(\mathbb{R}^N); \int_{\mathbb{R}^N} g |\varphi_1|^{p-2} \varphi_1 w dx = 0 \right\} \quad (1.8)$$

and $V = \text{span}\{\varphi_1\}$, we have from the simplicity of λ_1 ,

$$D^{1,p}(\mathbb{R}^N) = V \oplus W. \quad (1.9)$$

Then, since λ_1 is also isolated (see [11]), we have

$$\lambda_2 := \inf \left\{ \|w\|_{D^{1,p}}^p; w \in W, \int_{\mathbb{R}^N} g |w|^p dx = 1 \right\}, \quad (1.10)$$

which satisfies $\lambda_1 < \lambda_2$. In addition,

$$\int_{\mathbb{R}^N} |\nabla w|^p dx \geq \lambda_2 \int_{\mathbb{R}^N} g|w|^p dx \quad \forall w \in W. \quad (1.11)$$

Next, we make some basic assumptions on the function f . We assume that $f : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory function satisfying the growth condition

$$|f(x, u)| \leq a(x)|u|^{\sigma-1} + b(x), \quad (1.12)$$

with $1 < \sigma < p$, $a \geq 0$, $a \in L^\infty(\mathbb{R}^N) \cap L^{(p^*/\sigma)'}(\mathbb{R}^N)$, and $b \in L^{p^{**}}(\mathbb{R}^N)$. Some of our hypotheses are given upon the primitive $F(x, u) = \int_0^u f(x, s) ds$, namely, there exists $\gamma \in L^1(\mathbb{R}^N)$ such that

$$F(x, u) \geq \gamma(x) \quad \text{a.e. in } \mathbb{R}^N, \quad \forall u \in \mathbb{R}. \quad (1.13)$$

We also consider the following: there exist $\alpha \in L^\infty(\mathbb{R}^N) \cap L^1(\mathbb{R}^N) \cap L^{(p^*/\mu)'}(\mathbb{R}^N)$ and $\beta \in L^\infty(\mathbb{R}^N) \cap L^1(\mathbb{R}^N) \cap L^{p^{**}}(\mathbb{R}^N)$ satisfying

$$pF(x, u) - f(x, u)u \geq \alpha(x)|u|^\mu + \beta(x) \quad \text{a.e. in } \mathbb{R}^N, \quad \forall u \in \mathbb{R}, \quad (1.14)$$

and $1 < \mu \leq \sigma < p$.

Now we are in a position to state our results.

THEOREM 1.1. *Assume that (1.12) and (1.13) hold. If in addition*

$$\lim_{|u| \rightarrow \infty} F(x, u) = +\infty \quad \text{a.e. in } \mathbb{R}^N, \quad (1.15)$$

then for any $h \in L^{p^{**}}(\mathbb{R}^N)$ satisfying

$$\int_{\mathbb{R}^N} h(x)\varphi_1(x) dx = 0, \quad (1.16)$$

problem (1.1) has at least three solutions when λ is sufficiently close to λ_1 from left.

THEOREM 1.2. *Assume that (1.12) and (1.14) hold with $\alpha \geq \max\{a, g\}$. Assume further that $\lambda_1 \leq \lambda < \lambda_2$. Then for any $h \in L^{p^{**}}(\mathbb{R}^N)$ satisfying $|h| \leq \alpha$, problem (1.1) has at least one solution.*

Since (1.14) implies (1.15), under the hypotheses of Theorem 1.2, we get an extension of the original work of Mawhin and Schmitt [15] to the p -Laplacian in $\mathbb{R}^N$. We note that our results do not assume f bounded nor satisfying a sign condition. Theorem 1.2 is related to a class of double resonance problems

introduced in [6] for semilinear elliptic equations. It was not considered for the p -Laplacian, even in bounded domains. Condition (1.14) was early used in [1, 8, 10] for example, as an Ambrosetti-Rabinowitz type condition [2]. A simple example of g satisfying all the hypotheses of both theorems is

$$f(x, u) = \sigma a(x) |u|^{\sigma-2} u, \quad (1.17)$$

where $a \in L^\infty(\mathbb{R}^N) \cap L^1(\mathbb{R}^N) \cap L^{(p^*/\sigma)'}(\mathbb{R}^N)$ and $1 < \mu = \sigma < p - 1$.

The proofs of the theorems are given in Section 3. In Section 2, we present some preliminary results on the variational setting of the p -Laplacian equations in $D^{1,p}(\mathbb{R}^N)$ and the related Palais-Smale compactness.

2. Preliminaries

We begin with some standard facts upon the variational formulation of problem (1.1). Let $J_\lambda : D^{1,p}(\mathbb{R}^N) \rightarrow \mathbb{R}$ be the functional defined by

$$J_\lambda(u) = \int_{\mathbb{R}^N} \left[\frac{1}{p} |\nabla u(x)|^p - \frac{\lambda}{p} g(x) |u(x)|^p - F(x, u(x)) - h(x)u(x) \right] dx. \quad (2.1)$$

It is proved in do Ó [10], that J_λ is of class $C^1(\mathbb{R}^N)$ and

$$\begin{aligned} \langle J'_\lambda(u), \varphi \rangle &= \int_{\mathbb{R}^N} |\nabla u|^{p-2} \nabla u \cdot \nabla \varphi dx - \lambda \int_{\mathbb{R}^N} g |u|^{p-2} u \varphi dx \\ &\quad - \int_{\mathbb{R}^N} f(x, u) \varphi dx - \int_{\mathbb{R}^N} h \varphi dx, \end{aligned} \quad (2.2)$$

for all $\varphi \in D^{1,p}(\mathbb{R}^N)$. In addition, the critical points of J_λ are precisely the weak solutions of (1.1).

Next we recall a compactness result which is proved in [5].

LEMMA 2.1 (see [5]). *The functional*

$$u \mapsto \int_{\mathbb{R}^N} m(x) |u(x)|^q dx \quad (2.3)$$

is well defined and weakly continuous in $D^{1,p}(\mathbb{R}^N)$, for $1 \leq q < p^$ and $m \in L^{(p^*/q)'}(\mathbb{R}^N)$.*

As a consequence, under the conditions of the lemma, there exists $C > 0$ such that

$$\int_{\mathbb{R}^N} m(x) |u(x)|^q dx \leq \|m\|_{L^{(p^*/q)'}(\mathbb{R}^N)} \|u\|_{L^{p^*}}^q \leq C \|u\|_{D^{1,p}}^q. \quad (2.4)$$

LEMMA 2.2. Assume that (1.12) holds. Then the Nemytskii mapping,

$$u \mapsto f(x, u) \quad (2.5)$$

is compact from $D^{1,p}(\mathbb{R}^N)$ to $L^{p^*}(\mathbb{R}^N)$.

Proof. Put $r = p^{*'}$ and $q = (\sigma - 1)r$ so that $q < p^*$ and

$$\left(\frac{p^*}{\sigma}\right)' = r \left(\frac{p^*}{q}\right)'. \quad (2.6)$$

Then we get from (1.12) that $a^r \in L^{(p^*/q)'}(\mathbb{R}^N)$. Now let (u_n) be a sequence such that $u_n \rightharpoonup u$ weakly for some $u \in D^{1,p}(\mathbb{R}^N)$. Then from Lemma 2.1 we have $a^{r/q}u_n \rightarrow a^{r/q}u$ strongly in $L^q(\mathbb{R}^N)$. It follows that

$$a^{r/q}u_n \rightarrow a^{r/q}u, \quad |a^{r/q}u_n| \leq k \quad \text{a.e. in } \mathbb{R}^N, \quad (2.7)$$

for some $k \in L^q(\mathbb{R}^N)$. Hence, for all n and a.e. $x \in \mathbb{R}^N$,

$$|f(x, u_n(x))|^r \leq 2^r \left(a(x)^r |u_n(x)|^{(\sigma-1)r} + |b(x)|^r \right) \leq 2^r \left(|k(x)|^q + |b(x)|^r \right). \quad (2.8)$$

Since the last term is an integrable function, from Lebesgue theorem, we infer that $f(x, u_n) \rightarrow f(x, u)$ strongly in $L^r(\mathbb{R}^N)$. $\square$

Next, we do some remarks about the Palais-Smale condition for J_λ . We recall that J_λ is said to satisfy the Palais-Smale condition at level c , $(PS)_c$, if every sequence for which

$$J(u_n) \rightarrow c, \quad \|J'(u_n)\|_{(D^{1,p})^*} \rightarrow 0 \quad (2.9)$$

possesses a convergent subsequence. When J satisfies $(PS)_c$ for all $c \in \mathbb{R}^N$, we simply say that J satisfies the (PS) condition. In Theorem 1.2, we use a weaker version of the (PS) condition due to Cerami (cf. [4]). We say that J satisfies the Palais-Smale-Cerami condition, (PSC), if every sequence, for which

$$J(u_n) \text{ is bounded,} \quad \left(1 + \|u_n\|_{D^{1,p}}\right) \|J'(u_n)\|_{(D^{1,p})^*} \rightarrow 0, \quad (2.10)$$

possesses a convergent subsequence.

LEMMA 2.3. Assume that condition (1.12) holds. Then any bounded sequence satisfying (2.9) or (2.10) possesses a convergent subsequence.

Proof. Let (u_n) be a bounded sequence satisfying (2.9). Then, passing to a subsequence if necessary, there exists $u \in D^{1,p}(\mathbb{R}^N)$ such that $u_n \rightharpoonup u$ weakly in $D^{1,p}(\mathbb{R}^N)$ and also in $L^{p^*}(\mathbb{R}^N)$. Consequently,

$$\lim_{n \rightarrow \infty} \langle J'_\lambda(u_n), u_n - u \rangle = 0. \quad (2.11)$$

On the other hand, from [Lemma 2.2](#), we know that $f(x, u_n) \rightarrow f(x, u)$ strongly in $L^{p^{**}}(\mathbb{R}^N)$ and therefore,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} (f(x, u_n) + h)(u_n - u) dx = 0. \quad (2.12)$$

Noting that

$$\int_{\mathbb{R}^N} g |u_n|^{p-1} |u_n - u| dx \leq \left(\int_{\mathbb{R}^N} g |u_n|^p dx \right)^{1/p'} \left(\int_{\mathbb{R}^N} g |u_n - u|^p dx \right)^{1/p}, \quad (2.13)$$

and since $g \in L^{(p^*/p)'}$, [Lemma 2.1](#) implies that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} g |u_n|^{p-2} u_n (u_n - u) dx = 0. \quad (2.14)$$

Combining (2.11) with (2.12) and (2.14), we get

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u_n|^{p-2} \nabla u_n \cdot \nabla (u_n - u) dx = 0. \quad (2.15)$$

But since we also have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u|^{p-2} \nabla u \cdot \nabla (u_n - u) dx = 0, \quad (2.16)$$

it follows that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} (|\nabla u_n|^{p-2} \nabla u_n - |\nabla u|^{p-2} \nabla u) \cdot \nabla (u_n - u) dx = 0. \quad (2.17)$$

Then from a well-known argument based on the Clarkson inequality (cf. Tolksdorf [19]), we conclude that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} |\nabla u_n - \nabla u|^p dx = 0. \quad (2.18)$$

This completes the proof since (2.10) implies (2.9). $\square$

3. Proofs of Theorems 1.1 and 1.2

The proof of [Theorem 1.1](#) is based on the Ekeland's variational principle and the Ambrosetti-Rabinowitz Mountain-Pass theorem [2]. [Theorem 1.2](#) is proved using the saddle point theorem of Rabinowitz [17].

Proof of Theorem 1.1. We divide the proof in several steps.

Step 1 (the coerciveness of J_λ). Since $\lambda < \lambda_1$ and (1.12) holds, from (1.7) and (2.4) we get

$$J_\lambda(u) \geq \left(\frac{\lambda_1 - \lambda}{p\lambda_1} \right) \|u\|_{D^{1,p}}^p - C \|u\|_{D^{1,p}}^\sigma - C \|u\|_{D^{1,p}}, \quad (3.1)$$

where $C > 0$ denotes several constants. Then J_λ is coercive as a consequence of the assumption that $1 < \sigma < p$. This implies that any sequence satisfying (2.9) must be bounded, and therefore Lemma 2.2 implies that J_λ satisfies the $(PS)_c$ for all $c \in \mathbb{R}$. Similarly, from (1.11),

$$J_{\lambda_1}(w) \geq \left(\frac{\lambda_2 - \lambda_1}{p\lambda_2} \right) \|w\|_{D^{1,p}}^p - C\|w\|_{D^{1,p}}^\sigma - C\|w\|_{D^{1,p}}, \quad (3.2)$$

which shows that J_{λ_1} is coercive in W . Noting that $J_{\lambda_1} \leq J_\lambda$ for all $\lambda < \lambda_1$, we have that

$$m = \inf_W J_{\lambda_1} \leq \inf_W J_\lambda. \quad (3.3)$$

Step 2 (estimating J_λ in V). From (1.16) we have for $t \in \mathbb{R}$,

$$J_\lambda(t\varphi_1) = \left(\frac{\lambda_1 - \lambda}{p} \right) \int_{\mathbb{R}^N} |t\varphi_1(x)|^p dx - \int_{\mathbb{R}^N} F(x, t\varphi_1(x)) dx. \quad (3.4)$$

Now, from (1.13) there exist constants $R, C > 0$ such that

$$\int_{|x|>R} F(x, t\varphi_1(x)) dx \geq \int_{|x|>R} \gamma(x) dx \geq -C, \quad \forall t \in \mathbb{R}. \quad (3.5)$$

Choosing $t^+ > 0$ sufficiently large, we get from (1.15) that

$$\int_{|x| \leq R} F(x, t^+\varphi_1(x)) dx > -m + C. \quad (3.6)$$

Then we have

$$\int_{\mathbb{R}^N} F(x, t^+\varphi_1(x)) dx > -m, \quad (3.7)$$

so that

$$J_\lambda(t^+\varphi_1) \leq \left(\frac{\lambda_1 - \lambda}{p} \right) \int_{\mathbb{R}^N} |t^+\varphi_1(x)|^p dx + m. \quad (3.8)$$

Then for λ sufficiently near to λ_1 , $J_\lambda(t^+\varphi_1) < m$. The same conclusion holds for a $t^- < 0$.

Step 3 (the existence of the first two solutions). Put

$$\mathbb{O}^\pm = \{u \in D^{1,p}(\mathbb{R}^N); u = \pm t\varphi_1 + w \text{ with } t > 0, w \in W\}. \quad (3.9)$$

Then from Step 2, for λ sufficiently near to λ_1 ,

$$-\infty < \inf_{\mathbb{O}^\pm} I_\lambda < m. \quad (3.10)$$

Now let $u_n \in \mathbb{O}^+$ be a sequence satisfying (2.9) for $c < m$. Then from coerciveness of J_λ , (u_n) has a convergent subsequence, say, (u_n) itself. Noting that $W = \partial\mathbb{O}^+$ and $\inf_W J_\lambda \geq m$ (Step 1), we conclude that (u_n) converges to an interior point $u \in \mathbb{O}^+$. This means that J_λ satisfies the $(PS)_c$ condition inside $\mathbb{O}^+$ for all $c < m$. Then applying the Ekeland variational principle in $\mathbb{O}^+$, we see that J_λ has a critical point u^+ as a local minimum in $\mathbb{O}^+$. (See complete argument in [16].) Similarly, we obtain a critical point u^- of J_λ in $\mathbb{O}^-$. Taking into account that $\mathbb{O}^- \cap \mathbb{O}^+ = \emptyset$, the existence of two weak solutions of (1.1) is proved.

Step 4 (the third solution). To fix ideas, suppose that $J_\lambda(u^+) \leq J_\lambda(u^-)$. If u^- is not an isolated critical point, then J_λ has at least three solutions. Otherwise, putting

$$I(u) = J_\lambda(u + u^-) - J_\lambda(u^-), \quad e = u^+ - u^-, \quad (3.11)$$

we have that $I(0) = 0$, $I(e) \leq 0$, and there exist $r, \rho > 0$ such that $I(u) \geq \rho$ if $\|u\|_{D^{1,p}} = r$. Then, since $I' = J'_\lambda$ and I also satisfies the (PS) condition, from the Mountain-Pass theorem, the number

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} J_\lambda(\gamma(t)), \quad (3.12)$$

where

$$\Gamma = \{\gamma \in C([0,1], D^{1,p}(\mathbb{R}^N)); \gamma(0) = u^-, \gamma(1) = u^+\} \quad (3.13)$$

is a critical value of J_λ . Noting that all paths joining u^- to u^+ pass through W , we have $c \geq m$. Therefore we have obtained a third critical point of J_λ . The proof is now complete. $\square$

Proof of Theorem 1.2. The proof is based on the arguments from [8, 10].

Step 1 (the growth of F). We prove that for some $C_1, C_2 > 0$,

$$\int_{\mathbb{R}^N} F(x, t\varphi_1) dx \geq C_1 \|t\varphi_1\|_{D^{1,p}}^\mu - C_2. \quad (3.14)$$

In fact, from (1.14) we have

$$\frac{d}{du} \left(\frac{F(x, u)}{|u|^p} \right) \leq -\alpha(x)|u|^{\mu-p-2}u - \beta(x)|u|^{-p-2}u \quad (u > 0). \quad (3.15)$$

Integrating from $u > 0$ to $+\infty$, and noting that $F(x, \theta)/(\theta^p) \rightarrow 0$ as $\theta \rightarrow \infty$, we get

$$F(x, u) \geq \frac{\alpha(x)}{p-\mu} |u|^\mu + \frac{\beta(x)}{p}. \quad (3.16)$$

Since this inequality holds for $u < 0$, we have

$$\int_{\mathbb{R}^N} F(x, t\varphi_1) dx \geq C|t|^\mu - C_2, \quad (3.17)$$

and inequality (3.14) follows.

Step 2 (the (PSC) condition). Let (u_n) be a sequence satisfying (2.9). Then from Lemma 2.2, it suffices to prove that (u_n) is bounded. In fact, first we note that (u_n) satisfies

$$\begin{aligned} \langle J'_\lambda(u_n), u_n \rangle - pJ_\lambda(u_n) &= \int_{\mathbb{R}^N} [pF(x, u_n) - f(x, u_n)u_n + (p-1)hu_n] dx \\ &\geq \int_{\mathbb{R}^N} \alpha|u|^\mu dx + \int_{\mathbb{R}^N} \beta dx + (p-1) \int_{\mathbb{R}^N} hu_n dx. \end{aligned} \quad (3.18)$$

Now, since $|h| \leq \alpha$,

$$\left| \int_{\mathbb{R}^N} hu_n dx \right| \leq \int_{\mathbb{R}^N} \alpha |u_n| dx \leq \|\alpha\|_{L^1}^{1/\mu'} \left(\int_{\mathbb{R}^N} \alpha |u_n|^\mu dx \right)^{1/\mu}. \quad (3.19)$$

Then from the boundedness of $\langle J'_\lambda(u_n), u_n \rangle - pJ_\lambda(u_n)$, we deduce that

$$\int_{\mathbb{R}^N} \alpha |u_n|^\mu dx \leq C + C \left(\int_{\mathbb{R}^N} \alpha |u_n|^\mu dx \right)^{1/\mu}, \quad (3.20)$$

so that

$$\int_{\mathbb{R}^N} \alpha |u_n|^\mu dx \leq C. \quad (3.21)$$

Now we use an interpolation inequality. Since $0 < \mu < p < p^*$, there exists $t \in (0, 1)$ such that

$$1 = \frac{p(1-t)}{\mu} + \frac{pt}{p^*}. \quad (3.22)$$

Then from Hölder inequality,

$$\begin{aligned} \int_{\mathbb{R}^N} \alpha(x) |u|^p dx &= \int_{\mathbb{R}^N} (\alpha^{1/\mu} |u|)^{p(1-t)} (\alpha^{1/p^*} |u|)^{pt} dx \\ &\leq \left(\int_{\mathbb{R}^N} \alpha |u|^\mu dx \right)^{p(1-t)/\mu} \left(\int_{\mathbb{R}^N} \alpha |u|^{p^*} dx \right)^{pt/p^*}. \end{aligned} \quad (3.23)$$

Using (3.21) and (2.4),

$$\int_{\mathbb{R}^N} \alpha(x) |u_n(x)|^p dx \leq C \|u_n\|_{D^{1,p}}^{tp}. \quad (3.24)$$

Taking into account the boundedness of $J_\lambda(u_n)$,

$$\frac{1}{p} \|u_n\|_{D^{1,p}}^p \leq C + \frac{\lambda}{p} \int_{\mathbb{R}^N} g |u_n|^p dx + \int_{\mathbb{R}^N} F(x, u_n) dx + \int_{\mathbb{R}^N} h u_n dx, \quad (3.25)$$

and since $\alpha \geq \max\{a, g\}$,

$$\begin{aligned} \frac{1}{p} \|u_n\|_{D^{1,p}}^p &\leq C + \frac{\lambda}{p} \int_{\mathbb{R}^N} \alpha |u_n|^p dx + \frac{1}{\sigma} \int_{\mathbb{R}^N} \alpha |u_n|^\sigma dx \\ &\quad + \int_{\mathbb{R}^N} b |u_n| dx + \int_{\mathbb{R}^N} h u_n dx. \end{aligned} \quad (3.26)$$

Consequently, if $\mu = \sigma$ we have, from (3.24),

$$\|u_n\|_{D^{1,p}}^p \leq C \left(1 + \|u_n\|_{D^{1,p}}^{tp} + \|u_n\|_{D^{1,p}} \right). \quad (3.27)$$

Otherwise, we have $\mu < \sigma < p^*$, and as before, we get $s \in (0, 1)$ such that

$$\int_{\mathbb{R}^N} \alpha |u_n|^\sigma dx \leq C \|u_n\|_{D^{1,p}}^{s\sigma}. \quad (3.28)$$

Then

$$\|u_n\|_{D^{1,p}}^p \leq C \left(1 + \|u_n\|_{D^{1,p}}^{tp} + \|u_n\|_{D^{1,p}}^{s\sigma} + \|u_n\|_{D^{1,p}} \right). \quad (3.29)$$

In both cases, we see that $\|u_n\|_{D^{1,p}}$ is uniformly bounded.

Step 3 (the saddle point theorem). It is well known that the (PS) condition can be replaced by the (PSC) condition in the saddle point theorem of Rabinowitz (see [4, 17]). Then to conclude that J_λ has a critical point it suffices to show that

$$\lim_{\|v\|_{D^{1,p}} \rightarrow \infty} J_\lambda(v) = -\infty, \quad \lim_{\|w\|_{D^{1,p}} \rightarrow \infty} J_\lambda(w) = +\infty, \quad (3.30)$$

where $v \in V$ and $w \in W$, as defined in (1.9). Now, from (3.14),

$$J_\lambda(t\varphi_1) \leq -\left(\frac{\lambda - \lambda_1}{p\lambda_1}\right) \|t\varphi_1\|_{D^{1,p}}^p - C_1 \|t\varphi_1\|_{D^{1,p}}^\mu + C \|t\varphi_1\|_{D^{1,p}} + C_2. \quad (3.31)$$

Since $\lambda \geq \lambda_1$, the first part of (3.30) holds. Finally, since $\lambda < \lambda_2$, the argument in [Step 1](#) of the proof of [Theorem 1.1](#) implies the second statement of (3.30). The proof is now complete. $\square$

Note 3.1. Just before the completion of this paper, we noticed that P. De Nápoli and M. C. Mariani [9] studied problem (1.1) in the same framework of our [Theorem 1.1](#). However, they considered only the case $\lambda \rightarrow \lambda_1$ from left. Our assumptions on f are slightly more general.

Acknowledgments

T. F. Ma was partially supported by CNPq/Brazil. M. L. Pelicer was partially supported by CAPES-PICD/Brazil.

References

- [1] C. O. Alves, P. C. Carrião, and O. H. Miyagaki, *Existence and multiplicity results for a class of resonant quasilinear elliptic problems on $\mathbb{R}^N$* , *Nonlinear Anal.* **39** (2000), no. 1, Ser. A: Theory Methods, 99–110.
- [2] A. Ambrosetti and P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, *J. Functional Analysis* **14** (1973), 349–381.
- [3] M. Badiale and D. Lupo, *Some remarks on a multiplicity result by Mawhin and Schmitt*, *Acad. Roy. Belg. Bull. Cl. Sci. (5)* **65** (1989), no. 6-9, 210–224.
- [4] P. Bartolo, V. Benci, and D. Fortunato, *Abstract critical point theorems and applications to some nonlinear problems with “strong” resonance at infinity*, *Nonlinear Anal.* **7** (1983), no. 9, 981–1012.
- [5] A. K. Ben-Naoum, C. Troestler, and M. Willem, *Extrema problems with critical Sobolev exponents on unbounded domains*, *Nonlinear Anal.* **26** (1996), no. 4, 823–833.
- [6] H. Berestycki and D. G. de Figueiredo, *Double resonance in semilinear elliptic problems*, *Comm. Partial Differential Equations* **6** (1981), no. 1, 91–120.
- [7] R. Chiappinelli, J. Mawhin, and R. Nugari, *Bifurcation from infinity and multiple solutions for some Dirichlet problems with unbounded nonlinearities*, *Nonlinear Anal.* **18** (1992), no. 12, 1099–1112.
- [8] D. G. Costa and C. A. Magalhães, *Existence results for perturbations of the p -Laplacian*, *Nonlinear Anal.* **24** (1995), no. 3, 409–418.
- [9] P. De Nápoli and M. C. Mariani, *Three solutions for quasilinear equations in $\mathbb{R}^n$ near resonance*, *Proceedings of the USA-Chile Workshop on Nonlinear Analysis (Viña del Mar-Valparaíso, 2000)*, Southwest Texas State University, Texas, 2001, pp. 131–140.
- [10] J. M. B. do Ó, *Solutions to perturbed eigenvalue problems of the p -Laplacian in $\mathbb{R}^N$* , *Electron. J. Differential Equations* **1997** (1997), no. 11, 1–15.
- [11] P. Drábek and Y. X. Huang, *Bifurcation problems for the p -Laplacian in $\mathbb{R}^N$* , *Trans. Amer. Math. Soc.* **349** (1997), no. 1, 171–188.
- [12] J. Fleckinger, R. F. Manásevich, N. M. Stavrakakis, and F. de Thélin, *Principal eigenvalues for some quasilinear elliptic equations on $\mathbb{R}^N$* , *Adv. Differential Equations* **2** (1997), no. 6, 981–1003.
- [13] T. F. Ma, M. Ramos, and L. Sanchez, *Multiple solutions for a class of nonlinear boundary value problems near resonance: a variational approach*, *Nonlinear Anal.* **30** (1997), no. 6, 3301–3311.
- [14] T. F. Ma and L. Sanchez, *Three solutions of a quasilinear elliptic problem near resonance*, *Math. Slovaca* **47** (1997), no. 4, 451–457.
- [15] J. Mawhin and K. Schmitt, *Nonlinear eigenvalue problems with the parameter near resonance*, *Ann. Polon. Math.* **51** (1990), 241–248.
- [16] N. Mizoguchi, *Existence of nontrivial solutions of partial differential equations with discontinuous nonlinearities*, *Nonlinear Anal.* **16** (1991), no. 11, 1025–1034.
- [17] P. H. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, vol. 65, Conference Board of the Mathematical Sciences, Amer. Math. Soc., Rhode Island, 1986.

- [18] L. Sanchez, *Boundary value problems for some fourth order ordinary differential equations*, Appl. Anal. **38** (1990), no. 3, 161–177.
- [19] P. Tolksdorf, *On quasilinear boundary value problems in domains with corners*, Nonlinear Anal. **5** (1981), no. 7, 721–735.

TO FU MA: DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE ESTADUAL DE MARINGÁ,
87020-900 MARINGÁ, PR, BRAZIL

E-mail address: matofu@uem.br

MAURÍCIO LUCIANO PELICER: DEPARTAMENTO DE CIÊNCIAS, UNIVERSIDADE ESTADUAL
DE MARINGÁ, 87360-000 GOIOERÊ, PR, BRAZIL

E-mail address: mlpelicer@uem.br