A REMARK ON THE APPROXIMATE FIXED-POINT PROPERTY

TADEUSZ KUCZUMOW

Received 30 November 2001

We give an example of an unbounded, convex, and closed set C in the Hilbert space l^2 with the following two properties: (i) C has the approximate fixed-point property for nonexpansive mappings, (ii) C is not contained in a block for every orthogonal basis in l^2 .

1. Introduction

In [6], Goebel and the author observed that some unbounded sets in Hilbert spaces have the approximate fixed-point property for nonexpansive mappings. Namely, they proved that every closed convex set C , which is contained in a block, has the approximate fixed-point property for nonexpansive mappings (AFPP). This result was extended by Ray [14] to all linearly bounded subsets of l_p , $1 < p < \infty$. Next, he proved that a closed convex subset C of a real Hilbert space has the fixed-point property for nonexpansive mappings if and only if it is bounded [15]. The first result of Ray [14] was generalized by Reich [16] (for other results of this type see [1, 2, 4, 5, 7, 8, 9, 10, 11, 12, 13, 17, 19]). Reich [16] proved the following remarkable theorem: a closed, convex subset of a reflexive Banach space has the AFPP if and only if it is linearly bounded. Next, Shafrir [18] introduced the notion of a directionally bounded set. Using this concept, he proved two important theorems [18].

(1) A convex subset C of a Banach space X has the AFPP if and only if C is directionally bounded.

(2) For a Banach space X , the following two conditions are equivalent: (i) X is reflexive; (ii) every closed, convex, and linearly bounded subset C of X is directionally bounded.

Therefore, the following statements are equivalent: (a) X is reflexive; (b) a closed, convex subset C of X has the AFPP if and only if C is linearly bounded. This result is strictly connected with the above-mentioned Reich theorem [16].

Now, it is worth to note that, recently, there is a return to study the AFPP First, Espínola and Kirk [3] published a paper about the AFPP in the product spaces. They proved that the product space $D = (M \times C)_\infty$ has the AFPP for nonexpansive mappings whenever M is a metric space which has the AFPP for such mappings and C is a bounded, convex subset of a Banach space. Next, Wiśnicki wrote a paper about a common approximate fixed-point sequence for two commuting nonexpansive mappings (see [20] for details). Therefore, the author decided to publish an example of a set which is closely related to the AFPP. Namely, it is obvious that every blockable set in l^2 is linearly bounded, but there are linearly bounded sets in l^2 which are not contained in any block with respect to an arbitrary basis. This was mentioned in [6] but never published. The aim of this paper is to show the construction of such a set.

2. Preliminaries

Throughout this paper, l^2 is real, $\langle \cdot, \cdot \rangle$ denotes the scalar product in l^2 , and $\{e_n\}$ is the standard basis in l^2 .

For any nonempty set $K \subset l^2$, the closed convex hull of K is denoted by $\text{conv} K$.

Let C be a nonempty subset of a Banach space X . A mapping $T : C \rightarrow C$ is said to be nonexpansive if for each $x, y \in C$,

$$\|T(x) - T(y)\| \leq \|x - y\|. \quad (2.1)$$

A convex subset C of a Banach space X has the approximate fixed-point property (AFPP) if each nonexpansive $T : C \rightarrow C$ satisfies

$$\inf \{\|x - T(x)\| : x \in C\} = 0. \quad (2.2)$$

It is obvious that bounded convex sets always have the AFPP.

A set $K \subset l^2$ is said to be a block in the orthogonal basis $\{\tilde{e}_n\}$ if K is of the form

$$K = \{x \in l^2 : |\langle x, \tilde{e}_n \rangle| \leq M_n, n = 1, 2, \dots\}, \quad (2.3)$$

where $\{M_n\}$ is a sequence of positive reals.

The set $C \subset l^2$ is called a block set if there exists a block $K \subset l^2$ such that C is a subset of K .

A subset C of a Banach space X is linearly bounded if C has bounded intersections with all lines in X .

3. The construction

Let $\{k_n\}_{n=2}^\infty$ and $\{l_n\}_{n=2}^\infty$ be two sequences of positive reals such that

$$\sum_{n=2}^\infty \frac{k_n}{l_n} < +\infty, \quad \lim_n k_n = +\infty. \quad (3.1)$$

For example, we may take $k_n = n$ and $l_n = n^3$ for $n = 2, 3, \dots$. Next, we set

$$a_n = k_n e_1 + l_n e_n, \quad b_n = -k_n e_1 + l_n e_n, \quad (3.2)$$

for $n = 2, 3, \dots$, and finally,

$$C = \text{conv} \{x \in l^2 : \exists n \geq 2 (x = a_n \vee x = b_n)\}. \quad (3.3)$$

THEOREM 3.1. *If*

$$x = \sum_{n=1}^\infty c_n e_n = c_1 e_1 + \sum_{n=2}^\infty d_n l_n e_n = c_1 e_1 + \bar{x} \quad (3.4)$$

is an element of the set C, then

$$d_n \geq 0 \quad (3.5)$$

for $n = 2, 3, \dots$,

$$\sum_{n=2}^\infty d_n \leq 1, \quad (3.6)$$

and there exist sequences $\{\alpha_n\}_{n=2}^\infty$ and $\{\beta_n\}_{n=2}^\infty$ such that

$$c_1 = \sum_{n=2}^\infty (\alpha_n k_n - \beta_n k_n), \quad \alpha_n, \beta_n \geq 0, \quad \alpha_n + \beta_n = d_n, \quad (3.7)$$

for $n = 2, 3, \dots$. Additionally, there exists a positive constant $M_{\bar{x}}$ such that

$$0 \leq (\alpha_n + \beta_n) k_n = d_n k_n \leq M_{\bar{x}} \frac{k_n}{l_n} \quad (3.8)$$

for $n = 2, 3, \dots$.

Proof. Set

$$\bar{x} = \sum_{n=2}^\infty c_n e_n = \sum_{n=2}^\infty d_n l_n e_n. \quad (3.9)$$

Observe that, there exists a sequence $\{x_j\}_{j=1}^{\infty}$ such that

$$x = \lim_j x_j \quad (3.10)$$

with

$$\begin{aligned} x_j &= \sum_{n=2}^{\infty} (\alpha_{nj}a_n + \beta_{nj}b_n) \\ &= \sum_{n=2}^{\infty} (\alpha_{nj}k_n - \beta_{nj}k_n)e_1 + \sum_{n=2}^{\infty} (\alpha_{nj}l_n + \beta_{nj}l_n)e_n \\ &= \sum_{n=2}^{\infty} (\alpha_{nj}k_n - \beta_{nj}k_n)e_1 + \bar{x}_j \in C, \end{aligned} \quad (3.11)$$

where

$$\bar{x}_j = \sum_{n=2}^{\infty} (\alpha_{nj}l_n + \beta_{nj}l_n)e_n, \quad \alpha_{nj}, \beta_{nj} \geq 0, \quad \sum_{n=2}^{\infty} (\alpha_{nj} + \beta_{nj}) = 1. \quad (3.12)$$

Without loss of generality, we can assume that $\{\alpha_{nj}\}_{j=1}^{\infty}$ and $\{\beta_{nj}\}_{j=1}^{\infty}$ tend to α_n and β_n , respectively, for $n = 2, 3, \dots$. Hence, we have

$$c_1 = \sum_{n=2}^m (\alpha_n k_n - \beta_n k_n) + \lim_j \sum_{n=m+1}^{\infty} (\alpha_{nj} k_n - \beta_{nj} k_n) \quad (3.13)$$

for each $m \geq 2$. On the other hand,

$$\bar{x} = \lim_j \bar{x}_j = \lim_j \sum_{n=2}^{\infty} (\alpha_{nj}l_n + \beta_{nj}l_n)e_n \quad (3.14)$$

and, therefore, there exists a constant $0 < M_{\bar{x}} < +\infty$ such that

$$\alpha_{nj}l_n + \beta_{nj}l_n \leq M_{\bar{x}} \quad (3.15)$$

for all $n \geq 2$ and $j \in \mathbb{N}$. This implies that

$$\begin{aligned} 0 \leq \alpha_{nj}k_n + \beta_{nj}k_n &= (\alpha_{nj}l_n + \beta_{nj}l_n) \frac{k_n}{l_n} \leq M_{\bar{x}} \frac{k_n}{l_n}, \\ 0 \leq (\alpha_n + \beta_n)k_n &= d_n k_n \leq M_{\bar{x}} \frac{k_n}{l_n}, \end{aligned} \quad (3.16)$$

for all j, n , and finally,

$$\begin{aligned} \sup_j \left| \sum_{n=m+1}^{\infty} (\alpha_{nj}k_n - \beta_{nj}k_n) \right| &\leq \sup_j \sum_{n=m+1}^{\infty} (\alpha_{nj}k_n + \beta_{nj}k_n) \\ &\leq \sum_{n=m+1}^{\infty} M_{\bar{x}} \frac{k_n}{l_n} = M_{\bar{x}} \sum_{n=m+1}^{\infty} \frac{k_n}{l_n} \xrightarrow{m \rightarrow \infty} 0. \end{aligned} \quad (3.17)$$

Combining (3.13) with (3.17), we conclude that

$$c_1 = \sum_{n=2}^{\infty} (\alpha_n k_n - \beta_n k_n). \quad (3.18)$$

This completes the proof. $\square$

THEOREM 3.2. *The set C is linearly bounded but is not a block set in any orthogonal basis in l^2 .*

Proof. First, we show that C is not a block set in any orthogonal basis,

$$\{\tilde{e}_i\}_{i=1}^{\infty} = \left\{ \sum_{n=1}^{\infty} c_{in} e_n \right\}_{i=1}^{\infty} \quad (3.19)$$

in l^2 . Indeed, there exists i_0 such that $c_{i_0 1} \neq 0$. Since we have

$$\max(|\langle a_n, \tilde{e}_{i_0} \rangle|, |\langle b_n, \tilde{e}_{i_0} \rangle|) = k_n |c_{i_0 1}| + l_n |c_{i_0 n}| \quad (3.20)$$

for every $n \geq 2$, these two facts imply that

$$\sup \{ |\langle x, \tilde{e}_{i_0} \rangle| : x \in C \} = +\infty. \quad (3.21)$$

Therefore, C is not a block set in $\{\tilde{e}_i\}_{i=1}^{\infty}$.

Now, we prove that the set C is linearly bounded. We begin with the following simple observation:

$$\sup \{ |\langle x, e_n \rangle| : x \in C \} \leq l_n \quad (3.22)$$

for $n = 2, 3, \dots$. Next, if $x \in C$ is of the form

$$x = \sum_{n=1}^{\infty} c_n e_n = c_1 e_1 + \sum_{n=2}^{\infty} d_n l_n e_n = c_1 e_1 + \bar{x}, \quad (3.23)$$

then, by Theorem 3.1, we see that

$$d_n \geq 0 \quad (3.24)$$

for $n = 2, 3, \dots$,

$$\sum_{n=2}^{\infty} d_n \leq 1, \quad (3.25)$$

and there exist sequences $\{\alpha_n\}_{n=2}^\infty$ and $\{\beta_n\}_{n=2}^\infty$ such that

$$c_1 = \sum_{n=2}^{\infty} (\alpha_n k_n - \beta_n k_n), \quad \alpha_n, \beta_n \geq 0, \quad \alpha_n + \beta_n = d_n, \quad (3.26)$$

for $n = 2, 3, \dots$. Additionally, there exists a positive constant $M_{\bar{x}}$ such that

$$0 \leq (\alpha_n + \beta_n) k_n = d_n k_n \leq M_{\bar{x}} \frac{k_n}{l_n} \quad (3.27)$$

for $n = 2, 3, \dots$. Hence, we obtain

$$|c_1| = \left| \sum_{n=2}^{\infty} (\alpha_n k_n - \beta_n k_n) \right| \leq \sum_{n=2}^{\infty} (\alpha_n + \beta_n) k_n \leq M_{\bar{x}} \sum_{n=2}^{\infty} \frac{k_n}{l_n}. \quad (3.28)$$

Then, it follows from (3.22) and (3.28) that an intersection of C with any line $\{y + tv : t \in \mathbb{R}\}$, where $y, v \in l^2$ and $v \neq 0$, is either empty or bounded which completes the proof. $\square$

References

- [1] A. Canetti, G. Marino, and P. Pietramala, *Fixed point theorems for multivalued mappings in Banach spaces*, Nonlinear Anal. **17** (1991), no. 1, 11–20.
- [2] A. Carbone and G. Marino, *Fixed points and almost fixed points of nonexpansive maps in Banach spaces*, Riv. Mat. Univ. Parma (4) **13** (1987), 385–393.
- [3] R. Espínola and W. A. Kirk, *Fixed points and approximate fixed points in product spaces*, Taiwanese J. Math. **5** (2001), no. 2, 405–416.
- [4] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge Studies in Advanced Mathematics, vol. 28, Cambridge University Press, Cambridge, 1990.
- [5] ———, *Classical theory of nonexpansive mappings*, Handbook of Metric Fixed Point Theory (W. A. Kirk and B. Sims, eds.), Kluwer Academic Publishers, Dordrecht, 2001, pp. 49–91.
- [6] K. Goebel and T. Kuczumow, *A contribution to the theory of nonexpansive mappings*, Bull. Calcutta Math. Soc. **70** (1978), no. 6, 355–357.
- [7] K. Goebel and S. Reich, *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 83, Marcel Dekker, New York, 1984.
- [8] W. A. Kirk, *Fixed point theory for nonexpansive mappings*, Fixed Point Theory (Sherbrooke, Que., 1980), Lecture Notes in Math., vol. 886, Springer, Berlin, 1981, pp. 484–505.
- [9] W. A. Kirk and W. O. Ray, *Fixed-point theorems for mappings defined on unbounded sets in Banach spaces*, Studia Math. **64** (1979), no. 2, 127–138.
- [10] G. Marino, *Fixed points for multivalued mappings defined on unbounded sets in Banach spaces*, J. Math. Anal. Appl. **157** (1991), no. 2, 555–567.
- [11] G. Marino and P. Pietramala, *Fixed points and almost fixed points for mappings defined on unbounded sets in Banach spaces*, Atti Sem. Mat. Fis. Univ. Modena **40** (1992), no. 1, 1–9.

- [12] J. L. Nelson, K. L. Singh, and J. H. M. Whitfield, *Normal structures and nonexpansive mappings in Banach spaces*, Nonlinear Analysis, World Scientific Publishing, Singapore, 1987, pp. 433–492.
- [13] S. Park, *Best approximations and fixed points of nonexpansive maps in Hilbert spaces*, Numer. Funct. Anal. Optim. **18** (1997), no. 5-6, 649–657.
- [14] W. O. Ray, *Nonexpansive mappings on unbounded convex domains*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **26** (1978), no. 3, 241–245.
- [15] ———, *The fixed point property and unbounded sets in Hilbert space*, Trans. Amer. Math. Soc. **258** (1980), no. 2, 531–537.
- [16] S. Reich, *The almost fixed point property for nonexpansive mappings*, Proc. Amer. Math. Soc. **88** (1983), no. 1, 44–46.
- [17] J. Schu, *A fixed point theorem for nonexpansive mappings on star-shaped domains*, Z. Anal. Anwendungen **10** (1991), no. 4, 417–431.
- [18] I. Shafirir, *The approximate fixed point property in Banach and hyperbolic spaces*, Israel J. Math. **71** (1990), no. 2, 211–223.
- [19] T. E. Williamson, *A geometric approach to fixed points of non-self-mappings $T : D \rightarrow X$* , Fixed Points and Nonexpansive Mappings (Cincinnati, Ohio, 1982), Contemp. Math., vol. 18, American Mathematical Society, Rhode Island, 1983, pp. 247–253.
- [20] A. Wiśnicki, *On a problem of common approximate fixed points*, preprint, 2001.

Tadeusz Kuczumow: Instytut Matematyki, Uniwersytet M. Curie-Skłodowskiej (UMCS), 20-031 Lublin, Poland; Instytut Matematyki PWSZ, 20-120 Chełm, Poland
E-mail address: tadek@golem.umcs.lublin.pl