A QUASI-LINEAR PARABOLIC SYSTEM OF CHEMOTAXIS

TAKASI SENBA AND TAKASI SUZUKI

Received 15 December 2004; Accepted 21 January 2005

We consider a quasi-linear parabolic system with respect to unknown functions u and v on a bounded domain of n -dimensional Euclidean space. We assume that the diffusion coefficient of u is a positive smooth function $A(u)$, and that the diffusion coefficient of v is a positive constant. If $A(u)$ is a positive constant, the system is referred to as so-called Keller-Segel system. In the case where the domain is a bounded domain of two-dimensional Euclidean space, it is shown that some solutions to Keller-Segel system blow up in finite time. In three and more dimensional cases, it is shown that solutions to so-called Nagai system blow up in finite time. Nagai system is introduced by Nagai. The diffusion coefficients of Nagai system are positive constants. In this paper, we describe that solutions to the quasi-linear parabolic system exist globally in time, if the positive function $A(u)$ rapidly increases with respect to u .

Copyright © 2006 T. Senba and T. Suzuki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Our purpose is to show the well-posedness of a system of parabolic equations proposed in mathematical biology. Its origin is in Keller and Segel [9], describing the chemotactic aggregation of cellular slime molds which move preferentially toward relatively high concentrations of a chemical secreted by the amoebae themselves. Here, we study the simplified system,

$$\begin{aligned}u_t &= \nabla \cdot (A(u)\nabla u - \chi u \nabla v) \quad \text{in } \Omega \times (0, \infty), \\ \tau v_t &= d\Delta v - av + bu \quad \text{in } \Omega \times (0, \infty), \\ \frac{\partial u}{\partial \nu} &= \frac{\partial v}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, \infty), \\ u(\cdot, 0) &= u_0, \quad v(\cdot, 0) = v_0 \quad \text{in } \Omega,\end{aligned}\tag{1.1}$$

2 A quasi-linear parabolic system of chemotaxis

where $\Omega \subset \mathbf{R}^N$ ($N = 1, 2, 3, \dots$) is a bounded domain with smooth boundary $\partial\Omega$. u_0 and v_0 are smooth non-negative functions on $\overline{\Omega}$ satisfying

$$u_0 \not\equiv 0 \quad \text{in } \Omega, \quad \frac{\partial u_0}{\partial \nu} = \frac{\partial v_0}{\partial \nu} = \frac{\partial \Delta v_0}{\partial \nu} = 0 \quad \text{on } \partial\Omega. \quad (1.2)$$

τ, χ, d, a and b are positive constants, and $A(\cdot)$ is a smooth function defined on $[0, \infty)$.

If $A(\cdot)$ is a positive constant, this system describes the mean field of many self-gravitating particles, and the cases $\tau > 0$ and $\tau = 0$ are referred to as Keller-Segel system and Nagai system, respectively. In this case, if $N = 2$, there is a threshold of the initial value in L^1 norm for existence of the solution globally in time, denoted by c_* , so that if $\|u_0\|_1 < c_*$, then the solution exists globally in time and is bounded ([2, 5, 11, 14]), while blowup of the solution occurs in finite time in the case of $\|u_0\|_1 > c_*$ ([8, 11, 12, 17]). This phenomenon was conjectured by [4] and gave a motivation of our previous works on the mass quantization of collapses in the blowup solution. See [19, 20] for more details. On the other hand, in [13, 16], Nagai and the first author studied

$$\begin{aligned} u_t &= \nabla \cdot (\nabla u - \chi u \nabla \phi(v)) \quad \text{in } \Omega \times (0, \infty), \\ 0 &= d\Delta v - av + bu \quad \text{in } \Omega \times (0, \infty), \end{aligned} \quad (1.3)$$

and gave necessary and sufficient conditions for the existence of the solution globally in time, assuming that the sensitivity function $\phi = \phi(v)$ takes the form $\phi(v) = v^p$ for $p > 0$ or $\phi(v) = \log v$. Also, Biler [3] studied the same problem for more general ϕ . However, system (1.1) has not been studied so much and in this paper, we show that the solution exists globally in time if

$$\inf_{0 \leq u} A(u) > 0, \quad \liminf_{u \rightarrow \infty} \frac{A(u)}{u^{\tilde{\gamma}}} > 0 \quad (1.4)$$

hold for some $\tilde{\gamma} > \max((N-2)/N, 0)$. Those conditions imply

$$A(u) \geq \alpha u^\gamma + \beta \quad (u \geq 0) \quad (1.5)$$

for some $\alpha > 0, \beta > 0, \gamma \in (\max(0, (N-2)/N), \min(\tilde{\gamma}, 1))$. Henceforth, we put $\tau = d = a = b = 1$ for simplicity.

To state the main theorem, we need a few notations. First, given $T \in (0, \infty]$, we set $Q_T = \Omega \times (0, T)$ and take the function spaces $H^\ell(\overline{\Omega})$ and $H^{\ell, \ell/2}(\overline{Q_T})$ as in [10], where ℓ is a non-integral positive number. That is, $H^\ell(\overline{\Omega})$ denotes the Banach space of continuous functions $\{w(x)\}$ defined on $\overline{\Omega}$, provided with continuous derivatives up to order $[\ell]$ and the finite

$$|w|_\Omega^{(\ell)} = \langle w \rangle_\Omega^{(\ell)} + \sum_{j=0}^{[\ell]} \langle w \rangle_\Omega^{(j)} = \sum_{|\delta|=[\ell]} \sup_{x', x \in \Omega} \frac{|D_x^\delta w(x) - D_x^\delta w(x')|}{|x - x'|^{\ell - [\ell]}} + \sum_{j=0}^{[\ell]} \sum_{|\delta|=j} \sup_{x \in \Omega} |D_x^\delta w(x)|, \quad (1.6)$$

where $\delta = (\delta_1, \delta_2, \dots, \delta_N)$, $|\delta| = \delta_1 + \delta_2 + \dots + \delta_N$, and $D_x^\delta = D_{x_1}^{\delta_1} D_{x_2}^{\delta_2} \dots D_{x_N}^{\delta_N}$. Next,

$H^{\ell, \ell/2}(\overline{Q_T})$ denotes the Banach space of continuous functions $\{w(x, t)\}$ defined on $\overline{Q_T}$, provided with continuous derivatives $\{D_t^r D_x^\delta w(x, t)\}$ for $2r + |\delta| < \ell$ and the finite

$$\begin{aligned}
 |w|_{Q_T}^{(\ell)} &= \langle w \rangle_{x, Q_T}^{(\ell)} + \langle w \rangle_{t, Q_T}^{(\ell/2)} + \sum_{j=0}^{[\ell]} \langle w \rangle_{Q_T}^{(j)} \\
 &= \sum_{2r+|\delta|=[\ell]} \sup_{(x', t), (x, t) \in Q_T} \frac{|D_t^r D_x^\delta w(x, t) - D_t^r D_x^\delta w(x', t)|}{|x - x'|^{\ell - [\ell]}} \\
 &\quad + \sum_{2r+|\delta|=[\ell]} \sup_{(x, t'), (x, t) \in Q_T} \frac{|D_t^r D_x^\delta w(x, t) - D_t^r D_x^\delta w(x, t')|}{|t - t'|^{(\ell - [\ell])/2}} \\
 &\quad + \sum_{j=0}^{[\ell]} \sum_{2r+|\delta|=j} \sup_{(x, t) \in Q_T} |D_t^r D_x^\delta w(x, t)|.
 \end{aligned} \tag{1.7}$$

Then, the first theorem guarantees the local well-posedness of (1.1).

THEOREM 1.1 (time-local solution). *If (1.2) holds and*

$$\inf_{u \geq 0} A(u) = \beta > 0 \tag{1.8}$$

in (1.1), then there is $T > 0$ such that (1.1) has a unique classical solution (u, v) in Q_T satisfying

$$|u|_{Q_T}^{2+\theta} < +\infty, \quad |v|_{Q_T}^{2+\theta} < +\infty \tag{1.9}$$

for some $\theta \in (0, 1)$.

The main theorem is now stated as follows.

THEOREM 1.2 (time-global solution). *If (1.2) and (1.4) hold in (1.1), then it has a unique classical solution (u, v) in Q_∞ satisfying*

$$|u|_{Q_\infty}^{2+\theta} < +\infty, \quad |v|_{Q_\infty}^{2+\theta} < +\infty \tag{1.10}$$

for some $\theta \in (0, 1)$.

In Section 2, we consider some linear system related to (1.1). By using some estimates of the solutions and a standard contraction mapping principle, we show Theorem 1.1.

In Section 3, we introduce Lyapunov function of (1.1). By using the Lyapunov function, a maximal regularity in [6] and Moser's technique in [1], we show Theorem 1.2.

2. Time local solution

We take $T > 0$, $p > N + 2$, and $\ell \in (0, 1 - (N + 2)/p)$. Given

$$u \in H^{1+\ell, \ell/2}(\overline{Q_T}) \equiv \left\{ w \in H^{\ell, \ell/2}(\overline{Q_T}) \mid \nabla_x u \in H^{\ell, \ell/2}(\overline{Q_T})^N \right\}, \tag{2.1}$$

4 A quasi-linear parabolic system of chemotaxis

we apply [10, Theorem IV.5.3] and get the unique solution

$$(U, v) \in H^{3+\ell, (3+\ell)/2}(\overline{Q_T})^2 \quad (2.2)$$

to the system

$$\begin{aligned} U_t &= \nabla \cdot (A(u) \nabla U - U \nabla v) \quad \text{in } Q_T, \\ v_t &= \Delta v - v + u \quad \text{in } Q_T, \\ \frac{\partial U}{\partial \nu} &= \frac{\partial v}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T), \\ U(\cdot, 0) &= u_0, \quad v(\cdot, 0) = v_0 \quad \text{in } \Omega. \end{aligned} \quad (2.3)$$

Thus, we can introduce the operator $\mathcal{F}$ on $\{w \in H^{1+\ell, \ell/2}(\overline{Q_T}) \mid w \geq 0 \text{ in } \overline{Q_T}\}$ by

$$\mathcal{F}u = U. \quad (2.4)$$

Let

$$\begin{aligned} B(M, T) \equiv \Big\{ u \in H^{1+\ell, \ell/2}(\overline{Q_T}) \cap C([0, T]; W^{2,p}(\Omega)) \mid u \geq 0 \text{ in } \overline{Q_T}, \\ \partial u / \partial \nu = 0 \text{ on } \partial\Omega \times [0, T], \max_{0 \leq t \leq T} (\|\Delta u\|_p^p + \|u\|_p^p)^{1/p} \leq M \Big\} \end{aligned} \quad (2.5)$$

for

$$M = 2^{1/p} \left(\|\Delta u_0\|_p^p + \|u_0\|_p^p \right)^{1/p}, \quad (2.6)$$

where $\|\cdot\|_q$ and $\|\cdot\|_{m,q}$ denote the standard norms in $L^p(\Omega)$ and $W^{m,p}(\Omega)$, respectively.

Henceforth, C_i ($i = 1, 2, 3, 4, 5$) denote positive constants determined by M and T , which are monotone increasing in T . First, we show the following.

LEMMA 2.1. *If $u \in B(M, T)$, then the solution v to*

$$v_t = \Delta v - v + u \quad \text{in } Q_T, \quad \frac{\partial v}{\partial \nu} = 0 \quad \text{on } \partial\Omega \times (0, T) \quad (2.7)$$

satisfies that

$$\max_{0 \leq t \leq T} \left(\|\nabla \Delta v(\cdot, t)\|_p + \|v(\cdot, t)\|_p \right) \leq C_1. \quad (2.8)$$

Proof. Let $\Delta_{N,q}$ be the Laplacian with homogeneous Neumann boundary condition, realized as an m -accretive operator in $L^q(\Omega)$, where $q \in (1, \infty)$. Its domain is given by

$$D(\Delta_{N,q}) \equiv \left\{ w \in W^{2,q}(\Omega) \mid \frac{\partial w}{\partial \nu} = 0 \text{ on } \partial\Omega \right\}, \quad (2.9)$$

and henceforth we will write it as Δ_N for simplicity. Then, $-\Delta_N + 1$ generates the analytic

semi-group $\{e^{t(\Delta_N-1)}\}_{t \geq 0}$ ([7]), and it holds that

$$v(\cdot, t) = e^{(\Delta_N-1)t} v_0 + \int_0^t e^{(\Delta_N-1)(t-s)} u(\cdot, s) ds. \quad (2.10)$$

On the other hand, we have

$$\|(-\Delta_N + 1)^{3/2} v_0\|_p \leq C_2 \quad (2.11)$$

from the assumption, and if $u \in B(M, T)$, then it holds that $u(\cdot, t) \in D(\Delta_N)$ and

$$\|(-\Delta_N + 1)u(\cdot, t)\|_p \leq \left(\|\Delta u(\cdot, t)\|_p^p + \|u(\cdot, t)\|_p^p \right)^{1/p} \leq M \quad (2.12)$$

for $t \in [0, T]$. Thus, we have from (2.10), (2.11), and (2.12) that

$$\begin{aligned} \|(-\Delta_N + 1)^{3/2} v(\cdot, t)\|_p &\leq C_2 + \int_0^t \|(-\Delta_N + 1)^{1/2} e^{(\Delta_N-1)(t-s)} (-\Delta_N + 1) u(\cdot, s)\|_p ds \\ &\leq C_2 + \int_0^t \frac{C_3}{\sqrt{t-s}} M ds \leq C_2 + \frac{1}{2} \sqrt{T} C_3 M, \end{aligned} \quad (2.13)$$

and in particular,

$$\|\nabla \Delta v(\cdot, t)\|_p \leq C_4 \quad (2.14)$$

follows for $t \in [0, T]$. It also holds that

$$\|v(\cdot, t)\|_p \leq C_5 \quad (2.15)$$

and the proof is complete. $\square$

Now, we show the following.

LEMMA 2.2. *There exists $T_1 > 0$ such that*

$$\mathcal{FB}(M, T) \subset B(M, T) \quad (2.16)$$

holds for any $T \in (0, T_1]$.

Proof. Henceforth, C_i ($i = 6, 7, 8, \dots, 20$) denote positive constants depending only on M . We have $W^{2,p}(\Omega) \subset W^{1,\infty}(\Omega)$ and it holds that

$$\sup_{0 \leq t \leq S} \left(\|u(\cdot, t)\|_{1,\infty} + \|v(\cdot, t)\|_{1,\infty} \right) \leq C_6 \quad (2.17)$$

for $u \in B(M, S)$ if $S \in (0, 1]$.

6 A quasi-linear parabolic system of chemotaxis

From the first equation of (2.3), we obtain that

$$\begin{aligned}
\frac{1}{p} \frac{d}{dt} \int_{\Omega} |U|^p dx &= \int_{\Omega} \nabla \cdot (A(u) \nabla U - U \nabla v) |U|^{p-2} U dx \\
&= -(p-1) \int_{\Omega} A(u) |U|^{p-2} |\nabla U|^2 dx + (p-1) \int_{\Omega} |U|^{p-2} U \nabla v \cdot \nabla U dx \\
&\leq -\beta(p-1) \int_{\Omega} |U|^{p-2} |\nabla U|^2 dx + (p-1) \int_{\Omega} |U|^{p-1} |\nabla v| |\nabla U| dx \\
&\leq -\frac{1}{2} \beta(p-1) \int_{\Omega} |U|^{p-2} |\nabla U|^2 dx + C_7 \int_{\Omega} |\nabla v|^2 |U|^p dx.
\end{aligned} \tag{2.18}$$

We have $W^{2,p}(\Omega) \subset H^{1+\ell}(\overline{\Omega})$, and hence Lemma 2.1 implies the boundedness of ∇v . Therefore, it follows that

$$\frac{d}{dt} \int_{\Omega} |U|^p dx \leq C_8 \int_{\Omega} |U|^p dx. \tag{2.19}$$

The first equation of (2.3) implies also that

$$\begin{aligned}
\frac{1}{p(p-1)} \frac{d}{dt} \int_{\Omega} |\Delta U|^p dx &= \frac{1}{p-1} \int_{\Omega} |\Delta U|^{p-2} \Delta U \cdot \Delta U_t dx \\
&= -\frac{1}{p-1} \int_{\Omega} \nabla U_t \cdot \nabla (|\Delta U|^{p-2} \Delta U) dx \\
&= -\int_{\Omega} \nabla U_t \cdot (|\Delta U|^{p-2} \Delta \nabla U) dx \\
&= -\int_{\Omega} [\nabla (\nabla \cdot (A(u) \nabla U - U \nabla v)) \cdot \nabla \Delta U] |\Delta U|^{p-2} dx.
\end{aligned} \tag{2.20}$$

Here, we have

$$\begin{aligned}
&\nabla \{ \nabla \cdot (A(u) \nabla U - U \nabla v) \} \\
&= A(u) \nabla \Delta U + (\Delta U) \nabla A(u) + \nabla (\nabla A(u) \cdot \nabla U) \\
&\quad - \nabla (\nabla U \cdot \nabla v) - (\Delta v) \nabla U - U \nabla \Delta v \\
&= A(u) \nabla \Delta U + (\Delta U) \nabla A(u) + \sum_{i=1}^N \frac{\partial \nabla A(u)}{\partial x_i} \frac{\partial U}{\partial x_i} \\
&\quad + \sum_{i=1}^N \frac{\partial A(u)}{\partial x_i} \frac{\partial \nabla U}{\partial x_i} - \sum_{i=1}^N \frac{\partial \nabla U}{\partial x_i} \frac{\partial v}{\partial x_i} - \sum_{i=1}^N \frac{\partial U}{\partial x_i} \frac{\partial \nabla v}{\partial x_i} \\
&\quad - (\Delta v) \nabla U - U \nabla \Delta v \\
&= I + II + III + IV - V - VI - VII - VIII
\end{aligned} \tag{2.21}$$

and those terms are estimated as follows:

$$\begin{aligned}
& - \int_{\Omega} (I' \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \leq -\beta \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx, \\
& - \int_{\Omega} (II' \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \\
& \quad = - \int_{\Omega} (\nabla A(u) \cdot \nabla \Delta U) |\Delta U|^{p-2} \Delta U dx \\
& \quad \leq C_6 \|A'\|_{L^\infty([0, C_6])} \cdot \int_{\Omega} |\nabla \Delta U| |\Delta U|^{p-1} dx \\
& \quad \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_9 \int_{\Omega} |\Delta U|^p dx, \\
& - \int_{\Omega} (III' \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \\
& \quad \leq \|A'\|_{L^\infty([0, C_6])} \int_{\Omega} \sum_{i,j=1}^N \left| \frac{\partial^2 u}{\partial x_i \partial x_j} \right| |\nabla U| |\nabla \Delta U| |\Delta U|^{p-2} dx \\
& \quad \quad + \|A''\|_{L^\infty([0, C_6])} \int_{\Omega} |\nabla u|^2 |\nabla U| |\nabla \Delta U| |\Delta U|^{p-2} dx \\
& \quad \leq C_{10} (\|\Delta u\|_p^p + \|u\|_p^p)^{1/p} \|\nabla U\|_\infty \\
& \quad \quad \cdot \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\Delta U\|_p^{(p-2)/2} \\
& \quad \quad + C_{10} C_6^2 \|\nabla U\|_p \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\Delta U\|_p^{(p-2)/2} \\
& \quad \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_{11} (\|\Delta U\|_p^p + \|U\|_p^p), \\
& - \int_{\Omega} (IV' \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \\
& \quad \leq \|A'\|_{L^\infty([0, C_6])} \int_{\Omega} |\nabla u| \left(\sum_{i,j=1}^N \left| \frac{\partial^2 U}{\partial x_i \partial x_j} \right| \right) |\nabla \Delta U| |\Delta U|^{p-2} dx \\
& \quad \leq C_6 \|A'\|_{L^\infty([0, C_6])} \left\{ \int_{\Omega} \left(\sum_{i,j=1}^N \left| \frac{\partial^2 U}{\partial x_i \partial x_j} \right| \right)^p dx \right\}^{1/p} \\
& \quad \quad \cdot \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\Delta U\|_p^{(p-2)/2} \\
& \quad \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_{12} (\|\Delta U\|_p^p + \|U\|_p^p), \\
& - \int_{\Omega} (-V \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \\
& \quad \leq \int_{\Omega} |\nabla v| \left(\sum_{i,j=1}^N \left| \frac{\partial^2 U}{\partial x_i \partial x_j} \right| \right) |\nabla \Delta U| |\Delta U|^{p-2} dx
\end{aligned} \tag{2.22}$$

$$\begin{aligned}
& \leq C_6 \left\{ \int_{\Omega} \left(\sum_{i,j=1}^N \left| \frac{\partial^2 U}{\partial x_i \partial x_j} \right| \right)^p dx \right\}^{1/p} \\
& \quad \cdot \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\Delta U\|_p^{(p-2)/2} \\
& \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_{13} (\|\Delta U\|_p^p + \|U\|_p^p), \\
& - \int_{\Omega} (-VI \cdot \nabla \Delta U) |\nabla \Delta U|^{p-2} dx \\
& \leq \int_{\Omega} |\nabla U| \left(\sum_{i,j=1}^N \left| \frac{\partial^2 v}{\partial x_i \partial x_j} \right| \right) |\nabla \Delta U| |\Delta U|^{p-2} dx \\
& \leq C_{14} \|v\|_{2,p} \|\nabla U\|_{\infty} \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\Delta U\|_p^{(p-2)/p} \\
& \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_{15} (\|\Delta U\|_p^p + \|U\|_p^p), \\
& - \int_{\Omega} (-VII \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \\
& \leq \int_{\Omega} |\Delta v| |\nabla U| |\nabla \Delta U| |\Delta U|^{p-2} dx \\
& \leq C_{16} \|v\|_{2,p} \|\nabla U\|_{\infty} \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\Delta U\|_p^{(p-2)/2} \\
& \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_{17} (\|\Delta U\|_p^p + \|U\|_p^p), \\
& - \int_{\Omega} (-VIII \cdot \nabla \Delta U) |\Delta U|^{p-2} dx \\
& \leq \int_{\Omega} U |\nabla \Delta v| |\nabla \Delta U| |\Delta U|^{p-2} dx \\
& \leq \left\{ \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \right\}^{1/2} \|\nabla \Delta v\|_p \|\Delta U\|_p^{(p-2)/2} \|U\|_{\infty} \\
& \leq \frac{\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx + C_{18} (\|\Delta U\|_p^p + \|U\|_p^p).
\end{aligned} \tag{2.23}$$

Those inequalities are summarised as

$$\frac{1}{p(p-1)} \frac{d}{dt} \int_{\Omega} |\Delta U|^p dx + \frac{3\beta}{10} \int_{\Omega} |\nabla \Delta U|^2 |\Delta U|^{p-2} dx \leq C_{19} (\|\Delta U\|_p^p + \|U\|_p^p), \tag{2.24}$$

and therefore, we have

$$\frac{d}{dt} (\|\Delta U\|_p^p + \|U\|_p^p) \leq C_{20} (\|\Delta U\|_p^p + \|U\|_p^p) \tag{2.25}$$

by (2.19). Taking

$$T_1 = \min \left(\frac{\log 2}{C_{20}}, S \right), \quad (2.26)$$

we have for $T \in (0, T_1]$ that

$$\sup_{0 \leq t \leq T} \left(\|\Delta U(\cdot, t)\|_p^p + \|U(\cdot, t)\|_p^p \right) \leq e^{C_{20}T} \left(\|\Delta u_0\|_p^p + \|u_0\|_p^p \right) \leq \frac{1}{2} M^p e^{C_{20}T} \leq M^p, \quad (2.27)$$

and the proof is complete. $\square$

Next, we show the following.

LEMMA 2.3. *There is $T_2 \in (0, T_1]$ satisfying*

$$\max_{0 \leq t \leq T_2} \|\mathcal{F}u_1(\cdot, t) - \mathcal{F}u_2(\cdot, t)\|_p^p \leq \frac{1}{2} \max_{0 \leq t \leq T_2} \|u_1(\cdot, t) - u_2(\cdot, t)\|_p^p \quad (2.28)$$

for any $u_1, u_2 \in B(M, T_2)$.

Proof. We take $T \in (0, T_1]$ and $u_i \in B(M, T)$ for $i = 1, 2$. Let U_i and v_i be the solutions to (2.3) for $u = u_i$. Then, it holds that

$$v_2(\cdot, t) - v_1(\cdot, t) = \int_0^t e^{(s-t)(\Delta_N - 1)} (u_1(\cdot, s) - u_2(\cdot, s)) ds \quad (2.29)$$

and we have

$$\|\nabla(v_2(\cdot, t) - v_1(\cdot, t))\|_p \leq C_{21} \sqrt{t} \sup_{0 \leq s \leq t} \|u_2(\cdot, s) - u_1(\cdot, s)\|_p. \quad (2.30)$$

Next, we note

$$\begin{aligned} \frac{1}{p} \frac{d}{dt} \|U_2(\cdot, t) - U_1(\cdot, t)\|_p^p &= \int_{\Omega} (U_{2t} - U_{1t}) |U_2 - U_1|^{p-2} (U_2 - U_1) dx \\ &= \int_{\Omega} \{ \nabla \cdot [(A(u_2) - A(u_1)) \nabla U_2] + \nabla \cdot [A(u_1) \nabla (U_2 - U_1)] \\ &\quad - \nabla \cdot [(U_2 - U_1) \nabla v_2] - \nabla \cdot [U_1 \nabla (v_2 - v_1)] \} |U_2 - U_1|^{p-2} (U_2 - U_1) dx \\ &= -(p-1) \int_{\Omega} (A(u_2) - A(u_1)) [\nabla U_2 \cdot \nabla (U_2 - U_1)] |U_2 - U_1|^{p-2} dx \\ &\quad - (p-1) \int_{\Omega} A(u_1) |\nabla (U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx \\ &\quad + (p-1) \int_{\Omega} [\nabla v_2 \cdot \nabla (U_2 - U_1)] |U_2 - U_1|^{p-2} (U_2 - U_1) dx \\ &\quad + (p-1) \int_{\Omega} U_1 [\nabla (v_2 - v_1) \cdot \nabla (U_2 - U_1)] |U_2 - U_1|^{p-2} dx \\ &= -IX - X + XI + XII. \end{aligned} \quad (2.31)$$

Those terms are estimated by Lemmas 2.1 and 2.2 in the following way:

$$\begin{aligned}
|IX| &\leq (p-1)\|A'\|_{L^\infty([0,C_6])}\|u_2 - u_1\|_p\|\nabla U_2\|_\infty \\
&\quad \cdot \left\{ \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx \right\}^{1/2} \|U_2 - U_1\|_p^{(p-2)/2} \\
&\leq \frac{(p-1)\beta}{4} \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx \\
&\quad + C_{22}(\|U_2 - U_1\|_p^p + \|u_2 - u_1\|_p^p), \\
&\quad -X \leq -\beta(p-1) \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx, \\
|XI| &\leq (p-1)\|\nabla v_2\|_\infty \left\{ \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx \right\}^{1/2} \cdot \|U_2 - U_1\|_p^{p/2} \\
&\leq \frac{\beta(p-1)}{4} \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx + C_{23}\|U_2 - U_1\|_p^p, \\
|XII| &\leq (p-1)\|U_1\|_\infty \left\{ \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx \right\}^{1/2} \\
&\quad \cdot \|U_2 - U_1\|_p^{(p-2)/2} \|\nabla(v_2 - v_1)\|_p \\
&\leq \frac{(p-1)\beta}{4} \int_\Omega |\nabla(U_2 - U_1)|^2 |U_2 - U_1|^{p-2} dx \\
&\quad + C_{24}(\|U_2 - U_1\|_p^p + \|\nabla(v_2 - v_1)\|_p^p).
\end{aligned} \tag{2.32}$$

Thus, we obtain

$$\frac{1}{p} \frac{d}{dt} \|U_2 - U_1\|_p^p \leq C_{25} \left(\|U_2 - U_1\|_p^p + \|u_2 - u_1\|_p^p + \|\nabla(v_2 - v_1)\|_p^p \right), \tag{2.33}$$

and hence it follows from (2.30) that

$$\begin{aligned}
&\sup_{0 \leq t \leq T} \|U_2(\cdot, t) - U_1(\cdot, t)\|_p^p \\
&\leq pC_{25}T \left(\sup_{0 \leq t \leq T} \|U_2(\cdot, t) - U_1(\cdot, t)\|_p^p + \sup_{0 \leq t \leq T} \|u_2(\cdot, t) - u_1(\cdot, t)\|_p^p \right) \\
&\quad + T^{(p/2)+1} pC_{25}C_{21}^p \sup_{0 \leq t \leq T} \|u_2(\cdot, t) - u_1(\cdot, t)\|_p^p.
\end{aligned} \tag{2.34}$$

Therefore, for $T_2 > 0$ in

$$T_2 \leq T_1, \quad pC_{25}T_2 + T_2^{(p/2)+1} pC_{25}C_{21}^p \leq \frac{1}{4}, \tag{2.35}$$

it holds that

$$\sup_{0 \leq t \leq T_2} \|U_2(\cdot, t) - U_1(\cdot, t)\|_p^p \leq \frac{1}{2} \sup_{0 \leq t \leq T_2} \|u_2(\cdot, t) - u_1(\cdot, t)\|_p^p, \quad (2.36)$$

and the proof is complete. $\square$

Now, we can give the following.

Proof of Theorem 1.1. By Lemma 2.2 (with $T = T_2$) and Lemma 2.3, we have a positive constant C_{26} such that

$$\begin{aligned} \|U_t\|_p &\leq \|A(u)\Delta U\|_p + \|A'(u)\nabla u \cdot \nabla U\|_p \\ &\quad + \|\nabla U \cdot \nabla v\|_p + \|U\Delta v\|_p \leq C_{26} \end{aligned} \quad (2.37)$$

for $u \in B(M, T_2)$. Then [10, Lemma 3.3] guarantees the existence of $\ell \in (0, 1)$ such that

$$|\mathcal{F}u|_{Q_{T_2}}^{(\ell)} + |\nabla \mathcal{F}u|_{Q_{T_2}}^{(\ell)} \leq C_{27} \quad (2.38)$$

for $u \in B(M, T_2)$. Because the set

$$\mathcal{H} \equiv \left\{ u \in B(M, T_2) \mid |\mathcal{F}u|_{Q_{T_2}}^{(\ell)} + |\nabla \mathcal{F}u|_{Q_{T_2}}^{(\ell)} \leq C_{27} \right\}, \quad (2.39)$$

is compact in $X = C([0, T_2]; L^p(\Omega))$, we can apply the standard contraction mapping principle and get a unique fix point of $\mathcal{F}$ on $B = B(M, T_2) \subset X$. This fixed point induces the classical solution $(u, v) \in H^{2+\ell, 1+\ell/2}(\overline{Q_{T_2}})^2$ to (1.1) on Q_{T_2} . Conversely, if (u, v) is a classical solution to (1.1) and if $T_2 > 0$ is taken smaller, then, we can regard (u, v) as a fixed point of $\mathcal{F}$ on $B(M, T_2)$. This implies the uniqueness of the classical solution to (1.1) in Q_{T_2} , and the proof is complete. $\square$

3. Time global solution

To prove Theorem 1.2, we take

$$W(u, v) = \int_{\Omega} \left\{ B(u) - uv + \frac{1}{2}(|\nabla v|^2 + v^2) \right\} dx \quad (3.1)$$

for

$$B(u) = \int_1^u b(s) ds \quad \text{with } b(u) = \int_1^u \frac{A(s)}{s} ds \quad (3.2)$$

and show the following.

LEMMA 3.1. *If (u, v) is the classical solution to (1.1), then it holds that*

$$\frac{d}{dt} W(u, v) + \int_{\Omega} \left(|v_t|^2 + u |\nabla(b(u) - v)|^2 \right) dx = 0. \quad (3.3)$$

Proof. From the first equation of (1.1) we have

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} (B(u) - uv) dx + \int_{\Omega} uv_t dx &= \int_{\Omega} u_t (b(u) - v) dx \\ &= - \int_{\Omega} u |\nabla (b(u) - v)|^2 dx = 0, \end{aligned} \quad (3.4)$$

and the second equation implies that

$$\int_{\Omega} uv_t dx = \int_{\Omega} (v_t - \Delta v + v) v_t dx = \int_{\Omega} |v_t|^2 dx + \frac{1}{2} \frac{d}{dt} \int_{\Omega} (|\nabla v|^2 + v^2) dx. \quad (3.5)$$

Thus, the conclusion follows from (3.4) and (3.5). $\square$

We have

$$\begin{aligned} b(u) &\geq \frac{\alpha}{\gamma} (u^\gamma - 1) + \beta \log u \quad (u \geq 1) \\ b(u) &\leq \frac{\alpha}{\gamma} (u^\gamma - 1) + \beta \log u \quad (0 < u \leq 1) \end{aligned} \quad (3.6)$$

for α, β, γ given in (1.5), and hence it follows that

$$B(u) \geq B_0(u) \equiv \frac{\alpha}{\gamma(1+\gamma)} (u^{1+\gamma} - (1+\gamma)u + \gamma) + \beta(u \log u - u + 1) \quad (3.7)$$

for $u \in (0, \infty)$. Therefore, we have $W_0(u, v) \leq W(u, v)$ for $W_0 = W_0(u, v)$ defined by

$$W_0(u, v) \equiv \int_{\Omega} \left\{ B_0(u) - uv + \frac{1}{2} (|\nabla v|^2 + v^2) \right\} dx. \quad (3.8)$$

Henceforth, $T_{\max}$ denotes the maximal existence time of the classical solution (u, v) to (1.1), and C_i ($i = 28, 29, 30, \dots, 40$) are positive constants independent of $T_{\max}$. We have

$$\|u(\cdot, t)\|_1 = \|u_0\|_1 \quad (3.9)$$

and the following lemma is a consequence of the estimate on the fundamental solution to the heat equation, described in [21]. The proof is immediate and is omitted.

LEMMA 3.2. *If p is in $[1, N/(N-2))$ and $[1, \infty)$ for $N \geq 3$ and $N = 1, 2$, respectively, then it follows that*

$$\sup_{0 \leq t < T_{\max}} \|v(\cdot, t)\|_p \leq C_{28}. \quad (3.10)$$

Now, we show the following.

LEMMA 3.3. *If (u, v) is the classical solution to (1.1), then it holds that*

$$\begin{aligned} \sup_{0 \leq t < T_{\max}} W_0(u(\cdot, t), v(\cdot, t)) &\leq C_{29} \\ \sup_{0 \leq t < T_{\max}} \left(\|u(\cdot, t)\|_{1+\gamma}^{1+\gamma} + \|v(\cdot, t)\|_{1,2}^2 \right) &\leq C_{29}. \end{aligned} \quad (3.11)$$

Proof. First, we have

$$\sup_{0 \leq t < T_{\max}} W_0(u(\cdot, t), v(\cdot, t)) \leq \sup_{0 \leq t < T_{\max}} W(u(\cdot, t), v(\cdot, t)) \leq W(u_0, v_0) \quad (3.12)$$

with

$$\int_{\Omega} uv \, dx \leq \|u\|_{1+\gamma} \|v\|_{(1+\gamma)/\gamma}. \quad (3.13)$$

Then, Gagliardo-Nirenberg's inequality is indicated as

$$\|w\|_q \leq K_{p,q} (\|\nabla w\|_2^2 + \|w\|_2^2)^{\theta/2} \|w\|_p^{1-\theta} \quad (3.14)$$

for $w \in H^1(\Omega)$, where $p, q \in [1, \infty]$, and $\theta \in (0, 1)$ are in

$$-\frac{N}{q} = \theta\{1 - (N/2)\} - (1 - \theta)\frac{N}{p}. \quad (3.15)$$

In the case of $N \geq 3$, we take $q = (\gamma + 1)/\gamma$, $\theta = 2\gamma/(\gamma + 1)$ and $p = N(1 - \gamma)/(2\gamma)$ to adjust (3.15). We have $p \in [1, N/(N - 2))$ from the assumption, and (3.14) assures $C_\varepsilon > 0$ for any $\varepsilon > 0$ satisfying

$$\varepsilon \|\nabla v\|_2^2 \geq \|v\|_{(\gamma+1)/\gamma}^{(\gamma+1)/\gamma} - C_\varepsilon \quad (3.16)$$

by Lemma 3.2. The case $N = 1, 2$ is easier to guarantee (3.16). Therefore, we have from (3.12) that

$$W(u_0, v_0) \geq \int_{\Omega} \left\{ \frac{\alpha}{2\gamma(\gamma+1)} u^{1+\gamma} + \frac{1}{4} (|\nabla v|^2 + v^2) \right\} dx - C_{30}. \quad (3.17)$$

This implies the second inequality, and the proof is complete. $\square$

LEMMA 3.4. *If $q \geq (3 + \gamma)/2$ and*

$$\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{q+(\gamma-1)/2} \leq C_{32}, \quad (3.18)$$

then it holds that

$$\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{2q-1} \leq C_{33}. \quad (3.19)$$

Proof. From the first equation of (1.1) we have

$$\begin{aligned} \int_{\Omega} u_t u^{2q-2} dx &= \int_{\Omega} \nabla \cdot (A(u) \nabla u - u \nabla v) u^{2q-2} dx \\ &\leq -(2q-2) \int_{\Omega} (\alpha u^\gamma + \beta) u^{2q-3} |\nabla u|^2 dx + (2q-2) \int_{\Omega} u^{2q-2} \nabla u \cdot \nabla v dx \\ &= -(2q-2) \int_{\Omega} (\alpha u^\gamma + \beta) u^{2q-3} |\nabla u|^2 dx - \frac{2q-2}{2q-1} \int_{\Omega} u^{2q-1} \Delta v dx. \end{aligned} \quad (3.20)$$

On the other hand, the second equation of (1.1) implies

$$(e^{t/(4q)}v)_t = \left(\Delta - 1 + \frac{1}{4q}\right)(e^{t/(4q)}v) + e^{t/(4q)}u \quad (3.21)$$

and the maximal regularity theorem [6] guarantees

$$\begin{aligned} & \int_0^t \|(e^{s/(4q)}v)_s\|_{2q}^{2q} ds + \int_0^t \left\| \left(\Delta - 1 + \frac{1}{4q}\right)(e^{s/(4q)}v) \right\|_{2q}^{2q} ds \\ & \leq C_{34} \left\{ \int_0^t \left\| \left(\Delta - 1 + \frac{1}{4q}\right)e^{\{\Delta - 1 + (1/(4q))\}s} v_0 \right\|_{2q}^{2q} ds + \int_0^t \|e^{s/(4q)}u\|_{2q}^{2q} ds \right\} \end{aligned} \quad (3.22)$$

with C_{34} independent of $t \geq 0$. This implies

$$\int_0^t e^{s/2} \|\Delta v\|_{2q}^{2q} ds \leq C_{35} \left\{ t \|v_0\|_{2,2q}^{2q} + \int_0^t e^{s/2} \|u\|_{2q}^{2q} ds \right\}. \quad (3.23)$$

We can apply (3.14) for $q = r$ and $p = 1$ if $\theta = N(1 - (1/r))/(1 + (N/2)) \in (0, 1)$, or equivalently $1 < r < 2N/(N - 2)$. Putting

$$k = (2q + \gamma - 1)/2, \quad r = 2q/k, \quad (3.24)$$

we have

$$r = \frac{4q}{2q + \gamma - 1} \in \left[2, 2 + \frac{1 - \gamma}{1 + \gamma}\right], \quad (3.25)$$

and therefore, that inequality is applicable to $w = u^k$, $q = r$ and $p = 1$:

$$\|u^k\|_r \leq K_r \left(\|\nabla u^k\|_2^2 + \|u^k\|_2^2 \right)^{\theta/2} \|u^k\|_1^{1-\theta}. \quad (3.26)$$

Here, we have

$$K = \max_{2 \leq r \leq 2 + (1-\gamma)/(1+\gamma)} K_r < +\infty, \quad r\theta = \frac{2N}{N+2}(r-1) < 2, \quad (3.27)$$

and any $\varepsilon > 0$ admits $C_\varepsilon > 0$ such that

$$\begin{aligned} \|u^k\|_r^r & \leq K^r \left(\|\nabla u^k\|_2^2 + \|u^k\|_2^2 \right)^{r\theta/2} \|u^k\|_1^{r(1-\theta)} \\ & \leq \varepsilon \left(\|\nabla u^k\|_2^2 + \|u^k\|_2^2 \right) + C_\varepsilon \|u^k\|_1^{r(1-\theta)(2/(r\theta))'}, \end{aligned} \quad (3.28)$$

where

$$\begin{aligned} \left(\frac{2}{r\theta}\right)' &= \frac{2/(r\theta)}{(2/(r\theta)) - 1} = \frac{N+2}{(N+2) - N(r-1)} \\ &\leq \begin{cases} 3(1+\gamma)/(1+3\gamma) & \text{if } N=1 \\ (1+\gamma)/\gamma & \text{if } N=2 \\ (N+2)(N-1)/(N-2) & \text{if } N \geq 3 \end{cases} \\ &= \kappa \end{aligned} \quad (3.29)$$

with

$$\Lambda = \Lambda(q) = r(1-\theta) \left(\frac{2}{r\theta}\right)' = \frac{2N - (N-2)r}{2N+2-Nr}. \quad (3.30)$$

Thus, we have

$$\begin{aligned} C_{35} \int_0^t e^{s/2} \|u\|_{2q}^{2q} ds &\leq (q-1)(2q-1)\alpha \int_0^t \int_{\Omega} e^{s/2} u^{2q+\gamma-3} |\nabla u|^2 dx ds \\ &\quad + C_{36} \int_0^t e^{s/2} \left(\|u\|_{q+(y-1)/2}^{q+(y-1)/2} + 1 \right)^{\Lambda} ds. \end{aligned} \quad (3.31)$$

On the other hand, by (3.26), (3.29) and $\|u^k\|_r^r = \|u\|_{2q}^{2q}$, we have

$$\begin{aligned} &-(q-1)(2q-1) \int_{\Omega} \alpha u^{2q+\gamma-3} |\nabla u|^2 dx + (2q-2) \int_{\Omega} u^{2q-1} |\Delta v| dx \\ &\leq -\frac{(q-1)(2q-1)}{k^2} \alpha \|\nabla u^k\|_2^2 + (2q-2) \|u\|_{2q}^{2q-1} \|\Delta v\|_{2q} \\ &\leq -\frac{1}{2} \|u\|_{2q-1}^{2q-1} + \|\Delta v\|_{2q}^{2q} + C_{37} \left(\|u\|_{q+(y-1)/2}^{q+(y-1)/2} + 1 \right)^{\Lambda}. \end{aligned} \quad (3.32)$$

From (3.32) and (3.20) we obtain

$$\begin{aligned} &\frac{d}{dt} \|u\|_{2q-1}^{2q-1} + (q-1)(2q-1)\alpha \int_{\Omega} u^{2q+\gamma-3} |\nabla u|^2 dx \\ &\leq -(q-1)(2q-1)\alpha \int_{\Omega} u^{2q+\gamma-3} |\nabla u|^2 dx + (2q-2) \int_{\Omega} u^{2q-1} |\Delta v| dx \\ &\leq -\frac{1}{2} \|u\|_{2q-1}^{2q-1} + \|\Delta v\|_{2q}^{2q} + C_{37} \left(\|u\|_{q+(y-1)/2}^{q+(y-1)/2} + 1 \right)^{\Lambda}. \end{aligned} \quad (3.33)$$

Multiplying $e^{t/2}$ for the above inequality, integrating over $[0, t]$ and using (3.23) and (3.31), we now get that

$$\begin{aligned} &e^{t/2} \|u\|_{2q-1}^{2q-1} + (q-1)(2q-1)\alpha \int_0^t e^{s/2} \int_{\Omega} u^{2q+\gamma-3} |\nabla u|^2 dx ds \\ &\leq \int_0^t e^{s/2} \|\Delta v\|_{2q}^{2q} ds + C_{38} e^{t/2} (L_{q+(y-1)/2} + 1)^{\Lambda} \end{aligned} \quad (3.34)$$

or

$$\|u(\cdot, t)\|_{2q-1}^{2q-1} \leq C_{39}(L_{q+(\gamma-1)/2} + 1)^\Lambda, \quad (3.35)$$

where

$$L_{q+(\gamma-1)/2} = \sup_{0 \leq t < T_{\max}} \|u\|_{q+(\gamma-1)/2}^{q+(\gamma-1)/2}. \quad (3.36)$$

Thus, the proof is complete. $\square$

Now, we show the uniform boundedness of the solution.

PROPOSITION 3.5. *It holds that*

$$\sup_{0 \leq t < T_{\max}} (\|u(\cdot, t)\|_\infty + \|v(\cdot, t)\|_\infty) \leq C_{40}. \quad (3.37)$$

Proof. By means of Lemmas 3.3 and 3.4, we have

$$\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_q < +\infty \quad \text{for any } q \geq 1 + \gamma. \quad (3.38)$$

We have $D((-\Delta_{N,2N+1} + 1)^{3/4}) \subset W^{1,\infty}(\Omega)$, and hence it holds that

$$\begin{aligned} \|\nabla v(\cdot, t)\|_\infty &\leq C_{41} \|(-\Delta_N + 1)^{3/4} v(\cdot, t)\|_{2N+1} \\ &\leq C_{41} \left(\|(-\Delta_N + 1)^{3/4} e^{-t(\Delta_N - 1)} v_0\|_{2N+1} \right. \\ &\quad \left. + \int_0^t \|(-\Delta_N + 1)^{3/4} e^{-(t-s)(\Delta_N - 1)} u(\cdot, s)\|_{2N+1} ds \right) \\ &\leq C_{42} \left(\|(-\Delta_N + 1)^{3/4} v_0\|_{2N+1} + \int_0^t (t-s)^{-3/4} e^{-(t-s)} \|u(\cdot, s)\|_{2N+1} ds \right) \\ &\leq C_{43} \left(\|v_0\|_{2,2N+1} + \sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{2N+1} \right) \leq C_{44}. \end{aligned} \quad (3.39)$$

Taking $q \geq 1 + \gamma$ and recalling (3.24), we have

$$\begin{aligned} &\frac{1}{2q-1} \frac{d}{dt} \int_\Omega u^{2q-1} dx + (2q-2) \int_\Omega (\alpha u^\gamma + \beta) u^{2q-3} |\nabla u|^2 dx \\ &\leq (2q-2) \int_\Omega u^{2q-2} \nabla u \cdot \nabla v dx \\ &\leq (2q-2) \|\nabla v\|_\infty \left(\int_\Omega u^{2q+\gamma-3} |\nabla u|^2 dx \right)^{1/2} \cdot \left(\int_\Omega u^{2q-\gamma-1} dx \right)^{1/2} \\ &\leq \alpha(q-1) \int_\Omega u^{2q+\gamma-3} |\nabla u|^2 dx + \frac{q-1}{\alpha} \|\nabla v\|_\infty^2 \int_\Omega u^{2q-\gamma-1} dx \\ &\leq \alpha(q-1) \int_\Omega u^{2q+\gamma-3} |\nabla u|^2 dx + \frac{q-1}{\alpha} \|\nabla v\|_\infty^2 \left(\|u\|_{2q}^{2q} + |\Omega| \right), \end{aligned} \quad (3.40)$$

or

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} u^{2q-1} dx + \frac{\alpha(q-1)(2q-1)}{k^2} \int_{\Omega} |\nabla u^k|^2 dx \\ & \leq \frac{(q-1)(2q-1)}{\alpha} \|\nabla v\|_{\infty}^2 \left(\|u\|_{2q}^{2q} + |\Omega| \right). \end{aligned} \quad (3.41)$$

Here, we have from (3.24), (3.26) and (3.39) that

$$\begin{aligned} & \left(\frac{(q-1)(2q-1)}{\alpha} \|\nabla v\|_{\infty}^2 + 1 \right) \left(\|u\|_{2q}^{2q} + |\Omega| \right) \\ & \leq C_{44} q^2 \left\{ \int_{\Omega} |\nabla u^k|^2 dx + \int_{\Omega} u^{2k} dx \right\}^{r\theta/2} \cdot \left\{ \int_{\Omega} u^k dx \right\}^{r(1-\theta)} + C_{44} q^2 \\ & \leq \frac{\alpha(q-1)(2q-1)}{k^2} \left\{ \int_{\Omega} |\nabla u^k|^2 dx + \int_{\Omega} u^{2k} dx \right\} \\ & \quad + C_{45} q^{2(2/(r\theta))'} \left\{ \int_{\Omega} u^k dx \right\}^{r(1-\theta)(2/(r\theta))'} + C_{44} q^2. \end{aligned} \quad (3.42)$$

In other words,

$$\begin{aligned} & \left(\frac{(q-1)(2q-1)}{\alpha} \|\nabla v\|_{\infty}^2 + 1 \right) \left(\|u\|_{2q}^{2q} + |\Omega| \right) \\ & \leq \frac{\alpha(q-1)(2q-1)}{k^2} \left\{ \int_{\Omega} |\nabla u^k|^2 dx + \|u\|_{2q+\gamma-1}^{2q+\gamma-1} \right\} \\ & \quad + C_{45} q^{2(2/(r\theta))'} \left\{ \int_{\Omega} u^k dx \right\}^{r(1-\theta)(2/(r\theta))'} + C_{44} q^2. \end{aligned} \quad (3.43)$$

Here and henceforce, C_i ($i = 41, 42, 43, \dots, 50$) denote positive constants independent of $q \geq 1 + \gamma$ and $T_{\max}$.

Combining this with (3.30) and (3.41), we get that

$$\frac{d}{dt} \|u\|_{2q-1}^{2q-1} + \|u\|_{2q}^{2q} \leq C_{46} q^{2\kappa} \left\{ \int_{\Omega} u^{q+((\gamma-1)/2)} dx + 1 \right\}^{\Lambda}. \quad (3.44)$$

Because of $\gamma \in (0, 1)$ and (3.29), we have

$$\left(\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{2q-1}^{2q-1} + 1 \right) \leq C_{47} q^{2\kappa} \left(\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_q^q + 1 \right)^{\Lambda}. \quad (3.45)$$

For $q = 2^{j-1} + 1$, we put

$$\alpha_j = \sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{2q-1}^{2q-1} + 1, \quad \Lambda_j = \max(2, \tilde{\Lambda}_j), \quad (3.46)$$

where $\tilde{\Lambda}_j = \Lambda(q) = \Lambda(2^{j-1} + 1)$.

18 A quasi-linear parabolic system of chemotaxis

Here, we have $\Lambda_j \leq 2(1 + C_{48}2^{-j})$. In use of $\Lambda_j \geq 2$, we apply (3.45) as $q = 2^{j-1} + 1$ for $j = 0, 1, 2, \dots$. Then, we get that

$$\begin{aligned} \log \alpha_j &\leq \log C_{47} + 2\kappa \log(2^{j-1} + 1) + \Lambda_j \log \alpha_{j-1} \\ &\leq \log C_{47} + 2\kappa j \log 2 + \Lambda_j \log \alpha_{j-1} \end{aligned} \quad (3.47)$$

or

$$\log \alpha_j + 2\kappa(j+2) \log 2 + \log C_{47} \leq \Lambda_j (\log \alpha_{j-1} + 2\kappa(j+1) \log 2 + \log C_{47}), \quad (3.48)$$

which implies

$$\log \alpha_j + 2\kappa(j+2) \log 2 + \log C_{47} \leq \left(\prod_{i=1}^j \Lambda_i \right) (\log \alpha_0 + 4\kappa \log 2 + \log C_{47}), \quad (3.49)$$

or

$$\alpha_j \leq (C_{47} 2^{4\kappa} \alpha_0)^{\prod_{i=1}^j \Lambda_i} (C_{47} 2^{2\kappa(j+2)})^{-1}. \quad (3.50)$$

Hence, we obtain

$$\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{2^{j+1}} \leq (C_{47} 2^{4\kappa} \alpha_0)^{(\prod_{i=1}^j \Lambda_i)/(2^{j+1})} (C_{47} 2^{2\kappa(j+3)})^{-1/(2^{j+1})}. \quad (3.51)$$

Here, we apply

$$\prod_{i=1}^j \Lambda_i \leq 2^j \exp \left(C_{48} \sum_{i=1}^j 2^{-i} \right) \leq 2^j e^{C_{48}} = 2^j C_{49}, \quad (3.52)$$

and get that

$$\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{\infty} \leq C_{50} \left(\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_2^2 + 1 \right)^{C_{49}}. \quad (3.53)$$

The desired conclusion follows from this inequality and (3.38) with $q = 2$. $\square$

Let us confirm that Proposition 3.5 says that $\sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_{\infty}$ and $\sup_{0 \leq t < T_{\max}} \|v(\cdot, t)\|_{\infty}$ are estimated from above by constants independent of $T_{\max}$. We recall $p > N + 2$ and $\ell \in (0, 1 - (N + 2)/p)$. Then, (3.23) with $2q = p$ means that

$$\int_0^t e^{s/2} \|\Delta v\|_p^p ds \leq C_{51} \left(t \|v_0\|_{2,p}^p + \int_0^t e^{s/2} \|u\|_p^p ds \right) \quad (3.54)$$

and hence we obtain

$$\int_0^t e^{(s-t)/2} \|\Delta v\|_p^p ds \leq C_{52} \left(t e^{-t/2} \|v_0\|_{2,p}^p + \sup_{0 \leq t < T_{\max}} \|u(\cdot, t)\|_p^p \right). \quad (3.55)$$

Here, C_i ($i = 51, 52$) are independent of $t \in [0, T_{\max})$.

We take $\tau \in (0, (1/2) \min(T_{\max}, 1))$ in $1 < T_{\max} - \tau$ without loss of generality. Moreover, we take $T \in [\tau, T_{\max} - 1)$. Henceforce, C_i ($i = 53, 54, 55, \dots, 58$) denote positive constants independent of $T \in [\tau, T_{\max})$ and $T_{\max}$. In use of the reflection with respect to $\partial\Omega$, we have a domain $\tilde{\Omega} \supset \bar{\Omega}$ with smooth boundary and the extension $(\tilde{u}, \tilde{v})$ of (u, v) defined on $\tilde{\Omega}$. Then, we can apply [10, Theorem 3.1] for $(\tilde{u}, \tilde{v})$, and find $\theta \in (0, 1)$ such that

$$|\tilde{u}|_{\Omega \times (T, T+1)}^{(\theta)} \leq C_{53}. \quad (3.56)$$

In fact, $\int_T^{T+1} \|\Delta v\|_p^p dt$ is estimated from above independent of T and $T_{\max}$ by (3.55), and this implies (3.56). Then, it holds that

$$|u|_{\Omega \times (T, T+1)}^{(\theta)} \leq C_{53}, \quad (3.57)$$

and the parabolic regularity guarantees that

$$|u|_{Q_{T_{\max}}}^{(\theta)} \leq C_{54}. \quad (3.58)$$

Then,

$$|v|_{Q_{T_{\max}}}^{(2+\theta)} \leq C_{55} \quad (3.59)$$

follows from [10, Theorem 5.3].

We take $\zeta \in C^\infty(\mathbf{R})$ in $0 \leq \zeta \leq 1$ and

$$\zeta(s) = \begin{cases} 1 & \text{if } \tau \leq s, \\ 0 & \text{if } s \leq 0, \end{cases} \quad (3.60)$$

and set $\zeta_T(s) = \zeta(s - T)$. The function $W = \mathcal{A}(u)$ defined by

$$\mathcal{A}(s) = \int_0^s A(s') ds' \quad (3.61)$$

satisfies that

$$W_t = A\Delta W - (\nabla v \cdot \nabla W + Au\Delta v) \quad (3.62)$$

and hence we can deduce for $W\zeta_T$ that

$$\begin{aligned} (W\zeta_T)_t &= A\Delta(W\zeta_T) - (\nabla v \cdot \nabla(W\zeta_T)) \\ &\quad + (W\zeta'_T - Au\Delta v\zeta_T) \quad \text{in } \Omega \times [T, T+1], \\ \frac{\partial}{\partial v}(W\zeta_T) &= 0 \quad \text{on } \partial\Omega \times [T, T+1], \\ W(\cdot, T)\zeta_T(T) &= 0 \quad \text{in } \Omega. \end{aligned} \quad (3.63)$$

Here, we regard u and v as coefficients, and apply [10, Theorem 5.3] by (3.58). Then, we obtain

$$|u|_{\Omega \times (T+\tau, T+1)}^{(2+\theta)} = |u\zeta_T|_{\Omega \times (T, T+1)}^{(2+\theta)} \leq C_{56}, \quad (3.64)$$

or

$$|u|_{\Omega \times (\tau, T_{\max})}^{(2+\theta)} \leq C_{57}. \quad (3.65)$$

On the other hand we have the parabolic regularity locally in time, so that we obtain

$$|u|_{Q_{T_{\max}}}^{(2+\theta)}, |v|_{Q_{T_{\max}}}^{(2+\theta)} \leq C_{58}. \quad (3.66)$$

Finally, if $T_{\max} < \infty$, the solution (u, v) is extended after $T_{\max}$, a contradiction. Hence it holds that $T_{\max} = \infty$. We have the uniformly bounded solution globally in time, and the proof of Theorem 1.1 is complete.

References

- [1] N. D. Alikakos, *L^p bounds of solutions of reaction-diffusion equations*, Communications in Partial Differential Equations **4** (1979), no. 8, 827–868.
- [2] P. Biler, *Local and global solvability of some parabolic systems modelling chemotaxis*, Advances in Mathematical Sciences and Applications **8** (1998), no. 2, 715–743.
- [3] ———, *Global solutions to some parabolic-elliptic systems of chemotaxis*, Advances in Mathematical Sciences and Applications **9** (1999), no. 1, 347–359.
- [4] S. Childress and J. K. Percus, *Nonlinear aspects of chemotaxis*, Mathematical Biosciences. An International Journal **56** (1981), no. 3–4, 217–237.
- [5] H. Gajewski and K. Zacharias, *Global behaviour of a reaction-diffusion system modelling chemotaxis*, Mathematische Nachrichten **195** (1998), 77–114.
- [6] M. Giga, Y. Giga, and H. Sohr, *L^p estimate for abstract linear parabolic equations*, Japan Academy. Proceedings. Series A. Mathematical Sciences **67** (1991), no. 6, 197–202.
- [7] D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Lecture Notes in Mathematics, vol. 840, Springer, Berlin, 1981.
- [8] M. A. Herrero and J. J. L. Velázquez, *A blow-up mechanism for a chemotaxis model*, Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Serie IV **24** (1997), no. 4, 633–683 (1998).
- [9] E. F. Keller and L. A. Segel, *Initiation of slime mold aggregation viewed as an instability*, Journal of Theoretical Biology **26** (1970), 399–415.
- [10] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'tseva, *Linear and quasi-linear equations of parabolic type*, Izdat. "Nauka", Moscow, 1968, English translation; American Mathematical Society, Rhode Island, 1968.
- [11] T. Nagai, *Blow-up of radially symmetric solutions to a chemotaxis system*, Advances in Mathematical Sciences and Applications **5** (1995), no. 2, 581–601.
- [12] ———, *Blowup of nonradial solutions to parabolic-elliptic systems modeling chemotaxis in two-dimensional domains*, Journal of Inequalities and Applications **6** (2001), no. 1, 37–55.
- [13] T. Nagai and T. Senba, *Global existence and blow-up of radial solutions to a parabolic-elliptic system of chemotaxis*, Advances in Mathematical Sciences and Applications **8** (1998), no. 1, 145–156.
- [14] T. Nagai, T. Senba, and K. Yoshida, *Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis*, Funkcialaj Ekvacioj Japana Matematika Societo. Funkcialaj Ekvacioj. Serio Internacia **40** (1997), no. 3, 411–433.
- [15] V. Nanjundiah, *Chemotaxis, signal relaying, and aggregation morphology*, Journal of Theoretical Biology **42** (1973), 63–105.
- [16] T. Senba, *Blow-up of radially symmetric solutions to some systems of partial differential equations modelling chemotaxis*, Advances in Mathematical Sciences and Applications **7** (1997), no. 1, 79–92.

- [17] T. Senba and T. Suzuki, *Local and norm behavior of blowup solutions to a parabolic system of chemotaxis*, Journal of the Korean Mathematical Society **37** (2000), no. 6, 929–941.
- [18] ———, *Chemotactic collapse in a parabolic-elliptic system of mathematical biology*, Advances in Differential Equations **6** (2001), no. 1, 21–50.
- [19] ———, *Applied Analysis*, Imperial College Press, London, 2004.
- [20] T. Suzuki, *Free Energy and Self-Interacting Particles*, Progress in Nonlinear Differential Equations and Their Applications, vol. 62, Birkhäuser Boston, Massachusetts, 2005.
- [21] H. Tanabe, *Functional Analytic Methods for Partial Differential Equations*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 204, Marcel Dekker, New York, 1997.

Takasi Senba: Department of Applied Mathematics, Faculty of Technology, Miyazaki University,
1-1 Gakuen Kibanadai Nishi, Miyazaki-shi 889-2192, Japan
E-mail address: t0f302u@cc.miyazaki-u.ac.jp

Takasi Suzuki: Department of Mathematical Science, Graduate School of Engineering Science,
Osaka University, 1-3 Machikaneyama-machi, Toyonaka-shi 560-8531, Japan
E-mail address: suzuki@sigmath.es.osaka-u.ac.jp