COMMON FIXED POINTS OF ONE-PARAMETER NONEXPANSIVE SEMIGROUPS IN STRICTLY CONVEX BANACH SPACES

TOMONARI SUZUKI

Received 20 December 2003; Accepted 10 July 2005

One of our main results is the following convergence theorem for one-parameter nonexpansive semigroups: let C be a bounded closed convex subset of a Hilbert space E , and let $\{T(t) : t \in \mathbb{R}_+\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Fix $u \in C$ and $t_1, t_2 \in \mathbb{R}_+$ with $t_1 < t_2$. Define a sequence $\{x_n\}$ in C by $x_n = (1 - \alpha_n)/(t_2 - t_1) \int_{t_1}^{t_2} T(s)x_n ds + \alpha_n u$ for $n \in \mathbb{N}$, where $\{\alpha_n\}$ is a sequence in $(0, 1)$ converging to 0. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

Copyright © 2006 Tomonari Suzuki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let C be a closed convex subset of a Banach space E , and let T be a *nonexpansive mapping* on C , that is, $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. We know that T has a fixed point in the case that E is uniformly convex and C is bounded; see Browder [4], Göhde [10], and Kirk [15]. We denote by $F(T)$ the set of fixed points of T .

Let $\{T(t) : t \in \mathbb{R}_+\}$ be a *strongly continuous semigroup of nonexpansive mappings* (*nonexpansive semigroup*, in short) on a closed convex subset C of a Banach space E , that is,

- (i) for each $t \in \mathbb{R}_+$, $T(t)$ is a nonexpansive mapping on C ;
- (ii) $T(s+t) = T(s) \circ T(t)$ for all $s, t \in \mathbb{R}_+$;
- (iii) for each $x \in C$, the mapping $t \mapsto T(t)x$ from $\mathbb{R}_+$ into C is strongly continuous.

We also know that $\{T(t) : t \in \mathbb{R}_+\}$ has a common fixed point in the case that E is uniformly convex and C is bounded; see Browder [4]. Bruck [7] prove the following theorem.

THEOREM 1.1 (Bruck [7]). *Suppose a closed convex subset C of a Banach space has the fixed point property for nonexpansive mappings, and C is either weakly compact, or bounded and separable. Then for any commuting family S of nonexpansive mappings on C , the set of common fixed points of S is a nonempty nonexpansive retract of C .*

2 Nonexpansive semigroup in SC

This theorem yields that $\{T(t) : t \in \mathbb{R}_+\}$ has a common fixed point in the case that C has the fixed point property, and that C is weakly compact, or bounded and separable.

Several authors have studied about convergence theorems for nonexpansive semigroups; see [1, 2, 13, 16, 19, 21, 22] and others. For example, the following theorem is a corollary of Theorem 8 in [19].

THEOREM 1.2 (Shioji and Takahashi [19]). *Let C be a bounded closed convex subset of a Hilbert space E . Let $\{T(t) : t \in \mathbb{R}_+\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Let $\{\alpha_n\}$ and $\{t_n\}$ be sequences of real numbers satisfying $0 < \alpha_n < 1$, $\lim_n \alpha_n = 0$, $t_n > 0$ and $\lim_n t_n = \infty$. Fix $u \in C$ and define a sequence $\{x_n\}$ in C by*

$$x_n = \frac{1 - \alpha_n}{t_n} \int_0^{t_n} T(s)x_n ds + \alpha_n u \quad (1.1)$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

Also, Suzuki[21] proved the following theorem.

THEOREM 1.3 (Suzuki [21]). *Let E , C , $\{T(t) : t \in \mathbb{R}_+\}$ be as in Theorem 1.2. Let $\{\alpha_n\}$ and $\{t_n\}$ be sequences of real numbers satisfying $0 < \alpha_n < 1$, $t_n > 0$ and $\lim_n t_n = \lim_n \alpha_n/t_n = 0$. Fix $u \in C$ and define a sequence $\{x_n\}$ in C by*

$$x_n = (1 - \alpha_n)T(t_n)x_n + \alpha_n u \quad (1.2)$$

for $n \in \mathbb{N}$. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

We note that in these theorems, real sequences $\{t_n\}$ converge to 0 and ∞ . So, it is natural to study convergence theorems under the assumption that $\{t_n\}$ is a constant sequence. In this paper, motivated by Theorems 1.2 and 1.3, we consider such type of convergence theorems to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

2. Preliminaries

Throughout this paper we denote by $\mathbb{R}$ the set of real numbers, by $\mathbb{R}_+$ the set of nonnegative real numbers, and by $\mathbb{N}$ the set of positive integers. For a Banach space E , we also denote by E^* the dual space of E .

We recall that a Banach space E is called strictly convex if $\|x + y\|/2 < 1$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $x \neq y$. We know the following lemma.

LEMMA 2.1. *Let E be a Banach space. Then the following are equivalent:*

- (i) E is strictly convex;
- (ii) $\|\lambda x + (1 - \lambda)y\| < 1$ for all $\lambda \in (0, 1)$ and $x, y \in E$ with $\|x\| = \|y\| = 1$ and $x \neq y$;
- (iii) if $\|x\| = \|y\| = \|\lambda x + (1 - \lambda)y\|$ for some $\lambda \in (0, 1)$, then $x = y$.

A Banach space E is called uniformly convex if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $\|x + y\|/2 < 1 - \delta$ for all $x, y \in E$ with $\|x\| = \|y\| = 1$ and $\|x - y\| \geq \varepsilon$. It is clear that a uniformly convex Banach space is strictly convex. The norm of E is called Fréchet differentiable if for each $x \in E$ with $\|x\| = 1$, $\lim_{t \rightarrow 0} (\|x + ty\| - \|x\|)/t$ exists and is attained uniformly in $y \in E$ with $\|y\| = 1$. A Banach space E is said to have the Opial property [17]

if for each weakly convergent sequence $\{x_n\}$ in E with weak limit z , $\liminf_n \|x_n - z\| < \liminf_n \|x_n - y\|$ for all $y \in E$ with $y \neq z$. All Hilbert spaces, all finite dimensional Banach spaces and ℓ^p ($1 \leq p < \infty$) have the Opial property. Gossez and Lami Dozo[11] prove that every weakly compact convex subset of a Banach space with the Opial property has normal structure. We also know that every separable Banach space can be equivalently renormed so that it has the Opial property; see [23].

3. Common fixed points

In this section, we give our main results. The following proposition plays an important role in this paper.

PROPOSITION 3.1. *Let C be a closed convex subset of a strictly convex Banach space E . Let $\tau_\infty > 0$ and let $\{T(t) : t \in [0, \tau_\infty)\}$ be a family of mappings on C satisfying the following:*

- (i) *for each $t \in [0, \tau_\infty)$, $T(t)$ is nonexpansive;*
- (ii) *there exists a strictly increasing sequence $\{\tau_n\}$ in $[0, \tau_\infty)$ such that $\tau_1 = 0$, $\{\tau_n\}$ converges to τ_∞ , and mappings $t \mapsto T(t)x$ are weakly continuous on $[\tau_n, \tau_{n+1})$ for all $x \in C$ and $n \in \mathbb{N}$.*

Suppose that

$$\bigcap_{t \in [0, \tau_\infty)} F(T(t)) \neq \emptyset. \quad (3.1)$$

Then

$$\bigcap_{t \in [0, \tau_\infty)} F(T(t)) = F(S), \quad (3.2)$$

where S is a nonexpansive mapping on C defined by

$$Sx = \frac{1}{\tau_\infty} \int_0^{\tau_\infty} T(s)x \, ds \quad (3.3)$$

for all $x \in C$.

Remark 3.2. We do not assume $\{T(\cdot)\}$ is a nonexpansive semigroup.

Proof. Fix $f \in E^*$. Then the functions $t \mapsto f(T(t)x)$ from $[\tau_n, \tau_{n+1})$ into $\mathbb{R}$ are continuous on $[\tau_n, \tau_{n+1})$ for $x \in C$ and $n \in \mathbb{N}$. So, the functions $t \mapsto f(T(t)x)$ from $[0, \tau_\infty)$ into $\mathbb{R}$ are measurable for $x \in C$. We also have $\{T(t)x : t \in [0, \tau_\infty)\}$ is separable for each $x \in C$. Fix $w \in \bigcap_{t \in [0, \tau_\infty)} F(T(t))$. Since

$$\begin{aligned} \|T(t)x\| &= \|T(t)x\| - \|T(t)w\| + \|w\| \leq \|T(t)x - T(t)w\| + \|w\| \\ &\leq \|x - w\| + \|w\|, \end{aligned} \quad (3.4)$$

for $x \in C$ and $t \in [0, \tau_\infty)$, we have that the mappings $t \mapsto T(t)x$ are Bochner integrable for all $x \in C$ and hence S is well-defined. Using the separation theorem, we can easily prove

4 Nonexpansive semigroup in SC

that S is a mapping on C . Since

$$\begin{aligned}\|Sx - Sy\| &= \left\| \frac{1}{\tau_\infty} \int_0^{\tau_\infty} (T(s)x - T(s)y) ds \right\| \\ &\leq \frac{1}{\tau_\infty} \int_0^{\tau_\infty} \|T(s)x - T(s)y\| ds \\ &\leq \frac{1}{\tau_\infty} \int_0^{\tau_\infty} \|x - y\| ds = \|x - y\|\end{aligned}\tag{3.5}$$

for $x, y \in C$, S is nonexpansive. Therefore S is a nonexpansive mapping on C . It is obvious that $\bigcap_{t \in [0, \tau_\infty)} F(T(t)) \subset F(S)$. We assume that $z \in F(S) \setminus \bigcap_{t \in [0, \tau_\infty)} F(T(t))$. Then there exists $t_1 \in [0, \tau_\infty)$ such that $T(t_1)z \neq z$. Fix $g \in E^*$ with

$$\|g\| = 1, \quad g(T(t_1)z - z) = \|T(t_1)z - z\|.\tag{3.6}$$

For some $m \in \mathbb{N}$, t_1 belongs to $[\tau_m, \tau_{m+1})$. From the assumption (ii), there exists $t_2 \in (t_1, \tau_{m+1})$ such that

$$g(T(t)z - z) > \frac{1}{2} \|T(t_1)z - z\|\tag{3.7}$$

for all $t \in [t_1, t_2)$. Define nonexpansive mappings S_1 and S_2 on C by

$$\begin{aligned}S_1x &= \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} T(s)x ds, \\ S_2x &= \frac{1}{\tau_\infty - t_2 + t_1} \left(\int_0^{t_1} T(s)x ds + \int_{t_2}^{\tau_\infty} T(s)x ds \right)\end{aligned}\tag{3.8}$$

for all $x \in C$. We note that

$$Sx = \frac{t_2 - t_1}{\tau_\infty} S_1x + \frac{\tau_\infty - t_2 + t_1}{\tau_\infty} S_2x\tag{3.9}$$

for all $x \in C$. We have

$$\begin{aligned}g(S_1z - Sz) &= g\left(\frac{1}{t_2 - t_1} \int_{t_1}^{t_2} T(s)z ds - z\right) \\ &= g\left(\frac{1}{t_2 - t_1} \int_{t_1}^{t_2} (T(s)z - z) ds\right) \\ &= \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} g(T(s)z - z) ds \\ &\geq \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \frac{1}{2} \|T(t_1)z - z\| ds \\ &= \frac{1}{2} \|T(t_1)z - z\| > 0.\end{aligned}\tag{3.10}$$

Hence

$$g(S_2z - Sz) = \frac{t_2 - t_1}{\tau_\infty - t_2 + t_1} g(Sz - S_1z) < 0.\tag{3.11}$$

Therefore $S_1 z \neq S_2 z$. Fix $w \in \bigcap_{t \in [0, \tau_\infty)} F(T(t))$. Then we note that $S_1 w = S_2 w = w$. We have

$$\begin{aligned}
 \|z - w\| &= \|S_z - w\| = \left\| \frac{t_2 - t_1}{\tau_\infty} S_1 z + \frac{\tau_\infty - t_2 + t_1}{\tau_\infty} S_2 z - w \right\| \\
 &\leq \frac{t_2 - t_1}{\tau_\infty} \|S_1 z - w\| + \frac{\tau_\infty - t_2 + t_1}{\tau_\infty} \|S_2 z - w\| \\
 &= \frac{t_2 - t_1}{\tau_\infty} \|S_1 z - S_1 w\| + \frac{\tau_\infty - t_2 + t_1}{\tau_\infty} \|S_2 z - S_2 w\| \\
 &\leq \frac{t_2 - t_1}{\tau_\infty} \|z - w\| + \frac{\tau_\infty - t_2 + t_1}{\tau_\infty} \|z - w\| = \|z - w\|
 \end{aligned} \tag{3.12}$$

and hence

$$\|S_z - w\| = \|S_1 z - w\| = \|S_2 z - w\|. \tag{3.13}$$

This contradicts the strict convexity of E . Therefore, $F(S) \subset \bigcap_{t \in [0, \tau_\infty)} F(T(t))$. This completes the proof. $\square$

As a direct consequence of Proposition 3.1, we can prove the following, which was proved by Bruck [6]; see also [20].

COROLLARY 3.3 (Bruck [6]). *Let C be a closed convex subset of a strictly convex Banach space E . Let $\{T_n : n \in \mathbb{N}\}$ be a sequence of nonexpansive mappings on C . Suppose $\bigcap_{n=1}^\infty F(T_n)$ is nonempty. Let $\{\alpha_n\}$ be a sequence of positive numbers with $\sum_{n=1}^\infty \alpha_n = 1$. Define a nonexpansive mapping S on C by*

$$Sx = \sum_{n=1}^\infty \alpha_n T_n x \tag{3.14}$$

for $x \in C$. Then $F(S) = \bigcap_{n=1}^\infty F(T_n)$ holds.

Proof. Define a strictly increasing sequence $\{\tau_n\}$ in $[0, 1)$ by $\tau_1 = 0$ and

$$\tau_n = \sum_{k=1}^{n-1} \alpha_k \tag{3.15}$$

for $n \in \mathbb{N}$ with $n \geq 2$. We note that $\lim_n \tau_n = 1$. Define a family $\{T(t) : t \in [0, 1)\}$ of nonexpansive mappings as follows: If $\tau_n \leq t < \tau_{n+1}$, then

$$T(t)x = T_n x \tag{3.16}$$

for all $x \in C$. Then we note that

$$Sx = \sum_{n=1}^\infty \alpha_n T_n x = \sum_{n=1}^\infty \int_{\tau_n}^{\tau_{n+1}} T(s)x \, ds = \int_0^1 T(s)x \, ds = \frac{1}{1} \int_0^1 T(s)x \, ds \tag{3.17}$$

6 Nonexpansive semigroup in SC

for $x \in C$ and

$$\bigcap_{n=1}^{\infty} F(T_n) = \bigcap_{t \in [0,1)} F(T(t)). \quad (3.18)$$

So, by Proposition 3.1, we obtain the desired result. $\square$

As another direct consequence of Proposition 3.1, we obtain the following proposition.

PROPOSITION 3.4. *Let C be a closed convex subset of a strictly convex Banach space E . Let $\tau > 0$ and let $\{T(t) : t \in [0, \tau)\}$ be a family of mappings on C satisfying the following:*

- (i) *for each $t \in [0, \tau)$, $T(t)$ is nonexpansive;*
- (ii) *mappings $t \mapsto T(t)x$ are weakly continuous on $[0, \tau)$ for all $x \in C$.*

Suppose that

$$\bigcap_{t \in [0, \tau)} F(T(t)) \neq \emptyset. \quad (3.19)$$

Then

$$\bigcap_{t \in [0, \tau)} F(T(t)) = F(S), \quad (3.20)$$

where S is a nonexpansive mapping on C defined by

$$Sx = \frac{1}{\tau} \int_0^{\tau} T(s)x \, ds \quad (3.21)$$

for all $x \in C$.

Now, we prove one of our main results.

THEOREM 3.5. *Let C be a closed convex subset of a strictly convex Banach space E and let $\{T(t) : t \in \mathbb{R}_+\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Suppose that*

$$\bigcap_{t \in \mathbb{R}_+} F(T(t)) \neq \emptyset. \quad (3.22)$$

Fix $t_1, t_2 \in \mathbb{R}_+$ with $t_1 < t_2$, and define a nonexpansive mapping S on C by

$$Sx = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} T(s)x \, ds \quad (3.23)$$

for all $x \in C$. Then

$$\bigcap_{t \in \mathbb{R}_+} F(T(t)) = F(S) \quad (3.24)$$

holds.

Proof. It is clear that $\bigcap_{t \in \mathbb{R}_+} F(T(t)) \subset F(S)$. Fix $w \in F(S)$. By Proposition 3.4, we have

$$\bigcap_{t \in [t_1, t_2]} F(T(t)) = F(S). \quad (3.25)$$

So, $T(t)w = w$ for $t \in [t_1, t_2]$. Hence, for every $t \in [0, (t_2 - t_1)/2]$, we have

$$T(t)w = T(t) \circ T(t_1)w = T(t + t_1)w = w. \quad (3.26)$$

Let $t \in \mathbb{R}_+$ be fixed. Then there exist $m \in \mathbb{N} \cup \{0\}$ and $u \in [0, (t_2 - t_1)/2]$ such that $t = u + m(t_2 - t_1)/2$. We have

$$T(t)w = T\left(u + m \frac{t_2 - t_1}{2}\right)w = T(u) \circ T\left(\frac{t_2 - t_1}{2}\right)^m w = T(u)w = w, \quad (3.27)$$

where $T((t_2 - t_1)/2)^0$ is the identity mapping on C . Therefore w is a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$. This completes the proof. $\square$

Similarly we can prove the following theorem.

THEOREM 3.6. *Let C be a closed convex subset of a strictly convex Banach space E and let $\{T_n(t) : t \in \mathbb{R}_+, n \in \mathbb{N}\}$ be a sequence of strongly continuous semigroups of nonexpansive mappings on C . Let $\{U_n : n \in \mathbb{N}\}$ be a sequence of nonexpansive mappings on C . Suppose that*

$$\bigcap_{n=1}^{\infty} \bigcap_{t \in \mathbb{R}_+} F(T_n(t)) \cap \bigcap_{n=1}^{\infty} F(U_n) \neq \emptyset. \quad (3.28)$$

Let $\{t_n\}$, $\{u_n\}$, $\{\alpha_n\}$ and $\{\beta_n\}$ be real sequences such that $0 \leq t_n < u_n$, $\alpha_n > 0$ and $\beta_n > 0$ for all $n \in \mathbb{N}$, and $\sum_{n=1}^{\infty} \alpha_n + \sum_{n=1}^{\infty} \beta_n = 1$. Define a nonexpansive mapping S on C by

$$Sx = \sum_{n=1}^{\infty} \frac{\alpha_n}{u_n - t_n} \int_{t_n}^{u_n} T_n(s)x \, ds + \sum_{n=1}^{\infty} \beta_n U_n x \quad (3.29)$$

for all $x \in C$. Then

$$\bigcap_{n=1}^{\infty} \bigcap_{t \in \mathbb{R}_+} F(T_n(t)) \cap \bigcap_{n=1}^{\infty} F(U_n) = F(S). \quad (3.30)$$

holds.

We recall that a closed convex subset C of a Banach space E is said to have the fixed point property for nonexpansive mappings (FPP, in short) if for every bounded closed convex subset D of C , every nonexpansive mapping on D has a fixed point. So, by the results of Browder [4] and Göhde [10], every uniformly convex Banach space has FPP. Also, by Kirk's fixed point theorem [15], every weakly compact convex subset with normal structure has FPP.

As a direct consequence of Theorem 3.6, we obtain the following corollary.

8 Nonexpansive semigroup in SC

COROLLARY 3.7. *Let $E, C, \{\{T_n(t) : t \in \mathbb{R}_+\} : n \in \mathbb{N}\}, \{U_n : n \in \mathbb{N}\}, \{t_n\}, \{u_n\}, \{\alpha_n\}$, and $\{\beta_n\}$ be as in Theorem 3.6. Assume that C is weakly compact and has FPP, and*

$$T_m(s) \circ T_n(t) = T_n(t) \circ T_m(s), \quad U_m \circ U_n = U_n \circ U_m, \quad U_m \circ T_n(t) = T_n(t) \circ U_m \quad (3.31)$$

for all $s, t \in \mathbb{R}_+$ and $m, n \in \mathbb{N}$. Define a nonexpansive mapping S on C as in Theorem 3.6. Then

$$\bigcap_{n=1}^{\infty} \bigcap_{t \in \mathbb{R}_+} F(T_n(t)) \cap \bigcap_{n=1}^{\infty} F(U_n) = F(S) \neq \emptyset. \quad (3.32)$$

holds.

4. Convergence theorems

Using Theorem 3.5, we can prove many convergence theorems to a common fixed point of nonexpansive semigroups. In this section, we state some of them.

From the result of Ishikawa [14], we obtain the following theorem see also Edelstein [8].

THEOREM 4.1. *Let C be a compact convex subset of a strictly convex Banach space E . Let $\{T(t) : t \in \mathbb{R}_+\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Fix $t_1, t_2 \in \mathbb{R}_+$ with $t_1 < t_2$. Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and*

$$x_{n+1} = \frac{\alpha_n}{t_2 - t_1} \int_{t_1}^{t_2} T(s)x_n ds + (1 - \alpha_n)x_n \quad (4.1)$$

for $n \in \mathbb{N}$, where $\{\alpha_n\}$ is a sequence in $[0, 1]$ satisfying $\sum_{n=1}^{\infty} \alpha_n = \infty$ and $\limsup_n \alpha_n < 1$. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

From the results of Edelstein and O'Brien [9], and Reich [18], we obtain the following theorem.

THEOREM 4.2. *Let E be a Banach space. Suppose either of the following holds:*

- (i) E is strictly convex and has the Opial property; or
- (ii) E is uniformly convex and its norm is Fréchet differentiable.

Let C be a weakly compact convex subset of E , and let $\{T(t) : t \in \mathbb{R}_+\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Fix $t_1, t_2 \in \mathbb{R}_+$ with $t_1 < t_2$. Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and

$$x_{n+1} = \frac{\alpha}{t_2 - t_1} \int_{t_1}^{t_2} T(s)x_n ds + (1 - \alpha)x_n \quad (4.2)$$

for $n \in \mathbb{N}$, where α is a constant number in $(0, 1)$. Then $\{x_n\}$ converges weakly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

We note that

$$x \mapsto (1 - \alpha)Tx + \alpha u \quad (4.3)$$

is a contractive mapping if T is a nonexpansive mapping and $\alpha \in (0, 1)$. By the Banach contraction principle [3], such mappings have a unique fixed point. From the results of Browder [5], and Wittmann [24], we obtain the following theorem; see also [12]. Compare Theorem 4.3 with Theorems 1.2 and 1.3.

THEOREM 4.3. *Let C be a bounded closed convex subset of a Hilbert space E , and let $\{T(t) : t \in \mathbb{R}_+\}$ be a strongly continuous semigroup of nonexpansive mappings on C . Fix $u \in C$ and $t_1, t_2 \in \mathbb{R}_+$ with $t_1 < t_2$. Define a sequence $\{x_n\}$ in C by*

$$x_n = \frac{1 - \alpha_n}{t_2 - t_1} \int_{t_1}^{t_2} T(s)x_n ds + \alpha_n u \quad (4.4)$$

for $n \in \mathbb{N}$, where $\{\alpha_n\}$ is a sequence in $(0, 1)$ converging to 0. Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

THEOREM 4.4. *Let $E, C, \{T(t) : t \in \mathbb{R}_+\}$, u, t_1 and t_2 be as in Theorem 4.3. Define a sequence $\{x_n\}$ in C by $x_1 \in C$ and*

$$x_{n+1} = \frac{1 - \alpha_n}{t_2 - t_1} \int_{t_1}^{t_2} T(s)x_n ds + \alpha_n u \quad (4.5)$$

for $n \in \mathbb{N}$, where $\{\alpha_n\}$ is a sequence in $[0, 1]$ satisfying the following:

$$\lim_{n \rightarrow \infty} \alpha_n = 0; \quad \sum_{n=1}^{\infty} \alpha_n = \infty; \quad \sum_{n=1}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty. \quad (4.6)$$

Then $\{x_n\}$ converges strongly to a common fixed point of $\{T(t) : t \in \mathbb{R}_+\}$.

Acknowledgment

The author is supported in part by Grants-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology.

References

- [1] S. Atsushiba and W. Takahashi, *Strong convergence theorems for one-parameter nonexpansive semigroups with compact domains*, Fixed Point Theory and Applications (Y. J. Cho, J. K. Kim, and S. M. Kang, eds.), vol. 3, Nova Science, New York, 2002, pp. 15–31.
- [2] J.-B. Baillon, *Quelques propriétés de convergence asymptotique pour les semi-groupes de contractions impaires*, Comptes Rendus de l'Académie des Sciences. Série A-B **283** (1976), no. 3, A75–A78.
- [3] S. Banach, *Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales*, Fundamenta Mathematicae **3** (1922), 133–181.
- [4] F. E. Browder, *Nonexpansive nonlinear operators in a Banach space*, Proceedings of the Academy of Natural Sciences of Philadelphia U.S.A **54** (1965), 1041–1044.
- [5] ———, *Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces*, Archive for Rational Mechanics and Analysis **24** (1967), no. 1, 82–90.
- [6] R. E. Bruck Jr., *Properties of fixed-point sets of nonexpansive mappings in Banach spaces*, Transactions of the American Mathematical Society **179** (1973), 251–262.

- [7] ———, *A common fixed point theorem for a commuting family of nonexpansive mappings*, Pacific Journal of Mathematics **53** (1974), no. 1, 59–71.
- [8] M. Edelstein, *A remark on a theorem of M. A. Krasnoselski*, The American Mathematical Monthly **73** (1966), no. 5, 509–510.
- [9] M. Edelstein and R. C. O'Brien, *Nonexpansive mappings, asymptotic regularity and successive approximations*, The Journal of the London Mathematical Society. Second Series **17** (1978), no. 3, 547–554.
- [10] D. Göhde, *Zum Prinzip der kontraktiven Abbildung*, Mathematische Nachrichten **30** (1965), 251–258.
- [11] J.-P. Gossez and E. Lami Dozo, *Some geometric properties related to the fixed point theory for nonexpansive mappings*, Pacific Journal of Mathematics **40** (1972), no. 3, 565–573.
- [12] B. Halpern, *Fixed points of nonexpanding maps*, American Mathematical Society. Bulletin **73** (1967), 957–961.
- [13] N. Hirano, *Nonlinear ergodic theorems and weak convergence theorems*, Journal of the Mathematical Society of Japan **34** (1982), no. 1, 35–46.
- [14] S. Ishikawa, *Fixed points and iteration of a nonexpansive mapping in a Banach space*, Proceedings of the American Mathematical Society **59** (1976), no. 1, 65–71.
- [15] W. A. Kirk, *A fixed point theorem for mappings which do not increase distances*, The American Mathematical Monthly **72** (1965), no. 9, 1004–1006.
- [16] I. Miyadera and K. Kobayasi, *On the asymptotic behaviour of almost-orbits of nonlinear contraction semigroups in Banach spaces*, Nonlinear Analysis **6** (1982), no. 4, 349–365.
- [17] Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, American Mathematical Society. Bulletin **73** (1967), 591–597.
- [18] S. Reich, *Weak convergence theorems for nonexpansive mappings in Banach spaces*, Journal of Mathematical Analysis and Applications **67** (1979), no. 2, 274–276.
- [19] N. Shioji and W. Takahashi, *Strong convergence theorems for asymptotically nonexpansive semigroups in Hilbert spaces*, Nonlinear Analysis **34** (1998), no. 1, 87–99.
- [20] T. Suzuki, *Convergence theorems to common fixed points for infinite families of nonexpansive mappings in strictly convex Banach spaces*, Nihonkai Mathematical Journal **14** (2003), no. 1, 43–54.
- [21] ———, *On strong convergence to common fixed points of nonexpansive semigroups in Hilbert spaces*, Proceedings of the American Mathematical Society **131** (2003), no. 7, 2133–2136.
- [22] T. Suzuki and W. Takahashi, *Strong convergence of Mann's type sequences for one-parameter nonexpansive semigroups in general Banach spaces*, Journal of Nonlinear and Convex Analysis **5** (2004), no. 2, 209–216.
- [23] D. van Dulst, *Equivalent norms and the fixed point property for nonexpansive mappings*, The Journal of the London Mathematical Society. Second Series **25** (1982), no. 1, 139–144.
- [24] R. Wittmann, *Approximation of fixed points of nonexpansive mappings*, Archiv der Mathematik **58** (1992), no. 5, 486–491.

Tomonari Suzuki: Department of Mathematics, Kyushu Institute of Technology, Sensuicho, Tobata, Kitakyushu 804-8550, Japan
E-mail address: suzuki-t@mns.kyutech.ac.jp