ON CALCULATION OF THE RELATIVE INDEX OF A FIXED POINT IN THE NONDEGENERATE CASE

A. V. GUMINSKAYA AND P. P. ZABREIKO

Received 26 June 2005; Accepted 1 July 2005

The paper is devoted to the calculation of the index of a zero and the asymptotic index of a linear completely continuous nonnegative operator. Also the case of a nonlinear completely continuous operator A whose domain and image are situated in a closed convex set Q of a Banach space is considered. For this case, we formulate the rules for calculating the index of an arbitrary fixed point and the asymptotic index under the assumption that the corresponding linearizations exist and the operators of derivative do not have eigenvectors with eigenvalue 1 in some wedges.

Copyright © 2006 A. V. Guminskaya and P. P. Zabreiko. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let Q be a closed convex set in a Banach space and let $A : Q \rightarrow Q$ be a completely continuous operator. In [4] the calculation problem of a fixed point index of a vector field $I - A$ was formulated. In the simplest case, when Q is a cone, this problem was investigated in the articles by Isaenko [3], Mukhamadiev and Sabirov [5], and Pokornyi [6] (see also references in [8]). Later, in [1, 2] Dancer presented the general formula for the fixed point index of a completely continuous operator A with its domain and image in an arbitrary closed convex set. However, the case of an asymptotic index was not considered. Note that for the case of a cone this problem was earlier considered in the articles by Pokornyi [6] and Pokornyi and Astaf'eva [7]. The present paper concerns the cases of a wedge and an arbitrary closed convex set. In the latter case, the calculation of a fixed point index is reduced to the index calculation with respect to a specially constructed wedge. We also show that for the infinity singular point one needs to take the wedge

$$W_\infty = \{h \in X : x + th \in Q \ (x \in Q, 0 \leq t < \infty)\}. \quad (1.1)$$

2. Index of a linear operator

Let X be a Banach space, W a wedge in X (this means that W is a closed subset of X such that $W + W \subseteq W$ and $\lambda W \subset W$ for $\lambda \geq 0$), and A a linear completely continuous operator such that

$$AW \subseteq W. \quad (2.1)$$

Let $L = W \cap (-W)$. Then L is the maximal subspace which is contained in W . From (2.1) and the linearity of A it follows that the inclusion

$$A(L) \subseteq L \quad (2.2)$$

holds.

Consider a quotient space X/L and a quotient mapping $[\cdot] : X \rightarrow X/L$. It is easy to check that the image $\widehat{W}$ of W under the quotient mapping $[\cdot]$ is a cone in X/L .

From (2.2) it follows that the operator A induces a linear mapping $\widehat{A}$ of the quotient space X/L into itself such that $\widehat{A}[x] = [Ax]$. And by (2.1) we have that the cone $\widehat{W}$ is invariant under the operator $\widehat{A}$, that is, $\widehat{A}$ is nonnegative in the quotient space X/L .

THEOREM 2.1. *Let A be a linear completely continuous operator, acting in a Banach space X , and let W be a wedge that is invariant under the operator A . Suppose that $Ax \neq x$ for $x \in W$, $x \neq 0$. Then $\rho(\widehat{A}) \neq 1$ and*

$$\text{ind}(0, I - A; W) = \text{ind}(\infty, I - A; W) = \begin{cases} (-1)^{\beta(A|_L)} & \text{if } \rho(\widehat{A}) < 1, \\ 0 & \text{if } \rho(\widehat{A}) > 1, \end{cases} \quad (2.3)$$

where $A|_L$ is the restriction of the operator A to the space L , $\beta(A|_L)$ is the sum of multiplicities of eigenvalues of $A|_L$, greater than 1, and $\rho(\widehat{A})$ is the positive spectral radius of the operator $\widehat{A}$.

Proof. If $Ax \neq x$ for $x \in W$, $x \neq 0$, then zero and infinity singular points of the vector field $\Phi = I - A$ are isolated in W . Hence the relative indices $\text{ind}(0, I - A; W)$ and $\text{ind}(\infty, I - A; W)$ are well posed. In this case, by the definition of index at infinity (see, e.g., [4]), since the operator A has no more fixed points in W , it follows that $\text{ind}(\infty, I - A; W) = \text{ind}(0, I - A; W)$.

To calculate the index $\text{ind}(0, \Phi; W)$ we will consider two possible cases: when the spectral radius $\rho(\widehat{A})$ of the operator $\widehat{A}$ is less than 1 and when it is greater than 1. The case $\rho(\widehat{A}) = 1$ is impossible. Indeed, if this is not true, then there exists an element $[x]^* \neq 0$ of the quotient space X/L such that $\widehat{A}[x]^* = [x]^*$. In other words, there exist $u \in W$, $u \notin L$, and $y \in L$ such that the equality $u - Au = y$ holds. Let us explore the solvability in L of $(I - A)w = y$. If it had a solution $w \in L$, then the vector $u - w$ would be an eigenvector of the operator A , corresponding to the eigenvalue 1 and would be in W , which would contradict our assumptions. If we supposed that the equation had no solutions in L , then the operator $I - A$ would be invertible in L , which is impossible.

In the case $\rho(\widehat{A}) < 1$, let us show that the vector field $\Phi x = x - Ax$ is linearly relatively homotopic on $S_W = \{x \in W : \|x\| = 1\}$ to the field $\Phi_1 x = x - AQ(x)$, where $Q : X \rightarrow L$ is

a projection (in general, nonlinear) of X on L (see [4, Theorem 18.1]). To prove this fact assume the converse, that is, that

$$x = (1 - \lambda)Ax + \lambda AQx \quad (2.4)$$

has a solution for some $x \in S_W$ and $\lambda \in [0, 1]$. Now if we rewrite this equation for the operator $\hat{A}$ and recall (2.2), we get

$$[x] = (1 - \lambda)\hat{A}[x] \quad ([x] \in \hat{S}_W, \lambda \in [0, 1]), \quad (2.5)$$

where $\hat{S}_W = \{[x] \in \hat{W} : \|[x]\| = 1\}$.

Under our assumption on the spectral radius of the operator $\hat{A}$ this yields $[x] = 0$. In this case, we obtain $x = Qx$ and then (2.4) implies $x = Ax$ for $x \in S_W$, which contradicts the assumption of our theorem.

Thus the relative index $\text{ind}(0, I - A; W)$ is equal to the relative index $\text{ind}(0, I - AQ; W)$, which actually is the Leray-Schauder zero fixed point index of the restriction $A|_L$ of the operator A to the subspace L and we can calculate it by the well-known formula (see, e.g., [4, Theorem 21.1])

$$\text{ind}(0, I - AQ; W) = (-1)^{\beta(A|_L)}, \quad (2.6)$$

where $\beta(A|_L)$ is the sum of multiplicities of eigenvalues of $A|_L$, greater than 1.

Now assume that $\rho(\hat{A}) > 1$. In this case, there exists an element $[x]^*$ in the cone $\hat{W}$ such that $\hat{A}[x]^* = \rho[x]^*$ ($\rho > 1$). In other words, there exist an element $x^* \notin L$ of the wedge W (we assume that $\|x^*\| = 1$) and an element z of the subspace L such that $Ax^* = \rho x^* + z$.

Show that the vector field $\Phi x = x - Ax$ is linearly relatively homotopic on S_W to the field $\Phi_2 x = x - cx^*$, where the constant c will be defined later. Let us show that

$$x = (1 - \lambda)Ax + \lambda cx^* \quad (2.7)$$

has no solutions for $x \in S_W, \lambda \in [0, 1]$.

For $\lambda = 0$, (2.7) coincides with $x = Ax$. The latter equation has no solutions for $x \in S_W$. For $0 < \lambda \leq 1$ from (2.7), it follows that there exists a real $t > 0$ such that

$$x \geq tx^*. \quad (2.8)$$

We claim that there exists the maximal of such reals: $\xi = \max_{x \in S_W, x \geq tx^*} t$. We argue by contradiction. The inequality $x^* \leq x/t$ implies that if t tends to infinity, then $x^* \leq 0$. On the other hand, x^* is an element of the wedge W , thus $x^* \geq 0$. Hence we get that $x^* \in L$, which contradicts its choice.

Further, (2.7) and (2.8) imply

$$x \geq (1 - \lambda)\xi \rho x^* + \lambda cx^* = ((1 - \lambda)\xi \rho + \lambda c)x^*. \quad (2.9)$$

Choose $c > \xi$ and observe that for this c the inequality $(1 - \lambda)\xi \rho + \lambda c > \xi$ holds. Indeed, for $\lambda = 0$ this follows from $\rho > 1$. For $\lambda = 1$ this follows from our assumption for c . For other $\lambda \in (0, 1)$ the inequality holds as the result of two previous cases.

Figure 2.1

Therefore for the chosen c , the real ξ is not maximal among t such that $x \geq tx^*$. This contradiction proves the nondegeneracy of the linear homotopy connecting vector fields Φ and Φ_2 . Consequently, $\text{ind}(0, \Phi; W) = \text{ind}(0, \Phi_2; W)$.

To calculate $\text{ind}(0, \Phi_2; W)$ we will use the corollary of the Hahn-Banach theorem. According to it there exists a functional $l \in X^*$ such that $l(W) \geq 0$ and $l(x^*) = 1$ (then $\|l\| \geq 1$). Since (2.8), we obtain $\xi \leq l(x) \leq \|l\|$ for $x \in S_W$. And our assumption of c yields $c > \|l\| \geq 1$. Now we can show that the vector field $\Phi_2 x = x - cx^*$ is nondegenerate on $\text{cl}B_W = \{x \in W : \|x\| \leq 1\}$. Indeed,

$$\|x - cx^*\| \geq \|cx^*\| - 1 = c - 1 > 0. \quad (2.10)$$

Hence, by the relative rotation property, $\text{ind}(0, \Phi_2; W) = 0$. This completes the proof. $\square$

In applications there usually exist a complement X_1 of the linear hull $L(W) = W - W$ of the wedge W to X and a complement X_2 of a maximal subspace $X_3 = W \cap (-W)$ in W to $L(W)$. Then X can be presented as the direct sum of subspaces (see Figure 2.1)

$$X = X_1 \dot{+} X_2 \dot{+} X_3. \quad (2.11)$$

From (2.1) and the linearity of the operator A it follows that the inclusions

$$A(X_2 \dot{+} X_3) \subseteq X_2 \dot{+} X_3, \quad A(X_3) \subseteq X_3 \quad (2.12)$$

hold.

Assume that an intersection $W \cap X_2$ is not empty. Then it is easy to prove that this set is a cone K in X_2 . It generates the order relation in X_2 by the following rule: $x \leq y$ if $y - x \in K$. The cone K can be set as $K = \{x \in X_2 : x \geq 0\}$.

It can be proved that under such decomposition of the space X , the wedge W is invariant under a linear operator A if and only if A is determined by the matrix

$$A = \begin{pmatrix} a_{11} & 0 & 0 \\ a_{21} & a_{22} & 0 \\ a_{31} & a_{32} & a_{33} \end{pmatrix}, \quad (2.13)$$

where $a_{22} \geq 0$. Furthermore, one can show that zero and infinity singular points of such operator A are isolated in W if and only if 1 is not an eigenvalue of the operators a_{22} and a_{33} . In this case, Theorem 2.1 can be formulated in the following way.

THEOREM 2.2. *Let A be a linear completely continuous operator, acting in a Banach space X , and let W be a wedge that is invariant under the operator A . Then A can be defined by matrix (2.13). If 1 is not an eigenvalue of the operators a_{22} and a_{33} , then zero and infinity singular points of the operator A are isolated in W and*

$$\text{ind}(0, I - A; W) = \text{ind}(\infty, I - A; W) = \begin{cases} (-1)^{\beta(a_{33})} & \text{if } \rho(a_{22}) < 1, \\ 0 & \text{if } \rho(a_{22}) > 1, \end{cases} \quad (2.14)$$

where $\rho(a_{22})$ is the spectral radius of the operator a_{22} and $\beta(a_{33})$ is the sum of multiplicities of eigenvalues of the operator a_{33} , greater than 1.

3. Index of a nonlinear operator

Let A be a nonlinear operator and let Q be a closed convex set of a Banach space X that is invariant under the operator A . In this part, we discuss the relative fixed point index calculations of A under the assumption that A has a Fréchet derivative at its fixed point. Recall that an operator A is called *differentiable at the point x_0 with respect to Q* , if

$$\lim_{\substack{x \in Q, \\ x \rightarrow x_0}} \frac{\|Ax - Ax_0 - A'(x_0)x\|}{\|x\|} = 0, \quad (3.1)$$

and *differentiable at infinity with respect to Q* , if

$$\lim_{\substack{x \in Q, \\ x \rightarrow \infty}} \frac{\|Ax - A'(\infty)x\|}{\|x\|} = 0, \quad (3.2)$$

where $A'(x_0)$ and $A'(\infty)$ are linear operators.

Let Q be an arbitrary closed convex set of a Banach space X . Assume that Q is invariant under a completely continuous operator A . Let x_0 be a fixed point of the operator A and let A have a Fréchet derivative at the point x_0 with respect to Q .

The following lemmas show that in the case when the fixed point x_0 is not the infinity singular point some wedge W_{x_0} comprising the set Q is invariant under the mapping of the derivative $A'(x_0)$, whereas in the case of the infinity singular point some wedge W_∞ lying in the set Q is invariant under the mapping of derivative $A'(\infty)$.

LEMMA 3.1. *Let A be a completely continuous operator, acting in a Banach space X , and let $Q \subseteq X$ be a closed convex set that is invariant under A . Let $x_0 \in Q$ be a fixed point of the operator A and let A have Fréchet derivative at x_0 with respect to Q . Then the wedge*

$$W_{x_0} = \text{cl}\{h = t(x - x_0) : t \geq 0, x \in Q\} \quad (3.3)$$

is invariant under the mapping of the derivative $A'(x_0)$.

Proof. To prove that W_{x_0} is actually a wedge it is sufficient to show that W_{x_0} is closed under addition of its elements and their multiplication by nonnegative reals. If $h_1, h_2 \in W_{x_0}$, then $h_1 = t_1(x_1 - x_0)$, $h_2 = t_2(x_2 - x_0)$, where $x_1, x_2 \in Q$ and $0 \leq t_1, t_2 < \infty$. Thus for any α and β for which $0 \leq \alpha, \beta < \infty$, and $\alpha t_1 + \beta t_2 > 0$, we have

$$\begin{aligned} \alpha h_1 + \beta h_2 &= \alpha t_1(x_1 - x_0) + \beta t_2(x_2 - x_0) \\ &= (\alpha t_1 + \beta t_2) \left(\frac{\alpha t_1}{\alpha t_1 + \beta t_2}(x_1 - x_0) + \frac{\beta t_2}{\alpha t_1 + \beta t_2}(x_2 - x_0) \right) \\ &= (\alpha t_1 + \beta t_2) \left(\frac{\alpha t_1}{\alpha t_1 + \beta t_2}x_1 + \frac{\beta t_2}{\alpha t_1 + \beta t_2}x_2 - x_0 \right). \end{aligned} \quad (3.4)$$

An element

$$\frac{\alpha t_1}{\alpha t_1 + \beta t_2}x_1 + \frac{\beta t_2}{\alpha t_1 + \beta t_2}x_2 \quad (3.5)$$

is a convex combination of elements $x_1, x_2 \in Q$ and therefore it is in Q . Thus (3.4) implies $\alpha h_1 + \beta h_2 \in W_{x_0}$. If $\alpha t_1 + \beta t_2 = 0$, then $\alpha h_1 + \beta h_2 = 0$ and thus $\alpha h_1 + \beta h_2 \in W_{x_0}$.

Let h be an arbitrary nonzero element of W_{x_0} . Then there exist $x \in Q$ and $t > 0$ such that $h = t(x - x_0)$. From the differentiability of the operator A at x_0 and the linearity of the operator $A'(x_0)$ it follows that

$$t(Ax - x_0) = A'(x_0)h + t\omega\left(\frac{h}{t}\right). \quad (3.6)$$

Since $A(Q) \subseteq Q$, the element in the left-hand side of this equality is in W_{x_0} . Taking the limit as $t \rightarrow \infty$, by the closedness of W_{x_0} , we get $A'(x_0)h \in W_{x_0}$. This completes the proof. $\square$

LEMMA 3.2. *Let A be a completely continuous operator, acting in a Banach space X , and let $Q \subseteq X$ be a closed convex unbounded set that is invariant under A . Let A have Fréchet derivative at infinity with respect to Q . Then the wedge*

$$W_\infty = \{h \in X : x + th \in Q(x \in Q, 0 \leq t < \infty)\} \quad (3.7)$$

is invariant under the mapping of the derivative $A'(\infty)$.

Proof. To prove that W_∞ is invariant under the operator $A'(\infty)$, it suffices to show that there exists an element $x_* \in Q$ such that $x_* + tA'(\infty)h \in Q$ for any $h \in W_\infty$ and $0 \leq t < \infty$. Let $x_* \in Q$, then for any $\lambda > 0$, $t \geq 0$, $h \in W_\infty$ we have $x_* + \lambda th \in Q$. Since Q is invariant

under the operator A , we get $A'(\infty)x_* + \lambda t A'(\infty)h + w(x_* + \lambda th) \in Q$ for any $\lambda \geq 0, t \geq 0, h \in W_\infty$. Then, by the convexity of Q , for any $\lambda \geq 0$ we have

$$\left(1 - \frac{1}{\lambda}\right)x_* + \frac{1}{\lambda}A'(\infty)x_* + tA'(\infty)h + \frac{1}{\lambda}w(x_* + \lambda th) \in Q. \quad (3.8)$$

Taking the limit as $\lambda \rightarrow \infty$, by the closedness of Q , we get $x_* + tA'(\infty)h \in Q$ for any $t \geq 0, h \in W_\infty$. This completes the proof. $\square$

The following theorem specifies the main result of Dancer [2, Theorem 1].

THEOREM 3.3. *Let A be a completely continuous operator, acting in a Banach space X , and let $Q \subseteq X$ be a closed convex set that is invariant under A . Let $x_0 \in \partial Q$ be a fixed point of the operator A and let A have Fréchet derivative at x_0 with respect to Q . Then the wedge*

$$W_{x_0} = \text{cl} \{h = t(x - x_0) : t \geq 0, x \in Q\} \quad (3.9)$$

is invariant under the mapping of the derivative $A'(x_0)$.

If $A'(x_0)x \neq x$ for $x \in W_{x_0}, x \neq 0$, then the fixed point x_0 of the vector field $\Phi = I - A$ is isolated in Q and

$$\text{ind}(x_0, I - A; Q) = \text{ind}(0, I - A'(x_0); W_{x_0}). \quad (3.10)$$

Proof. Without loss of generality it can be assumed that $x_0 = 0$ (in the opposite case, the whole argument needs to be made for the operator $A(x_0 + x) - x_0$).

From the differentiability of the operator A at the point 0 it follows that there exists a linear operator $B = A'(0)$ such that

$$Ax = Bx + w(x), \quad (3.11)$$

where the operator w meets the condition

$$\lim_{\substack{x \in Q, \\ x \rightarrow 0}} \frac{\|w(x)\|}{\|x\|} = 0. \quad (3.12)$$

If $Bx \neq x$ for $x \in W_0, x \neq 0$, there exists a positive real $c > 0$ such that for any $x \in W_0$ the inequality

$$\|x - Bx\| \geq c\|x\| \quad (3.13)$$

holds.

Choose a real $r > 0$ such that inequalities

$$\|Ax - Bx\| \leq \frac{c}{2}\|x\|, \quad \frac{cr}{(c + 2\|B\|)\rho_r} - 1 > 0 \quad (3.14)$$

hold for $x \in Q, \|x\| \leq r$, where $\rho_r = \sup_{x \in W_0, \|x\|=r} \rho(x, Q)$ and $\rho(x, Q)$ denotes the distance from the point x to the set Q .

8 On calculation of the relative index

Show that on the intersection of the sphere $S_r = \{x \in X : \|x\| = r\}$ of radius r with the set W_0 the vector field $\Phi' = I - B$ is linearly homotopic to the field $\Phi_\alpha x = x - AP_\alpha x$ where α meets the condition

$$0 < \alpha < \frac{cr}{(c + 2\|B\|)\rho_r} - 1, \quad (3.15)$$

and P_α is a projection (in general, nonlinear) on Q that has the following property:

$$\|x - P_\alpha x\| \leq (1 + \alpha)\rho(x, Q) \quad (x \in X) \quad (3.16)$$

(the existence of such projection follows from [4, Theorem 18.1]).

To prove this, consider the linear deformation

$$\Phi(\lambda, x) = x - (1 - \lambda)Bx - \lambda AP_\alpha x \quad (x \in S_r \cap W_0, \lambda \in [0, 1]) \quad (3.17)$$

that connects vector fields Φ' and Φ_α . From the convexity of W_0 and the invariance of the sets W_0 and $Q \subseteq W_0$ under operators B and A , respectively, it follows that the element $(1 - \lambda)Bx + \lambda AP_\alpha x$ is in W_0 for any $x \in S_r \cap W_0$ and $\lambda \in [0, 1]$.

The nondegeneracy of the deformation $\Phi(\lambda, x)$ for $x \in S_r \cap W_0, \lambda \in [0, 1]$ follows from the inequalities

$$\begin{aligned} & \|x - (1 - \lambda)Bx - \lambda AP_\alpha x\| \\ & \geq \|x - Bx\| - \lambda \|AP_\alpha x - Bx\| \\ & \geq \|x - Bx\| - \|AP_\alpha x - BP_\alpha x\| - \|Bx - BP_\alpha x\| \\ & \geq cr - \frac{c}{2}(r + (1 + \alpha)\rho_r) - \|B\|(1 + \alpha)\rho_r = \frac{cr}{2} - \left(\frac{c}{2} + \|B\|\right)(1 + \alpha)\rho_r > 0. \end{aligned} \quad (3.18)$$

Hence the vector fields Φ' and Φ_α are homotopic on $S_r \cap W_0$. Thus, by the first property of the relative rotation,

$$\text{ind}(0, I - B; W_0) = \text{ind}(0, I - AP_\alpha; W_0). \quad (3.19)$$

By definition, the relative index $\text{ind}(0, I - AP_\alpha; W_0)$ is equal to the relative rotation $\gamma(I - AP_\alpha, B_r \cap W_0; W_0)$ of the vector field $I - AP_\alpha$ on the boundary of an open set $B_r \cap W_0$, where $B_r = \{x \in X : \|x\| < r\}$. By the additivity property of rotation,

$$\gamma(I - AP_\alpha, B_r \cap W_0; W_0) = \gamma(I - AP_\alpha, B_r \cap Q; W_0) + \gamma(I - AP_\alpha, B_r \cap (W_0 \setminus Q); W_0). \quad (3.20)$$

From the fact that AP_α has no fixed points beyond Q , it follows that $\gamma(I - AP_\alpha, B_r \cap (W_0 \setminus Q); W_0) = 0$. On the other hand, the relative rotation $\gamma(I - AP_\alpha, B_r \cap Q; W_0)$ can be considered as the rotation $\gamma(I - A, B_r \cap Q; Q)$ of the vector field $I - A$ on the boundary of the open set $B_r \cap Q$ with respect to Q . By the definition of relative index, this rotation coincides with $\text{ind}(0, I - A; Q)$. This completes the proof. $\square$

As it appears, the analogous statement is true for the case of asymptotic index.

THEOREM 3.4. *Let A be a completely continuous operator, acting in a Banach space X , and let $Q \subseteq X$ be a closed convex unbounded set that is invariant under A . Let A have Fréchet derivative at infinity with respect to Q . Then the wedge*

$$W_\infty = \{h \in X : x + th \in Q \ (x \in Q, 0 \leq t < \infty)\} \quad (3.21)$$

is invariant under the mapping of the derivative $A'(\infty)$.

If $W_\infty \neq \{0\}$ and $A'(\infty)x \neq x$ for $x \in W_\infty$, $x \neq 0$, then the infinity singular point of the vector field $\Phi = I - A$ is isolated in Q and

$$\text{ind}(\infty, I - A; Q) = \text{ind}(0, I - A'(\infty); W_\infty). \quad (3.22)$$

Proof. From the differentiability of the operator A at infinity it follows that there exists linear operator $B = A'(\infty)$ such that

$$Ax = Bx + w(x), \quad (3.23)$$

where the operator w meets the condition

$$\lim_{\substack{x \in Q, \\ x \rightarrow \infty}} \frac{\|w(x)\|}{\|x\|} = 0. \quad (3.24)$$

If $Bx \neq x$ for $x \in W_\infty$, $x \neq 0$, then there exists a positive real $c > 0$ such that for all $x \in W_\infty$ the inequality

$$\|x - Bx\| \geq c\|x\| \quad (3.25)$$

holds.

Choose a real $R > 0$ such that inequalities

$$\|Ax - Bx\| \leq \frac{c}{2}\|x\|, \quad \frac{cR}{2(1+c+\|B\|)\rho_R} - 1 > 0 \quad (3.26)$$

hold for all $x \in Q$, $\|x\| \geq R$, where $\rho_R = \sup_{x \in Q, \|x\|=R} \rho(x, W_\infty)$ and $\rho(x, W_\infty)$ denotes the distance from the point x to the set W_∞ .

Show that on the intersection of the sphere $S_R = \{x \in X : \|x\| = R\}$ of radius R with the set Q the vector field $\Phi = I - A$ is linearly homotopic to the field $\Phi_\alpha x = x - BP_\alpha x$, where a real α meets the condition

$$0 < \alpha < \frac{cR}{2(1+c+\|B\|)\rho_R} - 1, \quad (3.27)$$

and P_α is a projection (in general, nonlinear) on W_∞ that has the following property:

$$\|x - P_\alpha x\| \leq (1 + \alpha)\rho(x, W_\infty) \quad (x \in X). \quad (3.28)$$

Consider the linear deformation

$$\Phi(\lambda, x) = x - (1 - \lambda)BP_\alpha x - \lambda Ax \quad (x \in S_R \cap Q, \lambda \in [0, 1]) \quad (3.29)$$

that connects vector fields Φ and Φ_α . Since the convexity of Q and the invariance of the sets $W_\infty \subseteq Q$ and Q under the operators B and A , respectively, it follows that the element $(1 - \lambda)BP_\alpha x + \lambda Ax$ is in Q for any $x \in S_R \cap Q$ and $\lambda \in [0, 1]$.

The nondegeneracy of $\Phi(\lambda, x)$ for $x \in S_R \cap Q$, $\lambda \in [0, 1]$ follows from the inequalities

$$\begin{aligned}
 & \|x - (1 - \lambda)BP_\alpha x - \lambda Ax\| \\
 & \geq \|x - BP_\alpha x\| - \lambda \|Ax - BP_\alpha x\| \\
 & \geq \|P_\alpha x - BP_\alpha x\| - \|x - P_\alpha x\| - \|Ax - Bx\| - \|Bx - BP_\alpha x\| \\
 & \geq c(R - (1 + \alpha)\rho_R) - (1 + \alpha)\rho_R - \frac{cR}{2} - \|B\|(1 + \alpha)\rho_R \\
 & = \frac{cR}{2} - (1 + c + \|B\|)(1 + \alpha)\rho_R > 0.
 \end{aligned} \tag{3.30}$$

Hence the vector fields Φ and Φ_α are homotopic on $S_R \cap Q$. Thus, by the first property of the relative rotation,

$$\text{ind}(\infty, I - A; Q) = \text{ind}(\infty, I - BP_\alpha; Q). \tag{3.31}$$

By definition, the relative index $\text{ind}(\infty, I - BP_\alpha; Q)$ is equal to the relative rotation $\gamma(I - BP_\alpha, B_R \cap Q; Q)$ of the vector field $I - BP_\alpha$ on the boundary of an open set $B_R \cap Q$, where $B_R = \{x \in X : \|x\| < R\}$. By the additivity property of rotation,

$$\gamma(I - BP_\alpha, B_R \cap Q; Q) = \gamma(I - BP_\alpha, B_R \cap W_\infty; Q) + \gamma(I - BP_\alpha, B_R \cap (Q \setminus W_\infty); Q). \tag{3.32}$$

From the fact that the operator BP_α has no fixed points beyond W_∞ , it follows that the relative rotation $\gamma(I - BP_\alpha, B_R \cap (Q \setminus W_\infty); Q)$ is equal to zero. Finally, the relative rotation $\gamma(I - BP_\alpha, B_R \cap W_\infty; Q)$ can be considered as the rotation $\gamma(I - B, B_R \cap W_\infty; W_\infty)$ of the vector field $I - B$ on the boundary of open set $B_R \cap W_\infty$ with respect to W_∞ . By the definition of relative index and since the operator B has no nonzero fixed points in W_∞ , the latter rotation coincides with $\text{ind}(0, I - B; W_\infty)$. This completes the proof. $\square$

References

- [1] E. N. Dancer, *On the indices of fixed points of mappings in cones and applications*, Journal of Mathematical Analysis and Applications **91** (1983), no. 1, 131–151.
- [2] ———, *Positivity of maps and applications*, Topological Nonlinear Analysis: Degree, Singularity, and Variations, Progr. Nonlinear Differential Equations Appl., vol. 15, Birkhäuser Boston, Massachusetts, 1995, pp. 303–340.
- [3] Y. Y. Isaenko, *The fixed point index of linear positive operators*, Trudy Aspirantov Voronezhskogo Universiteta (Proceedings of Postgraduate Students of Voronezh State University) (1969), 54–58 (Russian).
- [4] M. A. Krasnosel'skiĭ and P. P. Zabreiko, *Geometrical Methods of Nonlinear Analysis*, Nauka, Moscow, 1975, translated into English: Springer, Berlin, 1984.
- [5] E. M. Mukhamadiev and T. Sabirov, *On the computation of the relative index of the zero of a vector field*, Trudy Seminara po Funktsional'nomu Analizu (Proceedings of Seminar on Functional Analysis) **12** (1969), 116–121 (Russian).

- [6] Y. V. Pokornyi, *Relative indices of positive operators*, Trudy Matematicheskogo Fakul'teta Voronezhskogo Universiteta (Proceedings of Department of Mathematics of Voronezh State University) **4** (1970), 79–89 (Russian).
- [7] Y. V. Pokornyi and I. N. Astaf'eva, *On the index at zero and at infinity for positive nonlinear operators*, Uspehi Matematicheskikh Nauk **26** (1971), no. 3, 189 (Russian).
- [8] P. P. Zabrejko, *Rotation of vector fields: definition, basic properties, and calculation*, Topological Nonlinear Analysis, II: Degree, Singularity, and Variations (Frascati, 1995), Progr. Nonlinear Differential Equations Appl., vol. 27, Birkhäuser Boston, Massachusetts, 1997, pp. 445–601.

A. V. Guminskaya: Mechanics and Mathematics Faculty, Belarusian State University,
Independence Avenue 4, 220050 Minsk, Belarus
E-mail address: aguminskaya@yandex.ru

P. P. Zabreiko: Mechanics and Mathematics Faculty, Belarusian State University,
Independence Avenue 4, 220050 Minsk, Belarus
E-mail address: zabreiko@mail.ru