Research Article

q -Genocchi Numbers and Polynomials Associated with Fermionic p -Adic Invariant Integrals on $\mathbb{Z}_p$

Leechae Jang¹ and Taekyun Kim²

¹ Department of Mathematics and Computer Science, KonKuk University, Chungju 380-701, South Korea

² Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, South Korea

Correspondence should be addressed to Leechae Jang, leechae.jang@kku.ac.kr

Received 3 April 2008; Accepted 22 April 2008

Recommended by Ferhan Atici

The main purpose of this paper is to present a systemic study of some families of multiple Genocchi numbers and polynomials. In particular, by using the fermionic p -adic invariant integral on $\mathbb{Z}_p$, we construct p -adic Genocchi numbers and polynomials of higher order. Finally, we derive the following interesting formula: $G_{n+k,q}^{(k)}(x) = 2^k k! \binom{n+k}{k} \sum_{l=0}^{\infty} \sum_{d_0+d_1+\dots+d_k=k-1, d_i \in \mathbb{N}} (-1)^l (l+x)^n$, where $G_{n+k,q}^{(k)}(x)$ are the q -Genocchi polynomials of order k .

Copyright © 2008 L. Jang and T. Kim. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$, and $\mathbb{C}_p$ will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field and the p -adic completion of the algebraic closure of $\mathbb{Q}_p$. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p|_p = p^{v_p(p)} = 1/p$. When one talks of q -extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, then we assume that $|1 - q|_p < 1$, see [1–6].

In $\mathbb{C}$, the ordinary Euler polynomials are defined as

$$\frac{2}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (|t| < \pi). \quad (1.1)$$

In the case $x = 0$, $E_n(0) = E_n$ are called Euler numbers, see [1–13]. Let $\delta_{0,n}$ be the Kronecker symbol. From (1.1) we derive the following relation:

$$E_0 = 1, \quad (E + 1)^n + E_n = 2\delta_{0,n}, \quad n \in \mathbb{N}, \quad (1.2)$$

(cf. [7–13]). Here, we use the technique method notation by replacing E^n by E_n ($n \geq 0$), symbolically. The first few are $1, -1/2, 0, 1/4, \dots$, and $E_{2k} = 0$ for $k = 1, 2, \dots$. A sequence consisting of the Genocchi numbers G_n satisfies the following relations:

$$\frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad (|t| < \pi), \quad (1.3)$$

see [11, 12]. It satisfies $G_1 = 1$, $G_3 = G_5 = G_7 = \dots = G_{2k+1} = 0$, $k = 1, 2, 3, \dots$, and even coefficients are given by

$$G_{2n} = 2(1 - 2^{2n})B_{2n} = 2nE_{2n-1}(0), \quad (1.4)$$

where B_n is Bernoulli numbers. The first few Genocchi numbers for even integers are $-1, 1, -3, 17, -155, 2073, \dots$. The first few prime Genocchi numbers are -3 and 17 , which occur at $n = 6$ and 8 . There are no others with $n < 10^5$. We now define the Genocchi polynomials as follows:

$$\frac{2t}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}, \quad (|t| < \pi). \quad (1.5)$$

Thus, we note that

$$G_n(x) = \sum_{l=0}^n \binom{n}{l} G_l x^{n-l}. \quad (1.6)$$

In this paper, we use the following notations: $[x]_q = (1 - q^x)/(1 - q)$ and $[x]_{-q} = (1 + (-q)^x)/(1 + q)$. Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the p -adic q -deformed fermionic integral on $\mathbb{Z}_p$ is defined as

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \quad (1.7)$$

see [1–4]. The fermionic p -adic invariant integral on $\mathbb{Z}_p$ can be obtained as $q \rightarrow 1$. That is,

$$I_{-1}(f) = \lim_{q \rightarrow 1} I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x). \quad (1.8)$$

From (1.8), we easily derive the following integral equation related to fermionic invariant p -adic integral on $\mathbb{Z}_p$:

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0), \quad (1.9)$$

where $f_1(x) = f(x+1)$, see [5].

The purpose of this paper is to present a systemic study of some families of multiple Genocchi numbers and polynomials by using the fermionic multivariate p -adic invariant integral on $\mathbb{Z}_p$. In addition, we will investigate some interesting identities related to Genocchi numbers and polynomials.

2. Genocchi numbers associated with fermionic p -adic invariant integral on $\mathbb{Z}_p$

From (1.9) we can derive

$$\begin{aligned} t \int_{\mathbb{Z}_p} e^{xt} d\mu_{-1}(x) &= \frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \\ t \int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) &= \frac{2t}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}, \end{aligned} \quad (2.1)$$

where $G_n(x)$ are Genocchi polynomials. It is easy to check that

$$t \int_{\mathbb{Z}_p} e^{(2x+1)t} d\mu_{-1}(x) = \frac{2t}{e^t + e^{-t}} = t \operatorname{sech} t = \frac{1}{2} \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} 2^l G_l \right) \frac{t^n}{n!}. \quad (2.2)$$

By comparing the coefficient on both sides in (2.1), we easily see that

$$\int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(x) = \frac{G_{n+1}(x)}{n+1}. \quad (2.3)$$

Therefore, we obtain the following proposition.

Proposition 2.1. For $k \in \mathbb{Z}_+$,

- (i) $\int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(x) = G_{n+1}(x)/(n+1)$ (Witt's formula for Genocchi polynomials);
- (ii) $\int_{\mathbb{Z}_p} e^{(2x+1)t} d\mu_{-1}(x) = (1/(n+1))((1/2)\sum_{l=0}^{n+1} \binom{n+1}{l} 2^l G_l)$, where $\binom{n}{l} = (n(n-1)\cdots(n-l+1))/l!$.

Let $\mathfrak{O}_{\mathbb{C}_p} = \{x \in \mathbb{C}_p \mid |x|_p \leq 1\}$ be the integer ring of $\mathbb{C}_p$. We note that $i = (-1)^{1/2} \in \mathfrak{O}_{\mathbb{C}_p}$. By using Taylor expansion, we see that

$$e^{ix} = \sum_{n=0}^{\infty} i^n \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} + i \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}. \quad (2.4)$$

In the p -adic number field, $\sin x$ and $\cos x$ are defined as

$$\begin{aligned} \sin x &= \frac{x}{1!} - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \cdots, \\ \cos x &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \cdots. \end{aligned} \quad (2.5)$$

From (2.4) and (2.5), we derive

$$e^{ix} = \cos x + i \sin x. \quad (2.6)$$

This is equivalent to

$$\cos x = \frac{e^{ix} + e^{-ix}}{2}, \quad \sin x = \frac{e^{ix} - e^{-ix}}{2i}. \quad (2.7)$$

By (2.7), we easily see that

$$\sec t = \frac{2}{e^{it} + e^{-it}} = \int_{\mathbb{Z}_p} e^{(2x+1)it} d\mu_{-1}(x) = \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} (2x+1)^n d\mu_{-1}(x) \frac{i^n t^n}{n!}. \quad (2.8)$$

It is not difficult to show that $\int_{\mathbb{Z}_p} (2x+1)^{2n+1} d\mu_{-1}(x) = 0$ for $n \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. From (2.8), we note that

$$\sec t = \sum_{n=0}^{\infty} i^n \int_{\mathbb{Z}_p} (2x+1)^n d\mu_{-1}(x) \frac{t^n}{n!} = \sum_{n=0}^{\infty} (-1)^n \int_{\mathbb{Z}_p} (2x+1)^{2n} d\mu_{-1}(x) \frac{t^{2n}}{(2n)!}. \quad (2.9)$$

Thus, we have

$$t \sec t = \frac{1}{2} \sum_{n=0}^{\infty} (-1)^n \left(\sum_{l=0}^{2n+1} \binom{2n+1}{l} 2^l G_l \right) \frac{t^{2n+1}}{(2n+1)!}. \quad (2.10)$$

Now we consider the fermionic multivariate p -adic invariant integral on $\mathbb{Z}_p$ as follows:

$$t^k \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} e^{(x_1+x_2+\cdots+x_k)t} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) = 2^k \underbrace{\frac{t^k}{(e^t+1) \cdots (e^t+1)}}_{k\text{-times}} = \sum_{n=0}^{\infty} G_n^{(k)} \frac{t^n}{n!}, \quad (2.11)$$

where $G_n^{(k)}$ are the n th Genocchi number of order k . By comparing the coefficient on both sides in (2.11), we see that $G_0^{(k)} = G_1^{(k)} = \cdots = G_n^{(k)} = 0$, and

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} (x_1 + x_2 + \cdots + x_k)^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) (n+k)_k = G_{n+k}^{(k)}, \quad (2.12)$$

where $(n+k)_k$ is the Jordan factor which is defined by $(n+k)_k = (n+k) \cdots (n+1)$. Thus, we note that

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} (x_1 + x_2 + \cdots + x_k)^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) = \frac{G_{n+k}^{(k)}}{\binom{n+k}{n} n!}, \quad (2.13)$$

for $k \in \mathbb{N}$, $n \in \mathbb{Z}_+$.

Theorem 2.2. For $n \in \mathbb{Z}_+$, $k \in \mathbb{N}$,

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} (x_1 + x_2 + \cdots + x_k)^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) = \frac{G_{n+k}^{(k)}}{\binom{n+k}{n} n!}. \quad (2.14)$$

The multinomial coefficient is well known as

$$(x_1 + x_2 + \cdots + x_k)^n = \sum_{\substack{l_1 + \cdots + l_k = n \\ l_1, \dots, l_k \geq 0}} \binom{n}{l_1, \dots, l_k} x_1^{l_1} x_2^{l_2} \cdots x_k^{l_k}. \quad (2.15)$$

Therefore, we obtain the following corollary.

Corollary 2.3. For $n \in \mathbb{Z}_+$, $k \in \mathbb{N}$,

$$\sum_{\substack{l_1 + \cdots + l_k = n \\ l_1, \dots, l_k \geq 0}} \binom{n}{l_1, \dots, l_k} \left(\frac{G_{l_1+1}}{l_1+1} \right) \left(\frac{G_{l_2+1}}{l_2+1} \right) \cdots \left(\frac{G_{l_k+1}}{l_k+1} \right) = \frac{G_{n+k}^{(k)}}{\binom{n+k}{n} n!}. \quad (2.16)$$

For $q \in \mathbb{C}_p$ with $|1 - q| < 1$, it is not difficult to show that

$$t \int_{\mathbb{Z}_p} q^x e^{xt} d\mu_{-1}(x) = \frac{2t}{qe^t + 1}. \quad (2.17)$$

Now, we define the q -extension of the Genocchi numbers as follows:

$$\frac{2t}{qe^t + 1} = \sum_{n=0}^{\infty} G_{n,q} \frac{t^n}{n!}. \quad (2.18)$$

By (2.17) and (2.18), we easily see that

$$\frac{G_{n+1,q}}{n+1} = \int_{\mathbb{Z}_p} q^x x^n d\mu_{-1}(x). \quad (2.19)$$

With the same motivation to construct the Genocchi polynomials of higher order, we can consider the q -extension of higher-order Genocchi numbers as follows:

$$\begin{aligned} & t^k \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} q^{x_2 + 2x_3 + \cdots + (k-1)x_k} e^{(x+x_1+x_2+\cdots+x_k)t} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) \\ &= \frac{t^k 2^k}{\underbrace{(e^t + 1)(qe^t + 1) \cdots (q^{k-1}e^t + 1)}_{k\text{-times}}} e^{xt} = \sum_{n=0}^{\infty} G_{n,q}^{(k)} \frac{t^n}{n!}, \end{aligned} \quad (2.20)$$

where $G_{n,q}^{(k)}$ are the q -Genocchi polynomials of order k . The basic q -natural numbers are defined as

$$[n]_q = \frac{1 - q^n}{1 - q} = 1 + q + q^2 + \cdots + q^{n-1}, \quad n \in \mathbb{N}. \quad (2.21)$$

The q -factorial of n is defined as

$$[n]_q! = [n]_q [n-1]_q \cdots [2]_q [1]_q = (1 + q + \cdots + q^{n-1}) \cdots (1 + q) \cdot 1. \quad (2.22)$$

The q -binomial coefficient is also defined as

$$\binom{n}{k}_q = \frac{[n]_q!}{[n-k]_q![k]_q!} = \frac{[n]_q[n-1]_q \cdots [n-k+1]_q}{[k]_q!}. \quad (2.23)$$

Note that $\lim_{q \rightarrow 1} \binom{n}{k}_q = \binom{n}{k} = (n(n-1) \cdots (n-k+1))/k!$. The q -binomial coefficient satisfies the following recursion formula:

$$\binom{n+1}{k}_q = \binom{n}{k-1}_q + q^k \binom{n}{k}_q = q^{n-k} \binom{n}{k-1}_q + \binom{n}{k}_q. \quad (2.24)$$

From this recursion formula, we can derive

$$\binom{n}{k}_q = \sum_{\substack{d_0+d_1+\cdots+d_k=k-1 \\ d_i \in \mathbb{N}}} q^{0 \cdot d_0 + 1 \cdot d_1 + \cdots + k \cdot d_k}. \quad (2.25)$$

The q -binomial expansion is given by

$$\begin{aligned} \prod_{i=1}^n (a + bq^{i-1}) &= \sum_{k=0}^n \binom{n}{k}_q q^{\binom{k}{2}} a^{n-k} b^k, \\ \prod_{i=1}^n (1 - bq^{i-1})^{-1} &= \sum_{k=0}^{\infty} \binom{n+k-1}{k}_q b^k. \end{aligned} \quad (2.26)$$

By (2.20) and (2.26), we see that

$$\begin{aligned} & t^k \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} q^{x_2+2x_3+\cdots+(k-1)x_k} e^{(x+x_1+x_2+\cdots+x_k)t} d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) \\ &= t^k 2^k \prod_{i=1}^k (1 - (-q^{i-1})e^t)^{-1} e^{xt} \\ &= t^k 2^k \sum_{l=0}^{\infty} \binom{k+l-1}{l}_q (-1)^l e^{(l+x)t} \\ &= t^k \sum_{n=0}^{\infty} \left(2^k \sum_{l=0}^{\infty} \binom{k+l-1}{l}_q (-1)^l (l+x)^n \right) \frac{t^n}{n!}. \end{aligned} \quad (2.27)$$

Therefore, we obtain the following theorem.

Theorem 2.4. For $n \in \mathbb{Z}_+$, $k \in \mathbb{N}$, we have

$$\begin{aligned} & \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} q^{x_2+2x_3+\cdots+(k-1)x_k} (x_1 + x_2 + \cdots + x_k)^n d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) \\ &= 2^k \sum_{l=0}^{\infty} \binom{k+l-1}{l}_q (-1)^l (l+x)^n. \end{aligned} \quad (2.28)$$

By (2.20), it is not difficult to show that

$$\begin{aligned}
 G_{n+k,q}^{(k)}(x) &= k! \binom{n+k}{k} \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} q^{x_2+2x_3+\cdots+(k-1)x_k} \times (x + x_1 + x_2 + \cdots + x_k)^k \\
 &\quad \times d\mu_{-1}(x_1) \cdots d\mu_{-1}(x_k) (n+k)_k, \\
 G_{0,q}^{(k)} &= G_{1,q}^{(k)} = \cdots = G_{n,q}^{(k)} = 0,
 \end{aligned} \tag{2.29}$$

where $n = 0, 1, 2, \dots$. Therefore, we obtain the following corollary.

Corollary 2.5. For $n \in \mathbb{Z}_+$, $k \in \mathbb{N}$,

$$\frac{G_{n+k,q}^{(k)}(x)}{k! \binom{n+k}{k}} = 2^k \sum_{l=0}^{\infty} \binom{k+l-1}{l}_q (-1)^l (l+x)^n. \tag{2.30}$$

Corollary 2.6. For $n \in \mathbb{Z}_+$, $k \in \mathbb{N}$,

$$G_{n+k,q}^{(k)}(x) = 2^k k! \binom{n+k}{k} \sum_{l=0}^{\infty} \sum_{\substack{d_0+d_1+\cdots+d_k=k-1 \\ d_i \in \mathbb{N}}} q^{0 \cdot d_0 + 1 \cdot d_1 + \cdots + k \cdot d_k} (-1)^l (l+x)^n. \tag{2.31}$$

Acknowledgment

The present research has been conducted by the research grant of Kwangwoon University in 2008.

References

- [1] T. Kim, “ q -Euler numbers and polynomials associated with p -adic q -integrals,” *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 15–27, 2007.
- [2] T. Kim, “ q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients,” *Russian Journal Mathematical Physics*, vol. 15, no. 1, pp. 51–57, 2008.
- [3] T. Kim, “ q -Bernoulli numbers associated with q -stirling numbers,” *Advances in Difference Equations*, vol. 2008, Article ID 743295, 10 pages, 2008.
- [4] T. Kim, “The modified q -Euler numbers and polynomials,” *Advanced Studies in Contemporary Mathematics*, vol. 16, pp. 161–170, 2008.
- [5] T. Kim, “Euler numbers and polynomials associated with zeta functions,” *Abstract and Applied Analysis*, vol. 2008, Article ID 581582, 13 pages, 2008.
- [6] T. Ernst, “Examples of a q -umbral calculus,” *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 1, pp. 1–22, 2008.
- [7] H. Ozden, I. N. Cangul, and Y. Simsek, “Multivariate interpolation functions of higher order q -Euler numbers and their applications,” *Abstract and Applied Analysis*, vol. 2008, Article ID 390857, 16 pages, 2008.
- [8] H. Ozden, Y. Simsek, S.-H. Rim, and I. N. Cangul, “A note on p -adic q -Euler measure,” *Advanced Studies in Contemporary Mathematics*, vol. 14, no. 2, pp. 233–239, 2007.
- [9] H. Ozden, I. N. Cangul, and Y. Simsek, “Remarks on sum of products of (h, q) -twisted Euler polynomials and numbers,” *Journal of Inequalities and Applications*, vol. 2008, Article ID 816129, 8 pages, 2008.

- [10] Y. Simsek, V. Kurt, and D. Kim, "New approach to the complete sum of products of the twisted (h, q) -Bernoulli numbers and polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 44–56, 2007.
- [11] M. Cenkci, M. Can, and V. Kurt, " q -extensions of Genocchi numbers," *Journal of the Korean Mathematical Society*, vol. 43, no. 1, pp. 183–198, 2006.
- [12] M. Cenkci, Y. Simsek, and V. Kurt, "Further remarks on multiple p -adic q - L -function of two variables," *Advanced Studies in Contemporary Mathematics*, vol. 14, no. 1, pp. 49–68, 2007.
- [13] T. Machide, "Sums of products of Kronecker's double series," *Journal of Number Theory*, vol. 128, no. 4, pp. 820–834, 2008.