Research Article

Spectral Singularities of Sturm-Liouville Problems with Eigenvalue-Dependent Boundary Conditions

Elgiz Bairamov and Nihal Yokus

Department of Mathematics, Ankara University, 06100 Tandoğan, Ankara, Turkey

Correspondence should be addressed to Elgiz Bairamov, bairamov@science.ankara.edu.tr

Received 25 June 2009; Accepted 20 August 2009

Recommended by Ağacık Zafer

Let L denote the operator generated in $L_2(\mathbb{R}_+)$ by Sturm-Liouville equation $-y'' + q(x)y = \lambda^2 y$, $x \in \mathbb{R}_+ = [0, \infty)$, $y'(0)/y(0) = \alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2$, where q is a complex-valued function and $\alpha_i \in \mathbb{C}$, $i = 0, 1, 2$ with $\alpha_2 \neq 0$. In this article, we investigate the eigenvalues and the spectral singularities of L and obtain analogs of Naimark and Pavlov conditions for L .

Copyright © 2009 E. Bairamov and N. Yokus. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let L_0 denote Sturm-Liouville operator generated in $L_2(\mathbb{R}_+)$ by the differential expression

$$l_0(y) := -y'' + q(x)y, \quad x \in \mathbb{R}_+, \quad (1.1)$$

and the boundary condition $y(0) = 0$, where $q : \mathbb{R}_+ \rightarrow \mathbb{C}$. Since q is a complex-valued function, the operator L_0 is a non-selfadjoint. The spectral analysis of L_0 has been investigated by Naimark [1]. He proved that some of the poles of the kernel of resolvent of L_0 are not the eigenvalues of the operator. He also showed that those poles (which are called spectral singularities by Schwartz [2]) are on the continuous spectrum. Moreover, he has shown the spectral singularities play an important role in the spectral analysis of L_0 , and if

$$\int_0^\infty e^{\varepsilon x} |q(x)| dx < \infty, \quad \varepsilon > 0, \quad (\text{N})$$

then the eigenvalues and the spectral singularities are of a finite number and each of them is of a finite multiplicity.

One very important step in the spectral analysis of L_0 was taken by Pavlov [3]. He studied the dependence of the structure of the eigenvalues and the spectral singularities of L_0 on the behavior of potential function at infinity. He also proved that if

$$\sup_{x \in \mathbb{R}_+} \left[e^{\varepsilon \sqrt{x}} |q(x)| \right] < \infty, \quad \varepsilon > 0, \quad (\text{P})$$

then the eigenvalues and the spectral singularities are of a finite number and each of them is of a finite multiplicity.

Conditions (N) and (P) are called Naimark and Pavlov conditions for L_0 , respectively.

Lyance showed that the spectral singularities play an important role in the spectral analysis of L_0 [4, 5]. He also investigated the effect of the spectral singularities in the spectral expansion.

The spectral singularities of non-selfadjoint operator generated in $L_2(\mathbb{R}_+)$ by (1.1) and the boundary condition

$$\int_0^\infty K(x)y(x)dx + \alpha y'(0) - \beta y(0) = 0 \quad (1.2)$$

was investigated in detail by Krall [6, 7].

Some problems of spectral theory of differential operator and some other types of operators with spectral singularities were studied by some authors [8–14]. Note that in all papers the boundary conditions are not depending on the spectral parameter.

In a recent series of papers, Binding et al. and Browne [15–18] have studied the spectral theory of regular Sturm-Liouville operators with boundary conditions depending on the spectral parameter.

Let L denote the operator generated in $L_2(\mathbb{R}_+)$ by

$$-y'' + q(x)y = \lambda^2 y, \quad x \in \mathbb{R}_+, \quad (1.3)$$

$$\frac{y'(0)}{y(0)} = \alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2, \quad (1.4)$$

where q is a complex-valued function, $\alpha_i \in \mathbb{C}, i = 0, 1, 2$, with $\alpha_2 \neq 0$. In this paper, we investigate the eigenvalues and the spectral singularities of L . In particular, we show that the analogs of Naimark and Pavlov conditions for L are

$$\begin{aligned} q \in AC(\mathbb{R}_+), \quad \lim_{x \rightarrow \infty} q(x) = 0, \quad \int_0^\infty e^{\varepsilon x} |q'(x)| dx < \infty, \quad \varepsilon > 0, \\ q \in AC(\mathbb{R}_+), \quad \lim_{x \rightarrow \infty} q(x) = 0, \quad \sup_{x \in \mathbb{R}_+} \left[e^{\varepsilon \sqrt{x}} |q'(x)| \right] < \infty, \quad \varepsilon > 0, \end{aligned} \quad (1.5)$$

respectively, where $AC(\mathbb{R}_+)$ denotes the class of complex-valued absolutely continuous functions on $\mathbb{R}_+$.

2. Jost Functions of (1.3)-(1.4)

Under the condition

$$\int_0^\infty x|q(x)|dx < \infty, \quad (2.1)$$

(1.3) has a solution $e(x, \lambda)$ satisfying

$$\lim_{x \rightarrow \infty} e(x, \lambda) e^{-i\lambda x} = 1, \quad \lambda \in \overline{\mathbb{C}}_+, \quad (2.2)$$

where $\overline{\mathbb{C}}_+ = \{\lambda : \lambda \in \mathbb{C}, \operatorname{Im} \lambda \geq 0\}$. The solution $e(x, \lambda)$ is called Jost solution of (1.3). Note that Jost solution has a representation [19]

$$e(x, \lambda) = e^{i\lambda x} + \int_x^\infty K(x, t) e^{i\lambda t} dt, \quad \lambda \in \overline{\mathbb{C}}_+, \quad (2.3)$$

where $K(x, t)$ is the solution of the integral equation

$$K(x, t) = \frac{1}{2} \int_{(x+t)/2}^\infty q(s) ds + \frac{1}{2} \int_x^{(x+t)/2} \int_{t+x-s}^{t+s-x} q(s) K(s, u) du ds + \frac{1}{2} \int_{(x+t)/2}^\infty \int_s^{t+s-x} q(s) K(s, u) du ds, \quad (2.4)$$

and $K(x, t)$ are continuously differentiable with respect to their arguments. We also have

$$|K(x, t)| \leq cw \left(\frac{x+t}{2} \right), \quad (2.5)$$

$$|K_x(x, t)|, |K_t(x, t)| \leq \frac{1}{4} \left| q \left(\frac{x+t}{2} \right) \right| + cw \left(\frac{x+t}{2} \right),$$

where $w(x) = \int_x^\infty |q(s)| ds$ and $c > 0$ is a constant.

Let

$$E^+(\lambda) := e'(0, \lambda) - (\alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2) e(0, \lambda), \quad \lambda \in \overline{\mathbb{C}}_+, \quad (2.6)$$

$$E^-(\lambda) := e'(0, -\lambda) - (\alpha_0 + \alpha_1 \lambda + \alpha_2 \lambda^2) e(0, -\lambda), \quad \lambda \in \overline{\mathbb{C}}_-,$$

where $\overline{\mathbb{C}}_- = \{\lambda : \lambda \in \mathbb{C}, \operatorname{Im} \lambda \leq 0\}$. Therefore, E^+ and E^- are analytic in $\mathbb{C}_+ = \{\lambda : \lambda \in \mathbb{C}, \operatorname{Im} \lambda > 0\}$ and $\mathbb{C}_- = \{\lambda : \lambda \in \mathbb{C}, \operatorname{Im} \lambda < 0\}$, respectively, and continuous up to real axis. The functions E^+ and E^- are called Jost functions of L .

Let us denote the eigenvalues and the spectral singularities of L by $\sigma_d(L)$ and $\sigma_{ss}(L)$, respectively. It is evident that

$$\sigma_d(L) = \{\lambda : \lambda \in \mathbb{C}_+, E^+(\lambda) = 0\} \cup \{\lambda : \lambda \in \mathbb{C}_-, E^-(\lambda) = 0\}, \quad (2.7)$$

$$\begin{aligned} \sigma_{ss}(L) &= \{\lambda : \lambda \in \mathbb{R}^*, E^+(\lambda) = 0\} \cup \{\lambda : \lambda \in \mathbb{R}^*, E^-(\lambda) = 0\}, \\ &\{\lambda : \lambda \in \mathbb{R}^*, E^+(\lambda) = 0\} \cap \{\lambda : \lambda \in \mathbb{R}^*, E^-(\lambda) = 0\} = \emptyset, \end{aligned} \quad (2.8)$$

where $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$.

Definition 2.1. The multiplicity of a zero E^+ (or E^-) in $\overline{\mathbb{C}}_+$ (or $\overline{\mathbb{C}}_-$) is defined as the multiplicity of the corresponding eigenvalue and spectral singularity of L .

In order to investigate the quantitative properties of the eigenvalues and the spectral singularities of L , we need to discuss the quantitative properties of the zeros of E^+ and E^- in $\overline{\mathbb{C}}_+$ and $\overline{\mathbb{C}}_-$, respectively.

Define

$$M_1^\pm = \{\lambda : \lambda \in \mathbb{C}_\pm, E^\pm(\lambda) = 0\}, \quad M_2^\pm = \{\lambda : \lambda \in \mathbb{R}^*, E^\pm(\lambda) = 0\}, \quad (2.9)$$

then by (2.7), we have

$$\sigma_d(L) = M_1^+ \cup M_1^-, \quad \sigma_{ss}(L) = M_2^+ \cup M_2^-. \quad (2.10)$$

Now, let us assume that

$$q \in AC(\mathbb{R}_+), \quad \lim_{x \rightarrow \infty} q(x) = 0, \quad \int_0^\infty x^3 |q'(x)| dx < \infty. \quad (2.11)$$

Theorem 2.2. Under condition (2.11), the functions E^+ and E^- have the representations

$$E^+(\lambda) = -\alpha_2 \lambda^2 + \beta^+ \lambda + \delta^+ + \int_0^\infty f^+(t) e^{i\lambda t} dt, \quad \lambda \in \overline{\mathbb{C}}_+, \quad (2.12)$$

$$E^-(\lambda) = -\alpha_2 \lambda^2 + \beta^- \lambda + \delta^- + \int_0^\infty f^-(t) e^{-i\lambda t} dt, \quad \lambda \in \overline{\mathbb{C}}_-, \quad (2.13)$$

where $\beta^\pm, \delta^\pm \in \mathbb{C}$, and $f^\pm \in L_1(\mathbb{R}_+)$.

Proof. Using (2.3), (2.4), and (2.6), we get (2.12), where

$$\begin{aligned} \beta^+ &= i - \alpha_1 - i\alpha_2 K(0, 0), \\ \delta^+ &= -K(0, 0) - \alpha_0 - i\alpha_1 K(0, 0) + \alpha_2 K_t(0, 0), \\ f^+(t) &= K_x(0, t) - \alpha_0 K(0, t) - i\alpha_1 K_t(0, t) + \alpha_2 K_{tt}(0, t). \end{aligned} \quad (2.14)$$

From (2.4), we see that

$$|K_{\#}(0, t)| \leq c \left[t \left| q\left(\frac{t}{2}\right) \right| + \left| q'\left(\frac{t}{2}\right) \right| + tw\left(\frac{t}{2}\right) + w_1\left(\frac{t}{2}\right) \right] \quad (2.15)$$

holds, where $w_1(t) = \int_t^\infty w(s)ds$ and $c > 0$ is a constant. It follows from (2.5), (2.14), and (2.15) that $f^+ \in L_1(\mathbb{R}_+)$. In a similar way, we obtain (2.13). $\square$

Theorem 2.3. *Under condition (2.11), we have the following.*

- (i) *The set of $\sigma_d(L)$ is bounded and has at most a countable number of elements, and its limit points can lie only in a bounded subinterval of the real axis.*
- (ii) *The set of $\sigma_{ss}(L)$ is bounded and its linear Lebesgue measure is zero.*

Proof. From (2.14) and (2.15), we see that

$$\begin{aligned} E^+(\lambda) &= -\alpha_2 \lambda^2 + \beta^+ \lambda + \delta^+ + o(1), \quad \lambda \in \overline{\mathbb{C}}_+, |\lambda| \rightarrow \infty, \\ E^-(\lambda) &= -\alpha_2 \lambda^2 + \beta^- \lambda + \delta^- + o(1), \quad \lambda \in \overline{\mathbb{C}}_-, |\lambda| \rightarrow \infty. \end{aligned} \quad (2.16)$$

Using (2.10), (2.16), and the uniqueness theorem of analytic functions [20], we get (i) and (ii). $\square$

3. Naïmark and Pavlov Conditions for L

We will denote the set of all limit points of M_1^+ and M_1^- by M_3^+ and M_3^- , respectively, and the set of all zeros of E^+ and E^- with infinity multiplicity in $\overline{\mathbb{C}}_+$ and $\overline{\mathbb{C}}_-$, by M_4^+ and M_4^- , respectively. It is obvious that

$$M_3^\pm \subset M_2^\pm, \quad M_4^\pm \subset M_2^\pm, \quad M_3^\pm \subset M_4^\pm, \quad (3.1)$$

and the linear Lebesgue measures of $M_3^\pm$ and $M_4^\pm$ are zero.

Theorem 3.1. *If*

$$q \in AC(\mathbb{R}_+), \quad \lim_{x \rightarrow \infty} q(x) = 0, \quad \int_0^\infty e^{\varepsilon x} |q'(x)| dx < \infty, \quad \varepsilon > 0, \quad (3.2)$$

then the operator L has a finite number of eigenvalues and spectral singularities, and each of them is of a finite multiplicity.

Proof. From (2.5), (2.14), (2.15), and (3.2), we find that

$$|f^+(t)| \leq ce^{-(\varepsilon/2)t}, \quad (3.3)$$

where $c > 0$ is a constant. By (2.12) and (3.3), we observe that the function E^+ has an analytic continuation to the half-plane $\text{Im } \lambda > -\varepsilon/4$. So we get that $M_4^+ = \emptyset$. It follows from (3.1)

that $M_3^+ = \emptyset$. Therefore the sets M_1^+ and M_2^+ have a finite number of elements with a finite multiplicity. We obtain similar results for the sets M_1^- and M_2^- . By (2.10) we have the proof of the theorem. $\square$

Now let us assume that

$$q \in AC(\mathbb{R}_+), \quad \lim_{x \rightarrow \infty} q(x) = 0, \quad \sup_{x \in \mathbb{R}_+} \left[e^{\varepsilon \sqrt{x}} |q'(x)| \right] < \infty, \quad \varepsilon > 0. \quad (3.4)$$

Hence, we have the following lemma.

Lemma 3.2. *It holds that $M_4^+ = M_4^- = \emptyset$.*

Proof. From (2.12) and (3.4), we find that the function E^+ is analytic in $\mathbb{C}_+$, and all of its derivatives are continuous in $\overline{\mathbb{C}}_+$. For a sufficiently large $T > 0$, we have

$$\left| \frac{d^k}{d\lambda^k} E^+(\lambda) \right| \leq A_k, \quad \lambda \in \overline{\mathbb{C}}_+, \quad |\lambda| \leq T, \quad k = 0, 1, 2, \dots, \quad (3.5)$$

where

$$A_k = 2^k c \int_0^\infty t^k e^{-(\varepsilon/2)\sqrt{t}} dt, \quad k = 0, 1, 2, \dots, \quad (3.6)$$

and $c > 0$ is a constant. Since the function E^+ is not equal to zero identically, then by Pavlov's theorem, M_4^+ satisfies

$$\int_0^h \ln A(s) d\mu(M_4^+, s) > -\infty, \quad (3.7)$$

where $A(s) = \inf_k (A_k s^k / k!)$, $\mu(M_4^+, s)$ is the linear Lebesgue measure of s -neighborhood of M_4^+ , [3]. Now, we obtain the following estimates for A_k :

$$A_k \leq B b^k k^k k!, \quad (3.8)$$

where B and b are constants depending on c and ε . From (3.8), we get that

$$A(s) \leq B \inf_k \left(b^k s^k k^k \right) \leq B \exp \left(-b^{-1} e^{-1} s^{-1} \right). \quad (3.9)$$

Now, (3.7) yields that

$$\int_0^h \frac{1}{s} d\mu(M_4^+, s) < \infty. \quad (3.10)$$

However, (3.10) holds for an arbitrary s , if and only if $\mu(M_4^+, s) = 0$ or $M_4^+ = \emptyset$. In a similar way we can prove that $M_4^- = \emptyset$. $\square$

Theorem 3.3. *Under condition (3.4), the operator L has a finite number of eigenvalues and spectral singularities, and each of them is of a finite multiplicity.*

Proof. To be able to prove the theorem, we have to show that the functions E^+ and E^- have a finite number of zeros with finite multiplicities in $\overline{\mathbb{C}}_+$ and $\overline{\mathbb{C}}_-$, respectively. We give the proof for E^+ .

From Lemma 3.2 and (3.1), we find that $M_3^+ = \emptyset$. So the bounded sets M_1^+ and M_2^+ have no limit points, that is, the function E^+ has only a finite number of zeros in $\overline{\mathbb{C}}_+$. Since $M_4^+ = \emptyset$, these zeros are of finite multiplicity. $\square$

Acknowledgment

This work was supported by TUBITAK.

References

- [1] M. A. Naïmark, "Investigation of the spectrum and the expansion in eigenfunctions of a non-selfadjoint differential operator of the second order on a semi-axis," *American Mathematical Society Translations Series 2*, vol. 16, pp. 103–193, 1960.
- [2] J. Schwartz, "Some non-selfadjoint operators," *Communications on Pure and Applied Mathematics*, vol. 13, pp. 609–639, 1960.
- [3] B. S. Pavlov, "The non-selfadjoint Schrödinger operator," *Topics in Mathematical Physics*, vol. 1, pp. 87–110, 1967.
- [4] V. E. Lyance, "A differential operator with spectral singularities. I," *American Mathematical Society Translations Series 2*, vol. 60, pp. 185–225, 1967.
- [5] V. E. Lyance, "A differential operator with spectral singularities. II," *American Mathematical Society Translations Series 2*, vol. 60, pp. 227–283, 1967.
- [6] A. M. Krall, "A nonhomogeneous eigenfunction expansion," *Transactions of the American Mathematical Society*, vol. 117, pp. 352–361, 1965.
- [7] A. M. Krall, "The adjoint of a differential operator with integral boundary conditions," *Proceedings of the American Mathematical Society*, vol. 16, pp. 738–742, 1965.
- [8] M. Adivar and E. Bairamov, "Spectral properties of non-selfadjoint difference operators," *Journal of Mathematical Analysis and Applications*, vol. 261, no. 2, pp. 461–478, 2001.
- [9] M. Adivar and E. Bairamov, "Difference equations of second order with spectral singularities," *Journal of Mathematical Analysis and Applications*, vol. 277, no. 2, pp. 714–721, 2003.
- [10] E. Bairamov, Ö. Çakar, and A. M. Krall, "An eigenfunction expansion for a quadratic pencil of a Schrödinger operator with spectral singularities," *Journal of Differential Equations*, vol. 151, no. 2, pp. 268–289, 1999.
- [11] E. Bairamov, Ö. Çakar, and A. M. Krall, "Non-selfadjoint difference operators and Jacobi matrices with spectral singularities," *Mathematische Nachrichten*, vol. 229, pp. 5–14, 2001.
- [12] E. Bairamov and A. O. Çelebi, "Spectrum and spectral expansion for the non-selfadjoint discrete Dirac operators," *The Quarterly Journal of Mathematics. Oxford. Second Series*, vol. 50, no. 200, pp. 371–384, 1999.
- [13] A. M. Krall, E. Bairamov, and Ö. Çakar, "Spectrum and spectral singularities of a quadratic pencil of a Schrödinger operator with a general boundary condition," *Journal of Differential Equations*, vol. 151, no. 2, pp. 252–267, 1999.
- [14] A. M. Krall, E. Bairamov, and Ö. Çakar, "Spectral analysis of non-selfadjoint discrete Schrödinger operators with spectral singularities," *Mathematische Nachrichten*, vol. 231, pp. 89–104, 2001.

- [15] P. A. Binding, P. J. Browne, and B. A. Watson, "Transformations between Sturm-Liouville problems with eigenvalue dependent and independent boundary conditions," *The Bulletin of the London Mathematical Society*, vol. 33, no. 6, pp. 749–757, 2001.
- [16] P. A. Binding, P. J. Browne, and B. A. Watson, "Sturm-Liouville problems with boundary conditions rationally dependent on the eigenparameter. I," *Proceedings of the Edinburgh Mathematical Society*, vol. 33, pp. 749–757, 2001.
- [17] P. A. Binding, P. J. Browne, and B. A. Watson, "Sturm-Liouville problems with boundary conditions rationally dependent on the eigenparameter. II," *Journal of Computational and Applied Mathematics*, vol. 148, no. 1, pp. 147–168, 2002.
- [18] W. J. Code and P. J. Browne, "Sturm-Liouville problems with boundary conditions depending quadratically on the eigenparameter," *Journal of Mathematical Analysis and Applications*, vol. 309, no. 2, pp. 729–742, 2005.
- [19] V. A. Marchenko, *Sturm-Liouville Operators and Applications*, vol. 22 of *Operator Theory: Advances and Applications*, Birkhäuser, Basel, Switzerland, 1986.
- [20] E. P. Dolzhenko, "Boundary value uniqueness theorems for analytic functions," *Mathematical Notes*, vol. 26, pp. 437–442, 1979.