Research Article

On an Integral-Type Operator Acting between Bloch-Type Spaces on the Unit Ball

Stevo Stević¹ and Sei-Ichiro Ueki²

¹ Mathematical Institute of the Serbian Academy of Sciences, Knez Mihailova 36/III, 11000 Beograd, Serbia

² Faculty of Engineering, Ibaraki University, Hitachi 316-8511, Japan

Correspondence should be addressed to Stevo Stević, sstevic@ptt.rs

Received 30 January 2010; Accepted 18 March 2010

Academic Editor: Narcisa C. Apreutesei

Copyright © 2010 S. Stević and S.-I. Ueki. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $\mathbb{B}$ denote the open unit ball of $\mathbb{C}^n$. For a holomorphic self-map φ of $\mathbb{B}$ and a holomorphic function g in $\mathbb{B}$ with $g(0) = 0$, we define the following integral-type operator: $I_{\varphi}^s f(z) = \int_0^1 \Re f(\varphi(tz)) g(tz) (dt/t)$, $z \in \mathbb{B}$. Here $\Re f$ denotes the radial derivative of a holomorphic function f in $\mathbb{B}$. We study the boundedness and compactness of the operator between Bloch-type spaces $\mathcal{B}_{\omega}$ and $\mathcal{B}_{\mu}$, where ω is a normal weight function and μ is a weight function. Also we consider the operator between the little Bloch-type spaces $\mathcal{B}_{\omega,0}$ and $\mathcal{B}_{\mu,0}$.

1. Introduction

Let $\mathbb{B}$ denote the open unit ball of the n -dimensional complex vector space $\mathbb{C}^n$ and $H(\mathbb{B})$ the space of all holomorphic functions on $\mathbb{B}$. For $f \in H(\mathbb{B})$ with the Taylor expansion $f(z) = \sum_{|\gamma| \geq 0} a_{\gamma} z^{\gamma}$, let

$$\Re f(z) = \sum_{|\gamma| \geq 0} |\gamma| a_{\gamma} z^{\gamma} \quad (1.1)$$

be the radial derivative of f , where $\gamma = (\gamma_1, \dots, \gamma_n)$ is a multi-index, $|\gamma| = \gamma_1 + \dots + \gamma_n$, and $z^{\gamma} = z_1^{\gamma_1} \dots z_n^{\gamma_n}$. It is well known that

$$\Re f(z) = \sum_{j=1}^n z_j \frac{\partial f}{\partial z_j}(z) = \langle \nabla f(z), \bar{z} \rangle, \quad (1.2)$$

where ∇ is the usual gradient on $\mathbb{C}^n$.

Let φ be a holomorphic self-map of $\mathbb{B}$ and $g \in H(\mathbb{B})$ with $g(0) = 0$. Then φ and g define an operator I_φ^g on $H(\mathbb{B})$ as follows:

$$I_\varphi^g f(z) = \int_0^1 \Re f(\varphi(tz)) g(tz) \frac{dt}{t}, \quad f \in H(\mathbb{B}), z \in \mathbb{B}. \quad (1.3)$$

The following important formula involving $\Re$ and $I_\varphi^g f$ was proved, for example, in [1]

$$\Re \left[I_\varphi^g f \right] (z) = \Re f(\varphi(z)) g(z), \quad z \in \mathbb{B}. \quad (1.4)$$

Motivated by papers [2, 3], operators I_φ^g were introduced by the first author of the present paper and Zhu in [1, 4–6], where its boundedness and compactness from the α -Bloch space, the Zygmund space, the mixed-norm space, and the generalized weighted Bergman space into the Bloch-type space on the unit ball are studied. In our previous work [7], we studied the boundedness and compactness of I_φ^g acting between weighted-type spaces. For related operators on $\mathbb{C}^n$ see, for example, [8–21] and the references therein.

Let ω be a strictly positive continuous function on $\mathbb{B}$ (*weight*). If $\omega(z) = \omega(|z|)$ for every $z \in \mathbb{B}$, we call it *radial weight*. A weight ω is called *normal* ([9, 22]) if it is radial and there are a and b , $0 < a < b < \infty$ such that $\omega(r)/(1-r)^a$ is decreasing on $[0, 1)$, $\omega(r)/(1-r)^b$ is increasing on $[0, 1)$,

$$\lim_{r \rightarrow 1} \frac{\omega(r)}{(1-r)^a} = 0, \quad \lim_{r \rightarrow 1} \frac{\omega(r)}{(1-r)^b} = \infty. \quad (1.5)$$

A radial weight ω is called *typical* if it is nonincreasing with respect to $|z|$ and $\omega(z) \rightarrow 0$ as $|z| \rightarrow 1^-$. If ω is normal, then by the monotonicity of $\omega(r)/(1-r)^a$, for $0 \leq r_1 < r < 1$, we have that

$$\omega(r) = (1-r)^a \frac{\omega(r)}{(1-r)^a} < (1-r)^a \frac{\omega(r_1)}{(1-r_1)^a} < \omega(r_1), \quad (1.6)$$

that is, ω is decreasing on $[0, 1)$. On the other hand, from the first equality in (1.5), we have that for any $\varepsilon > 0$, there is a $\delta > 0$ such that

$$0 < \omega(r) < \varepsilon(1-r)^a, \quad (\delta < r < 1), \quad (1.7)$$

which implies $\lim_{r \rightarrow 1^-} \omega(r) = 0$. Hence every normal weight ω is also typical.

For a weight ω , the *associated weight* $\tilde{\omega}$ ([23]) is defined by

$$\tilde{\omega}(z) := \frac{1}{\sup \left\{ |f(z)| : f \in H_\omega^\infty, \|f\|_{H_\omega^\infty} \leq 1 \right\}}, \quad z \in \mathbb{B}. \quad (1.8)$$

Here H_ω^∞ denotes the *weighted-type space* consisting of all $f \in H(\mathbb{B})$ with

$$\|f\|_{H_\omega^\infty} = \sup_{z \in \mathbb{B}} \omega(z) |f(z)| < \infty \quad (1.9)$$

(see, e.g., [23, 24]). Associated weights assist us in studying of weighted-type spaces of holomorphic functions. It is known that associated weights are also continuous, $0 < \omega \leq \tilde{\omega}$, and for each $z \in \mathbb{B}$, we can find an $f_z \in H_\omega^\infty$, $\|f_z\|_{H_\omega^\infty} \leq 1$ such that $f_z(z) = 1/\tilde{\omega}(z)$. Let $H_{\omega,0}^\infty$ be the *little weighted-type space*, that is, the space of all $f \in H(\mathbb{B})$ such that $\omega(z)|f(z)| \rightarrow 0$ as $|z| \rightarrow 1^-$. If ω is typical, then the unit ball $B_{H_\omega^\infty}$ is the closure of $B_{H_{\omega,0}^\infty}$ for the compact open topology. Hence we have

$$\tilde{\omega}(z) = \frac{1}{\sup\{|f(z)| : f \in H_{\omega,0}^\infty, \|f\|_{H_\omega^\infty} \leq 1\}} \quad (1.10)$$

and so for each $z \in \mathbb{B}$, we can choose an $f_z \in B_{H_{\omega,0}^\infty}$ such that $f_z(z) = 1/\tilde{\omega}(z)$. A weight ω is called *essential* if it satisfies that $\tilde{\omega} \leq C\omega$ for some positive constant C . By the arguments in [25], we see that a normal weight function is also essential. For some examples of essential weights, see, for example, [25]. Related results can also be found in [22, 26].

The *Bloch-type space* $\mathcal{B}_\omega$ is the space of all holomorphic functions f on $\mathbb{B}$ such that

$$b_\omega(f) = \sup_{z \in \mathbb{B}} \omega(z) |\Re f(z)| < \infty, \quad (1.11)$$

where ω is a weight (see, e.g., [20]). The *little Bloch-type space* $\mathcal{B}_{\omega,0}$ consists of all $f \in H(\mathbb{B})$ such that

$$\lim_{|z| \rightarrow 1^-} \omega(z) |\Re f(z)| = 0. \quad (1.12)$$

Both spaces $\mathcal{B}_\omega$ and $\mathcal{B}_{\omega,0}$ are Banach spaces with the norm

$$\|f\|_{\mathcal{B}_\omega} = |f(0)| + b_\omega(f), \quad (1.13)$$

and $\mathcal{B}_{\omega,0}$ is a closed subspace of $\mathcal{B}_\omega$. When $\omega(r) = 1 - r^2$, the space $\mathcal{B}_\omega$ is a classical Bloch space.

The purpose of this paper is to characterize the boundedness and compactness of the operators $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ and $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$.

Throughout this paper, we assume that φ is a holomorphic self-map of $\mathbb{B}$ and $g \in H(\mathbb{B})$ with $g(0) = 0$. Furthermore, some constants are denoted by C ; they are positive and may differ from one occurrence to the other. The notation $a \leq b$ means that there exists a positive constant C such that $a \leq Cb$. Moreover, if both $a \leq b$ and $b \leq a$ hold, then one says that $a \asymp b$.

2. Auxiliary Results

Here we formulate and prove some auxiliary results which are used in the proofs of the main ones.

The following lemma was proved in [20, Theorem 2.1].

Lemma 2.1. *Let ω be a normal weight function and $f \in H(\mathbb{B})$. Then $f \in \mathcal{B}_\omega$ if and only if $\sup_{z \in \mathbb{B}} \omega(z)|\nabla f(z)| < \infty$ and it holds that*

$$\|f\|_{\mathcal{B}_\omega} \asymp |f(0)| + \sup_{z \in \mathbb{B}} \omega(z)|\nabla f(z)|. \quad (2.1)$$

Moreover, $f \in \mathcal{B}_{\omega,0}$ if and only if $\lim_{|z| \rightarrow 1^-} \omega(z)|\nabla f(z)| = 0$.

As an application of Lemma 2.1, we have the following result.

Lemma 2.2. *Let ω be a normal weight function and $f \in \mathcal{B}_\omega$. Then $f \in \mathcal{B}_{\omega,0}$ if and only if it holds that $\lim_{r \rightarrow 1^-} \|f_r - f\|_{\mathcal{B}_\omega} = 0$, where $f_r(z) = f(rz)$.*

Proof. Take an $f \in \mathcal{B}_{\omega,0}$. For a fixed $\varepsilon > 0$, by Lemma 2.1, we can choose a $\delta_0 \in (0, 1)$ such that

$$\omega(z)|\nabla f(z)| < \frac{\varepsilon}{2} \quad (2.2)$$

for any $z \in \mathbb{B} \setminus \delta_0^2 \overline{\mathbb{B}}$. Since $(\partial f_r / \partial z_j)(z) = r(\partial f / \partial z_j)(rz)$ for $j \in \{1, \dots, n\}$, $r \in (0, 1)$, and $z \in \mathbb{B}$, we have

$$\begin{aligned} \|f_r - f\|_{\mathcal{B}_\omega} &\asymp \sup_{z \in \mathbb{B}} \omega(z)|r\nabla f(rz) - \nabla f(z)| \\ &\leq \sup_{z \in \mathbb{B} \setminus \delta_0 \overline{\mathbb{B}}} \omega(z)|r\nabla f(rz) - \nabla f(z)| \\ &\quad + \sup_{z \in \delta_0 \overline{\mathbb{B}}} \omega(z)|r\nabla f(rz) - \nabla f(z)|. \end{aligned} \quad (2.3)$$

Since

$$\max_{|z| \leq \delta_0} |r\nabla f(rz) - \nabla f(z)| \longrightarrow 0, \quad \text{as } r \longrightarrow 1^-, \quad (2.4)$$

we see that the second term in (2.3) converges to 0 as $r \rightarrow 1^-$.

If $r \in (\delta_0, 1)$ and $z \in \mathbb{B} \setminus \delta_0 \overline{\mathbb{B}}$, then by (2.2) we have

$$\omega(rz)|\nabla f(rz)| < \frac{\varepsilon}{2}. \quad (2.5)$$

By (1.6) we have that $\omega(z) \leq \omega(rz)$ for $r, |z| \in [0, 1)$.

Hence we have

$$\sup_{z \in \mathbb{B} \setminus \delta_0 \overline{\mathbb{B}}} \omega(z)|r\nabla f(rz) - \nabla f(z)| \leq \sup_{z \in \mathbb{B} \setminus \delta_0 \overline{\mathbb{B}}} \omega(rz)|\nabla f(rz)| + \sup_{z \in \mathbb{B} \setminus \delta_0 \overline{\mathbb{B}}} \omega(z)|\nabla f(z)| < \varepsilon, \quad (2.6)$$

for all $r \in (\delta_0, 1)$. This proves that $\lim_{r \rightarrow 1^-} \|f_r - f\|_{\mathcal{B}_\omega} = 0$ whenever $f \in \mathcal{B}_{\omega,0}$.

Conversely, the normality of ω implies that for any $\varepsilon > 0$ we have

$$\omega(z)|\nabla f(rz)| \leq \varepsilon(1-|z|)^a \sup_{|w| \leq r} |\nabla f(w)| \longrightarrow 0, \quad \text{as } |z| \longrightarrow 1^-, \quad (2.7)$$

so that $f_r \in \mathcal{B}_{\omega,0}$ for any $r \in (0,1)$. On the other hand, by the assumption $\lim_{r \rightarrow 1^-} \|f_r - f\|_{\mathcal{B}_\omega} = 0$, we have that for every $\varepsilon > 0$ there is an $r_1 \in (0,1)$ such that

$$\|f_r - f\|_{\mathcal{B}_\omega} < \varepsilon, \quad (2.8)$$

for $r \in (r_1, 1)$.

By letting $|z| \rightarrow 1^-$ in the following inequality, which easily follows from Lemma 2.1:

$$\omega(z)|\nabla f(z)| \preceq \omega(z)|\nabla f(rz)| + \|f_r - f\|_{\mathcal{B}_\omega}, \quad (2.9)$$

then using (2.7) and (2.8), we get $f \in \mathcal{B}_{\omega,0}$, as claimed. $\square$

Corollary 2.3. *Let ω be a normal weight function. Then the set of all holomorphic polynomials is dense in $\mathcal{B}_{\omega,0}$.*

Proof. For the homogeneous expansion $f = \sum_{k=0}^{\infty} F_k$ of an $f \in \mathcal{B}_{\omega,0}$, we set $P^j = \sum_{k=0}^j F_k$ for each $j \in \mathbb{N}$. Since $P^j \rightarrow f$ uniformly on compact subsets of $\mathbb{B}$ as $j \rightarrow \infty$, we see that $\Re[P_r^j] \rightarrow \Re[f_r]$ uniformly on $\mathbb{B}$ for any $r \in (0,1)$. Moreover, we have

$$\begin{aligned} \|P_r^j - f\|_{\mathcal{B}_\omega} &\leq \|P_r^j - f_r\|_{\mathcal{B}_\omega} + \|f_r - f\|_{\mathcal{B}_\omega} \\ &\leq \sup_{z \in r\mathbb{B}} \omega(z) \sup_{z \in \mathbb{B}} |\Re[P_r^j](z) - \Re[f_r](z)| + \|f_r - f\|_{\mathcal{B}_\omega}. \end{aligned} \quad (2.10)$$

Combining this with Lemma 2.2, we get the desired result. $\square$

The following lemma can be found in [1, Lemma 3]. Its proof is similar to the proof of the corresponding one-dimensional result in [27], for the case of the little Bloch space $\mathcal{B}_{(1-r),0}$. Hence we omit the proof.

Lemma 2.4. *A closed subset K in $\mathcal{B}_{\omega,0}$ is compact if and only if it is bounded and*

$$\lim_{|z| \rightarrow 1^-} \sup_{f \in K} \omega(z)|\Re f(z)| = 0. \quad (2.11)$$

The following lemma is very useful for estimating the norm of the Bloch-type space.

Lemma 2.5. *Assume that m is a positive integer and ω is normal. Then for every $f \in H(\mathbb{B})$,*

$$\sup_{z \in \mathbb{B}} \omega(z)|f(z)| \asymp |f(0)| + \sup_{z \in \mathbb{B}} (1-|z|)^m \omega(z)|\Re^m f(z)|. \quad (2.12)$$

Proof. For the details of the proof, we can refer [9] or [28]. $\square$

3. The Boundedness of Operator I_φ^g

In this section we consider the boundedness of the operator $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ or $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$.

Theorem 3.1. *Let ω be a normal weight function and μ a weight function. Then the following conditions are equivalent:*

- (a) $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is bounded;
- (b) $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_\mu$ is bounded;
- (c) φ and g satisfy

$$\sup_{z \in \mathbb{B}} \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))} < \infty. \quad (3.1)$$

Moreover, if $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is bounded, then

$$\|I_\varphi^g\|_{\mathcal{B}_\omega \rightarrow \mathcal{B}_\mu} \asymp \sup_{z \in \mathbb{B}} \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))}. \quad (3.2)$$

Proof. The implication (a) $\Rightarrow$ (b) is clear, so we only prove (b) $\Rightarrow$ (c) and (c) $\Rightarrow$ (a).

(b) $\Rightarrow$ (c): assume that $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_\mu$ is bounded and fix $z \in \mathbb{B}$. We may assume that $\varphi(z) \neq 0$. For $w := \varphi(z)$, there exists $h_w \in H_{\omega,0}^\infty$ such that $\|h_w\|_{H_\omega^\infty} \leq 1$ and $h_w(w) = 1/\tilde{\omega}(w)$. We define the function f_w as follows:

$$f_w(v) = \int_0^1 h_w(tv) \frac{\langle tv, w \rangle}{|w|} \frac{dt}{t}, \quad v \in \mathbb{B}. \quad (3.3)$$

Since $\Re f_w(v) = h_w(v)(\langle v, w \rangle / |w|)$, we see that $f_w \in \mathcal{B}_{\omega,0}$ and $\|f_w\|_{\mathcal{B}_\omega} \leq 1$. Hence, by (1.4), we have

$$\|I_\varphi^g\|_{\mathcal{B}_\omega \rightarrow \mathcal{B}_\mu} \geq \|I_\varphi^g f_w\|_{\mathcal{B}_\mu} \geq \mu(z)|g(z)||\Re f_w(\varphi(z))| = \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))}, \quad (3.4)$$

and so condition (3.1) is true.

(c) $\Rightarrow$ (a): we assume (3.1) and take an $f \in \mathcal{B}_\omega$. Since ω is an essential weight (due to its normality), (1.4) gives

$$\begin{aligned} \mu(z)|\Re[I_\varphi^g f](z)| &= \mu(z)|g(z)||\Re f(\varphi(z))| \\ &\leq \mu(z)|g(z)||\varphi(z)| \frac{\tilde{\omega}(\varphi(z))|\nabla f(\varphi(z))|}{\tilde{\omega}(\varphi(z))} \\ &\leq \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))} \sup_{w \in \mathbb{B}} \omega(w)|\nabla f(w)|, \end{aligned} \quad (3.5)$$

for any $z \in \mathbb{B}$. By Lemma 2.1, we have $\sup_{w \in \mathbb{B}} \omega(w) |\nabla f(w)| \leq \|f\|_{\mathcal{B}_\omega}$, and so we obtain

$$\|I_\varphi^g f\|_{\mathcal{B}_\mu} \leq \sup_{z \in \mathbb{B}} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} \|f\|_{\mathcal{B}_\omega}. \quad (3.6)$$

This implies that $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is bounded. The relation (3.2) follows from (3.4) and (3.6). This completes the proof. $\square$

Theorem 3.2. *Let ω be a normal weight function and μ a weight function. Then the following conditions are equivalent:*

- (a) $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is bounded;
- (b) φ and g satisfy

$$\lim_{|z| \rightarrow 1^-} \mu(z) |g(z)| |\varphi(z)| = 0, \quad \sup_{z \in \mathbb{B}} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} < \infty. \quad (3.7)$$

Proof. (a) $\Rightarrow$ (b): as in the proof of Theorem 3.1, for fixed $z \in \mathbb{B}$ and $w = \varphi(z)$, we see that φ and g satisfy the condition

$$\sup_{z \in \mathbb{B}} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} < \infty. \quad (3.8)$$

On the other hand, since the normality of ω implies that the function $\pi_j(z) := z_j$ ($1 \leq j \leq n$) belongs to $\mathcal{B}_{\omega,0}$, we obtain that $\mu(z) |g(z)| |\varphi_j(z)| \rightarrow 0$ for each j , and so $\mu(z) |g(z)| |\varphi(z)| \rightarrow 0$ as $|z| \rightarrow 1^-$.

(b) $\Rightarrow$ (a): the assumption $\lim_{|z| \rightarrow 1^-} \mu(z) |g(z)| |\varphi(z)| = 0$ shows that $I_\varphi^g p \in \mathcal{B}_{\mu,0}$ for any polynomial p . For each $f \in \mathcal{B}_{\omega,0}$, by Corollary 2.3, we can choose a sequence of polynomials $\{p_j\}_{j \in \mathbb{N}}$ such that $\|f - p_j\|_{\mathcal{B}_\omega} \rightarrow 0$ as $j \rightarrow \infty$. Furthermore, the assumption

$$\sup_{z \in \mathbb{B}} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} < \infty \quad (3.9)$$

shows that $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_\mu$ is bounded by Theorem 3.1. Thus we obtain

$$0 \leq \|I_\varphi^g f - I_\varphi^g p_j\|_{\mathcal{B}_\mu} \leq \|I_\varphi^g\|_{\mathcal{B}_\omega \rightarrow \mathcal{B}_\mu} \|f - p_j\|_{\mathcal{B}_\omega} \rightarrow 0 \quad (\text{as } j \rightarrow \infty). \quad (3.10)$$

Since $I_\varphi^g f \in \mathcal{B}_\mu$, $\{I_\varphi^g p_j\}_{j \in \mathbb{N}} \subset \mathcal{B}_{\mu,0}$, and $\mathcal{B}_{\mu,0}$ is closed in $\mathcal{B}_\mu$, we have $I_\varphi^g f \in \mathcal{B}_{\mu,0}$ for any $f \in \mathcal{B}_{\omega,0}$. Hence $I_\varphi^g(\mathcal{B}_{\omega,0}) \subseteq \mathcal{B}_{\mu,0}$ which means that $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is bounded. The proof is accomplished. $\square$

The following corollary is an immediate consequence of Theorems 3.1 and 3.2.

Corollary 3.3. *Let ω be a normal weight function and μ a weight function. Then $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is bounded if and only if $\lim_{|z| \rightarrow 1^-} \mu(z) |g(z)| |\varphi(z)| = 0$ and $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is bounded.*

4. The Compactness of Operator I_φ^g

In this section we characterize the compactness of $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ or $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$. To do this, we need the following standard lemma (see, e.g., [13, Lemma 3]).

Lemma 4.1. *Let ω and μ be weight functions. Suppose that the operator $I_\varphi^g : \mathcal{B}_\omega$ (or $\mathcal{B}_{\omega,0}$) $\rightarrow \mathcal{B}_\mu$ is bounded. Then $I_\varphi^g : \mathcal{B}_\omega$ (or $\mathcal{B}_{\omega,0}$) $\rightarrow \mathcal{B}_\mu$ is compact if and only if for every bounded sequence $\{f_j\}_{j \in \mathbb{N}}$ in $\mathcal{B}_\omega$ (or $\mathcal{B}_{\omega,0}$) which converges to 0 uniformly on compact subsets of $\mathbb{B}$, $\|I_\varphi^g f_j\|_{\mathcal{B}_\mu} \rightarrow 0$ as $j \rightarrow \infty$.*

Theorem 4.2. *Let ω and μ be weight functions. Suppose that φ is a holomorphic self-map of $\mathbb{B}$ such that $\|\varphi\|_\infty < 1$ and the operator $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is bounded. Then $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is compact. Here $\|\varphi\|_\infty$ denotes the supremum $\sup_{z \in \mathbb{B}} |\varphi(z)|$.*

Proof. Since $\|\varphi\|_\infty < 1$, we see that $|\varphi(z)| \leq r$ for some $r \in (0, 1)$ and any $z \in \mathbb{B}$. From the proof of Theorem 3.1, we see that the boundedness of $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ implies

$$M := \sup_{z \in \mathbb{B}} \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))} < \infty. \quad (4.1)$$

Thus we obtain that

$$\sup_{z \in \mathbb{B}} \mu(z)|g(z)||\varphi(z)| \leq M \sup_{w \in r\bar{\mathbb{B}}} \tilde{\omega}(w) < \infty. \quad (4.2)$$

Take a bounded sequence $\{f_j\}_{j \in \mathbb{N}}$ in $\mathcal{B}_\omega$ such that $f_j \rightarrow 0$ uniformly on compact subsets of $\mathbb{B}$ as $j \rightarrow \infty$. By (1.4), we have

$$\begin{aligned} \|I_\varphi^g f_j\|_{\mathcal{B}_\mu} &= \sup_{z \in \mathbb{B}} \mu(z)|g(z)||\Re f_j(\varphi(z))| \\ &\leq \sup_{z \in \mathbb{B}} \mu(z)|g(z)||\varphi(z)||\nabla f_j(\varphi(z))| \\ &\leq M \sup_{w \in r\bar{\mathbb{B}}} \tilde{\omega}(w) \sup_{w \in r\bar{\mathbb{B}}} |\nabla f_j(w)|. \end{aligned} \quad (4.3)$$

Since $\partial f_j / \partial z_k$ ($1 \leq k \leq n$) also converges to 0 uniformly on $r\bar{\mathbb{B}}$ as $j \rightarrow \infty$, (4.2) and (4.3) show that $\|I_\varphi^g f_j\|_{\mathcal{B}_\mu} \rightarrow 0$ as $j \rightarrow \infty$. From Lemma 4.1, it follows that $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is compact, and so we get the assertions. $\square$

Lemma 4.3. *Suppose that ω is a weight function. Then there exists a sequence $\{f_k\}_{k \in \mathbb{N}}$ in the closed unit ball of $\mathcal{B}_\omega$ such that $f_k \rightarrow 0$ uniformly on compact subsets of $\mathbb{B}$ as $k \rightarrow \infty$.*

Proof. Let $\{w_k\}_{k \in \mathbb{N}} \subset \mathbb{B}$ with $|w_k| \rightarrow 1^-$ as $k \rightarrow \infty$. For each w_k , there exists $h_k := h_{w_k} \in H_\omega^\infty$ such that $\|h_k\|_{H_\omega^\infty} \leq 1$ and $h_k(w_k) = 1/\tilde{\omega}(w_k)$. We define f_k as follows:

$$f_k(z) = \int_0^1 h_k(tz) \left\{ \frac{\langle tz, w_k \rangle}{|w_k|} \right\}^{1/(1-|w_k|)} \frac{dt}{t}, \quad z \in \mathbb{B}. \quad (4.4)$$

Since $f_k(0) = 0$ and $|\Re f_k(z)| \leq |h_k(z)|$, we have $\{f_k\}_{k \in \mathbb{N}} \subset \mathcal{B}_\omega$ and $\|f_k\|_{\mathcal{B}_\omega} \leq 1$ for each $k \in \mathbb{N}$. For any compact subset $\mathcal{K}$ of $\mathbb{B}$, we can choose an $r \in (0, 1)$ such that $\mathcal{K} \subset r\overline{\mathbb{B}}$. Hence we obtain that for any $z \in \mathcal{K}$

$$|f_k(z)| \leq \int_0^1 |h_k(tz)| t^{|w_k|/(1-|w_k|)} dt \leq \|h_k\|_{H_\omega^\infty} \int_0^1 \frac{1}{\omega(tz)} t^{|w_k|/(1-|w_k|)} dt \leq \max_{w \in r\overline{\mathbb{B}}} \frac{1}{\omega(w)} (1 - |w_k|). \quad (4.5)$$

From the above inequality, it follows that f_k converges to 0 uniformly on compact subsets of $\mathbb{B}$ as $k \rightarrow \infty$. This completes the proof. $\square$

Remark 4.4. If we assume that ω is typical in Lemma 4.3, then we can choose $h_k \in H_{\omega,0}^\infty$. In this case, hence, we see that f_k belongs to $\mathcal{B}_{\omega,0}$ for each $k \in \mathbb{N}$.

Theorem 4.5. *Let ω be a normal weight function and μ a weight function. Suppose that the operator $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is bounded and $\|\varphi\|_\infty = 1$. Then the following conditions are equivalent:*

- (a) $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is compact;
- (b) $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_\mu$ is compact;
- (c) φ and g satisfy

$$\lim_{|\varphi(z)| \rightarrow 1^-} \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))} = 0. \quad (4.6)$$

Proof. (a) $\Rightarrow$ (b): this implication is obvious.

(b) $\Rightarrow$ (c): take a sequence $\{z_k\}_{k \in \mathbb{N}}$ in $\mathbb{B}$ with $|\varphi(z_k)| \rightarrow 1^-$ as $k \rightarrow \infty$ and put $w_k = \varphi(z_k)$ for each k . Then, by Remark 4.4 after Lemma 4.3, there exists a sequence $\{f_k\}_{k \in \mathbb{N}}$ in $\mathcal{B}_{\omega,0}$ such that $\sup_{k \in \mathbb{N}} \|f_k\|_{\mathcal{B}_\omega} \leq 1$ and $f_k \rightarrow 0$ uniformly on compact subsets of $\mathbb{B}$ as $k \rightarrow \infty$. By Lemma 4.1, the compactness of $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_\mu$ implies that $\|I_\varphi^g f_k\|_{\mathcal{B}_\mu} \rightarrow 0$ as $k \rightarrow \infty$.

On the other hand, (1.4) gives $\Re[I_\varphi^g f_k](z) = \Re f_k(\varphi(z))g(z)$, and so we have

$$\|I_\varphi^g f_k\|_{\mathcal{B}_\mu} \geq \mu(z_k) |\Re f_k(\varphi(z_k))| |g(z_k)| \geq \mu(z_k) |\Re f_k(\varphi(z_k))| |g(z_k)| |\varphi(z_k)|. \quad (4.7)$$

From the construction (4.4) of f_k , we obtain

$$\Re f_k(\varphi(z_k)) = \frac{|\varphi(z_k)|^{1/(1-|\varphi(z_k)|)}}{\tilde{\omega}(\varphi(z_k))}, \quad (4.8)$$

for each $k \in \mathbb{N}$. Combining this with (4.7), we have

$$\|I_\varphi^g f_k\|_{\mathcal{B}_\mu} \geq \frac{\mu(z_k) |g(z_k)| |\varphi(z_k)|}{\tilde{\omega}(\varphi(z_k))} |\varphi(z_k)|^{1/(1-|\varphi(z_k)|)}. \quad (4.9)$$

Letting $k \rightarrow \infty$, we have

$$\lim_{k \rightarrow \infty} \frac{\mu(z_k) |g(z_k)| |\varphi(z_k)|}{\tilde{\omega}(\varphi(z_k))} = 0, \quad (4.10)$$

for any sequence $\{z_k\}_{k \in \mathbb{N}}$ with $|\varphi(z_k)| \rightarrow 1^-$. This proves that (4.6) is true.

(c) $\Rightarrow$ (a): we will prove the following estimate:

$$\|I_\varphi^g\|_e \leq \limsup_{|\varphi(z)| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))}. \quad (4.11)$$

Here $\|I_\varphi^g\|_e$ denotes the essential norm of $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$, namely,

$$\|I_\varphi^g\|_e = \inf \left\{ \|I_\varphi^g + \mathcal{K}\|_{\mathcal{B}_\omega \rightarrow \mathcal{B}_\mu} \mid \mathcal{K} : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu \text{ is compact} \right\}. \quad (4.12)$$

Now we take a sequence $\{r_l\}_{l \in \mathbb{N}} \subset (0, 1)$ which increasingly converges to 1 and put

$$I_{r_l \varphi}^g f(z) = \int_0^1 \Re f(r_l \varphi(tz)) g(tz) \frac{dt}{t}. \quad (4.13)$$

Since $\|r_l \varphi\|_\infty \leq r_l < 1$, Theorem 4.2 implies that $I_{r_l \varphi}^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is compact for each $l \in \mathbb{N}$. For any $f \in \mathcal{B}_\omega$ with $\|f\|_{\mathcal{B}_\omega} \leq 1$, from (1.4) it follows that

$$\begin{aligned} \|I_\varphi^g f - I_{r_l \varphi}^g f\|_{\mathcal{B}_\omega} &= \sup_{z \in \mathbb{B}} \mu(z) |g(z)| |\Re f(\varphi(z)) - \Re f(r_l \varphi(z))| \\ &\leq \sup_{R < |\varphi(z)| < 1} \mu(z) |g(z)| |\Re f(\varphi(z)) - \Re f(r_l \varphi(z))| \\ &\quad + \sup_{|\varphi(z)| \leq R} \mu(z) |g(z)| |\Re f(\varphi(z)) - \Re f(r_l \varphi(z))|, \end{aligned} \quad (4.14)$$

for some fixed $R \in (0, 1)$. The essentiality of ω and Lemma 2.1 give

$$\begin{aligned} \mu(z) |g(z)| |\Re f(\varphi(z))| &\leq \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} \tilde{\omega}(\varphi(z)) |\nabla f(\varphi(z))| \\ &\leq \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} \sup_{w \in \mathbb{B}} \omega(w) |\nabla f(w)| \\ &\leq \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))}. \end{aligned} \quad (4.15)$$

Similarly, we also have

$$\mu(z)|g(z)||\Re f(r_l\varphi(z))| \leq \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(r_l\varphi(z))}, \quad (4.16)$$

for each $l \in \mathbb{N}$. The normality of ω implies that

$$\frac{\omega(r_l\varphi(z))}{(1-|r_l\varphi(z)|)^a} \geq \frac{\omega(\varphi(z))}{(1-|\varphi(z)|)^a}, \quad (4.17)$$

for each $l \in \mathbb{N}$ and some $a > 0$, and so by the essentiality,

$$\frac{\tilde{\omega}(r_l\varphi(z))}{(1-|r_l\varphi(z)|)^a} \geq \frac{\tilde{\omega}(\varphi(z))}{(1-|\varphi(z)|)^a}. \quad (4.18)$$

Thus (4.16) and (4.18) give

$$\mu(z)|g(z)||\Re f(r_l\varphi(z))| \leq \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))}, \quad (4.19)$$

for each $l \in \mathbb{N}$. By (4.15) and (4.19), we obtain

$$\sup_{R < |\varphi(z)| < 1} \mu(z)|g(z)||\Re f(\varphi(z)) - \Re f(r_l\varphi(z))| \leq \sup_{R < |\varphi(z)| < 1} \frac{\mu(z)|g(z)||\varphi(z)|}{\tilde{\omega}(\varphi(z))}. \quad (4.20)$$

When $|\varphi(z)| \leq R$, by using the mean value theorem, we have

$$\begin{aligned} \mu(z)|g(z)||\Re f(\varphi(z)) - \Re f(r_l\varphi(z))| &\leq (1-r_l)\mu(z)|g(z)||\varphi(z)| \sup_{|w| \leq R} |\nabla [\Re f](w)| \\ &\leq \frac{1-r_l}{1-R} \max_{w \in \mathbb{B}} \frac{1}{\omega(w)} \sup_{z \in \mathbb{B}} \mu(z)|g(z)||\varphi(z)| \\ &\quad \times \sup_{w \in \mathbb{B}} \omega(w)(1-|w|) |\nabla [\Re f](w)|. \end{aligned} \quad (4.21)$$

Since $\omega(w)(1-|w|)$ is also normal, by Lemmas 2.1 and 2.5, we have

$$\begin{aligned} \sup_{w \in \mathbb{B}} \omega(w)(1-|w|) |\nabla [\Re f](w)| &\asymp \sup_{w \in \mathbb{B}} \omega(w)(1-|w|) |\Re^2 f(w)| \\ &\asymp \sup_{w \in \mathbb{B}} \omega(w) |\Re f(w)|. \end{aligned} \quad (4.22)$$

Hence we obtain

$$\sup_{|\varphi(z)| \leq R} \mu(z) |g(z)| |\Re f(\varphi(z)) - \Re f(r_l \varphi(z))| \leq \frac{1-r_l}{1-R} \max_{w \in r\mathbb{B}} \frac{1}{\omega(w)} \sup_{z \in \mathbb{B}} \mu(z) |g(z)| |\varphi(z)|. \quad (4.23)$$

Since the boundedness of $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ implies $\sup_{z \in \mathbb{B}} \mu(z) |g(z)| |\varphi(z)| < \infty$, letting $l \rightarrow \infty$ in the above inequality, we have

$$\sup_{\|f\|_{\mathcal{B}_\omega} \leq 1} \sup_{|\varphi(z)| \leq R} \mu(z) |g(z)| |\Re f(\varphi(z)) - \Re f(r_l \varphi(z))| \rightarrow 0. \quad (4.24)$$

By using (4.14), (4.20), and (4.24) and letting $R \rightarrow 1^-$, we obtain the desired estimate

$$\|I_\varphi^g\|_e \leq \limsup_{|\varphi(z)| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))}. \quad (4.25)$$

So if condition (4.6) is true, then $\|I_\varphi^g\|_e = 0$, which means that $I_\varphi^g : \mathcal{B}_\omega \rightarrow \mathcal{B}_\mu$ is compact. Our proof is accomplished. $\square$

Theorem 4.6. *Let ω be a normal weight function and μ a weight function. Suppose that the operator $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is bounded. Then $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is compact if and only if*

$$\lim_{|z| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} = 0. \quad (4.26)$$

Proof. Suppose that (4.26) holds. For any $f \in \mathcal{B}_{\omega,0}$, by Lemma 2.1 and (1.4), we have

$$\begin{aligned} \mu(z) \left| \Re \left[I_\varphi^g f \right] (z) \right| &= \mu(z) |\Re f(\varphi(z))| |g(z)| \\ &\leq \mu(z) |g(z)| |\varphi(z)| |\nabla f(\varphi(z))| \\ &\leq \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} \omega(\varphi(z)) |\nabla f(\varphi(z))| \\ &\leq \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} \|f\|_{\mathcal{B}_\omega}. \end{aligned} \quad (4.27)$$

Combining this with (4.26), we obtain

$$\lim_{|z| \rightarrow 1^-} \sup_{\|f\|_{\mathcal{B}_\omega} \leq 1} \mu(z) \left| \Re \left[I_\varphi^g f \right] (z) \right| = 0. \quad (4.28)$$

Hence it follows from Lemma 2.4 that $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is compact.

Conversely, we assume that $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_{\mu,0}$ is compact. By Theorem 3.2, we see that

$$\lim_{|z| \rightarrow 1^-} \mu(z) |g(z)| |\varphi(z)| = 0. \quad (4.29)$$

Thus this implies (4.26) if $\|\varphi\|_\infty < 1$.

Now assume $\|\varphi\|_\infty = 1$. We claim that

$$\limsup_{|z| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} = \limsup_{|\varphi(z)| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))}. \quad (4.30)$$

Further assume that $\{z_k\}_{k \in \mathbb{N}}$ is a sequence in $\mathbb{B}$ such that

$$\limsup_{|z| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} = \lim_{k \rightarrow \infty} \frac{\mu(z_k) |g(z_k)| |\varphi(z_k)|}{\tilde{\omega}(\varphi(z_k))}. \quad (4.31)$$

If $\sup_{k \in \mathbb{N}} |\varphi(z_k)| < 1$, then from this and (4.29) we have that both limits in (4.30) are equal to zero. If $\sup_{k \in \mathbb{N}} |\varphi(z_k)| = 1$, then there is a subsequence $\{\varphi(z_{k_l})\}_{l \in \mathbb{N}}$ such that $|\varphi(z_{k_l})| \rightarrow 1^-$ as $l \rightarrow \infty$. Hence we have

$$\limsup_{|z| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))} = \lim_{l \rightarrow \infty} \frac{\mu(z_{k_l}) |g(z_{k_l})| |\varphi(z_{k_l})|}{\tilde{\omega}(\varphi(z_{k_l}))} \leq \limsup_{|\varphi(z)| \rightarrow 1^-} \frac{\mu(z) |g(z)| |\varphi(z)|}{\tilde{\omega}(\varphi(z))}, \quad (4.32)$$

and so (4.30) holds.

Since $I_\varphi^g : \mathcal{B}_{\omega,0} \rightarrow \mathcal{B}_\mu$ is also compact, by Theorem 4.5, we see that the second limit in (4.30) is equal to zero, so that (4.26) holds. This completes the proof. $\square$

Acknowledgment

This work was supported by JSPS Grant-in-Aid for Young Scientists (Start-up: no. 20840004) and Ibaraki University.

References

- [1] S. Stević, "On an integral operator from the Zygmund space to the Bloch-type space on the unit ball," *Glasgow Mathematical Journal*, vol. 51, no. 2, pp. 275–287, 2009.
- [2] S. Li and S. Stević, "Generalized composition operators on Zygmund spaces and Bloch type spaces," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1282–1295, 2008.
- [3] S. Stević, "Generalized composition operators from logarithmic Bloch spaces to mixed-norm spaces," *Utilitas Mathematica*, vol. 77, pp. 167–172, 2008.
- [4] S. Stević, "On an integral operator between Bloch-type spaces on the unit ball," *Bulletin des Sciences Mathématiques*. In press.
- [5] S. Stević, "Products of integral-type operators and composition operators from a mixed norm space to Bloch-type spaces," *Siberian Mathematical Journal*, vol. 50, no. 4, pp. 726–736, 2009.
- [6] X. Zhu, "Generalized composition operators from generalized weighted Bergman spaces to Bloch type spaces," *Journal of the Korean Mathematical Society*, vol. 46, no. 6, pp. 1219–1232, 2009.

- [7] S. Stević and S.-I. Ueki, "Integral-type operators acting between weighted-type spaces on the unit ball," *Applied Mathematics and Computation*, vol. 215, no. 7, pp. 2464–2471, 2009.
- [8] D.-C. Chang, S. Li, and S. Stević, "On some integral operators on the unit polydisk and the unit ball," *Taiwanese Journal of Mathematics*, vol. 11, no. 5, pp. 1251–1285, 2007.
- [9] Z. Hu, "Extended Cesàro operators on mixed norm spaces," *Proceedings of the American Mathematical Society*, vol. 131, no. 7, pp. 2171–2179, 2003.
- [10] S. G. Krantz and S. Stević, "On the iterated logarithmic Bloch space on the unit ball," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 5-6, pp. 1772–1795, 2009.
- [11] S. Stević, "On an integral operator on the unit ball in $\mathbb{C}^n$," *Journal of Inequalities and Applications*, vol. 2005, no. 1, pp. 81–88, 2005.
- [12] S. Stević, "Boundedness and compactness of an integral operator on a weighted space on the polydisc," *Indian Journal of Pure and Applied Mathematics*, vol. 37, no. 6, pp. 343–355, 2006.
- [13] S. Stević, "Boundedness and compactness of an integral operator in a mixed norm space on the polydisk," *Sibirskii Matematicheskii Zhurnal*, vol. 48, no. 3, pp. 694–706, 2007.
- [14] S. Stević, "On a new operator from H^∞ to the Bloch-type space on the unit ball," *Utilitas Mathematica*, vol. 77, pp. 257–263, 2008.
- [15] S. Stević, "On a new integral-type operator from the weighted Bergman space to the Bloch-type space on the unit ball," *Discrete Dynamics in Nature and Society*, vol. 2008, Article ID 154263, 14 pages, 2008.
- [16] S. Stević, "On a new operator from the logarithmic Bloch space to the Bloch-type space on the unit ball," *Applied Mathematics and Computation*, vol. 206, no. 1, pp. 313–320, 2008.
- [17] S. Stević, "On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball," *Journal of Mathematical Analysis and Applications*, vol. 354, no. 2, pp. 426–434, 2009.
- [18] S. Stević, "On an integral-type operator from logarithmic Bloch-type and mixed-norm spaces to Bloch-type spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 12, pp. 6323–6342, 2009.
- [19] J. Xiao, "Riemann-Stieltjes operators on weighted Bloch and Bergman spaces of the unit ball," *Journal of the London Mathematical Society*, vol. 70, no. 1, pp. 199–214, 2004.
- [20] X. Tang, "Extended Cesàro operators between Bloch-type spaces in the unit ball of $\mathbb{C}^n$," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 1199–1211, 2007.
- [21] W. Yang, "On an integral-type operator between Bloch-type spaces," *Applied Mathematics and Computation*, vol. 215, no. 3, pp. 954–960, 2009.
- [22] S. Stević and E. Wolf, "Differences of composition operators between weighted-type spaces of holomorphic functions on the unit ball of $\mathbb{C}^n$," *Applied Mathematics and Computation*, vol. 215, no. 5, pp. 1752–1760, 2009.
- [23] K. D. Bierstedt and W. H. Summers, "Biduals of weighted Banach spaces of analytic functions," *Journal of Australian Mathematical Society. Series A*, vol. 54, no. 1, pp. 70–79, 1993.
- [24] S. Stević, "Norm of weighted composition operators from Bloch space to H_μ^∞ on the unit ball," *Ars Combinatoria*, vol. 88, pp. 125–127, 2008.
- [25] K. D. Bierstedt, J. Bonet, and J. Taskinen, "Associated weights and spaces of holomorphic functions," *Studia Mathematica*, vol. 127, no. 2, pp. 137–168, 1998.
- [26] M. Lindström and E. Wolf, "Essential norm of the difference of weighted composition operators," *Monatshefte für Mathematik*, vol. 153, no. 2, pp. 133–143, 2008.
- [27] K. Madigan and A. Matheson, "Compact composition operators on the Bloch space," *Transactions of the American Mathematical Society*, vol. 347, no. 7, pp. 2679–2687, 1995.
- [28] S. Stević and S.-I. Ueki, "Weighted composition operators and integral-type operators between weighted Hardy spaces on the unit ball," *Discrete Dynamics in Nature and Society*, vol. 2009, Article ID 952831, 21 pages, 2009.