Research Article

Best Possible Inequalities between Generalized Logarithmic Mean and Classical Means

Yu-Ming Chu¹ and Bo-Yong Long²

¹ Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China

² College of Mathematics Science, Anhui University, Hefei 230039, China

Correspondence should be addressed to Yu-Ming Chu, chuyuming2005@yahoo.com.cn

Received 26 November 2009; Revised 16 February 2010; Accepted 16 March 2010

Academic Editor: Lance Littlejohn

Copyright © 2010 Y.-M. Chu and B.-Y. Long. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We answer the question: for $\alpha, \beta, \gamma \in (0, 1)$ with $\alpha + \beta + \gamma = 1$, what are the greatest value p and the least value q , such that the double inequality $L_p(a, b) < A^\alpha(a, b)G^\beta(a, b)H^\gamma(a, b) < L_q(a, b)$ holds for all $a, b > 0$ with $a \neq b$? Here $L_p(a, b)$, $A(a, b)$, $G(a, b)$, and $H(a, b)$ denote the generalized logarithmic, arithmetic, geometric, and harmonic means of two positive numbers a and b , respectively.

1. Introduction

For $p \in \mathbb{R}$ the generalized logarithmic mean $L_p(a, b)$ of two positive numbers a and b with $a \neq b$ is defined by

$$L_p(a, b) = \begin{cases} \left[\frac{a^{p+1} - b^{p+1}}{(p+1)(a-b)} \right]^{1/p}, & p \neq 0, p \neq -1, \\ \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{1/(b-a)}, & p = 0, \\ \frac{b-a}{\log b - \log a}, & p = -1. \end{cases} \quad (1.1)$$

It is well known that $L_p(a, b)$ is continuous and strictly increasing with respect to $p \in \mathbb{R}$ for fixed $a, b > 0$ with $a \neq b$. Recently, the generalized logarithmic mean has been

the subject of intensive research. Many remarkable inequalities and monotonicity results for the generalized logarithmic mean can be found in the literature [1–9]. It might be surprising that the generalized logarithmic mean has applications in physics, economics, and even in meteorology [10–13].

Let $A(a, b) = (a + b)/2$, $I(a, b) = (1/e)(b^b/a^a)^{1/(b-a)}$, $L(a, b) = (b - a)/(\log b - \log a)$, $G(a, b) = \sqrt{ab}$, and $H(a, b) = 2ab/(a + b)$ be the arithmetic, identric, logarithmic, geometric, and harmonic means of two positive numbers a and b with $a \neq b$, respectively. Then

$$\begin{aligned} \min\{a, b\} < H(a, b) < G(a, b) = L_{-2}(a, b) < L(a, b) = L_{-1}(a, b) \\ < I(a, b) = L_0(a, b) < A(a, b) = L_1(a, b) < \max\{a, b\}. \end{aligned} \quad (1.2)$$

For $p \in \mathbb{R}$, the p th power mean $M_p(a, b)$ of two positive numbers a and b with $a \neq b$ is defined by

$$M_p(a, b) = \left(\frac{a^p + b^p}{2} \right)^{1/p} \quad (p \neq 0), \quad M_0(a, b) = (ab)^{1/2}. \quad (1.3)$$

In [14], Alzer and Janous established the following sharp double inequality (see also [15, page 350]):

$$M_{\log 2 / \log 3}(a, b) < \frac{2}{3}A(a, b) + \frac{1}{3}G(a, b) < M_{2/3}(a, b) \quad (1.4)$$

for all $a, b > 0$ with $a \neq b$.

For $\alpha \in (0, 1)$, Janous [16] found the greatest value p and the least value q such that

$$M_p(a, b) < \alpha A(a, b) + (1 - \alpha)G(a, b) < M_q(a, b) \quad (1.5)$$

for all $a, b > 0$ with $a \neq b$.

In [17–19], the authors presented bounds for $L(a, b)$ and $I(a, b)$ in terms of $G(a, b)$ and $A(a, b)$.

Theorem A. *For all positive real numbers a and b with $a \neq b$, we have*

$$\begin{aligned} L(a, b) &< \frac{1}{3}A(a, b) + \frac{2}{3}G(a, b), \\ \frac{1}{3}G(a, b) + \frac{2}{3}A(a, b) &< I(a, b). \end{aligned} \quad (1.6)$$

The proof of the following Theorem B can be found in [20].

Theorem B. For all positive real numbers a and b with $a \neq b$, we have

$$\begin{aligned}\sqrt{G(a,b)A(a,b)} &< \sqrt{L(a,b)I(a,b)} \\ &< \frac{1}{2}(L(a,b) + I(a,b)) \\ &< \frac{1}{2}(G(a,b) + A(a,b)).\end{aligned}\tag{1.7}$$

The following Theorems C–E were established by Alzer and Qiu in [21].

Theorem C. The inequalities

$$\alpha A(a,b) + (1-\alpha)G(a,b) < I(a,b) < \beta A(a,b) + (1-\beta)G(a,b)\tag{1.8}$$

hold for all positive real numbers a and b with $a \neq b$ if and only if $\alpha \leq 2/3$ and $\beta \geq 2/e = 0.73575 \dots$.

Theorem D. Let a and b be real numbers with $a \neq b$. If $0 < a, b \leq e$, then

$$[G(a,b)]^{A(a,b)} < [L(a,b)]^{I(a,b)} < [A(a,b)]^{G(a,b)}.\tag{1.9}$$

And if $a, b \geq e$, then

$$[A(a,b)]^{G(a,b)} < [I(a,b)]^{L(a,b)} < [G(a,b)]^{A(a,b)}.\tag{1.10}$$

Theorem E. For all positive real numbers a and b with $a \neq b$, we have

$$M_p(a,b) < \frac{1}{2}(L(a,b) + I(a,b))\tag{1.11}$$

with the best possible parameter $p = \log 2 / (1 + \log 2) = 0.40938 \dots$.

However, the following problem is still open: for $\alpha, \beta, \gamma \in (0, 1)$ with $\alpha + \beta + \gamma = 1$, what are the greatest value p and the least value q , such that the double inequality

$$L_p(a,b) < A^\alpha(a,b)G^\beta(a,b)H^\gamma(a,b) < L_q(a,b)\tag{1.12}$$

holds for all $a, b > 0$ with $a \neq b$? The purpose of this paper is to give the solution to this open problem.

2. Lemmas

In order to establish our main result, we need four lemmas, which we present in this section.

Lemma 2.1. *If $t > 1$, then*

$$\frac{t-1}{\log t} - \left[\frac{t(t+1)}{2} \right]^{1/3} > 0. \quad (2.1)$$

Proof. If $t > 1$, then it follows from [22, 3.6.16] that

$$\frac{t-1}{\log t} > \frac{t+t^{1/3}}{1+t^{1/3}}. \quad (2.2)$$

It is not difficult to verify that

$$\frac{t+t^{1/3}}{1+t^{1/3}} > \left[\frac{t(t+1)}{2} \right]^{1/3} \quad (2.3)$$

for $t > 1$.

Therefore, Lemma 2.1 follows from inequalities (2.2) and (2.3). $\square$

Lemma 2.2. *If $t > 1$, then*

$$\frac{t}{t-1} \log t - \frac{1}{6} \log t - \frac{2}{3} \log \frac{1+t}{2} - 1 > 0. \quad (2.4)$$

Proof. Let $f(t) = (t/(t-1)) \log t - (1/6) \log t - (2/3) \log((1+t)/2) - 1$. Then simple computations lead to

$$\lim_{t \rightarrow 1} f(t) = 0, \quad (2.5)$$

$$f'(t) = \frac{g(t)}{6t(t-1)^2(t+1)}, \quad (2.6)$$

where $g(t) = -6t(t+1) \log t + t^3 + 9t^2 - 9t - 1$,

$$g(1) = 0, \quad (2.7)$$

$$g'(t) = -6(2t+1) \log t + 3t^2 + 12t - 15, \quad (2.8)$$

$$g'(1) = 0, \quad (2.9)$$

$$g''(t) = \frac{h(t)}{t}, \quad (2.10)$$

where $h(t) = -12 \log t + 6t^2 - 6$,

$$g''(1) = h(1) = 0, \quad (2.11)$$

$$h'(t) = -12 \log t + 12t - 12, \quad (2.12)$$

$$h'(1) = 0, \quad (2.13)$$

$$h''(t) = \frac{12}{t}(t-1) > 0 \quad (2.14)$$

for $t > 1$.

Therefore, Lemma 2.2 follows from (2.5)–(2.7), (2.9)–(2.11), (2.13), and (2.14). $\square$

Lemma 2.3. *Let $t > 1$ and $g(t) = (1 - (1/2)\lambda)(3\lambda - 5)t^{3\lambda-2} + [(\lambda - 1)(3\lambda - 5) - 1]t^{3\lambda-3} - [(1/2)\lambda(3\lambda - 5) + 1]t^{3\lambda-4} + [(1/2)\lambda(3\lambda - 5) + 1]t^2 - [(\lambda - 1)(3\lambda - 5) - 1]t - (1 - (1/2)\lambda)(3\lambda - 5)$. Then*

(1) $g(t) > 0$ for $\lambda \in (0, 2/3) \cup (1, 4/3) \cup (5/3, 2)$ and

(2) $g(t) < 0$ for $\lambda \in (2/3, 1) \cup (4/3, 5/3)$.

Proof. Simple computations yield

$$\lim_{t \rightarrow 1} g(t) = 0, \quad (2.15)$$

$$\begin{aligned} g'(t) = & \left(1 - \frac{1}{2}\lambda\right)(3\lambda - 5)(3\lambda - 2)t^{3\lambda-3} + 3(\lambda - 1)[(\lambda - 1)(3\lambda - 5) - 1]t^{3\lambda-4} \\ & - (3\lambda - 4)\left[\frac{1}{2}\lambda(3\lambda - 5) + 1\right]t^{3\lambda-5} + [\lambda(3\lambda - 5) + 2]t \\ & - (\lambda - 1)(3\lambda - 5) + 1, \end{aligned} \quad (2.16)$$

$$g'(1) = 0, \quad (2.17)$$

$$\begin{aligned} g''(t) = & 3\left(1 - \frac{1}{2}\lambda\right)(\lambda - 1)(3\lambda - 5)(3\lambda - 2)t^{3\lambda-4} \\ & + 3(\lambda - 1)(3\lambda - 4)[(\lambda - 1)(3\lambda - 5) - 1]t^{3\lambda-5} \\ & - (3\lambda - 4)(3\lambda - 5)\left[\frac{1}{2}\lambda(3\lambda - 5) + 1\right]t^{3\lambda-6} + \lambda(3\lambda - 5) + 2, \end{aligned} \quad (2.18)$$

$$g''(1) = 0, \quad (2.19)$$

$$g'''(t) = t^{3\lambda-7}h(t), \quad (2.20)$$

where $h(t) = 3(1 - (1/2)\lambda)(\lambda - 1)(3\lambda - 5)(3\lambda - 4)(3\lambda - 2)t^2 + 3(\lambda - 1)(3\lambda - 4)(3\lambda - 5)[(\lambda - 1)(3\lambda - 5) - 1]t - 3(\lambda - 2)(3\lambda - 4)(3\lambda - 5)[(1/2)\lambda(3\lambda - 5) + 1]$,

$$h(1) = 0, \quad (2.21)$$

$$h'(t) = 81(2 - \lambda)\left(\lambda - \frac{2}{3}\right)(\lambda - 1)\left(\lambda - \frac{4}{3}\right)\left(\lambda - \frac{5}{3}\right)(t - 1). \quad (2.22)$$

(1) If $\lambda \in (0, 2/3) \cup (1, 4/3) \cup (5/3, 2)$, then (2.22) implies

$$h'(t) > 0 \quad (2.23)$$

for $t > 1$.

Therefore, Lemma 2.3(1) follows from (2.15), (2.17), (2.19), (2.20), (2.21), and (2.23).

(2) If $\lambda \in (2/3, 1) \cup (4/3, 5/3)$, then (2.22) leads to

$$h'(t) < 0 \quad (2.24)$$

for $t > 1$.

Therefore, Lemma 2.3(2) follows from (2.15), (2.17), (2.19), (2.20), (2.21), and (2.24). $\square$

Lemma 2.4. Let $t > 1$ and $G(t) = (1 - 2/\lambda)t^{3-2/\lambda} - (3 - 2/\lambda)t^{2-2/\lambda} + (3 - 2/\lambda)t + (2/\lambda) - 1$. Then

(1) $G(t) > 0$ for $\lambda \in (2/3, 1)$ and

(2) $G(t) < 0$ for $\lambda \in (0, 2/3) \cup (1, 2)$.

Proof. From the expression of $G(t)$, we clearly see that

$$\lim_{t \rightarrow 1} G(t) = 0, \quad (2.25)$$

$$G'(t) = \left(1 - \frac{2}{\lambda}\right) \left(3 - \frac{2}{\lambda}\right) t^{2-2/\lambda} - 2 \left(3 - \frac{2}{\lambda}\right) \left(1 - \frac{1}{\lambda}\right) t^{1-2/\lambda} + 3 - \frac{2}{\lambda}, \quad (2.26)$$

$$G'(1) = 0, \quad (2.27)$$

$$G''(t) = \frac{6}{\lambda^3} \left(\lambda - \frac{2}{3}\right) (\lambda - 1)(\lambda - 2) t^{-2/\lambda} (t - 1). \quad (2.28)$$

From (2.28), we have

$$G''(t) > 0 \quad (2.29)$$

for $\lambda \in (2/3, 1)$ and $t > 1$, and

$$G''(t) < 0 \quad (2.30)$$

for $\lambda \in (0, 2/3) \cup (1, 2)$ and $t > 1$.

Therefore, Lemma 2.4(1) follows from (2.25) and (2.27) together with (2.29), and Lemma 2.4(2) follows from (2.25) and (2.27) together with (2.30). $\square$

3. Main Result

Theorem 3.1. Let $a, b > 0$ with $a \neq b$ and $\alpha, \beta > 0$ with $\alpha + \beta < 1$. Then

- (1) $L_{6\alpha+3\beta-5}(a, b) = A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) = L_{-2/(2\alpha+\beta)}(a, b)$ for $2\alpha + \beta \in \{2/3, 1\}$;
 (2) $L_{6\alpha+3\beta-5}(a, b) > A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) > L_{-2/(2\alpha+\beta)}(a, b)$ for $2\alpha + \beta \in (0, 2/3) \cup (1, 2)$ and $L_{6\alpha+3\beta-5}(a, b) < A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) < L_{-2/(2\alpha+\beta)}(a, b)$ for $2\alpha + \beta \in (2/3, 1)$, and the parameters $6\alpha+3\beta-5$ and $-2/(2\alpha+\beta)$ cannot be improved in either case.

Proof. (1) We divide the proof into two cases.

Case 1. If $2\alpha + \beta = 2/3$, then simple computations lead to

$$\begin{aligned} L_{6\alpha+3\beta-5}(a, b) &= L_{-2/(2\alpha+\beta)}(a, b) = L_{-3}(a, b) \\ &= \left(\frac{2a^2b^2}{a+b} \right)^{1/3} = 2^{1-(2\alpha+\beta)}(ab)^{1-(2\alpha+\beta)/2}(a+b)^{2\alpha+\beta-1} \\ &= A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b). \end{aligned} \quad (3.1)$$

Case 2. If $2\alpha + \beta = 1$, then we clearly see that

$$\begin{aligned} L_{6\alpha+3\beta-5}(a, b) &= L_{-2/(2\alpha+\beta)}(a, b) = L_{-2}(a, b) \\ &= (ab)^{1/2} = A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b). \end{aligned} \quad (3.2)$$

(2) Without loss of generality, we assume that $a > b$ and we put $t = a/b > 1$ and $\lambda = 2\alpha + \beta$.

Firstly, we compare $A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$ with $L_{3\lambda-5}(a, b)$. We divide the proof into five cases.

Case 1. If $\lambda = 4/3$, then (1.1) leads to

$$\begin{aligned} L_{3\lambda-5}(a, b) - A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) \\ = \frac{a-b}{\log a - \log b} - \left(\frac{a+b}{2} \right)^\alpha (ab)^{\beta/2} \left(\frac{2ab}{a+b} \right)^{1-\alpha-\beta} = b \left[\frac{t-1}{\log t} - \left(\frac{t(t+1)}{2} \right)^{1/3} \right]. \end{aligned} \quad (3.3)$$

Therefore, $L_{3\lambda-5}(a, b) > A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$ follows from (3.3) and Lemma 2.1.

Case 2. If $\lambda = 5/3$, then from (1.1) we have

$$\log[L_{3\lambda-5}(a, b)] - \log[A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)] = \frac{t}{t-1} \log t - \frac{1}{6} \log t - \frac{2}{3} \log \frac{1+t}{2} - 1. \quad (3.4)$$

From (3.4) and Lemma 2.2, we clearly see that $L_{3\lambda-5}(a, b) > A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$.

Case 3. If $\lambda \in (0, 2/3) \cup (1, 4/3) \cup (5/3, 2)$, then (1.1) yields

$$\begin{aligned} & \log[L_{3\lambda-5}(a, b)] - \log[A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)] \\ &= \frac{1}{3\lambda-5} \log \frac{t^{3\lambda-4}-1}{(3\lambda-4)(t-1)} - (\lambda-1) \log \frac{1+t}{2} - \left(1 - \frac{\lambda}{2}\right) \log t. \end{aligned} \quad (3.5)$$

Let $f(t) = (1/(3\lambda-5)) \log((t^{3\lambda-4}-1)/(3\lambda-4)(t-1)) - (\lambda-1) \log((1+t)/2) - (1-\lambda/2) \log t$. Then simple computations lead to

$$\lim_{t \rightarrow 1} f(t) = 0, \quad (3.6)$$

$$f'(t) = \frac{g(t)}{(3\lambda-5)(t^{3\lambda-4}-1)(t^2-1)t}, \quad (3.7)$$

where $g(t) = (1 - (1/2)\lambda)(3\lambda-5)t^{3\lambda-2} + [(\lambda-1)(3\lambda-5)-1]t^{3\lambda-3} - [(1/2)\lambda(3\lambda-5)+1]t^{3\lambda-4} + [(1/2)\lambda(3\lambda-5)+1]t^2 - [(\lambda-1)(3\lambda-5)-1]t - (1 - (1/2)\lambda)(3\lambda-5)$.

We clearly see that

$$(3\lambda-5)(t^{3\lambda-4}-1) > 0. \quad (3.8)$$

Therefore, $L_{3\lambda-5}(a, b) > A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$ follows from (3.5)–(3.8) and Lemma 2.3(1).

Case 4. If $\lambda \in (2/3, 1)$, then (3.5)–(3.8) again hold, and from (3.5)–(3.8) and Lemma 2.3(2), we know that $L_{3\lambda-5}(a, b) < A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$.

Case 5. If $\lambda \in (4/3, 5/3)$, then (3.5)–(3.7) hold and

$$(3\lambda-5)(t^{3\lambda-4}-1) < 0. \quad (3.9)$$

Therefore, $L_{3\lambda-5}(a, b) > A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$ follows from (3.5)–(3.7) and (3.9) together with Lemma 2.3(2).

Secondly, we compare $A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)$ with $L_{-2/\lambda}(a, b)$.

From (1.1), we have

$$\begin{aligned} & \log[L_{-2/\lambda}(a, b)] - \log[A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b)] \\ &= -\frac{\lambda}{2} \log \frac{t^{1-\lambda/2}-1}{(1-\lambda/2)(t-1)} - (\lambda-1) \log \frac{1+t}{2} - \left(1 - \frac{\lambda}{2}\right) \log t. \end{aligned} \quad (3.10)$$

Let $F(t) = -\lambda/2 \log((t^{1-(\lambda/2)} - 1)/(1 - \lambda/2)(t - 1)) - (\lambda - 1) \log((1 + t)/2) - (1 - \lambda/2) \log t$. Then simple computations yield

$$\lim_{t \rightarrow 1} F(t) = 0, \quad (3.11)$$

$$F'(t) = \frac{\lambda G(t)}{2t(1 - t^{1-2/\lambda})(t^2 - 1)}, \quad (3.12)$$

where $G(t) = (1 - 2/\lambda)t^{3-2/\lambda} - (3 - 2/\lambda)t^{2-2/\lambda} + (3 - 2/\lambda)t + (2/\lambda) - 1$.

We clearly see that

$$1 - t^{1-2/\lambda} > 0 \quad (3.13)$$

for $\lambda \in (0, 2)$.

Therefore, $A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) < L_{-2/\lambda}(a, b)$ for $\lambda \in (2/3, 1)$ follows from (3.10)–(3.13) and Lemma 2.4(1), and $A^\alpha(a, b)G^\beta(a, b)H^{1-\alpha-\beta}(a, b) > L_{-2/\lambda}(a, b)$ for $\lambda \in (0, 2/3) \cup (1, 2)$ follows from (3.10)–(3.13) and Lemma 2.4(2).

At last, we prove that the parameters $-2/\lambda$ and $3\lambda - 5$ cannot be improved in either case.

The following two cases will complete the proof for the optimality of parameter $-2/\lambda$.

Case I. If $\lambda \in (0, 2/3) \cup (1, 2)$, then for any $\epsilon \in (0, 2/\lambda - 1)$, one has

$$\begin{aligned} & \lim_{t \rightarrow +\infty} \frac{L_{-2/\lambda+\epsilon}(1, t)}{A^\alpha(1, t)G^\beta(1, t)H^{1-\alpha-\beta}(1, t)} \\ &= \lim_{t \rightarrow +\infty} \left\{ \left[\frac{2/\lambda - 1 - \epsilon}{1 - t^{1+\epsilon-2/\lambda}} \left(1 - \frac{1}{t}\right) \right]^{\lambda/(2-\epsilon\lambda)} \left(\frac{1+t}{2t}\right)^{1-\lambda} t^{\epsilon\lambda^2/2(2-\epsilon\lambda)} \right\} \\ &= +\infty. \end{aligned} \quad (3.14)$$

Equation (3.14) implies that for any $\epsilon \in (0, 2/\lambda - 1)$, there exists a sufficiently large $T_1 = T_1(\epsilon, \alpha, \beta) > 1$, such that $L_{-2/\lambda+\epsilon}(1, t) > A^\alpha(1, t)G^\beta(1, t)H^{1-\alpha-\beta}(1, t)$ for $t \in (T_1, +\infty)$.

Case II. If $\lambda \in (2/3, 1)$, then for any $\epsilon > 0$, one has

$$\begin{aligned} & \lim_{t \rightarrow +\infty} \frac{A^\alpha(1, t)G^\beta(1, t)H^{1-\alpha-\beta}(1, t)}{L_{-2/\lambda-\epsilon}(1, t)} \\ &= \lim_{t \rightarrow +\infty} \left\{ \left(\frac{2t}{1+t}\right)^{1-\lambda} \left[\frac{1 - t^{1-2/\lambda-\epsilon}}{(2/\lambda + \epsilon - 1)(1 - 1/t)} \right]^{\lambda/(2+\epsilon\lambda)} t^{\epsilon\lambda^2/2(2+\epsilon\lambda)} \right\} \\ &= +\infty. \end{aligned} \quad (3.15)$$

Equation (3.15) implies that for any $\epsilon > 0$, there exists a sufficiently large $T_2 = T_2(\epsilon, \alpha, \beta) > 1$, such that $A^\alpha(1, t)G^\beta(1, t)H^{1-\alpha-\beta}(1, t) > L_{-2/\lambda-\epsilon}(1, t)$ for $t \in (T_2, +\infty)$.

The following seven cases will complete the proof for the optimality of parameter $3\lambda - 5$.

Case A. If $\lambda = 4/3$, then for any $\epsilon > 0$ and $x > 0$, one has

$$[L_{3\lambda-5-\epsilon}(1, 1+x)]^{1+\epsilon} - \left[A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x) \right]^{1+\epsilon} = \frac{g_1(x)}{(1+x)^\epsilon - 1}, \quad (3.16)$$

where $g_1(x) = \epsilon x(1+x)^\epsilon - (1+x/2)^{(1+\epsilon)/3}(1+x)^{(1+\epsilon)/3}[(1+x)^\epsilon - 1]$.

Upon letting $x \rightarrow 0$, the Taylor expansion leads to

$$g_1(x) = -\frac{1}{24}\epsilon^2(\epsilon+1)x^3 + o(x^3). \quad (3.17)$$

Equations (3.16) and (3.17) imply that for $\epsilon > 0$, there exists a sufficiently small $\delta_1 = \delta_1(\epsilon) > 0$, such that $L_{3\lambda-5-\epsilon}(1, 1+x) < A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_1)$.

Case B. If $\lambda = 5/3$, then for any $\epsilon \in (0, 1)$ and $x > 0$ ($x \rightarrow 0$), one has

$$\begin{aligned} & [L_{3\lambda-5-\epsilon}(1, 1+x)]^\epsilon - \left[A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x) \right]^\epsilon \\ &= \frac{(1-\epsilon)x}{(1+x)^{1-\epsilon} - 1} - (1+x)^{\epsilon/6} \left(1 + \frac{x}{2} \right)^{(2/3)\epsilon} \\ &= \frac{1}{(1+x)^{1-\epsilon} - 1} \left[-\frac{1}{24}(1-\epsilon)\epsilon^2 x^3 + o(x^3) \right]. \end{aligned} \quad (3.18)$$

Equation (3.18) implies that for any $\epsilon \in (0, 1)$, there exists a sufficiently small $\delta_2 = \delta_2(\epsilon) > 0$, such that $L_{3\lambda-5-\epsilon}(1, 1+x) < A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_2)$.

Case C. If $\lambda \in (0, 2/3)$, then for any $\epsilon > 0$ and $x > 0$ ($x \rightarrow 0$), one has

$$\begin{aligned} & [L_{3\lambda-5-\epsilon}(1, 1+x)]^{5+\epsilon-3\lambda} - \left[A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x) \right]^{5+\epsilon-3\lambda} \\ &= \frac{(4+\epsilon-3\lambda)x(1+x)^{4+\epsilon-3\lambda}}{(1+x)^{4+\epsilon-3\lambda} - 1} - \left[(1+x)^{1-\lambda/2} \left(1 + \left(\frac{x}{2} \right) \right)^{\lambda-1} \right]^{5+\epsilon-3\lambda} \\ &= \frac{1}{(1+x)^{4+\epsilon-3\lambda} - 1} \left[-\frac{\epsilon}{24}(5+\epsilon-3\lambda)(4+\epsilon-3\lambda)x^3 + o(x^3) \right]. \end{aligned} \quad (3.19)$$

Equation (3.19) implies that for any $\epsilon > 0$, there exists a sufficiently small $\delta_3 = \delta_3(\epsilon, \alpha, \beta) > 0$, such that $L_{3\lambda-5-\epsilon}(1, 1+x) < A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_3)$.

Case D. If $\lambda \in (2/3, 1)$, then for any $\epsilon \in (0, 4 - 3\lambda)$ and $x > 0$ ($x \rightarrow 0$), one has

$$\begin{aligned}
 & [L_{3\lambda-5+\epsilon}(1, 1+x)]^{5-3\lambda-\epsilon} - \left[A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x) \right]^{5-3\lambda-\epsilon} \\
 &= \frac{(4-3\lambda-\epsilon)x(1+x)^{4-3\lambda-\epsilon}}{(1+x)^{4-3\lambda-\epsilon}-1} - \left[\frac{(1+x)^{1-\lambda/2}}{(1+x/2)^{1-\lambda}} \right]^{5-3\lambda-\epsilon} \\
 &= \frac{(\epsilon/24)(5-3\lambda-\epsilon)(4-3\lambda-\epsilon)x^3 + o(x^3)}{\left[(1+x)^{4-3\lambda-\epsilon} - 1 \right] (1+x/2)^{(1-\lambda)(5-3\lambda-\epsilon)}}.
 \end{aligned} \tag{3.20}$$

Equation (3.20) implies that for any $\epsilon \in (0, 4 - 3\lambda)$, there exists a sufficiently small $\delta_4 = \delta_4(\epsilon, \alpha, \beta) > 0$, such that $L_{3\lambda-5+\epsilon}(1, 1+x) > A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_4)$.

Case E. If $\lambda \in (1, 4/3)$, then for any $\epsilon > 0$ and $x > 0$ ($x \rightarrow 0$), one has

$$\begin{aligned}
 & [L_{3\lambda-5-\epsilon}(1, 1+x)]^{5+\epsilon-3\lambda} - \left[A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x) \right]^{5+\epsilon-3\lambda} \\
 &= \frac{(4+\epsilon-3\lambda)x(1+x)^{4+\epsilon-3\lambda}}{(1+x)^{4+\epsilon-3\lambda}-1} - \left[(1+x)^{1-\lambda/2} \left(1 + \left(\frac{x}{2} \right) \right)^{\lambda-1} \right]^{5+\epsilon-3\lambda} \\
 &= \frac{1}{(1+x)^{4+\epsilon-3\lambda}-1} \left[-(\epsilon/24)(5+\epsilon-3\lambda)(4+\epsilon-3\lambda)x^3 + o(x^3) \right].
 \end{aligned} \tag{3.21}$$

Equation (3.21) implies that for any $\epsilon > 0$, there exists a sufficiently small $\delta_5 = \delta_5(\epsilon, \alpha, \beta) > 0$, such that $L_{3\lambda-5-\epsilon}(1, 1+x) < A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_5)$.

Case F. If $\lambda \in (4/3, 5/3)$, then for any $\epsilon \in (0, 3\lambda - 4)$ and $x > 0$ ($x \rightarrow 0$), one has

$$\begin{aligned}
 & [L_{3\lambda-5-\epsilon}(1, 1+x)]^{5+\epsilon-3\lambda} - \left[A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x) \right]^{5+\epsilon-3\lambda} \\
 &= \frac{(3\lambda-4-\epsilon)x}{(1+x)^{3\lambda-4-\epsilon}-1} - \left[(1+x)^{1-\lambda/2} \left(1 + \left(\frac{x}{2} \right) \right)^{\lambda-1} \right]^{5+\epsilon-3\lambda} \\
 &= \frac{1}{(1+x)^{3\lambda-4-\epsilon}-1} \left[-\frac{\epsilon}{24}(5+\epsilon-3\lambda)(3\lambda-4-\epsilon)x^3 + o(x^3) \right].
 \end{aligned} \tag{3.22}$$

Equation (3.22) implies that for any $\epsilon \in (0, 3\lambda - 4)$, there exists a sufficiently small $\delta_6 = \delta_6(\epsilon, \alpha, \beta) > 0$, such that $L_{3\lambda-5-\epsilon}(1, 1+x) < A^\alpha(1, 1+x)G^\beta(1, 1+x)H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_6)$.

Case G. If $\lambda \in (5/3, 2)$, then for any $\epsilon \in (0, 3\lambda - 5)$ and $x > 0$ ($x \rightarrow 0$), one has

$$\begin{aligned}
 & [L_{3\lambda-5-\epsilon}(1, 1+x)]^{3\lambda-5-\epsilon} - \left[A^\alpha(1, 1+x) G^\beta(1, 1+x) H^{1-\alpha-\beta}(1, 1+x) \right]^{3\lambda-5-\epsilon} \\
 &= \frac{(1+x)^{3\lambda-4-\epsilon} - 1}{(3\lambda-4-\epsilon)x} - \left[(1+x)^{1-\lambda/2} \left(1 + \left(\frac{x}{2} \right) \right)^{\lambda-1} \right]^{3\lambda-5-\epsilon} \\
 &= \frac{1}{(3\lambda-4-\epsilon)x} \left[-\frac{\epsilon}{24} (3\lambda-4-\epsilon)(3\lambda-5-\epsilon)x^3 + o(x^3) \right].
 \end{aligned} \tag{3.23}$$

Equation (3.23) implies that for any $\epsilon \in (0, 3\lambda - 5)$, there exists a sufficiently small $\delta_7 = \delta_7(\epsilon, \alpha, \beta) > 0$, such that $L_{3\lambda-5+\epsilon}(1, 1+x) < A^\alpha(1, 1+x) G^\beta(1, 1+x) H^{1-\alpha-\beta}(1, 1+x)$ for $x \in (0, \delta_7)$. $\square$

Acknowledgments

The authors wish to thank the anonymous referee for his very careful reading of the manuscript and fruitful comments and suggestions. This research is partly supported by N S Foundation of China (Grant 60850005), N S Foundation of Zhejiang Province (Grants D7080080 and Y607128), and Innovation Team Foundation of the Department of Education of Zhejiang Province (Grant T200924).

References

- [1] K. B. Stolarsky, "The power and generalized logarithmic means," *The American Mathematical Monthly*, vol. 87, no. 7, pp. 545–548, 1980.
- [2] C. E. M. Pearce and J. Pečarić, "Some theorems of Jensen type for generalized logarithmic means," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 40, no. 9-10, pp. 789–795, 1995.
- [3] B. Mond, C. E. M. Pearce, and J. Pečarić, "The logarithmic mean is a mean," *Mathematical Communications*, vol. 2, no. 1, pp. 35–39, 1997.
- [4] Ch.-P. Chen and F. Qi, "Monotonicity properties for generalized logarithmic means," *The Australian Journal of Mathematical Analysis and Applications*, vol. 1, no. 2, article 2, p. 499, 2004.
- [5] A. A. K. Abuhany, S. R. Salem, and I. M. Salman, "On Steffensen's integral inequality with applications," *Journal of Rajasthan Academy of Physical Sciences*, vol. 5, no. 1, pp. 1–12, 2006.
- [6] F. Qi, Sh.-X. Chen, and Ch.-P. Chen, "Monotonicity of ratio between the generalized logarithmic means," *Mathematical Inequalities & Applications*, vol. 10, no. 3, pp. 559–564, 2007.
- [7] H.-N. Shi, "Schur-convex functions related to hadamard-type inequalities," *Journal of Mathematical Inequalities*, vol. 1, no. 1, pp. 127–136, 2007.
- [8] Ch.-P. Chen, "The monotonicity of the ratio between generalized logarithmic means," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 86–89, 2008.
- [9] F. Qi, X.-A. Li, and Sh.-X. Chen, "Refinements, extensions and generalizations of the second Kershaw's double inequality," *Mathematical Inequalities & Applications*, vol. 11, no. 3, pp. 457–465, 2008.
- [10] P. Kahlig and J. Matkowski, "Functional equations involving the logarithmic mean," *Zeitschrift für Angewandte Mathematik und Mechanik*, vol. 76, no. 7, pp. 385–390, 1996.
- [11] A. O. Pittenger, "The logarithmic mean in n variables," *The American Mathematical Monthly*, vol. 92, no. 2, pp. 99–104, 1985.
- [12] N. S. Nadirashvili, "New isoperimetric inequalities in mathematical physics," in *Partial Differential Equations of Elliptic Type (Cortona, 1992)*, Sympos. Math., XXXV, pp. 197–203, Cambridge University Press, Cambridge, UK, 1994.
- [13] G. Pólya and G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Princeton University, Princeton, NJ, USA, 1951.

- [14] H. Alzer and W. Janous, "Solution of problem 8*," *Crux Mathematicorum with Mathematical Mayhem*, vol. 13, pp. 173–178, 1987.
- [15] P. S. Bullen, D. S. Mitrinović, and P. M. Vasić, *Means and Their Inequalities*, vol. 31, D. Reidel, Dordrecht, The Netherlands, 1988.
- [16] W. Janous, "A note on generalized heronian means," *Mathematical Inequalities & Applications*, vol. 4, no. 3, pp. 369–375, 2001.
- [17] B. C. Carlson, "The logarithmic mean," *The American Mathematical Monthly*, vol. 79, pp. 615–618, 1972.
- [18] B. E. Leach and M. C. Sholander, "Extended mean values II," *Journal of Mathematical Analysis and Applications*, vol. 92, no. 1, pp. 207–223, 1983.
- [19] J. Sándor, "A note on some inequalities for means," *Archiv der Mathematik*, vol. 56, no. 5, pp. 471–473, 1991.
- [20] H. Alzer, "Ungleichungen für mittelwerte," *Archiv der Mathematik*, vol. 47, no. 5, pp. 422–426, 1986.
- [21] H. Alzer and S.-L. Qiu, "Inequalities for means in two variables," *Archiv der Mathematik*, vol. 80, no. 2, pp. 201–215, 2003.
- [22] D. S. Mitrinović, *Analytic Inequalities*, Springer, New York, NY, USA, 1970.