Research Article

The Essential Norm of the Generalized Hankel Operators on the Bergman Space of the Unit Ball in C^n

Luo Luo and Yang Xuemei

Department of Mathematics, University of Science and Technology of China, Hefei, Anhui 230026, China

Correspondence should be addressed to Luo Luo, lluo@ustc.edu.cn

Received 26 August 2010; Revised 1 December 2010; Accepted 31 December 2010

Academic Editor: Simeon Reich

Copyright © 2010 L. Luo and Y. Xuemei. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In 1993, Peloso introduced a kind of operators on the Bergman space $A^2(B)$ of the unit ball that generalizes the classical Hankel operator. In this paper, we estimate the essential norm of the generalized Hankel operators on the Bergman space $A^p(B)$ ($p > 1$) of the unit ball and give an equivalent form of the essential norm.

1. Introduction

Let B be the open unit ball in C^n , m the Lebesgue measure on C^n normalized so that $m(B) = 1$, $H(B)$ denotes the class of all holomorphic functions on B . The Bergman space $A^2(B)$ is the Banach space of all holomorphic functions f on B such that $\int_B |f(z)|^2 dm(z) < \infty$. It is easy to show that $A^2(B)$ is a closed subspace of $L^2(B, dm)$.

There is an orthogonal projection of $L^2(B, dm)$ onto $A^2(B)$, denoted by P and

$$Pf(z) = \int_B K(z, w) f(w) dm(w), \quad (1.1)$$

where $K(z, w) = 1/(1 - \langle z, w \rangle)^{n+1}$ is the Bergman kernel on B .

For a function $f \in H(B)$, define the Hankel operator $H_f : A^2(B) \rightarrow A^2(B)^\perp$ with symbol f by

$$H_f g = (I - P)(\overline{f}g) = \int_B \overline{(f(z) - f(w))} K(z, w) g(w) dm(w), \quad (1.2)$$

where I is the identity operator.

Since the Hankel operator H_f is connected with the Toeplitz operator, the commutator, the Bloch space, and the Besov space, it has been extensively studied. Important papers in this context are [1, 2] for the case $n = 1$ and [3–5] for the case $n > 1$. It is known that H_f is bounded on $A^2(B)$ if and only if $f \in \beta(B)$ and H_f is compact $A^2(B)$ if and only if $f \in \beta_0(B)$, where

$$\begin{aligned}\beta(B) &= \left\{ f \in H(B) : \sup_{z \in B} (1 - |z|^2) |Rf(z)| < \infty \right\}, \\ \beta_0(B) &= \left\{ f \in H(B) : (1 - |z|^2) |Rf(z)| \rightarrow 0, \text{ as } |z| \rightarrow 1 \right\}.\end{aligned}\tag{1.3}$$

Rf is the radial derivative of f defined by

$$Rf(z) = \sum_{j=1}^n z_j \frac{\partial f(z)}{\partial z_j}.\tag{1.4}$$

$\beta(B)$ is called the Bloch space, and $\beta_0(B)$ is called the little Bloch space.

For $n = 1$, $f \in H(D)$ (D is the open unit disc), H_f is in the Schatten class S_p ($1 < p < \infty$) if and only if $f \in B_p(D)$; $H_f \in S_p$ ($0 < p \leq 1$) if and only if f is a constant, where

$$B_p(D) = \left\{ f \in H(D) : f'(z)(1 - |z|^2) \in L^p(d\lambda) \right\}, \quad p > 1,\tag{1.5}$$

and $d\lambda(z) = (1 - |z|^2)^{-2} dm(z)$ is the invariant volume measure on D , $B_p(D)$ is called the Besov space on D . This theorem expresses that there is a cutoff of H_f at $p = 1$.

For $n > 1$, $f \in H(B)$, $H_f \in S_p$ ($2n < p < \infty$) if and only if $f \in B_p(B)$, $H_f \in S_p$ ($0 < p \leq 2n$) if and only if f is a constant, where

$$B_p(B) = \left\{ f \in H(B) : (1 - |z|^2) Rf(z) \in L^p(d\lambda) \right\}, \quad p > n,\tag{1.6}$$

and $d\lambda(z) = (1 - |z|^2)^{-(n+1)} dm(z)$ is the invariant volume measure on B . $B_p(B)$ is called the Besov space on B . Then, the cutoff phenomenon of H_f appears at $p = 2n$. If $c(n)$ denotes the value of “cutoff,” then

$$c(n) = \begin{cases} 1, & n = 1, \\ 2n, & n > 1. \end{cases}\tag{1.7}$$

Obviously, $c(n)$ depends on the dimension n of the unit ball.

In 1993, Peloso [3] replaced $f(z) - f(w)$ with

$$\Delta_j f(w, z) = f(w) - \sum_{|\alpha| < j} \frac{D^\alpha f(z)}{\alpha!} (w - z)^\alpha\tag{1.8}$$

to define a kind of generalized Hankel operator:

$$\begin{aligned} H_{f,j}g(z) &= \int_B \overline{-\Delta_j f(w, z)} K(z, w) g(w) dm(w), \\ H'_{f,j}g(z) &= \int_B \overline{\Delta_j f(z, w)} K(z, w) g(w) dm(w). \end{aligned} \quad (1.9)$$

Here, $(D^\alpha f)(z) = (\partial^{| \alpha |} f(z)) / (\partial z_1^{\alpha_1} \cdots \partial z_n^{\alpha_n})$. Clearly, if $j = 1$, $H_{f,1}$ and $H'_{f,1}$ are just the classical Hankel operator H_f . He proved that $H_{f,j}$ has the same boundedness and compactness properties as H_f , but the Schatten class property of $H_{f,j}$ is different from that of H_f . If $n \geq 2$, $f \in H(B)$, $H_{f,j} \in S_p((2n/j) < p < \infty)$ if and only if $f \in B_p(B)$; if $0 < p \leq (2n/j)$, $H_{f,j} \in S_p$ if and only if f is a polynomial of degree at most $j - 1$. So the value of "cutoff" of $H_{f,j}$ is $2n/j$; this means that the cutoff constant $c(n)$ depends not only on the dimension but also on the degree of the polynomial

$$\sum_{|\alpha| < j} \frac{D^\alpha f(z)}{\alpha!} (w - z)^\alpha, \quad (1.10)$$

and we are able to lower the cutoff constant by increasing j .

The cutoff phenomenon expressed that the generalized Hankel operator $H_{f,j}$ defined by Peloso and the classical Hankel operator H_f are different.

In the present paper, we will consider the generalized Hankel operators $H_{f,j}$ defined by Peloso on the Bergman space $A^p(B)$ which is the Banach space of all holomorphic functions f on B such that $\int_B |f(z)|^p dm(z) < \infty$, for $p > 1$.

For $f(z) \in H(B)$, j is a positive integer, and we define the generalized Hankel operators $H_{f,j}$ and $H'_{f,j}$ of order j with symbol f by

$$\begin{aligned} H_{f,j}g(z) &= \int_B \overline{-\Delta_j f(w, z)} K(z, w) g(w) dm(w), \\ H'_{f,j}g(z) &= \int_B \overline{\Delta_j f(z, w)} K(z, w) g(w) dm(w), \end{aligned} \quad (1.11)$$

where $g \in A^p(B)$,

$$\begin{aligned} \Delta_j f(w, z) &= f(w) - \sum_{|\alpha| < j} \frac{D^\alpha f(z)}{\alpha!} (w - z)^\alpha, \\ \Delta_j f(z, w) &= f(w) - \sum_{|\alpha| < j} \frac{D^\alpha f(w)}{\alpha!} (z - w)^\alpha. \end{aligned} \quad (1.12)$$

Luo and Ji-Huai [6] studied the boundedness, compactness, and the Schatten class property of the generalized Hankel operator $H_{f,j}$ on the Bergman space $A^p(B)$ ($p > 1$), which extended the known results.

We will study the essential norm of this kind of generalized Hankel operators $H_{f,j}$ and $H'_{f,j}$. We recall that the essential norm of a bounded linear operator T is the distance from T to the compact operators; that is,

$$\|T\|_{\text{ess}} = \inf\{\|T - K\| : K \text{ is a compact operator}\}. \quad (1.13)$$

The essential norm of a bounded linear operator T is connected with the compactness of the operator T and the spectrum of the operator T .

We know that $\|T\|_{\text{ess}} = 0$ if and only if T is compact, so that estimates on $\|T\|_{\text{ess}}$ lead to conditions for T to be compact. Thus, we will obtain a different proof of the compactness of the generalized Hankel operators $H_{f,j}$ and $H'_{f,j}$.

Throughout the paper, C denotes a positive constant, whose value may change from one occurrence to the next one.

2. Preliminaries

For any fixed point $a \in B - \{0\}$, $z \in B$, define the Möbius transformation φ_a by

$$\varphi_a(z) = \frac{a - P_a(z) - s_a Q_a(z)}{1 - \langle z, a \rangle}, \quad (2.1)$$

where $s_a = \sqrt{1 - |a|^2}$ and P_a is the orthogonal projection from C^n onto the one-dimensional subspace $[a]$ generated by a , Q_a is the orthogonal projection from C^n onto $C^n \ominus [a]$. It is clear that

$$\begin{aligned} P_a(z) &= \frac{\langle z, a \rangle}{|a|^2} a, \quad z \in C^n, \\ Q_a(z) &= z - \frac{\langle z, a \rangle}{|a|^2} a, \quad z \in B. \end{aligned} \quad (2.2)$$

Lemma 2.1. *For every $a \in B$, φ_a has the following properties:*

- (1) $\varphi_a(0) = a$ and $\varphi_a(a) = 0$,
- (2) $\varphi_a \circ \varphi_a(z) = z$, $z \in B$,
- (3) $1/(1 - \langle \varphi_a(z), a \rangle) = (1 - \langle z, a \rangle)/(1 - |a|^2)$, $z \in B$.

Proof. The proofs can be found in [7]. □

Lemma 2.2. *For $s > -1$, t real, define*

$$I_t(z) = \int_B \frac{(1 - |w|^2)^s}{|1 - \langle z, w \rangle|^{n+1+s+t}} dm(w), \quad z \in B. \quad (2.3)$$

Then,

- (1) $t < 0$, $I_t(z)$ is bounded in B ,
- (2) $t = 0$, $I_t(z) \sim \log(1/(1 - |z|^2))$ as $|z| \rightarrow 1^-$,
- (3) $t > 0$, $I_t(z) \sim (1 - |z|^2)^{-t}$ as $|z| \rightarrow 1^-$.

Here, the notation $a(z) \sim b(z)$ means that the ratio $a(z)/b(z)$ has a positive finite limit as $|z| \rightarrow 1^-$.

Proof. This is in [7, Theorem 1.12]. $\square$

Lemma 2.3. Let $k_\xi(z) = K(z, \xi)/\|K_\xi\|_{L^p(dm)}$, where $K_\xi(z) = K(z, \xi) = 1/(1 - \langle z, \xi \rangle)^{n+1}$, then $k_\xi(z)$ has the following properties:

- (1) $\|k_\xi\|_{L^p(dm)} = 1$,
- (2) $k_\xi(z) \rightarrow 0$ at every point $z \in B$ as $|\xi| \rightarrow 1^-$.

Proof. It is obvious. $\square$

Lemma 2.4. Let $K_\xi(z) = K(z, \xi)$. Then, for any positive integer j ,

- (1) $H_{f,j}K_\xi = \overline{-\Delta_j f(\xi, \cdot)}K_\xi$,
- (2) $H'_{f,j}K_\xi = \overline{\Delta_j f(\cdot, \xi)}K_\xi$.

Proof. The proof is obtained by the definition of $H_{f,j}$ and $H'_{f,j}$ and the reproducing property of $K(z, \xi)$, through the direct computation to get them. $\square$

Lemma 2.5. Let j be any positive integer, $f \in H(B)$, and $0 < q < \infty$, then there is a constant C independent of f , such that

- (1) $(1 - |z|^2)^j |R^j f(z)| \leq C \{ \int_B |\Delta_j f(\varphi_z(w), z)|^q dm(w) \}^{1/q}$,
- (2) $(1 - |z|^2)^j |R^j f(z)| \leq C \{ \int_B |\Delta_j f(z, \varphi_z(w))|^q dm(w) \}^{1/q}$,

where $R^j f$ is the j th order radial derivative of f ,

$$R^j f(z) = \sum_{k=1}^{\infty} k^j f_k(z), \quad (2.4)$$

and $f(z) = \sum_{k=0}^{\infty} f_k(z)$ is the homogeneous expansion.

Proof. This is in [3, Proposition 3.2]. $\square$

Lemma 2.6. Let j be any positive integer, $f \in H(B)$, and $0 < \rho < 1$, $p > 1$, then

- (1) $\int_B |\Delta_j f(w, z)|^p ((1 - |w|^2)^{-\rho} / |1 - \langle z, w \rangle|^{n+1}) dm(w) \leq C(1 - |z|^2)^{-\rho} (\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)|)^p$,
- (2) $\int_B |\Delta_j f(z, w)|^p ((1 - |w|^2)^{-\rho} / |1 - \langle z, w \rangle|^{n+1}) dm(w) \leq C(1 - |z|^2)^{-\rho} (\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)|)^p$.

Proof. (1) Write $F(w, z)$ for $\Delta_j f(w, z)$. Using the change of variables $w = \varphi_z(\xi)$, we obtain

$$\begin{aligned} & \int_B |F(w, z)|^p \frac{(1 - |w|^2)^{-\rho}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) \\ &= \int_B |F(\varphi_z(\xi), z)|^p \frac{(1 - |\varphi_z(\xi)|^2)^{-\rho}}{|1 - \langle z, \varphi_z(\xi) \rangle|^{n+1}} \frac{(1 - |z|^2)^{n+1}}{|1 - \langle \xi, z \rangle|^{2(n+1)}} dm(\xi) \\ &= (1 - |z|^2)^{-\rho} \int_B |F(\varphi_z(\xi), z)|^p \frac{(1 - |\xi|^2)^{-\rho}}{|1 - \langle \xi, z \rangle|^{n+1-2\rho}} dm(\xi). \end{aligned} \quad (*)$$

Let

$$1 < q' < \min\left(\frac{1}{\rho}, \frac{n+1}{n+1-\rho}\right) \quad (2.5)$$

and set $q = q' / (q' - 1)$. Then, applying Hölder's inequality to $(*)$, we obtain

$$\begin{aligned} & \int_B |F(w, z)|^p \frac{(1 - |w|^2)^{-\rho}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) \\ & \leq (1 - |z|^2)^{-\rho} \left(\int_B |F(\varphi_z(\xi), z)|^{pq} dm(\xi) \right)^{1/q} \left(\int_B \frac{(1 - |\xi|^2)^{-\rho q'}}{|1 - \langle \xi, z \rangle|^{(n+1-2\rho)q'}} dm(\xi) \right)^{1/q'}. \end{aligned} \quad (2.6)$$

Because of our choice of q' , it follows that $-\rho q' > -1$ and $(n+1-2\rho)q' < n+1-\rho q'$. Now, Lemma 2.2 implies that

$$\int_B \frac{(1 - |\xi|^2)^{-\rho q'}}{|1 - \langle \xi, z \rangle|^{(n+1-2\rho)q'}} dm(\xi) \quad (2.7)$$

is bounded by a constant. Therefore, applying [3, Theorem 3.4], we get

$$\int_B |\Delta_j f(w, z)|^p \frac{(1 - |w|^2)^{-\rho}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) \leq C (1 - |z|^2)^{-\rho} \left(\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)| \right)^p. \quad (2.8)$$

(2) The proof of (2) is similar to that of (1). □

3. The Main Result and Its Proof

Theorem 3.1. *Let $f \in H(B)$, j any positive integer, $p > 1$, and the generalized Hankel operators $H_{f,j}$, $H'_{f,j}$ defined on $A^p(B)$ by*

$$\begin{aligned} H_{f,j}g(z) &= \int_B \overline{-\Delta_j f(w, z)} K(z, w) g(w) dm(w), \\ H'_{f,j}g(z) &= \int_B \overline{\Delta_j f(z, w)} K(z, w) g(w) dm(w). \end{aligned} \quad (3.1)$$

Suppose that $H_{f,j}$ and $H'_{f,j}$ are bounded on $A^p(B)$, then the following quantities are equivalent:

- (1) $\|H_{f,j}\|_{\text{ess}}$ and $\|H'_{f,j}\|_{\text{ess}}$,
- (2) $\overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2)^j |R^j f(z)|$,
- (3) $\overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2) |Rf(z)|$.

Particularly, $H_{f,j}$ and $H'_{f,j}$ are compact on $A^p(B)$ if and only if $\overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2) |Rf(z)| = 0$.

Proof. First, we will prove that $\|H_{f,j}\|_{\text{ess}} \geq C \overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2)^j |R^j f(z)|$. By the definition of $k_\xi(z)$ of Lemmas 2.3 and 2.4, we have

$$\begin{aligned} \|H_{f,j}k_\xi\|_{L^p(dm)}^p &= \int_B |H_{f,j}k_\xi(z)|^p dm(z) \\ &= \int_B \left| H_{f,j} \frac{K(z, \xi)}{\|K_\xi\|_{L^p(dm)}} \right|^p dm(z) \\ &= \frac{1}{\|K_\xi\|_{L^p(dm)}^p} \int_B |H_{f,j}K(z, \xi)|^p dm(z) \\ &= \frac{1}{\|K_\xi\|_{L^p(dm)}^p} \int_B |\Delta_j f(\xi, z)|^p |K(z, \xi)|^p dm(z) \\ &= \frac{1}{\|K_\xi\|_{L^p(dm)}^p} \cdot I, \end{aligned} \quad (3.2)$$

here $I = \int_B |\Delta_j f(\xi, z)|^p |K(z, \xi)|^p dm(z)$.

Use the change of variables $z = \varphi_\xi(\tau)$ in the integral I , and recall that

$$dm(z) = \left(\frac{1 - |\xi|^2}{|1 - \langle \tau, \xi \rangle|^2} \right)^{n+1} dm(\tau). \quad (3.3)$$

Thus

$$\begin{aligned}
I &= \int_B \frac{|\Delta_j f(\xi, \varphi_\xi(\tau))|^p}{|1 - \langle \varphi_\xi(\tau), \xi \rangle|^{p(n+1)}} \cdot \left(\frac{1 - |\xi|^2}{|1 - \langle \tau, \xi \rangle|^2} \right)^{n+1} dm(\tau) \\
&= \frac{1}{(1 - |\xi|^2)^{(n+1)(p-1)}} \int_B \frac{|\Delta_j f(\xi, \varphi_\xi(\tau))|^p}{|1 - \langle \tau, \xi \rangle|^{(2-p)(n+1)}} dm(\tau) \\
&\geq \frac{1}{(1 - |\xi|^2)^{(n+1)(p-1)}} \left(\int_B |\Delta_j f(\xi, \varphi_\xi(\tau))| dm(\tau) \right)^p \\
&\quad \times \left(\int_B \frac{1}{|1 - \langle \tau, \xi \rangle|^{(2-p)(n+1)/(1-p)}} dm(\tau) \right)^{1-p} \\
&\geq \frac{C}{(1 - |\xi|^2)^{(n+1)(p-1)}} \left(\int_B |\Delta_j f(\xi, \varphi_\xi(\tau))| dm(\tau) \right)^p \\
&\geq \frac{C}{(1 - |\xi|^2)^{(n+1)(p-1)}} \left[(1 - |\xi|^2)^j |R^j f(\xi)| \right]^p.
\end{aligned} \tag{3.4}$$

Here, we have used (3) of Lemma 2.1, Hölder's inequality for the indexes p and $p/(p-1)$, (1) of Lemma 2.2, and (2) of Lemma 2.5.

Therefore,

$$\begin{aligned}
\|H_{f,j} k_\xi\|_{L^p(dm)}^p &\geq \frac{1}{\|K_\xi\|_{L^p(dm)}^p} \frac{C}{(1 - |\xi|^2)^{(n+1)(p-1)}} \left[(1 - |\xi|^2)^j |R^j f(\xi)| \right]^p \\
&\geq C (1 - |\xi|^2)^{(n+1)(p-1)} \frac{1}{(1 - |\xi|^2)^{(n+1)(p-1)}} \left[(1 - |\xi|^2)^j |R^j f(\xi)| \right]^p \\
&= C \left[(1 - |\xi|^2)^j |R^j f(\xi)| \right]^p.
\end{aligned} \tag{3.5}$$

So $\|H_{f,j} k_\xi\|_{L^p(dm)} \geq C(1 - |\xi|^2)^j |R^j f(\xi)|$.

For any compact operator T , by (2) of Lemma 2.3, we have $\|Tk_\xi\|_{L^p(dm)} \rightarrow 0$ as $|\xi| \rightarrow 1^-$. Then,

$$\begin{aligned}
\|H_{f,j} - T\| &\geq \overline{\lim}_{|\xi| \rightarrow 1^-} \|(H_{f,j} - T)k_\xi\|_{L^p(dm)} \\
&\geq \overline{\lim}_{|\xi| \rightarrow 1^-} \left(\|H_{f,j}k_\xi\|_{L^p(dm)} - \|Tk_\xi\|_{L^p(dm)} \right) \\
&= \overline{\lim}_{|\xi| \rightarrow 1^-} \|H_{f,j}k_\xi\|_{L^p(dm)} \\
&\geq C \overline{\lim}_{|\xi| \rightarrow 1^-} (1 - |\xi|^2)^j |R^j f(\xi)|.
\end{aligned} \tag{3.6}$$

Thus, $\|H_{f,j}\|_{\text{ess}} \geq C \overline{\lim}_{|z| \rightarrow 1^-} (1 - |z|^2)^j |R^j f(z)|$.

Now, we will prove that $\|H_{f,j}\|_{\text{ess}} \leq C \overline{\lim}_{|z| \rightarrow 1^-} (1 - |z|^2)^j |R^j f(z)|$.

Write $F(z, w)$ for $-\Delta_j f(z, w)$. For $0 < \rho < 1$ and $g \in A^p(B)$, let $B(0, \rho)$ and $B(0; \rho, 1)$ denote the ball $|z| \leq \rho$ and the ring $\rho < |z| < 1$, respectively, then we have

$$\begin{aligned}
H_{f,j}g(z) &= \chi_{B(0,\rho)}(z) \int_B \overline{F(w, z)} K(z, w) g(w) dm(w) \\
&\quad + \chi_{B(0;\rho,1)}(z) \int_B \overline{F(w, z)} K(z, w) g(w) dm(w) \\
&= T_1 g(z) + T_2 g(z).
\end{aligned} \tag{3.7}$$

Here,

$$\begin{aligned}
T_1 g(z) &= \chi_{B(0,\rho)}(z) \int_B \overline{F(w, z)} K(z, w) g(w) dm(w), \\
T_2 g(z) &= \chi_{B(0;\rho,1)}(z) \int_B \overline{F(w, z)} K(z, w) g(w) dm(w).
\end{aligned} \tag{3.8}$$

We first show that T_1 is compact. Let $\{g_l\}$ be a sequence weakly converging to 0 and $p' = p/(p-1)$, by Hölder's inequality, then we have

$$\begin{aligned}
|T_1 g_l(z)|^p &= \left| \chi_{B(0,\rho)}(z) \int_B \overline{F(w, z)} K(z, w) g_l(w) dm(w) \right|^p \\
&\leq \chi_{B(0,\rho)}(z) \left(\int_B |F(w, z)| \frac{|g_l(w)|}{|1 - \langle z, w \rangle|^{n+1}} dm(w) \right)^p \\
&\leq \chi_{B(0,\rho)}(z) \left(\int_B \frac{|F(w, z)|^{p'} (1 - |w|^2)^{-1/p}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) \right)^{p/p'} \\
&\quad \times \int_B \frac{|g_l(w)|^p (1 - |w|^2)^{1/p'}}{|1 - \langle z, w \rangle|^{n+1}} dm(w).
\end{aligned} \tag{3.9}$$

By Lemma 2.6, we get

$$\begin{aligned}
 |T_1 g_l(z)|^p &\leq C \chi_{B(0,\rho)}(z) \left(\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)| \right)^p \\
 &\quad \times (1 - |z|^2)^{-1/p'} \int_B \frac{|g_l(w)|^p (1 - |w|^2)^{1/p'}}{|1 - \langle z, w \rangle|^{n+1}} dm(w).
 \end{aligned} \tag{3.10}$$

Thus,

$$\begin{aligned}
 \|T_1 g_l\|_{L^p(dm)}^p &= \int_B |T_1 g_l(z)|^p dm(z) \\
 &\leq C \int_{|z| < \rho} \left(\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)| \right)^p (1 - |z|^2)^{-1/p'} \\
 &\quad \times \int_B \frac{|g_l(w)|^p (1 - |w|^2)^{1/p'}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) dm(z) \\
 &\leq C (1 - |\rho|^2)^{-1/p'} \left(\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)| \right)^p \\
 &\quad \times \int \int_B \frac{|g_l(w)|^p (1 - |w|^2)^{1/p'}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) dm(z) \\
 &= C (1 - |\rho|^2)^{-1/p'} \left(\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)| \right)^p \\
 &\quad \cdot \int_B |g_l(w)|^p (1 - |w|^2)^{1/p'} \int_B \frac{1}{|1 - \langle z, w \rangle|^{n+1}} dm(z) dm(w) \\
 &\leq C (1 - |\rho|^2)^{-1/p'} \left(\sup_{z \in B} (1 - |z|^2)^j |R^j f(z)| \right)^p \\
 &\quad \times \int_B |g_l(w)|^p (1 - |w|^2)^{1/p'} \log(1 - |w|^2) dm(w) \\
 &\longrightarrow 0, \quad \text{as } l \longrightarrow \infty.
 \end{aligned} \tag{3.11}$$

So, T_1 is compact.

For $g \in A^p$ and $p' = p/(p-1)$, by Hölder's inequality,

$$\begin{aligned}
 |T_2 g(z)|^p &= \left| \chi_{B(0,\rho,1)}(z) \int_B \overline{F(w, z)} K(z, w) g(w) dm(w) \right|^p \\
 &\leq \left(\int_B \chi_{B(0,\rho,1)}(z) |F(w, z)| \frac{|g(w)|}{|1 - \langle z, w \rangle|^{n+1}} dm(w) \right)^p
 \end{aligned}$$

$$\begin{aligned}
&\leq \left(\int_B \chi_{B(0;\rho,1)}(z) \frac{|F(w,z)|^{p'} (1-|w|^2)^{-1/p}}{|1-\langle z,w \rangle|^{n+1}} dm(w) \right)^{p/p'} \\
&\quad \times \int_B \frac{|g(w)|^p (1-|w|^2)^{1/p'}}{|1-\langle z,w \rangle|^{n+1}} dm(w).
\end{aligned} \tag{3.12}$$

So

$$\begin{aligned}
\|T_2 g\|_{L^p(dm)}^p &= \int_B |T_2 g(z)|^p dm(z) \leq \int_B \left(\int_B \chi_{B(0;\rho,1)}(z) \frac{|F(w,z)|^{p'} (1-|w|^2)^{-1/p}}{|1-\langle z,w \rangle|^{n+1}} dm(w) \right)^{p/p'} \\
&\quad \cdot \int_B \frac{|g(w)|^p (1-|w|^2)^{1/p'}}{|1-\langle z,w \rangle|^{n+1}} dm(w) dm(z).
\end{aligned} \tag{3.13}$$

Change the variables $w = \varphi_z(\xi)$, let

$$1 < q' < \min\left(p, \frac{n+1}{n+1-1/p}\right), \tag{3.14}$$

and set $q = q'/(q' - 1)$, by Lemmas 2.1 and 2.2, then we obtain

$$\begin{aligned}
&\int_B \chi_{B(0;\rho,1)}(z) \frac{|F(w,z)|^{p'} (1-|w|^2)^{-1/p}}{|1-\langle z,w \rangle|^{n+1}} dm(w) \\
&= \int_B \chi_{B(0;\rho,1)}(z) \frac{|F(\varphi_z(\xi), z)|^{p'} (1-|\varphi_z(\xi)|^2)^{-1/p}}{|1-\langle z, \varphi_z(\xi) \rangle|^{n+1}} \frac{(1-|z|^2)^{n+1}}{|1-\langle \xi, z \rangle|^{2(n+1)}} dm(\xi) \\
&= \int_B \chi_{B(0;\rho,1)}(z) (1-|z|^2)^{-1/p} |F(\varphi_z(\xi), z)|^{p'} \frac{(1-|\xi|^2)^{-1/p}}{|1-\langle \xi, z \rangle|^{n+1-2/p}} dm(\xi) \\
&\leq (1-|z|^2)^{-1/p} \left(\int_B \chi_{B(0;\rho,1)}(z) |F(\varphi_z(\xi), z)|^{p'q} dm(\xi) \right)^{1/q} \\
&\quad \times \left(\int_B \frac{(1-|\xi|^2)^{-q'/p}}{|1-\langle \xi, z \rangle|^{(n+1-2/p)q}} dm(\xi) \right)^{1/q'} \\
&\leq C(1-|z|^2)^{-1/p} \left(\int_B \chi_{B(0;\rho,1)}(z) |F(\varphi_z(\xi), z)|^{p'q} dm(\xi) \right)^{1/q}.
\end{aligned} \tag{3.15}$$

By the same argument of [3, Theorem 3.4], we know that

$$\left(\int_B \chi_{B(0;\rho,1)}(z) |F(\varphi_z(\xi), z)|^{p'q} dm(\xi) \right)^{1/q} \leq C \left(\sup_{\rho < |z| < 1} (1 - |z|^2)^j |R^j f(z)| \right)^{p'}. \quad (3.16)$$

Applying Fubini's theorem and Lemma 2.2, we have

$$\begin{aligned} \|T_2 g\|_{L^p(dm)}^p &\leq C \left(\sup_{\rho < |z| < 1} (1 - |z|^2)^j |R^j f(z)| \right)^p \\ &\quad \times \int_B (1 - |z|^2)^{-1/p'} \int_B \frac{|g(w)|^p (1 - |w|^2)^{1/p'}}{|1 - \langle z, w \rangle|^{n+1}} dm(w) dm(z) \\ &\leq C \left(\sup_{\rho < |z| < 1} (1 - |z|^2)^j |R^j f(z)| \right)^p \|g\|_{L^p(dm)}^p. \end{aligned} \quad (3.17)$$

So

$$\|T_2\| \leq C \sup_{\rho < |z| < 1} (1 - |z|^2)^j |R^j f(z)|. \quad (3.18)$$

Thus, by the definition of the essential norm, we have

$$\|H_{f,j}\|_{\text{ess}} \leq \|T_1 + T_2\|_{\text{ess}} \leq \|T_2\| \leq C \sup_{\rho < |z| < 1} (1 - |z|^2)^j |R^j f(z)|. \quad (3.19)$$

As $\rho \rightarrow 1$, we have

$$\|H_{f,j}\|_{\text{ess}} \leq C \overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2)^j |R^j f(z)|. \quad (3.20)$$

Similarly, we get the equality of $\|H'_{f,j}\|_{\text{ess}}$ and $\overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2)^j |R^j f(z)|$.

By [7, Theorems 3.4 and 3.5], we obtain the equality of $\overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2)^j |R^j f(z)|$ and $\overline{\lim_{|z| \rightarrow 1^-}} (1 - |z|^2) |Rf(z)|$.

We complete the proof of Theorem 3.1. $\square$

Acknowledgments

The authors would like to thank the referee for the careful reading of the first version of this paper and for the several suggestions made for improvement. The paper was supported by the NNSF of China (no. 10771201) and the Natural Science Foundation of Anhui Province (no. 090416233).

References

- [1] J. Arazy, S. D. Fisher, and J. Peetre, "Hankel operators on weighted Bergman spaces," *American Journal of Mathematics*, vol. 110, no. 6, pp. 989–1054, 1988.
- [2] S. Axler, "The Bergman space, the Bloch space, and commutators of multiplication operators," *Duke Mathematical Journal*, vol. 53, no. 2, pp. 315–332, 1986.
- [3] M. M. Peloso, "Besov spaces, mean oscillation, and generalized Hankel operators," *Pacific Journal of Mathematics*, vol. 161, no. 1, pp. 155–184, 1993.
- [4] K. Stroethoff, "Compact Hankel operators on the Bergman spaces of the unit ball and polydisk in C^n ," *Journal of Operator Theory*, vol. 23, no. 1, pp. 153–170, 1990.
- [5] K. H. Zhu, "Hilbert-Schmidt Hankel operators on the Bergman space," *Proceedings of the American Mathematical Society*, vol. 109, no. 3, pp. 721–730, 1990.
- [6] L. Luo and S. Ji-Huai, "Generalized Hankel operators on the Bergman space of the unit ball," *Complex Variables. Theory and Application*, vol. 44, no. 1, pp. 55–71, 2001.
- [7] K. Zhu, *Spaces of Holomorphic Functions in the Unit ball*, vol. 226, Springer, New York, NY, USA, 2005.