Research Article

Existence and Nonexistence Results for Classes of Singular Elliptic Problem

Peng Zhang and Jia-Feng Liao

Department of Mathematics, Zunyi Normal College, Zunyi 563002, China

Correspondence should be addressed to Peng Zhang, gzzypd@sina.com

Received 14 January 2010; Accepted 19 July 2010

Academic Editor: Norimichi Hirano

Copyright © 2010 P. Zhang and J.-F. Liao. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The singular semilinear elliptic problem $-\Delta u + k(x)u^{-\gamma} = \lambda u^p$ in Ω , $u > 0$ in Ω , $u = 0$ on $\partial\Omega$, is considered, where Ω is a bounded domain with smooth boundary in R^N , $k \in C_{\text{loc}}^\alpha(\Omega) \cap C(\overline{\Omega})$, and γ, p, λ are three positive constants. Some existence or nonexistence results are obtained for solutions of this problem by the sub-supersolution method.

1. Introduction and Main Results

In this paper, we study the existence or the nonexistence of solutions to the following singular semilinear elliptic problem

$$\begin{aligned} -\Delta u + k(x)u^{-\gamma} &= \lambda u^p, & \text{in } \Omega, \\ u &> 0, & \text{in } \Omega, \\ u &= 0, & \text{on } \partial\Omega, \end{aligned} \tag{1.1}$$

where $\Omega \subset R^N$ ($N \geq 1$) is a bounded domain with $C^{2+\alpha}$ boundary for some $\alpha \in (0, 1)$, $k \in C_{\text{loc}}^\alpha(\Omega) \cap C(\overline{\Omega})$, and γ, p , and λ are three nonnegative constants. This problem arises in the study of non-Newtonian fluids, chemical heterogeneous catalysts, in the theory of heat conduction in electrically conducting materials (see [1–7] and their references).

Many authors have considered this problem. For examples, when $k(x) < 0$ in Ω , problem (1.1) was studied in [3, 8–11]; when $k(x) > 0$ in Ω , problem (1.1) was considered in [12–14]. Particularly, when $k(x) \equiv 1$, it has been established in Zhang [14] that there exists $\bar{\lambda} > 0$ such that problem (1.1) has at least one solution in $C^{2+\alpha}(\Omega) \cap C(\overline{\Omega})$ for all $\lambda > \bar{\lambda}$ and

has no solution in $C^2(\Omega) \cap C(\overline{\Omega})$ if $\lambda < \bar{\lambda}$. After that Shi and Yao in [13] have also obtained the same results with $k \in C^{2,\alpha}(\overline{\Omega})$ and $k(x) > 0$ in $\overline{\Omega}$. Recently, Ghergu and Rădulescu in [12] considered more general sublinear singular elliptic problem with $k \in C^\alpha(\overline{\Omega})$.

In this paper, we consider the case that $k \in C_{\text{loc}}^\alpha(\Omega) \cap C(\overline{\Omega})$, and k may have zeros in $\overline{\Omega}$. The following main results are obtained by the sub-supersolution method with restriction on the boundary in Cui [15].

Theorem 1.1. *Suppose that $k \in C_{\text{loc}}^\alpha(\Omega) \cap C(\overline{\Omega})$, $k \geq 0$, and $k \neq 0$. Assume that $0 < \gamma < 1$ and $0 < p < 1$, then there exists $\bar{\lambda} \in (0, \infty)$ such that problem (1.1) has at least one solution $u_\lambda \in C^{2+\alpha}(\Omega) \cap C(\overline{\Omega})$ and $u_\lambda^{-\gamma} \in L^1(\Omega)$ for all $\lambda > \bar{\lambda}$, and problem (1.1) has no solution in $C^2(\Omega) \cap C(\overline{\Omega})$ if $\lambda < \bar{\lambda}$. Moreover, problem (1.1) has a maximal solution v_λ which is increasing with respect to λ for all $\lambda > \bar{\lambda}$.*

Remark 1.2. Theorem 1.1 generalizes Theorem 1.2 in [13] in coefficient $k(x)$ of the singular term. Consequently, it also generalizes Theorem 1 in [14]. Moreover, there are functions k satisfying our Theorem 1.1 and not satisfying Theorem 1.2 in [13]. For example, let

$$k(x) = \begin{cases} -\frac{1}{\ln(|x - x_0|/(2d))}, & x \in \overline{\Omega} \setminus \{x_0\}, \\ 0, & x = x_0, \end{cases} \quad (1.2)$$

where $x_0 \in \partial\Omega$, and

$$d = \text{diam}(\Omega) \triangleq \max\{|x - y| \mid x, y \in \overline{\Omega}\}. \quad (1.3)$$

Certainly, this example does not satisfy Theorem 1.2 in [12] yet.

Theorem 1.3. *Suppose that $k \in C_{\text{loc}}^\alpha(\Omega) \cap C(\overline{\Omega})$ and $k(x) > 0$ in $\overline{\Omega}$. If $\gamma \geq 1$, problem (1.1) has no solution in $C^2(\Omega) \cap C(\overline{\Omega})$ for all $\lambda > 0$ and $p > 0$.*

Remark 1.4. Obviously, Theorem 1.3 is a generalization of Theorem 2 in [14]. There are also functions $k(x)$ satisfying our Theorem 1.3 and not satisfying Theorem 2 in [14] and Theorem 1.1 in [12]. For example, let

$$k(x) = \begin{cases} -\frac{1}{\ln(|x - x_0|/(2d))} + \varepsilon, & x \in \overline{\Omega} \setminus \{x_0\}, \\ \varepsilon, & x = x_0, \end{cases} \quad (1.4)$$

where $x_0 \in \partial\Omega$, ε is any positive constant and $d = \text{diam}(\Omega)$ is the diameter of Ω .

2. Proof of Theorems

Consider the more general semilinear elliptic problem

$$\begin{aligned} -\Delta u &= f(x, u), \quad \text{in } \Omega, \\ u &> 0, \quad \text{in } \Omega, \\ u &= 0, \quad \text{on } \partial\Omega, \end{aligned} \tag{2.1}$$

where the function $f(x, s)$ is locally Hölder continuous in $\Omega \times (0, \infty)$ and continuously differentiable with respect to the variable s . A function $\underline{u}$ is called to be a subsolution of problem (2.1) if $\underline{u} \in C^2(\Omega) \cap C(\overline{\Omega})$, and

$$\begin{aligned} -\Delta \underline{u} &\leq f(x, \underline{u}), \quad \text{in } \Omega, \\ \underline{u} &> 0, \quad \text{in } \Omega, \\ \underline{u} &= 0, \quad \text{on } \partial\Omega. \end{aligned} \tag{2.2}$$

A function $\overline{u}$ is called to be a supersolution of problem (2.1) if $\overline{u} \in C^2(\Omega) \cap C(\overline{\Omega})$, and

$$\begin{aligned} -\Delta \overline{u} &\geq f(x, \overline{u}), \quad \text{in } \Omega, \\ \overline{u} &> 0, \quad \text{in } \Omega, \\ \overline{u} &= 0, \quad \text{on } \partial\Omega. \end{aligned} \tag{2.3}$$

According to Lemma 3 in the study of Cui [15], we can easily have the following basic existence of classical solution to problem (2.1).

Lemma 2.1. *Let $f \in C_{\text{loc}}^\alpha(\Omega \times (0, \infty))$ be continuously differentiable with respect to the variable s . Suppose that problem (2.1) has a supersolution $\overline{u}$ and a subsolution $\underline{u}$ such that*

$$\underline{u}(x) \leq \overline{u}(x), \quad \text{in } \Omega, \tag{2.4}$$

then problem (2.1) has at least one solution $u \in C^{2+\alpha}(\Omega) \cap C(\overline{\Omega})$ satisfying

$$\underline{u}(x) \leq u(x) \leq \overline{u}(x), \quad \text{in } \overline{\Omega}. \tag{2.5}$$

Let λ_1 be the first eigenvalue of the eigenvalue problem

$$\begin{aligned} -\Delta u &= \lambda u, \quad \text{in } \Omega, \\ u &= 0, \quad \text{on } \partial\Omega, \end{aligned} \tag{2.6}$$

and $\varphi_1 > 0$ in Ω the corresponding eigenfunction. Then $\varphi_1 \in C^{2+\alpha}(\overline{\Omega})$. Moreover one has the following lemma.

Lemma 2.2 (see [10]). *One has*

$$\int_{\Omega} \varphi_1^r dx < \infty \quad (2.7)$$

if and only if $r > -1$.

Now we give the proof of our theorems.

Proof of Theorem 1.1. Let $p \in (0, 1)$, and let u^* denote the unique solution of

$$\begin{aligned} -\Delta u &= u^p, & \text{in } \Omega, \\ u &> 0, & \text{in } \Omega, \\ u &= 0, & \text{on } \partial\Omega, \end{aligned} \quad (2.8)$$

where u^* belongs to $C^2(\overline{\Omega})$ (see [16]). Then $u = \lambda^{1/(1-p)} u^*$ is a solution of

$$\begin{aligned} -\Delta u &= \lambda u^p, & \text{in } \Omega, \\ u &> 0, & \text{in } \Omega, \\ u &= 0, & \text{on } \partial\Omega, \end{aligned} \quad (2.9)$$

where $0 < p < 1$ and $\lambda > 0$. Then fix $\lambda > 0$ and set

$$\bar{u} = \lambda^{1/(1-p)} u^*, \quad (2.10)$$

thus we can easily obtain that $\bar{u}$ is a supersolution of problem (1.1).

Now, we want to find a subsolution of problem (1.1). Let

$$\underline{u} = M \varphi_1^{2/(1+\gamma)}, \quad (2.11)$$

where M is a positive constant; now we will prove that $\underline{u}$ is a subsolution of problem (1.1). By Hopf's maximum principle in [17], there exist $\delta > 0$ and $\varepsilon_0 > 0$ such that

$$\begin{aligned} |\nabla \varphi_1| &\geq \delta, & \text{on } \Omega \setminus \Omega', \\ \varphi_1 &\geq \delta, & \text{on } \Omega', \end{aligned} \quad (2.12)$$

where $\Omega' = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > \varepsilon_0\}$. On Ω' , we choose $M \geq M_1 \triangleq ((\|k\|_{\infty}(1 + \gamma))/\lambda_1 \delta^2)^{1/(1+\gamma)}$, then we have

$$\frac{k(x)}{M^{\gamma} \varphi_1^{2\gamma/(1+\gamma)}} \leq \frac{\lambda_1 M}{1 + \gamma} \varphi_1^{2/(1+\gamma)}, \quad (2.13)$$

where $\|k\|_\infty = \max\{|k(x)| \mid x \in \overline{\Omega}\}$ for $k \in C(\overline{\Omega})$. On $\Omega \setminus \Omega'$, we choose $M \geq M_2 \triangleq (\|k\|_\infty(1+\gamma)^2/2(1-\gamma)\delta^2)^{1/(1+\gamma)}$, then one obtains

$$\frac{k(x)}{M^\gamma \varphi_1^{2\gamma/(1+\gamma)}} \leq \frac{2(1-\gamma)M|\nabla \varphi_1|^2}{(1+\gamma)^2 \varphi_1^{2\gamma/(1+\gamma)}}. \quad (2.14)$$

Thus, we choose $M \geq \max\{M_1, M_2\}$, then fixing M , let $\lambda > \lambda' \triangleq (3\lambda_1 M^{1-p})/(1+\gamma) \|\varphi_1\|_\infty^{2(1-p)/(1+\gamma)}$, it follows from (2.13) and (2.14) that

$$\begin{aligned} -\Delta \underline{u} + k(x) \underline{u}_\lambda^{-\gamma} &= -M \Delta \varphi_1^{2/(1+\gamma)} + \frac{k(x)}{M^\gamma \varphi_1^{2\gamma/(1+\gamma)}} \\ &= -M \left(\frac{2(1-\gamma)}{(1+\gamma)^2} |\nabla \varphi_1|^2 \varphi_1^{-2\gamma/(1+\gamma)} + \frac{2}{1+\gamma} \varphi_1^{(1-\gamma)/(1+\gamma)} \Delta \varphi_1 \right) + \frac{k(x)}{M^\gamma \varphi_1^{2\gamma/(1+\gamma)}} \\ &= \frac{2\lambda_1 M}{1+\gamma} \varphi_1^{2/(1+\gamma)} + \frac{k(x)}{M^\gamma \varphi_1^{2\gamma/(1+\gamma)}} - \frac{2(1-\gamma)M|\nabla \varphi_1|^2}{(1+\gamma)^2 \varphi_1^{2\gamma/(1+\gamma)}} \\ &\leq \frac{3\lambda_1 M}{1+\gamma} \varphi_1^{2/(1+\gamma)} \\ &\leq \lambda \left(M \varphi_1^{2/(1+\gamma)} \right)^p \\ &= \lambda \underline{u}_\lambda^p. \end{aligned} \quad (2.15)$$

Thus we proved that $\underline{u} = M \varphi_1^{2/(1+\gamma)}$ is a subsolution of problem (1.1) for all $\lambda > \lambda'$. According to Lemma 4 in [14], there exists a positive constant C such that

$$\varphi_1(x) \leq C u^*(x), \quad \text{in } \overline{\Omega}. \quad (2.16)$$

Set $\lambda \geq \lambda'' \triangleq (MC \|\varphi_1\|_\infty^{(1-\gamma)/(1+\gamma)})^{1-p}$, then we have

$$\overline{u} = \lambda^{1/(1-p)} u^* \geq \underline{u} = M \varphi_1^{2/(1+\gamma)}, \quad \text{in } \Omega. \quad (2.17)$$

Thus we choose $\lambda^* = \max\{\lambda', \lambda''\}$; via Lemma 2.1, problem (1.1) has at least one solution $u_\lambda \in C^{2+\alpha}(\Omega) \cap C(\overline{\Omega})$ and satisfying

$$\underline{u}(x) \leq u_\lambda(x) \leq \overline{u}(x), \quad \text{in } \overline{\Omega}, \quad (2.18)$$

for all $\lambda \geq \lambda^*$.

Since $u_\lambda \geq M\varphi_1^{2/(1+\gamma)}$ in $\overline{\Omega}$ for all $\lambda \geq \lambda^*$ and $-2\gamma/(1+\gamma) > -1$, according to Lemma 2.2 one has

$$\int_{\Omega} u_\lambda^{-\gamma}(x) dx \leq \frac{1}{M^\gamma} \int_{\Omega} \varphi_1^{-2\gamma/(1+\gamma)}(x) dx < +\infty. \quad (2.19)$$

So we obtain $u_\lambda^{-\gamma} \in L^1(\Omega)$.

Let $\Omega_j = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > r/2j\}$, $j = 1, 2, 3, \dots$, and let u_j be the unique solution of

$$\begin{aligned} -\Delta u + k(x)u_{j-1}^{-\gamma} &= \lambda u_{j-1}^p, \quad \text{in } \Omega_j, \\ u &= u_{j-1}, \quad \text{on } \overline{\Omega} \setminus \Omega_j, \end{aligned} \quad (2.20)$$

for $j = 1, 2, 3, \dots$, and with $u_0 = \bar{u} = \lambda^{1/(1-p)} u^*$, where

$$r = \max_{x \in \Omega} \min_{y \in \partial\Omega} |x - y|. \quad (2.21)$$

We claim that u_j is nonincreasing with respect to j in $\overline{\Omega}$ for all $j \in N$. Indeed, since $\bar{u}$ is a supersolution of problem (1.1) for all $\lambda > 0$, then we have

$$\begin{aligned} -\Delta(u_0 - u_1) &= -\Delta u_0 + \Delta u_1 \\ &= -\Delta u_0 + k(x)u_0^{-\gamma} - \lambda u_0^p \\ &= -\Delta \bar{u} + k(x)\bar{u}^{-\gamma} - \lambda \bar{u}^p \\ &> 0, \end{aligned} \quad (2.22)$$

for all $x \in \Omega_1$. Since $u_1 = u_0$ in $\overline{\Omega} \setminus \Omega_1$, so by the maximum principle, one has $u_0 \geq u_1$ in $\overline{\Omega}$. So when $j = 0$ our claim is true. We assume that our claim is true when $j = n$; that is, $u_n \leq u_{n-1}$ in $\overline{\Omega}$. Then we obtain

$$\begin{aligned} -\Delta(u_n - u_{n+1}) &= -\Delta u_n + \Delta u_{n+1} \\ &= \lambda(u_{n-1}^p - u_n^p) + k(x)(u_n^{-\gamma} - u_{n-1}^{-\gamma}) \\ &> 0, \end{aligned} \quad (2.23)$$

for all $x \in \Omega_{n+1}$. Since $u_n = u_{n+1}$ in $\overline{\Omega} \setminus \Omega_{n+1}$, so by the maximum principle, one has $u_n \geq u_{n+1}$ in $\overline{\Omega}$. Thus by the induction, one obtains

$$u_{j+1} \leq u_j, \quad \text{in } \overline{\Omega}, \quad (2.24)$$

for all $j \in N$. Then by the monotonicity of u_j , we have

$$\begin{aligned} -\Delta u_j &= \lambda u_{j-1}^p - k(x) u_{j-1}^{-\gamma} \\ &\geq \lambda u_j^p - k(x) u_j^{-\gamma}, \end{aligned} \quad (2.25)$$

for all $x \in \Omega_j$ and $j \in N^+$. According to the definitions of u_j and u_0 , we obtain that u_j is a supersolution of problem (1.1) for all $j \in N^+$. Let u_λ be a classical solution of problem (1.1), thus one has

$$u_\lambda(x) \leq u_{j+1}(x) \leq u_j(x) \leq u_0(x), \quad \text{in } \overline{\Omega}. \quad (2.26)$$

Assume that $v_\lambda(x) = \lim_{j \rightarrow \infty} u_j(x)$ for all $x \in \overline{\Omega}$, then by standard elliptic arguments (see [17]) it follows that v_λ is a solution of problem (1.1), and $v_\lambda \geq u_\lambda$ in Ω for any u_λ . Therefore, v_λ is the maximal solution of problem (1.1). According to the above arguments, problem (1.1) has a maximal solution for $\lambda \geq \lambda^*$.

To complete the proof of Theorem 1.1, setting

$$\begin{aligned} \sigma &= \{\lambda > 0 \mid \text{problem (1.1) has at least one solution } u_\lambda\}, \\ \bar{\lambda} &= \inf \sigma, \end{aligned} \quad (2.27)$$

then $[\lambda^*, +\infty) \subset \sigma$, $\bar{\lambda} \leq \lambda^*$. It suffices to prove that if $\lambda_0 \in \sigma$, then $[\lambda_0, +\infty) \subset \sigma$; that is, assume that $\lambda > \lambda_0$, then problem (1.1) has at least one solution. Let u_{λ_0} be a solution of problem (1.1) corresponding to λ_0 , then u_{λ_0} is a subsolution of problem (1.1) with every fixed $\lambda > \lambda_0$. Since $\bar{u} = \lambda^{1/(1-p)} u^*$ is a supersolution of problem (1.1) for any $\lambda > 0$, then one has

$$\lambda^{1/(1-p)} u^* \geq \lambda_0^{1/(1-p)} u^* \geq u_{\lambda_0}, \quad \text{in } \Omega, \quad (2.28)$$

for all $\lambda > \lambda_0$. According to Lemma 2.1, problem (1.1) has at least one solution $u_\lambda \in C^{2+\alpha}(\Omega) \cap C(\overline{\Omega})$ for all $\lambda > \lambda_0$. Moreover,

$$u_{\lambda_0}(x) \leq u_\lambda(x) \leq \bar{u}(x), \quad \text{in } \Omega. \quad (2.29)$$

Consequently, the maximal solution v_λ of problem (1.1) is increasing with respect to λ for all $\lambda > \bar{\lambda}$. So the proof of Theorem 1.1 is completed. $\square$

Proof of Theorem 1.3. Suppose to the contrary that there exists $\lambda > 0$ such that problem (1.1) has one solution $u_\lambda \in C^2(\Omega) \cap C(\overline{\Omega})$. Let e be the unique solution of

$$\begin{aligned} -\Delta u &= 1, \quad \text{in } \Omega, \\ u &= 0, \quad \text{on } \partial\Omega, \end{aligned} \quad (2.30)$$

$e \in C^{2+\alpha}(\overline{\Omega})$. By the maximum principle, $e > 0$ in Ω . We claim that for any solution u_λ of problem (1.1), there exists a constant $M = M(\lambda) > 0$ such that

$$Me(x) > u_\lambda(x), \quad \text{in } \Omega. \quad (2.31)$$

Indeed, let $M = \lambda \|u_\lambda\|_\infty^p + 1$, then one obtains

$$\begin{aligned} -\Delta(Me - u_\lambda) &= -M\Delta e + \Delta u_\lambda \\ &= \lambda \|u_\lambda\|_\infty^p + 1 - \lambda u_\lambda^p(x) + k(x)u_\lambda^{-\gamma} \\ &> 0, \end{aligned} \quad (2.32)$$

for all $x \in \Omega$. Since $(Me - u_\lambda)|_{\partial\Omega} = 0$, by the maximum principle we have

$$Me(x) > u_\lambda(x), \quad \text{in } \Omega. \quad (2.33)$$

According to Lemma 4 in [14], there exists a positive constant C such that

$$e(x) \leq C\varphi_1(x), \quad \text{in } \Omega. \quad (2.34)$$

Since $\gamma \geq 1$, from Lemma 2.2, it follows that

$$\int_{\Omega} u_\lambda^{-\gamma}(x) dx \geq \frac{1}{(CM)^\gamma} \int_{\Omega} \varphi_1^{-\gamma}(x) dx = +\infty. \quad (2.35)$$

Thus we obtain

$$\int_{\Omega} u_\lambda^{-\gamma} dx = +\infty. \quad (2.36)$$

Set

$$\Omega_i = \left\{ x \in \Omega \mid \text{dist}(x, \partial\Omega) > \frac{r}{2i}, \quad i \in N^+ \right\}, \quad (2.37)$$

and $\Omega = \bigcup_{i=1}^{\infty} \Omega_i$, then $\Omega_i \subset \Omega$ and $u_\lambda \in C^2(\overline{\Omega_i})$, satisfying

$$-\Delta u_\lambda + k(x)u_\lambda^{-\gamma} = \lambda u_\lambda^p, \quad (2.38)$$

for all $x \in \overline{\Omega_i}$ and $i \in N^+$. Consequently, integrating (2.38) we have

$$-\int_{\Omega_i} \Delta u_\lambda dx + \int_{\Omega_i} k(x)u_\lambda^{-\gamma} dx = \lambda \int_{\Omega_i} u_\lambda^p dx \leq \lambda \int_{\Omega} u_\lambda^p dx, \quad (2.39)$$

noting that

$$\int_{\Omega_i} \Delta u_\lambda dx = \int_{\partial\Omega_i} \frac{\partial u_\lambda}{\partial n} ds, \quad (2.40)$$

where n denotes the outward normal to $\partial\Omega_i$. From (2.39) and (2.40), letting $i \rightarrow \infty$, one has

$$\int_{\Omega} k(x) u_\lambda^{-\gamma} dx - \int_{\partial\Omega} \frac{\partial u_\lambda}{\partial n} ds \leq \lambda \|u_\lambda\|_\infty^p |\Omega|, \quad (2.41)$$

where $|\Omega|$ denotes the Lebesgue measure of Ω . According to (2.36) and $k(x) > 0$ in $\bar{\Omega}$, one obtains

$$\int_{\partial\Omega} \frac{\partial u_\lambda}{\partial n} ds = +\infty. \quad (2.42)$$

But this is impossible, by Hopf's maximum principle, we have

$$\frac{\partial u_\lambda}{\partial n} < 0, \quad (2.43)$$

for all $x \in \partial\Omega$, where n denotes the outward normal to $\partial\Omega$ at x . Therefore Theorem 1.3 is true. $\square$

Acknowledgment

This paper is supported by NNSF of China under Grant 10771173 and the Natural Science Foundation of Education of Guizhou Province under Grant 2008067.

References

- [1] A. Callegari and A. Nachman, "Some singular, nonlinear differential equations arising in boundary layer theory," *Journal of Mathematical Analysis and Applications*, vol. 64, no. 1, pp. 96–105, 1978.
- [2] A. Nachman and A. Callegari, "A nonlinear singular boundary value problem in the theory of pseudoplastic fluids," *SIAM Journal on Applied Mathematics*, vol. 38, no. 2, pp. 275–281, 1980.
- [3] Y. S. Choi, A. C. Lazer, and P. J. McKenna, "Some remarks on a singular elliptic boundary value problem," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 32, no. 3, pp. 305–314, 1998.
- [4] J. I. Diaz, J.-M. Morel, and L. Oswald, "An elliptic equation with singular nonlinearity," *Communications in Partial Differential Equations*, vol. 12, no. 12, pp. 1333–1344, 1987.
- [5] W. Fulks and J. S. Maybee, "A singular nonlinear elliptic equations," *Osaka Journal of Mathematics*, vol. 12, no. 5, pp. 1–19, 1960.
- [6] C. A. Stuart, "Existence and approximation of solutions of non-linear elliptic equations," *Mathematische Zeitschrift*, vol. 147, no. 1, pp. 53–63, 1976.
- [7] M. M. Coclite and G. Palmieri, "On a singular nonlinear Dirichlet problem," *Communications in Partial Differential Equations*, vol. 14, no. 10, pp. 1315–1327, 1989.
- [8] M. G. Crandall, P. H. Rabinowitz, and L. Tartar, "On a Dirichlet problem with a singular nonlinearity," *Communications in Partial Differential Equations*, vol. 2, no. 2, pp. 193–222, 1977.
- [9] A. L. Edelson, "Entire solutions of singular elliptic equations," *Journal of Mathematical Analysis and Applications*, vol. 139, no. 2, pp. 523–532, 1989.

- [10] A. C. Lazer and P. J. McKenna, "On a singular nonlinear elliptic boundary-value problem," *Proceedings of the American Mathematical Society*, vol. 111, no. 3, pp. 721–730, 1991.
- [11] M. A. del Pino, "A global estimate for the gradient in a singular elliptic boundary value problem," *Proceedings of the Royal Society of Edinburgh. Section A*, vol. 122, no. 3-4, pp. 341–352, 1992.
- [12] M. Ghergu and V. Rădulescu, "Sublinear singular elliptic problems with two parameters," *Journal of Differential Equations*, vol. 195, no. 2, pp. 520–536, 2003.
- [13] J. Shi and M. Yao, "On a singular nonlinear semilinear elliptic problem," *Proceedings of the Royal Society of Edinburgh. Section A*, vol. 128, no. 6, pp. 1389–1401, 1998.
- [14] Z. Zhang, "On a Dirichlet problem with a singular nonlinearity," *Journal of Mathematical Analysis and Applications*, vol. 194, no. 1, pp. 103–113, 1995.
- [15] S. Cui, "Existence and nonexistence of positive solutions for singular semilinear elliptic boundary value problems," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 41, no. 1-2, pp. 149–176, 2000.
- [16] P.-L. Lions, "On the existence of positive solutions of semilinear elliptic equations," *SIAM Review*, vol. 24, no. 4, pp. 441–467, 1982.
- [17] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, vol. 224 of *Fundamental Principles of Mathematical Sciences*, Springer, Berlin, Germany, 2nd edition, 1983.