Research Article

Fixed Points and the Stability of an AQCQ-Functional Equation in Non-Archimedean Normed Spaces

Choonkil Park

*Department of Mathematics, Research Institute for Natural Sciences, Hanyang University,
Seoul 133-791, South Korea*

Correspondence should be addressed to Choonkil Park, baak@hanyang.ac.kr

Received 8 January 2010; Accepted 8 April 2010

Academic Editor: W. A. Kirk

Copyright © 2010 Choonkil Park. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Using fixed point method, we prove the generalized Hyers-Ulam stability of the following additive-quadratic-cubic-quartic functional equation $f(x+2y) + f(x-2y) = 4f(x+y) + 4f(x-y) - 6f(x) + f(2y) + f(-2y) - 4f(y) - 4f(-y)$ in non-Archimedean Banach spaces.

1. Introduction and Preliminaries

A *valuation* is a function $|\cdot|$ from a field K into $[0, \infty)$ such that 0 is the unique element having the 0 valuation, $|rs| = |r| \cdot |s|$, and the triangle inequality holds, that is,

$$|r + s| \leq |r| + |s|, \quad \forall r, s \in K. \quad (1.1)$$

A field K is called a *valued field* if K carries a valuation. The usual absolute values of $\mathbb{R}$ and $\mathbb{C}$ are examples of valuations.

Let us consider a valuation which satisfies a stronger condition than the triangle inequality. If the triangle inequality is replaced by

$$|r + s| \leq \max\{|r|, |s|\}, \quad \forall r, s \in K, \quad (1.2)$$

then the function $|\cdot|$ is called a *non-Archimedean valuation*, and the field is called a *non-Archimedean field*. Clearly $|1| = |-1| = 1$ and $|n| \leq 1$ for all $n \in \mathbb{N}$. A trivial example of

a non-Archimedean valuation is the function $|\cdot|$ taking everything except for 0 into 1 and $|0| = 0$.

Throughout this paper, we assume that the base field is a non-Archimedean field, hence call it simply a field.

Definition 1.1 (see [1]). Let X be a vector space over a field K with a non-Archimedean valuation $|\cdot|$. A function $\|\cdot\| : X \rightarrow [0, \infty)$ is said to be a *non-Archimedean norm* if it satisfies the following conditions:

- (i) $\|x\| = 0$ if and only if $x = 0$;
- (ii) $\|rx\| = |r|\|x\|$ ($r \in K, x \in X$);
- (iii) the strong triangle inequality

$$\|x + y\| \leq \max\{\|x\|, \|y\|\}, \quad \forall x, y \in X \quad (1.3)$$

holds. Then $(X, \|\cdot\|)$ is called a *non-Archimedean normed space*.

Definition 1.2. (i) Let $\{x_n\}$ be a sequence in a non-Archimedean normed space X . Then the sequence $\{x_n\}$ is called *Cauchy* if for a given $\varepsilon > 0$, there is a positive integer N such that

$$\|x_n - x_m\| \leq \varepsilon \quad (1.4)$$

for all $n, m \geq N$.

(ii) Let $\{x_n\}$ be a sequence in a non-Archimedean normed space X . Then the sequence $\{x_n\}$ is called *convergent* if for a given $\varepsilon > 0$, there are a positive integer N and an $x \in X$ such that

$$\|x_n - x\| \leq \varepsilon \quad (1.5)$$

for all $n \geq N$. Then we call $x \in X$ a limit of the sequence $\{x_n\}$, and denote it by $\lim_{n \rightarrow \infty} x_n = x$.

(iii) If every Cauchy sequence in X converges, then the non-Archimedean normed space X is called a *non-Archimedean Banach space*.

The stability problem of functional equations originated from a question of Ulam [2] concerning the stability of group homomorphisms. Hyers [3] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [4] for additive mappings and by Rassias [5] for linear mappings by considering an unbounded Cauchy difference. The paper of Rassias [5] has provided a lot of influence in the development of what we call *generalized Hyers-Ulam stability* or as *Hyers-Ulam-Rassias stability* of functional equations. A generalization of the Rassias theorem was obtained by Găvruta [6] by replacing the unbounded Cauchy difference by a general control function in the spirit of Rassias' approach.

The functional equation

$$f(x + y) + f(x - y) = 2f(x) + 2f(y) \quad (1.6)$$

is called a *quadratic functional equation*. In particular, every solution of the quadratic functional equation is said to be a *quadratic mapping*. A generalized Hyers-Ulam stability problem for the quadratic functional equation was proved by Skof [7] for mappings $f : X \rightarrow Y$, where X is a normed space and Y is a Banach space. Cholewa [8] noticed that the theorem of Skof is still true if the relevant domain X is replaced by an Abelian group. Czerwik [9] proved the generalized Hyers-Ulam stability of the quadratic functional equation.

In [10], Jun and Kim considered the following cubic functional equation:

$$f(2x + y) + f(2x - y) = 2f(x + y) + 2f(x - y) + 12f(x), \quad (1.7)$$

which is called a *cubic functional equation* and every solution of the cubic functional equation is said to be a *cubic mapping*. In [11], Lee et al. considered the following quartic functional equation:

$$f(2x + y) + f(2x - y) = 4f(x + y) + 4f(x - y) + 24f(x) - 6f(y), \quad (1.8)$$

which is called a *quartic functional equation* and every solution of the quartic functional equation is said to be a *quartic mapping*.

The stability problems of several functional equations have been extensively investigated by a number of authors and there are many interesting results concerning this problem (see [12–27]).

Let X be a set. A function $d : X \times X \rightarrow [0, \infty]$ is called a *generalized metric* on X if d satisfies

- (1) $d(x, y) = 0$ if and only if $x = y$;
- (2) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (3) $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in X$.

We recall a fundamental result in fixed point theory.

Theorem 1.3 (see [28, 29]). *Let (X, d) be a complete generalized metric space and let $J : X \rightarrow X$ be a strictly contractive mapping with Lipschitz constant $L < 1$. Then for each given element $x \in X$, either*

$$d(J^n x, J^{n+1} x) = \infty \quad (1.9)$$

for all nonnegative integers n or there exists a positive integer n_0 such that

- (1) $d(J^n x, J^{n+1} x) < \infty$, for all $n \geq n_0$;
- (2) the sequence $\{J^n x\}$ converges to a fixed point y^* of J ;
- (3) y^* is the unique fixed point of J in the set $Y = \{y \in X \mid d(J^{n_0} x, y) < \infty\}$;
- (4) $d(y, y^*) \leq (1/(1-L))d(y, Jy)$ for all $y \in Y$.

In 1996, Isac and Rassias [30] were the first to provide applications of stability theory of functional equations for the proof of new fixed point theorems with applications. By using fixed point methods, the stability problems of several functional equations have been extensively investigated by a number of authors (see [31–36]).

This paper is organized as follows: in Section 2, using the fixed point method, we prove the generalized Hyers-Ulam stability of the additive-quadratic-cubic-quartic functional equation

$$f(x+2y) + f(x-2y) = 4f(x+y) + 4f(x-y) - 6f(x) + f(2y) + f(-2y) - 4f(y) - 4f(-y) \quad (1.10)$$

in non-Archimedean Banach spaces for an odd case. In Section 3, using the fixed point method, we prove the generalized Hyers-Ulam stability of the additive-quadratic-cubic-quartic functional equation (1.10) in non-Archimedean Banach spaces for an even case.

Throughout this paper, assume that X is a non-Archimedean normed vector space and that Y is a non-Archimedean Banach space.

2. Generalized Hyers-Ulam Stability of the Functional Equation (1.10): An Odd Case

One can easily show that an odd mapping $f : X \rightarrow Y$ satisfies (1.10) if and only if the odd mapping $f : X \rightarrow Y$ is an additive-cubic mapping, that is,

$$f(x+2y) + f(x-2y) = 4f(x+y) + 4f(x-y) - 6f(x). \quad (2.1)$$

It was shown in Lemma 2.2 of [37] that $g(x) := f(2x) - 2f(x)$ and $h(x) := f(2x) - 8f(x)$ are cubic and additive, respectively, and that $f(x) = (1/6)g(x) - (1/6)h(x)$.

One can easily show that an even mapping $f : X \rightarrow Y$ satisfies (1.10) if and only if the even mapping $f : X \rightarrow Y$ is a quadratic-quartic mapping, that is,

$$f(x+2y) + f(x-2y) = 4f(x+y) + 4f(x-y) - 6f(x) + 2f(2y) - 8f(y). \quad (2.2)$$

It was shown in Lemma 2.1 of [38] that $g(x) := f(2x) - 4f(x)$ and $h(x) := f(2x) - 16f(x)$ are quartic and quadratic, respectively, and that $f(x) = (1/12)g(x) - (1/12)h(x)$.

For a given mapping $f : X \rightarrow Y$, we define

$$\begin{aligned} Df(x, y) := & f(x+2y) + f(x-2y) - 4f(x+y) - 4f(x-y) + 6f(x) \\ & - f(2y) - f(-2y) + 4f(y) + 4f(-y) \end{aligned} \quad (2.3)$$

for all $x, y \in X$.

We prove the generalized Hyers-Ulam stability of the functional equation $Df(x, y) = 0$ in non-Archimedean Banach spaces: an odd case.

Theorem 2.1. *Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with*

$$\varphi(x, y) \leq \frac{L}{|8|} \varphi(2x, 2y) \quad (2.4)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying

$$\|Df(x, y)\| \leq \varphi(x, y) \quad (2.5)$$

for all $x, y \in X$. Then there is a unique cubic mapping $C : X \rightarrow Y$ such that

$$\|f(2x) - 2f(x) - C(x)\| \leq \frac{L}{|8| - |8|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (2.6)$$

for all $x \in X$.

Proof. Letting $x = y$ in (2.5), we get

$$\|f(3y) - 4f(2y) + 5f(y)\| \leq \varphi(y, y) \quad (2.7)$$

for all $y \in X$.

Replacing x by $2y$ in (2.5), we get

$$\|f(4y) - 4f(3y) + 6f(2y) - 4f(y)\| \leq \varphi(2y, y) \quad (2.8)$$

for all $y \in X$.

By (2.7) and (2.8),

$$\begin{aligned} & \|f(4y) - 10f(2y) + 16f(y)\| \\ & \leq \max\{\|4(f(3y) - 4f(2y) + 5f(y))\|, \|f(4y) - 4f(3y) + 6f(2y) - 4f(y)\|\} \\ & \leq \max\{|4| \cdot \|f(3y) - 4f(2y) + 5f(y)\|, \|f(4y) - 4f(3y) + 6f(2y) - 4f(y)\|\} \\ & \leq \max\{|4|\varphi(y, y), \varphi(2y, y)\} \end{aligned} \quad (2.9)$$

for all $y \in X$.

Letting $y := x/2$ and $g(x) := f(2x) - 2f(x)$ for all $x \in X$, we get

$$\left\|g(x) - 8g\left(\frac{x}{2}\right)\right\| \leq \max\left\{|4|\varphi\left(\frac{x}{2}, \frac{x}{2}\right), \varphi\left(x, \frac{x}{2}\right)\right\} \quad (2.10)$$

for all $x \in X$.

Consider the set

$$S := \{g : X \rightarrow Y\}, \quad (2.11)$$

and introduce the generalized metric on S

$$d(g, h) = \inf\{\mu \in \mathbb{R}_+ : \|g(x) - h(x)\| \leq \mu(\max\{|4|\varphi(x, x), \varphi(2x, x), \forall x \in X\})\}, \quad (2.12)$$

where, as usual, $\inf \phi = +\infty$. It is easy to show that (S, d) is complete. (See the proof of Lemma 2.1 of [39].)

Now we consider the linear mapping $J : S \rightarrow S$ such that

$$Jg(x) := 8g\left(\frac{x}{2}\right) \quad (2.13)$$

for all $x \in X$.

Let $g, h \in S$ be given such that $d(g, h) = \varepsilon$. Then

$$\|g(x) - h(x)\| \leq \varepsilon \cdot \max\{|4|\varphi(x, x), \varphi(2x, x)\}| \quad (2.14)$$

for all $x \in X$. Hence

$$\|Jg(x) - Jh(x)\| = \left\| 8g\left(\frac{x}{2}\right) - 8h\left(\frac{x}{2}\right) \right\| \leq |8|\varepsilon \frac{L}{|8|} \max\{|4|\varphi(x, x), \varphi(2x, x)\}| \quad (2.15)$$

for all $x \in X$. So $d(g, h) = \varepsilon$ implies that $d(Jg, Jh) \leq L\varepsilon$. This means that

$$d(Jg, Jh) \leq Ld(g, h) \quad (2.16)$$

for all $g, h \in S$.

It follows from (2.10) that

$$\left\| g(x) - 8g\left(\frac{x}{2}\right) \right\| \leq \frac{L}{|8|} (\max\{|4|\varphi(x, x), \varphi(2x, x)\}|) \quad (2.17)$$

for all $x \in X$. So $d(g, Jg) \leq L/|8|$.

By Theorem 1.3, there exists a mapping $C : X \rightarrow Y$ satisfying the following.

(1) C is a fixed point of J , that is,

$$C\left(\frac{x}{2}\right) = \frac{1}{8}C(x) \quad (2.18)$$

for all $x \in X$. The mapping C is a unique fixed point of J in the set

$$M = \{h \in S : d(g, h) < \infty\}. \quad (2.19)$$

This implies that C is a unique mapping satisfying (2.18) such that there exists a $\mu \in (0, \infty)$ satisfying

$$\|g(x) - C(x)\| \leq \mu \cdot \max\{|4|\varphi(x, x), \varphi(2x, x)\}| \quad (2.20)$$

for all $x \in X$; since $g : X \rightarrow Y$ is odd, $C : X \rightarrow Y$ is an odd mapping.

(2) $d(J^n g, C) \rightarrow 0$ as $n \rightarrow \infty$. This implies the equality

$$\lim_{n \rightarrow \infty} 8^n g\left(\frac{x}{2^n}\right) = C(x) \quad (2.21)$$

for all $x \in X$.

(3) $d(g, C) \leq (1/(1-L))d(g, Jg)$, which implies the inequality

$$d(g, C) \leq \frac{L}{|8| - |8|L}. \quad (2.22)$$

This implies that the inequality (2.6) holds.

By (2.5),

$$\left\| 8^n Dg\left(\frac{x}{2^n}, \frac{y}{2^n}\right) \right\| \leq |8|^n \max\left\{ \varphi\left(\frac{2x}{2^n}, \frac{2y}{2^n}\right), |2|\varphi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) \right\} \quad (2.23)$$

for all $x, y \in X$ and all $n \in \mathbb{N}$. So

$$\left\| 8^n Dg\left(\frac{x}{2^n}, \frac{y}{2^n}\right) \right\| \leq |8|^n \frac{L^n}{|8|^n} \max\{\varphi(2x, 2y), |2|\varphi(x, y)\} \quad (2.24)$$

for all $x, y \in X$ and all $n \in \mathbb{N}$. So

$$\|DC(x, y)\| = 0 \quad (2.25)$$

for all $x, y \in X$. Thus the mapping $C : X \rightarrow Y$ is cubic, as desired. $\square$

Corollary 2.2. *Let θ and p be positive real numbers with $p < 3$. Let $f : X \rightarrow Y$ be an odd mapping satisfying*

$$\|Df(x, y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.26)$$

for all $x, y \in X$. Then there exists a unique cubic mapping $C : X \rightarrow Y$ such that

$$\|f(2x) - 2f(x) - C(x)\| \leq \max\{2 \cdot |4|, |2|^p + 1\} \frac{\theta}{|2|^p - |8|} \|x\|^p \quad (2.27)$$

for all $x \in X$.

Proof. The proof follows from Theorem 2.1 by taking

$$\varphi(x, y) := \theta(\|x\|^p + \|y\|^p) \quad (2.28)$$

for all $x, y \in X$. Then we can choose $L = |8|/|2|^p$ and we get the desired result. $\square$

Theorem 2.3. Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y) \leq |8|L\varphi\left(\frac{x}{2}, \frac{y}{2}\right) \quad (2.29)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.5). Then there is a unique cubic mapping $C : X \rightarrow Y$ such that

$$\|f(2x) - 2f(x) - C(x)\| \leq \frac{1}{|8| - |8|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (2.30)$$

for all $x \in X$.

Proof. It follows from (2.10) that

$$\left\|g(x) - \frac{1}{8}g(2x)\right\| \leq \frac{1}{|8|} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (2.31)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

Theorem 2.4. Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y) \leq \frac{L}{|2|}\varphi(2x, 2y) \quad (2.32)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.5). Then there is a unique additive mapping $A : X \rightarrow Y$ such that

$$\|f(2x) - 8f(x) - A(x)\| \leq \frac{L}{|2| - |2|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (2.33)$$

for all $x \in X$.

Proof. Letting $y := x/2$ and $g(x) := f(2x) - 8f(x)$ for all $x \in X$ in (2.9), we get

$$\left\|g(x) - 2g\left(\frac{x}{2}\right)\right\| \leq \max\left\{|4|\varphi\left(\frac{x}{2}, \frac{x}{2}\right), \varphi\left(x, \frac{x}{2}\right)\right\} \quad (2.34)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

Corollary 2.5. *Let θ and p be positive real numbers with $p < 1$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.26). Then there exists a unique additive mapping $C : X \rightarrow Y$ such that*

$$\|f(2x) - 8f(x) - A(x)\| \leq \max\{2 \cdot |4|, |2|^p + 1\} \frac{\theta}{|2|^p - |2|} \|x\|^p \quad (2.35)$$

for all $x \in X$.

Proof. The proof follows from Theorem 2.4 by taking

$$\varphi(x, y) := \theta(\|x\|^p + \|y\|^p) \quad (2.36)$$

for all $x, y \in X$. Then we can choose $L = |2|/|2|^p$ and we get the desired result. $\square$

Theorem 2.6. *Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with*

$$\varphi(x, y) \leq |2|L\varphi\left(\frac{x}{2}, \frac{y}{2}\right) \quad (2.37)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an odd mapping satisfying (2.5). Then there is a unique additive mapping $A : X \rightarrow Y$ such that

$$\|f(2x) - 8f(x) - A(x)\| \leq \frac{1}{|2| - |2|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (2.38)$$

for all $x \in X$.

Proof. It follows from (2.34) that

$$\left\|g(x) - \frac{1}{2}g(2x)\right\| \leq \frac{1}{|2|} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (2.39)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

3. Generalized Hyers-Ulam Stability of the Functional Equation (1.10): An Even Case

Now we prove the generalized Hyers-Ulam stability of the functional equation $Df(x, y) = 0$ in non-Archimedean Banach spaces: an even case.

Theorem 3.1. *Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with*

$$\varphi(x, y) \leq \frac{L}{|16|} \varphi(2x, 2y) \quad (3.1)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.5) and $f(0) = 0$. Then there is a unique quartic mapping $Q : X \rightarrow Y$ such that

$$\|f(2x) - 4f(x) - Q(x)\| \leq \frac{L}{|16| - |16|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (3.2)$$

for all $x \in X$.

Proof. Letting $x = y$ in (2.5), we get

$$\|f(3y) - 6f(2y) + 15f(y)\| \leq \varphi(y, y) \quad (3.3)$$

for all $y \in X$.

Replacing x by $2y$ in (2.5), we get

$$\|f(4y) - 4f(3y) + 4f(2y) + 4f(y)\| \leq \varphi(2y, y) \quad (3.4)$$

for all $y \in X$.

By (3.3) and (3.4),

$$\begin{aligned} & \|f(4y) - 20f(2y) + 64f(y)\| \\ & \leq \max\{\|4(f(3y) - 6f(2y) + 15f(y))\|, \|f(4y) - 4f(3y) + 4f(2y) + 4f(y)\|\} \\ & \leq \max\{|4| \cdot \|f(3y) - 6f(2y) + 15f(y)\|, \|f(4y) - 4f(3y) + 4f(2y) + 4f(y)\|\} \\ & \leq \max\{|4|\varphi(y, y), \varphi(2y, y)\} \end{aligned} \quad (3.5)$$

for all $y \in X$.

Letting $y := x/2$ and $g(x) := f(2x) - 4f(x)$ for all $x \in X$, we get

$$\left\|g(x) - 16g\left(\frac{x}{2}\right)\right\| \leq \max\left\{|4|\varphi\left(\frac{x}{2}, \frac{x}{2}\right), \varphi\left(x, \frac{x}{2}\right)\right\} \quad (3.6)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

Corollary 3.2. Let θ and p be positive real numbers with $p < 4$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.26) and $f(0) = 0$. Then there exists a unique quartic mapping $Q : X \rightarrow Y$ such that

$$\|f(2x) - 4f(x) - Q(x)\| \leq \max\{2 \cdot |4|, |2|^p + 1\} \frac{\theta}{|2|^p - |16|} \|x\|^p \quad (3.7)$$

for all $x \in X$.

Proof. The proof follows from Theorem 3.1 by taking

$$\varphi(x, y) := \theta(\|x\|^p + \|y\|^p) \quad (3.8)$$

for all $x, y \in X$. Then we can choose $L = |16|/|2|^p$ and we get the desired result. $\square$

Theorem 3.3. Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y) \leq |16|L\varphi\left(\frac{x}{2}, \frac{y}{2}\right) \quad (3.9)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.5) and $f(0) = 0$. Then there is a unique quartic mapping $Q : X \rightarrow Y$ such that

$$\|f(2x) - 4f(x) - Q(x)\| \leq \frac{1}{|16| - |16|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (3.10)$$

for all $x \in X$.

Proof. It follows from (3.6) that

$$\left\|g(x) - \frac{1}{16}g(2x)\right\| \leq \frac{1}{|16|} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (3.11)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

Theorem 3.4. Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y) \leq \frac{L}{|4|}\varphi(2x, 2y) \quad (3.12)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.5) and $f(0) = 0$. Then there is a unique quadratic mapping $T : X \rightarrow Y$ such that

$$\|f(2x) - 16f(x) - T(x)\| \leq \frac{L}{|4| - |4|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (3.13)$$

for all $x \in X$.

Proof. Letting $y := x/2$ and $g(x) := f(2x) - 16f(x)$ for all $x \in X$ in (3.5), we get

$$\left\|g(x) - 4g\left(\frac{x}{2}\right)\right\| \leq \max\left\{|4|\varphi\left(\frac{x}{2}, \frac{x}{2}\right), \varphi\left(x, \frac{x}{2}\right)\right\} \quad (3.14)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

Corollary 3.5. *Let θ and p be positive real numbers with $p < 2$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.26) and $f(0) = 0$. Then there exists a unique quadratic mapping $T : X \rightarrow Y$ such that*

$$\|f(2x) - 16f(x) - T(x)\| \leq \max\{2 \cdot |4|, |2|^p + 1\} \frac{\theta}{|2|^p - |4|} \|x\|^p \quad (3.15)$$

for all $x \in X$.

Proof. The proof follows from Theorem 3.4 by taking

$$\varphi(x, y) := \theta(\|x\|^p + \|y\|^p) \quad (3.16)$$

for all $x, y \in X$. Then we can choose $L = |4|/|2|^p$ and we get the desired result. $\square$

Theorem 3.6. *Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with*

$$\varphi(x, y) \leq |4|L\varphi\left(\frac{x}{2}, \frac{y}{2}\right) \quad (3.17)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be an even mapping satisfying (2.5) and $f(0) = 0$. Then there is a unique quadratic mapping $T : X \rightarrow Y$ such that

$$\|f(2x) - 16f(x) - T(x)\| \leq \frac{1}{|4| - |4|L} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (3.18)$$

for all $x \in X$.

Proof. It follows from (3.14) that

$$\left\|g(x) - \frac{1}{4}g(2x)\right\| \leq \frac{1}{|4|} \max\{|4|\varphi(x, x), \varphi(2x, x)\} \quad (3.19)$$

for all $x \in X$.

The rest of the proof is similar to the proof of Theorem 2.1. $\square$

For a given f , let $f_o(x) := ((f(x) - f(-x))/2)$ and $f_e(x) := ((f(x) + f(-x))/2)$. Then f_o is odd and f_e is even. Let $g_o(x) := f_o(2x) - 2f_o(x)$ and $h_o(x) := f_o(2x) - 8f_o(x)$. Then $f_o(x) = (1/6)g_o(x) - (1/6)h_o(x)$. Let $g_e(x) := f_e(2x) - 4f_e(x)$ and $h_e(x) := f_e(2x) - 16f_e(x)$. Then $f_e(x) = (1/12)g_e(x) - (1/12)h_e(x)$. Thus

$$f(x) = \frac{1}{6}g_o(x) - \frac{1}{6}h_o(x) + \frac{1}{12}g_e(x) - \frac{1}{12}h_e(x). \quad (3.20)$$

Theorem 3.7. Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y) \leq \frac{L}{|2|} \varphi(2x, 2y) \quad (3.21)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ and (2.5). Then there exist an additive mapping $A : X \rightarrow Y$, a quadratic mapping $T : X \rightarrow Y$, a cubic mapping $C : X \rightarrow Y$, and a quartic mapping $Q : X \rightarrow Y$ such that

$$\begin{aligned} & \left\| f(x) - \frac{1}{6}A(x) - \frac{1}{12}T(x) - \frac{1}{6}C(x) - \frac{1}{12}Q(x) \right\| \\ & \leq \max \left\{ \frac{L}{|6| \cdot |2|(1-L)}, \frac{L}{|12| \cdot |4|(1-L)}, \frac{L}{|6| \cdot |8|(1-L)}, \frac{L}{|12| \cdot |16|(1-L)} \right\} \\ & \quad \cdot \frac{1}{|2|} \max \{ |4|\varphi(x, x), \varphi(2x, x), |4|\varphi(-x, -x), \varphi(-2x, -x) \} \\ & \leq \frac{L}{|12| \cdot |16| \cdot |2|(1-L)} \cdot \max \{ |4|\varphi(x, x), \varphi(2x, x), |4|\varphi(-x, -x), \varphi(-2x, -x) \} \end{aligned} \quad (3.22)$$

for all $x \in X$.

Corollary 3.8. Let θ and p be positive real numbers with $p < 1$. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ and (2.5). Then there exist an additive mapping $A : X \rightarrow Y$, a quadratic mapping $T : X \rightarrow Y$, a cubic mapping $C : X \rightarrow Y$ and a quartic mapping $Q : X \rightarrow Y$, such that

$$\left\| f(x) - \frac{1}{6}A(x) - \frac{1}{12}T(x) - \frac{1}{6}C(x) - \frac{1}{12}Q(x) \right\| \leq \max \{ 2 \cdot |4|, |2|^p + 1 \} \cdot \frac{\theta}{|12|(|2|^p - |2|)} \|x\|^p \quad (3.23)$$

for all $x \in X$.

Proof. The proof follows from Theorem 3.7 by taking

$$\varphi(x, y) := \theta(\|x\|^p + \|y\|^p) \quad (3.24)$$

for all $x, y \in X$. Then we can choose $L = |2|/|2|^p$ and we get the desired result. $\square$

Theorem 3.9. Let $\varphi : X^2 \rightarrow [0, \infty)$ be a function such that there exists an $L < 1$ with

$$\varphi(x, y) \leq |16|L\varphi\left(\frac{x}{2}, \frac{y}{2}\right) \quad (3.25)$$

for all $x, y \in X$. Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ and (2.5). Then there exist an additive mapping $A : X \rightarrow Y$, a quadratic mapping $T : X \rightarrow Y$, a cubic mapping $C : X \rightarrow Y$ and a quartic mapping $Q : X \rightarrow Y$ such that

$$\begin{aligned} & \left\| f(x) - \frac{1}{6}A(x) - \frac{1}{12}T(x) - \frac{1}{6}C(x) - \frac{1}{12}Q(x) \right\| \\ & \leq \max \left\{ \frac{1}{|6| \cdot |2|(1-L)}, \frac{1}{|12| \cdot |4|(1-L)}, \frac{1}{|6| \cdot |8|(1-L)}, \frac{1}{|12| \cdot |16|(1-L)} \right\} \\ & \quad \cdot \frac{1}{|2|} \max \{ |4|\varphi(x, x), \varphi(2x, x), |4|\varphi(-x, -x), \varphi(-2x, -x) \} \\ & \leq \frac{1}{|12| \cdot |16| \cdot |2|(1-L)} \cdot \max \{ |4|\varphi(x, x), \varphi(2x, x), |4|\varphi(-x, -x), \varphi(-2x, -x) \} \end{aligned} \quad (3.26)$$

for all $x \in X$.

Acknowledgment

This work was supported by Basic Science Research Program through the National Research Foundation of Korea funded by the Ministry of Education, Science and Technology (NRF-2009-0070788).

References

- [1] M. S. Moslehian and G. Sadeghi, "A Mazur-Ulam theorem in non-Archimedean normed spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 69, no. 10, pp. 3405–3408, 2008.
- [2] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience, New York, NY, USA, 1960.
- [3] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [4] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [5] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [6] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [7] F. Skof, "Proprietà locali e approssimazione di operatori," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 53, pp. 113–129, 1983.
- [8] P. W. Cholewa, "Remarks on the stability of functional equations," *Aequationes Mathematicae*, vol. 27, no. 1-2, pp. 76–86, 1984.
- [9] S. Czerwik, "On the stability of the quadratic mapping in normed spaces," *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, vol. 62, pp. 59–64, 1992.
- [10] K.-W. Jun and H.-M. Kim, "The generalized Hyers-Ulam-Rassias stability of a cubic functional equation," *Journal of Mathematical Analysis and Applications*, vol. 274, no. 2, pp. 267–278, 2002.
- [11] S. H. Lee, S. M. Im, and I. S. Hwang, "Quartic functional equations," *Journal of Mathematical Analysis and Applications*, vol. 307, no. 2, pp. 387–394, 2005.
- [12] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, USA, 2002.
- [13] Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431–434, 1991.

- [14] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, vol. 34 of *Progress in Nonlinear Differential Equations and Their Applications*, Birkhäuser, Boston, Mass, USA, 1998.
- [15] S.-M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, Fla, USA, 2001.
- [16] C. Park, "Hyers-Ulam-Rassias stability of homomorphisms in quasi-Banach algebras," *Bulletin des Sciences Mathématiques*, vol. 132, no. 2, pp. 87–96, 2008.
- [17] C. Park, "Hyers-Ulam-Rassias stability of a generalized Apollonius-Jensen type additive mapping and isomorphisms between C^* -algebras," *Mathematische Nachrichten*, vol. 281, no. 3, pp. 402–411, 2008.
- [18] C. Park and J. Cui, "Generalized stability of C^* -ternary quadratic mappings," *Abstract and Applied Analysis*, vol. 2007, Article ID 23282, 6 pages, 2007.
- [19] C. Park and A. Najati, "Homomorphisms and derivations in C^* -algebras," *Abstract and Applied Analysis*, vol. 2007, Article ID 80630, 12 pages, 2007.
- [20] Th. M. Rassias, "Problem 16; 2, Report of the 27th International Symposium on Functional Equations," *Aequationes Mathematicae*, vol. 39, pp. 292–293; 309, 1990.
- [21] Th. M. Rassias, "On the stability of the quadratic functional equation and its applications," *Studia Universitatis Babeş-Bolyai. Mathematica*, vol. 43, no. 3, pp. 89–124, 1998.
- [22] Th. M. Rassias, "The problem of S. M. Ulam for approximately multiplicative mappings," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 352–378, 2000.
- [23] Th. M. Rassias, "On the stability of functional equations in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 1, pp. 264–284, 2000.
- [24] Th. M. Rassias, "On the stability of functional equations and a problem of Ulam," *Acta Applicandae Mathematicae*, vol. 62, no. 1, pp. 23–130, 2000.
- [25] Th. M. Rassias and P. Šemrl, "On the behavior of mappings which do not satisfy Hyers-Ulam stability," *Proceedings of the American Mathematical Society*, vol. 114, no. 4, pp. 989–993, 1992.
- [26] Th. M. Rassias and P. Šemrl, "On the Hyers-Ulam stability of linear mappings," *Journal of Mathematical Analysis and Applications*, vol. 173, no. 2, pp. 325–338, 1993.
- [27] Th. M. Rassias and K. Shibata, "Variational problem of some quadratic functionals in complex analysis," *Journal of Mathematical Analysis and Applications*, vol. 228, no. 1, pp. 234–253, 1998.
- [28] L. Cădariu and V. Radu, "Fixed points and the stability of Jensen's functional equation," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 4, no. 1, article 4, 2003.
- [29] J. B. Diaz and B. Margolis, "A fixed point theorem of the alternative, for contractions on a generalized complete metric space," *Bulletin of the American Mathematical Society*, vol. 74, pp. 305–309, 1968.
- [30] G. Isac and Th. M. Rassias, "Stability of ψ -additive mappings: applications to nonlinear analysis," *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 2, pp. 219–228, 1996.
- [31] L. Cădariu and V. Radu, "On the stability of the Cauchy functional equation: a fixed point approach," in *Iteration Theory (ECIT '02)*, vol. 346 of *Grazer Mathematische Berichte*, pp. 43–52, Karl-Franzens-Universität Graz, Graz, Austria, 2004.
- [32] L. Cădariu and V. Radu, "Fixed point methods for the generalized stability of functional equations in a single variable," *Fixed Point Theory and Applications*, vol. 2008, Article ID 749392, 15 pages, 2008.
- [33] M. Mirzavaziri and M. S. Moslehian, "A fixed point approach to stability of a quadratic equation," *Bulletin of the Brazilian Mathematical Society*, vol. 37, no. 3, pp. 361–376, 2006.
- [34] C. Park, "Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equations in Banach algebras," *Fixed Point Theory and Applications*, vol. 2007, Article ID 50175, 15 pages, 2007.
- [35] C. Park, "Generalized Hyers-Ulam stability of quadratic functional equations: a fixed point approach," *Fixed Point Theory and Applications*, vol. 2008, Article ID 493751, 9 pages, 2008.
- [36] V. Radu, "The fixed point alternative and the stability of functional equations," *Fixed Point Theory*, vol. 4, no. 1, pp. 91–96, 2003.
- [37] M. Eshaghi Gordji, S. Kaboli-Gharetapeh, C. Park, and S. Zolfaghari, "Stability of an additive-cubic-quartic functional equation," *Advances in Difference Equations*, vol. 2009, Article ID 395693, 20 pages, 2009.
- [38] M. Eshaghi Gordji, S. Abbaszadeh, and C. Park, "On the stability of a generalized quadratic and quartic type functional equation in quasi-Banach spaces," *Journal of Inequalities and Applications*, vol. 2009, Article ID 153084, 26 pages, 2009.
- [39] D. Mihet and V. Radu, "On the stability of the additive Cauchy functional equation in random normed spaces," *Journal of Mathematical Analysis and Applications*, vol. 343, no. 1, pp. 567–572, 2008.