Research Article

Strong Convergence Theorems for Equilibrium Problems and k -Strict Pseudocontractions in Hilbert Spaces

Dao-Jun Wen

*School of Mathematics and Statistics, Chongqing Technology and Business University,
Chongqing 400067, China*

Correspondence should be addressed to Dao-Jun Wen, wendaojun@ctbu.edu.cn

Received 23 March 2011; Revised 31 May 2011; Accepted 23 June 2011

Academic Editor: Ljubisa Kocinac

Copyright © 2011 Dao-Jun Wen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce a new iterative scheme for finding a common element of the set of solutions of an equilibrium problem and the set of common fixed point of a finite family of k -strictly pseudocontractive nonself-mappings. Strong convergence theorems are established in a real Hilbert space under some suitable conditions. Our theorems presented in this paper improve and extend the corresponding results announced by many others.

1. Introduction

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, respectively. Let K be a nonempty closed convex subset of H . Let F be a bifunction from $K \times K$ into $\mathbb{R}$, where $\mathbb{R}$ denotes the set of real numbers. We consider the following problem: Find $x \in K$ such that

$$F(x, y) \geq 0, \quad \forall y \in K, \quad (1.1)$$

which is called equilibrium problem. We use $EP(F)$ to denote the set of solution of the problem (1.1). Given a mapping $T : K \rightarrow H$, let $F(x, y) = \langle Tx, y - x \rangle$ for all $x, y \in K$. Then, $z \in EP(F)$ if and only if $\langle Tz, y - z \rangle \geq 0$ for all $y \in K$; that is, z is a solution of the variational inequality. Numerous problems in physics, optimization, and economics reduce to find a solution of (1.1). Some methods have been proposed to solve the equilibrium problem (see, e.g., [1–3]).

Recall that a nonself-mapping $T : K \rightarrow H$ is called a k -strict pseudocontraction if there exists a constant $k \in [0, 1)$ such that

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + k\|(I - T)x - (I - T)y\|^2, \quad \forall x, y \in K. \quad (1.2)$$

We use $F(T)$ to denote the fixed point set of the mapping T , that is, $F(T) := \{x \in K : Tx = x\}$. As $k = 0$, T is said to be nonexpansive, that is, $\|Tx - Ty\| \leq \|x - y\|$, for all $x, y \in K$. T is said to be pseudocontractive if $k = 1$ and is also said to be strongly pseudocontractive if there exists a positive constant $\lambda \in (0, 1)$ such that $T + \lambda I$ is pseudocontractive. Clearly, the class of k -strict pseudocontractions falls into the one between classes of nonexpansive mappings and pseudocontractions. We remark also that the class of strongly pseudocontractive mappings is independent of the class of k -strict pseudocontractions (see, e.g., [4, 5]).

Iterative methods for equilibrium problem and nonexpansive mappings have been extensively investigated; see, for example, [1–18] and the references therein. However, iterative methods for strict pseudocontractions are far less developed than those for nonexpansive mappings though Browder and Petryshyn [5] initiated their work in 1967; the reason is probably that the second term appearing in the right-hand side of (1.2) impedes the convergence analysis for iterative algorithms used to find a fixed point of the strict pseudocontraction T . On the other hand, strict pseudocontractions have more powerful applications than nonexpansive mappings do in solving inverse problems (see, e.g., [6]). Therefore, it is interesting to develop the theory of iterative methods for equilibrium problem and strict pseudocontractions.

In 2007, Acedo and Xu [12] proposed the following parallel algorithm for a finite family of k_i -strict pseudocontractions $\{T_i\}_{i=1}^N$ in Hilbert space H :

$$\forall x_0 \in K, \quad x_{n+1} = (1 - \alpha_n)x_n + \alpha_n \sum_{i=1}^N \lambda_i T_i x_n, \quad (1.3)$$

where $\{\alpha_n\} \subset (0, 1)$, and $\{\lambda_i\}_{i=1}^N$ is a finite sequence of positive numbers such that $\sum_{i=1}^N \lambda_i = 1$. They proved that the sequence $\{x_n\}$ defined by (1.3) converges weakly to a common fixed point of $\{T_i\}_{i=1}^N$ under some appropriate conditions. Moreover, by applying additional projections, they further proved that algorithm can be modified to have strong convergence.

Recently, S. Takahashi and W. Takahashi [13] studied the equilibrium problem and fixed point of nonexpansive self-mappings T in Hilbert spaces by a viscosity approximation methods for finding an element of $\text{EP}(F) \cap F(T)$. Very recently, by using the general approximation method, Qin et al. [14] obtained a strong convergence theorem for finding an element of $F(T)$. On the other hand, Ceng et al. [16] proposed an iterative scheme for finding an element of $\text{EP}(F) \cap F(T)$ and then obtained some weak and strong convergence theorems.

In this paper, inspired and motivated by research going in this area, we introduce a modified parallel iteration, which is defined in the following way:

$$F(u_n, y) + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in K,$$

$$\begin{aligned}
y_n &= \alpha_n u_n + (1 - \alpha_n) \sum_{i=1}^N \eta_i^{(n)} T_i u_n, \\
x_{n+1} &= \beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n) y_n, \quad n \geq 0,
\end{aligned} \tag{1.4}$$

where $u \in K$ is a given point, $\{T_i\}_{i=1}^N : K \rightarrow H$ is a finite family of k_i -strictly pseudocontractive nonself-mappings, $\{\eta_i^{(n)}\}_{i=1}^N$ is a finite sequences of positive numbers, $\{\alpha_n\}$, $\{\beta_n\}$, and $\{\gamma_n\}$ are some sequences in $(0,1)$.

Our purpose is not only to modify the parallel algorithm (1.3) to the case of equilibrium problems and common fixed point for a finite family of k_i -strictly pseudocontractive nonself-mappings, but also to establish strong convergence theorems in a real Hilbert space under some different conditions. Our theorems presented in this paper improve and extend the main results of [9, 12–14, 16].

2. Preliminaries

Let K be a nonempty closed and convex subset of a Hilbert space H . We use P_K to denote the metric or nearest point projection of H onto K ; that is, for $x \in H$, $P_K x$ is the only point in K such that $\|x - P_K x\| = \inf\{\|x - z\| : z \in K\}$. we write $x_n \rightharpoonup x$ and $x_n \rightarrow x$ indicate that the sequence $\{x_n\}$ convergence weakly and strongly to x , respectively.

It is well known that Hilbert space H satisfies Opial's condition [8], that is, for any sequence $\{x_n\}$ with $x_n \rightharpoonup x$ and every $y \in H$ with $y \neq x$, we have

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\|. \tag{2.1}$$

To study the equilibrium problem (1.1), we may assume that the bifunction F of $K \times K$ into $\mathbb{R}$ satisfies the following conditions.

- (A1) $F(x, x) = 0$ for all $x \in K$.
- (A2) F is monotone, that is, $F(x, y) + F(y, x) \leq 0$ for all $x, y \in K$.
- (A3) For each $x, y, z \in K$, $\lim_{t \rightarrow 0} F(tz + (1-t)x, y) \leq F(x, y)$.
- (A4) For each $x \in K$, $y \mapsto F(x, y)$ is convex and lower semi-continuous.

In order to prove our main results, we need the following Lemmas and Propositions.

Lemma 2.1 (see [1, 3]). *Let F be a bifunction from $K \times K$ into $\mathbb{R}$ satisfying (A1)–(A4). Then, for any $r > 0$ and $x \in H$, there exists $z \in K$ such that*

$$F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in K. \tag{2.2}$$

Further, if $T_r x = \{z \in K : F(z, y) + (1/r) \langle y - z, z - x \rangle \geq 0, \text{ for all } y \in K\}$, then the following holds.

- (1) T_r is single-valued.
- (2) T_r is firmly nonexpansive, that is, $\|T_r x - T_r y\|^2 \leq \langle T_r x - T_r y, x - y \rangle$, for all $x, y \in H$.
- (3) $F(T_r) = EP(F)$.
- (4) $EP(F)$ is closed and convex.

Lemma 2.2 (see [7]). Let $(E, \langle \cdot, \cdot \rangle)$ be an inner product space. Then, for all $x, y, z \in E$ and $\alpha, \beta, \gamma \in [0, 1]$ with $\alpha + \beta + \gamma = 1$, we have

$$\|\alpha x + \beta y + \gamma z\|^2 = \alpha \|x\|^2 + \beta \|y\|^2 + \gamma \|z\|^2 - \alpha\beta \|x - y\|^2 - \alpha\gamma \|x - z\|^2 - \beta\gamma \|y - z\|^2. \quad (2.3)$$

Lemma 2.3 (see [19]). Let $\{x_n\}$ and $\{z_n\}$ be bounded sequence in Banach space E , and let $\{\lambda_n\}$ be a sequence in $[0, 1]$ such that $0 < \liminf_{n \rightarrow \infty} \lambda_n \leq \limsup_{n \rightarrow \infty} \lambda_n < 1$. Suppose $x_{n+1} = \lambda_n x_n + (1 - \lambda_n)z_n$ and

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0, \quad \forall n \geq 0. \quad (2.4)$$

Then $\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0$.

Lemma 2.4 (see [2, 10]). Let $T : K \rightarrow H$ be a k -strict pseudocontraction. For $\lambda \in [k, 1)$, define $S : K \rightarrow H$ by $Sx = \lambda x + (1 - \lambda)Tx$ for each $x \in K$. Then, as $\lambda \in [k, 1)$, S is a nonexpansive mapping such that $F(S) = F(T)$.

Lemma 2.5 (see [10]). If $T : K \rightarrow H$ is a k -strict pseudocontraction, then the fixed point set $F(T)$ is closed convex so that the projection $P_{F(T)}$ is well defined.

Lemma 2.6 (see [9]). Let K be a nonempty bounded closed convex subset of H . Given $x \in H$ and $z \in K$, then $z = P_K x$ if and only if there holds the relation:

$$\langle x - z, z - y \rangle \geq 0, \quad \forall y \in K. \quad (2.5)$$

Lemma 2.7 (see [20]). Assume $\{a_n\}$ is a sequence of nonnegative real numbers such that

$$a_{n+1} \leq (1 - \gamma_n)a_n + \gamma_n \delta_n, \quad n \geq 0, \quad (2.6)$$

where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a real sequence such that

- (i) $\sum_{n=0}^{\infty} \gamma_n = \infty$,
- (ii) $\limsup_{n \rightarrow \infty} \delta_n \leq 0$ or $\sum_{n=0}^{\infty} |\gamma_n \delta_n| < \infty$.

Then $\lim_{n \rightarrow \infty} a_n = 0$.

Proposition 2.8 (see, e.g., Acedo and Xu [12]). Let K be a nonempty closed convex subset of Hilbert space H . Given an integer $N \geq 1$, assume that $\{T_i\}_{i=1}^N : K \rightarrow H$ is a finite family of k_i -strict pseudocontractions. Suppose that $\{\lambda_i\}_{i=1}^N$ is a positive sequence such that $\sum_{i=1}^N \lambda_i = 1$. Then $\sum_{i=1}^N \lambda_i T_i$ is a k -strict pseudocontraction with $k = \max\{k_i : 1 \leq i \leq N\}$.

Proposition 2.9 (see, e.g., Acedo and Xu [12]). Let $\{T_i\}_{i=1}^N$ and $\{\lambda_i\}_{i=1}^N$ be given as in Proposition 2.8 above. Then $F(\sum_{i=1}^N \lambda_i T_i) = \bigcap_{i=1}^N F(T_i)$.

3. Main Results

Theorem 3.1. Let K be a nonempty closed convex subset of Hilbert space H , and let F be a bifunction from $K \times K$ into $\mathbb{R}$ satisfying (A1)–(A4). Let $\{T_i\}_{i=1}^N : K \rightarrow H$ be a finite family of k_i -strict pseudocontractions such that $k = \max\{k_i : 1 \leq i \leq N\}$ and $\mathcal{F} = \bigcap_{i=1}^N F(T_i) \cap \text{EP}(F) \neq \emptyset$. Assume $\{\eta_i^{(n)}\}_{i=1}^N$ is a finite sequences of positive numbers such that $\sum_{i=1}^N \eta_i^{(n)} = 1$ for all $n \geq 0$. Given $u \in K$ and $x_0 \in K$, $\{\alpha_n\}$, $\{\beta_n\}$, and $\{\gamma_n\}$ are some sequences in $(0,1)$; the following control conditions are satisfied.

- (i) $k \leq \alpha_n \leq \lambda < 1$ for all $n \geq 0$ and $\lim_{n \rightarrow \infty} \alpha_n = \alpha$,
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=0}^{\infty} \beta_n = \infty$,
- (iii) $0 < \liminf_{n \rightarrow \infty} \gamma_n \leq \limsup_{n \rightarrow \infty} \gamma_n < 1$,
- (iv) $\lim_{n \rightarrow \infty} |\eta_i^{(n+1)} - \eta_i^{(n)}| = 0$.

Then the sequence $\{x_n\}$ generated by (1.4) converges strongly to $q \in \mathcal{F}$, where $q = P_{\mathcal{F}}u$.

Proof. From Lemma 2.1, we see that $\text{EP}(F) = F(T_r)$, and note that u_n can be rewritten as $u_n = T_r x_n$. Putting $A_n = \sum_{i=1}^N \eta_i^{(n)} T_i$, we have $A_n : K \rightarrow H$ is a k -strict pseudocontraction and $F(A_n) = \bigcap_{i=1}^N F(T_i)$ by Propositions 2.8 and 2.9, where $k = \max\{k_i : 1 \leq i \leq N\}$.

From (1.4), condition (i), and Lemma 2.2, taking a point $p \in \mathcal{F}$, we have

$$\begin{aligned}
 \|y_n - p\|^2 &= \|\alpha_n(u_n - p) + (1 - \alpha_n)(A_n u_n - p)\|^2 \\
 &= \alpha_n \|u_n - p\|^2 + (1 - \alpha_n) \|A_n u_n - p\|^2 - \alpha_n(1 - \alpha_n) \|u_n - A_n u_n\|^2 \\
 &\leq \alpha_n \|u_n - p\|^2 + (1 - \alpha_n) \left[\|u_n - p\|^2 + k \|u_n - A_n u_n\|^2 \right] \\
 &\quad - \alpha_n(1 - \alpha_n) \|u_n - A_n u_n\|^2 \\
 &= \|u_n - p\|^2 - (1 - \alpha_n)(\alpha_n - k) \|u_n - A_n u_n\|^2 \\
 &\leq \|T_r x_n - T_r p\|^2 \\
 &\leq \|x_n - p\|^2.
 \end{aligned} \tag{3.1}$$

Furthermore, we have

$$\|y_n - p\| \leq \|u_n - p\| \leq \|x_n - p\|. \tag{3.2}$$

It follows from (1.4) and (3.2) that

$$\begin{aligned}
\|x_{n+1} - p\| &= \|\beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n)y_n - p\| \\
&\leq \beta_n \|u - p\| + \gamma_n \|x_n - p\| + (1 - \beta_n - \gamma_n) \|y_n - p\| \\
&\leq \beta_n \|u - p\| + (1 - \beta_n) \|x_n - p\| \\
&\leq \max\{\|u - p\|, \|x_0 - p\|\}.
\end{aligned} \tag{3.3}$$

Consequently, sequence $\{x_n\}$ is bounded and so are $\{u_n\}$ and $\{y_n\}$.

Define a mapping $T_n x := \alpha_n x + (1 - \alpha_n)A_n x$ for each $x \in K$. Then $T_n : K \rightarrow H$ is nonexpansive. Indeed, by using (1.1), condition (i), and Lemma 2.2, we have for all $x, y \in K$ that

$$\begin{aligned}
\|T_n x - T_n y\|^2 &= \|\alpha_n(x - y) + (1 - \alpha_n)(A_n x - A_n y)\|^2 \\
&= \alpha_n \|x - y\|^2 + (1 - \alpha_n) \|A_n x - A_n y\|^2 \\
&\quad - \alpha_n(1 - \alpha_n) \|x - A_n x - (y - A_n y)\|^2 \\
&\leq \alpha_n \|x - y\|^2 + (1 - \alpha_n) \left[\|x - y\|^2 + k \|x - A_n x - (y - A_n y)\|^2 \right] \\
&\quad - \alpha_n(1 - \alpha_n) \|x - A_n x - (y - A_n y)\|^2 \\
&= \|x - y\|^2 - (1 - \alpha_n)(\alpha_n - k) \|x - A_n x - (y - A_n y)\|^2 \\
&\leq \|x - y\|^2,
\end{aligned} \tag{3.4}$$

which shows that $T_n : K \rightarrow H$ is nonexpansive.

Next we show that $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$. Setting $x_{n+1} = \gamma_n x_n + (1 - \gamma_n)z_n$, we have

$$\begin{aligned}
z_{n+1} - z_n &= \frac{x_{n+2} - \gamma_{n+1}x_{n+1}}{1 - \gamma_{n+1}} - \frac{x_{n+1} - \gamma_n x_n}{1 - \gamma_n} \\
&= \frac{\beta_{n+1}u + (1 - \beta_{n+1} - \gamma_{n+1})y_{n+1}}{1 - \gamma_{n+1}} - \frac{\beta_n u + (1 - \beta_n - \gamma_n)y_n}{1 - \gamma_n} \\
&= \frac{\beta_{n+1}}{1 - \gamma_{n+1}}(u - y_{n+1}) + (y_{n+1} - y_n) - \frac{\beta_n}{1 - \gamma_n}(u - y_n).
\end{aligned} \tag{3.5}$$

It follows that

$$\|z_{n+1} - z_n\| \leq \frac{\beta_{n+1}}{1 - \gamma_{n+1}} \|u - y_{n+1}\| + \|y_{n+1} - y_n\| + \frac{\beta_n}{1 - \gamma_n} \|u - y_n\|. \tag{3.6}$$

From (1.4), we have $y_n = T_n u_n$ and

$$\begin{aligned}
 \|y_{n+1} - y_n\| &\leq \|T_{n+1} u_{n+1} - T_{n+1} u_n\| + \|T_{n+1} u_n - T_n u_n\| \\
 &\leq \|u_{n+1} - u_n\| + \|\alpha_{n+1} u_n + (1 - \alpha_{n+1}) A_{n+1} u_n - [\alpha_n u_n + (1 - \alpha_n) A_n u_n]\| \\
 &\leq \|u_{n+1} - u_n\| + |\alpha_{n+1} - \alpha_n| \|u_n - A_n u_n\| + (1 - \alpha_{n+1}) \|A_{n+1} u_n - A_n u_n\| \\
 &\leq \|u_{n+1} - u_n\| + |\alpha_{n+1} - \alpha_n| \|u_n - A_n u_n\| + (1 - \alpha_{n+1}) \sum_{i=1}^N \left| \eta_i^{(n+1)} - \eta_i^{(n)} \right| \|T_i u_n\|.
 \end{aligned} \tag{3.7}$$

By Lemma 2.1, $u_n = T_r x_n$ and $u_{n+1} = T_r x_{n+1}$, we have

$$F(u_n, y) + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in K, \tag{3.8}$$

$$F(u_{n+1}, y) + \frac{1}{r} \langle y - u_{n+1}, u_{n+1} - x_{n+1} \rangle \geq 0, \quad \forall y \in K. \tag{3.9}$$

Putting $y = u_{n+1}$ in (3.8) and $y = u_n$ in (3.9), we obtain

$$\begin{aligned}
 F(u_n, u_{n+1}) + \frac{1}{r} \langle u_{n+1} - u_n, u_n - x_n \rangle &\geq 0, \\
 F(u_{n+1}, u_n) + \frac{1}{r} \langle u_n - u_{n+1}, u_{n+1} - x_{n+1} \rangle &\geq 0.
 \end{aligned} \tag{3.10}$$

So, from (A2) and $r > 0$, we have

$$\langle u_{n+1} - u_n, u_n - u_{n+1} + u_{n+1} - x_n - (u_{n+1} - x_{n+1}) \rangle \geq 0, \tag{3.11}$$

and hence

$$\|u_{n+1} - u_n\|^2 \leq \langle u_{n+1} - u_n, x_{n+1} - x_n \rangle, \tag{3.12}$$

which implies that

$$\|u_{n+1} - u_n\| \leq \|x_{n+1} - x_n\|. \tag{3.13}$$

Combining (3.6), (3.7), and (3.13), we have

$$\begin{aligned}
 \|z_{n+1} - z_n\| &\leq \frac{\beta_{n+1}}{1 - \gamma_{n+1}} \|u - y_{n+1}\| + \|x_{n+1} - x_n\| + \frac{\beta_n}{1 - \gamma_n} \|u - y_n\| \\
 &\quad + |\alpha_{n+1} - \alpha_n| \|u_n - A_n u_n\| + (1 - \alpha_{n+1}) \sum_{i=1}^N \left| \eta_i^{(n+1)} - \eta_i^{(n)} \right| \|T_i u_n\|.
 \end{aligned} \tag{3.14}$$

This together with (i), (ii) and (iv) imply that

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0. \quad (3.15)$$

Hence, by Lemma 2.3, we obtain

$$\lim_{n \rightarrow \infty} \|z_n - x_n\| = 0. \quad (3.16)$$

Consequently,

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = \lim_{n \rightarrow \infty} (1 - \gamma_n) \|z_n - x_n\| = 0. \quad (3.17)$$

On the other hand, by (1.4) and (iii), we have

$$\|x_{n+1} - y_n\| \leq \beta_n \|u - y_n\| + \gamma_n \|x_n - x_{n+1}\| + \gamma_n \|x_{n+1} - y_n\|, \quad (3.18)$$

which implies that

$$\|x_{n+1} - y_n\| \leq \frac{\beta_n}{1 - \gamma_n} \|u - y_n\| + \frac{\gamma_n}{1 - \gamma_n} \|x_n - x_{n+1}\|. \quad (3.19)$$

Combining (ii), (3.17), and (3.19), we have

$$\lim_{n \rightarrow \infty} \|x_{n+1} - y_n\| = 0. \quad (3.20)$$

Note that

$$\|x_n - y_n\| \leq \|x_n - x_{n+1}\| + \|x_{n+1} - y_n\|, \quad (3.21)$$

which together with (3.17) and (3.20) implies

$$\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0. \quad (3.22)$$

Moreover, for $p \in \mathcal{F} = \bigcap_{i=1}^N F(T_i) \cap \text{EP}(F)$, we have

$$\begin{aligned} \|u_n - p\|^2 &= \|T_r x_n - T_r p\|^2 \leq \langle x_n - p, u_n - p \rangle \\ &= \frac{1}{2} (\|x_n - p\|^2 + \|u_n - p\|^2 - \|x_n - u_n\|^2), \end{aligned} \quad (3.23)$$

and hence

$$\|u_n - p\|^2 \leq \|x_n - p\|^2 - \|x_n - u_n\|^2. \quad (3.24)$$

From Lemma 2.2, (3.2) and (3.24), we have

$$\begin{aligned}
\|x_{n+1} - p\|^2 &= \|\beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n) y_n - p\|^2 \\
&\leq \beta_n \|u - p\|^2 + \gamma_n \|x_n - p\|^2 + (1 - \beta_n - \gamma_n) \|y_n - p\|^2 \\
&\leq \beta_n \|u - p\|^2 + \gamma_n \|x_n - p\|^2 + (1 - \beta_n - \gamma_n) (\|x_n - p\|^2 - \|x_n - u_n\|^2) \\
&\leq \beta_n \|u - p\|^2 + \|x_n - p\|^2 - (1 - \beta_n - \gamma_n) \|x_n - u_n\|^2,
\end{aligned} \tag{3.25}$$

and hence

$$\begin{aligned}
(1 - \beta_n - \gamma_n) \|x_n - u_n\|^2 &\leq \beta_n \|u - p\|^2 + \|x_n - p\|^2 - \|x_{n+1} - p\|^2 \\
&\leq \beta_n \|u - p\|^2 + \|x_n - x_{n+1}\| (\|x_n - p\| + \|x_{n+1} - p\|).
\end{aligned} \tag{3.26}$$

By (ii) and (3.17), we obtain

$$\lim_{n \rightarrow \infty} \|x_n - u_n\| = 0. \tag{3.27}$$

It follows from (3.22) and (3.27) that

$$\lim_{n \rightarrow \infty} \|y_n - u_n\| = 0. \tag{3.28}$$

Define $S_n : K \rightarrow H$ by $S_n x = \alpha x + (1 - \alpha) A_n x$. Then, S_n is a nonexpansive with $F(S_n) = F(A_n)$ by Lemma 2.4. Note that $\lim_{n \rightarrow \infty} \alpha_n = \alpha \in [k, 1)$ by condition (i) and

$$\begin{aligned}
\|u_n - S_n u_n\| &\leq \|u_n - y_n\| + \|y_n - S_n u_n\| \\
&\leq \|u_n - y_n\| + \|\alpha_n u_n + (1 - \alpha_n) A_n u_n - [\alpha u_n + (1 - \alpha) A_n u_n]\| \\
&\leq \|u_n - y_n\| + |\alpha_n - \alpha| \|u_n - A_n u_n\|,
\end{aligned} \tag{3.29}$$

which combines with condition (i) and (3.28) yielding that

$$\lim_{n \rightarrow \infty} \|u_n - S_n u_n\| = 0. \tag{3.30}$$

We now show that $\limsup_{n \rightarrow \infty} \langle u - q, x_n - q \rangle \leq 0$, where $q = P_{\mathcal{F}} u$. To see this, we choose a subsequence $\{x_{n_i}\}$ of $\{x_n\}$ such that

$$\limsup_{n \rightarrow \infty} \langle u - q, x_n - q \rangle = \lim_{i \rightarrow \infty} \langle u - q, x_{n_i} - q \rangle. \tag{3.31}$$

Since $\{u_{n_i}\}$ is bounded, there exists a subsequence $\{u_{n_{i_j}}\}$ of $\{u_{n_i}\}$ converging weakly to u^* . Without loss of generality, we assume that $u_{n_i} \rightharpoonup u^*$ as $i \rightarrow \infty$. From (3.27), we obtain $x_{n_i} \rightarrow u^*$

as $i \rightarrow \infty$. Since K is closed and convex, K is weakly closed. So, we have $u^* \in K$ and $u^* \in F(S_n)$. Otherwise, from $u^* \neq S_n u^*$ and Opial's condition, we obtain

$$\begin{aligned} \liminf_{i \rightarrow \infty} \|u_{n_i} - u^*\| &< \liminf_{i \rightarrow \infty} \|u_{n_i} - S_{n_i} u^*\| \\ &\leq \liminf_{i \rightarrow \infty} (\|u_{n_i} - S_{n_i} u_{n_i}\| + \|S_{n_i} u_{n_i} - S_{n_i} u^*\|) \\ &\leq \liminf_{i \rightarrow \infty} \|u_{n_i} - u^*\|. \end{aligned} \quad (3.32)$$

This is a contradiction. Hence, we get $u^* \in F(S_n) = F(A_n)$. Moreover, by $u_n = T_r x_n$, we have

$$F(u_n, y) + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in K. \quad (3.33)$$

It follows from (A2) that

$$\frac{1}{r} \langle y - u_n, u_n - x_n \rangle \geq F(y, u_n). \quad (3.34)$$

Replacing n by n_i , we have

$$\frac{1}{r} \langle y - u_{n_i}, u_{n_i} - x_{n_i} \rangle \geq F(y, u_{n_i}). \quad (3.35)$$

Since $u_{n_i} - x_{n_i} \rightarrow 0$ and $u_{n_i} \rightharpoonup u^*$, it follows from (A4) that $F(y, u^*) \leq 0$ for all $y \in K$. Put $z_t = ty + (1-t)u^*$ for all $t \in (0, 1]$ and $y \in K$. Then, we have $z_t \in K$, and hence, $F(z_t, u^*) \leq 0$. By (A1) and (A4), we have

$$0 = F(z_t, z_t) \leq tF(z_t, y) + (1-t)F(z_t, u^*) \leq tF(z_t, y), \quad (3.36)$$

which implies $F(z_t, y) \geq 0$. From (A3), we have $F(u^*, y) \geq 0$ for all $y \in K$, and hence, $u^* \in \text{EP}(F)$. Therefore, $u^* \in F(S_n) \cap \text{EP}(F)$. From Lemma 2.6, we know that

$$\langle u - P_{\mathcal{F}} u, u^* - P_{\mathcal{F}} u \rangle \leq 0. \quad (3.37)$$

It follows from (3.31) and (3.37) that

$$\limsup_{n \rightarrow \infty} \langle u - q, x_n - q \rangle = \lim_{i \rightarrow \infty} \langle u - q, x_{n_i} - q \rangle = \langle u - q, u^* - q \rangle \leq 0. \quad (3.38)$$

Finally, we prove that $x_n \rightarrow q = P_{\mathcal{F}}u$ as $n \rightarrow \infty$. From (1.4) again, we have

$$\begin{aligned}
\|x_{n+1} - q\|^2 &= \langle \beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n)y_n - q, x_{n+1} - q \rangle \\
&= \beta_n \langle u - q, x_{n+1} - q \rangle + \gamma_n \langle x_n - q, x_{n+1} - q \rangle + (1 - \beta_n - \gamma_n) \langle y_n - q, x_{n+1} - q \rangle \\
&\leq \beta_n \langle u - q, x_{n+1} - q \rangle + \gamma_n \|x_n - q\| \|x_{n+1} - q\| + (1 - \beta_n - \gamma_n) \|y_n - q\| \|x_{n+1} - q\| \\
&\leq (1 - \beta_n) \|x_n - q\| \|x_{n+1} - q\| + \beta_n \langle u - q, x_{n+1} - q \rangle \\
&\leq \frac{1 - \beta_n}{2} (\|x_n - q\|^2 + \|x_{n+1} - q\|^2) + \beta_n \langle u - q, x_{n+1} - q \rangle \\
&\leq \frac{1 - \beta_n}{2} \|x_n - q\|^2 + \frac{1}{2} \|x_{n+1} - q\|^2 + \beta_n \langle u - q, x_{n+1} - q \rangle,
\end{aligned} \tag{3.39}$$

which implies that

$$\|x_{n+1} - q\|^2 \leq (1 - \beta_n) \|x_n - q\|^2 + 2\beta_n \langle u - q, x_{n+1} - q \rangle. \tag{3.40}$$

It follows from (3.38), (3.40), and Lemma 2.7 that $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$. This completes the proof. $\square$

As $N = 1$, that is, $A_n = T$ and $\eta_i^{(n)} \equiv 1$ in Theorem 3.1, we have the following results immediately.

Theorem 3.2. *Let K be a nonempty closed convex subset of Hilbert space H , and let F be a bifunction from $K \times K$ into $\mathbb{R}$ satisfying (A1)–(A4). Let $T : K \rightarrow H$ be a k -strict pseudocontractions such that $\mathcal{F} = F(T) \cap \text{EP}(F) \neq \emptyset$. Let $\{x_n\}$ be a sequence generated in the following manner:*

$$\begin{aligned}
F(u_n, y) + \frac{1}{r} \langle y - u_n, u_n - x_n \rangle &\geq 0, \quad \forall y \in K, \\
y_n &= \alpha_n u_n + (1 - \alpha_n) T u_n, \\
x_{n+1} &= \beta_n u + \gamma_n x_n + (1 - \beta_n - \gamma_n) y_n, \quad n \geq 0,
\end{aligned} \tag{3.41}$$

where $u \in K$ and $x_0 \in K$, $\{\alpha_n\}$, $\{\beta_n\}$, and $\{\gamma_n\}$ are some sequences in $(0, 1)$. If the following control conditions are satisfied:

- (i) $k \leq \alpha_n \leq \lambda < 1$ for all $n \geq 0$ and $\lim_{n \rightarrow \infty} \alpha_n = \alpha$,
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$ and $\sum_{n=0}^{\infty} \beta_n = \infty$,
- (iii) $0 < \liminf_{n \rightarrow \infty} \gamma_n \leq \limsup_{n \rightarrow \infty} \gamma_n < 1$,

then $\{x_n\}$ converges strongly to $q \in \mathcal{F}$, where $q = P_{\mathcal{F}}u$.

Acknowledgment

This work was supported by the National Science Foundation of China (11001287), the Natural Science Foundation Project of Chongqing (CSTC 2010BB9254), and the Science and Technology Research Project of Chongqing Municipal Education Commission (KJ 110701).

References

- [1] E. Blum and W. Oettli, "From optimization and variational inequalities to equilibrium problems," *The Mathematics Student*, vol. 63, no. 1–4, pp. 123–145, 1994.
- [2] A. Moudafi and M. Théra, "Proximal and dynamical approaches to equilibrium problems," in *Ill-Posed Variational Problems and Regularization Techniques*, vol. 477 of *Lecture Notes in Economics and Mathematical Systems*, pp. 187–201, Springer, Berlin, Germany, 1999.
- [3] P. L. Combettes and A. Hirstoaga, "Equilibrium programming in Hilbert spaces," *Journal of Nonlinear and Convex Analysis*, vol. 6, no. 1, pp. 117–136, 2005.
- [4] F. E. Browder, "Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces," *Archive for Rational Mechanics and Analysis*, vol. 24, pp. 82–90, 1967.
- [5] F. E. Browder and W. V. Petryshyn, "Construction of fixed points of nonlinear mappings in Hilbert space," *Journal of Mathematical Analysis and Applications*, vol. 20, pp. 197–228, 1967.
- [6] O. Scherzer, "Convergence criteria of iterative methods based on Landweber iteration for solving nonlinear problems," *Journal of Mathematical Analysis and Applications*, vol. 194, no. 3, pp. 911–933, 1995.
- [7] M. O. Osilike and D. I. Igbokwe, "Weak and strong convergence theorems for fixed points of pseudocontractions and solutions of monotone type operator equations," *Computers & Mathematics with Applications*, vol. 40, no. 4–5, pp. 559–567, 2000.
- [8] W. Takahashi and M. Toyoda, "Weak convergence theorems for nonexpansive mappings and monotone mappings," *Journal of Optimization Theory and Applications*, vol. 118, no. 2, pp. 417–428, 2003.
- [9] G. Marino and H.-K. Xu, "Weak and strong convergence theorems for strict pseudo-contractions in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 336–346, 2007.
- [10] H. Zhou, "Convergence theorems of fixed points for k -strict pseudo-contractions in Hilbert spaces," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 69, no. 2, pp. 456–462, 2008.
- [11] T.-H. Kim and H.-K. Xu, "Strong convergence of modified Mann iterations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 61, no. 1–2, pp. 51–60, 2005.
- [12] G. L. Acedo and H.-K. Xu, "Iterative methods for strict pseudo-contractions in Hilbert spaces," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 67, no. 7, pp. 2258–2271, 2007.
- [13] S. Takahashi and W. Takahashi, "Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 331, no. 1, pp. 506–515, 2007.
- [14] X. Qin, M. Shang, and S. M. Kang, "Strong convergence theorems of modified Mann iterative process for strict pseudo-contractions in Hilbert spaces," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 70, no. 3, pp. 1257–1264, 2009.
- [15] H. Zhang and Y. Su, "Strong convergence theorems for strict pseudo-contractions in q -uniformly smooth Banach spaces," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 70, no. 9, pp. 3236–3242, 2009.
- [16] L.-C. Ceng, S. Al-Homidan, Q. H. Ansari, and J.-C. Yao, "An iterative scheme for equilibrium problems and fixed point problems of strict pseudo-contraction mappings," *Journal of Computational and Applied Mathematics*, vol. 223, no. 2, pp. 967–974, 2009.
- [17] S. Plubtieng and R. Punpaeng, "A new iterative method for equilibrium problems and fixed point problems of nonexpansive mappings and monotone mappings," *Applied Mathematics and Computation*, vol. 197, no. 2, pp. 548–558, 2008.
- [18] X. Qin, Y. J. Cho, and S. M. Kang, "Viscosity approximation methods for generalized equilibrium problems and fixed point problems with applications," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 1, pp. 99–112, 2010.

- [19] T. Suzuki, "Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter non-expansive semigroups without Bochner integrals," *Journal of Mathematical Analysis and Applications*, vol. 305, no. 1, pp. 227–239, 2005.
- [20] H. K. Xu, "An iterative approach to quadratic optimization," *Journal of Optimization Theory and Applications*, vol. 116, no. 3, pp. 659–678, 2003.