Research Article

Some Identities on the q -Bernoulli Numbers and Polynomials with Weight 0

T. Kim, J. Choi, and Y. H. Kim

Division of General Education-Mathematics, Kwangju University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to J. Choi, jeschoi@kw.ac.kr

Received 16 August 2011; Accepted 27 September 2011

Academic Editor: Josef Diblík

Copyright © 2011 T. Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently, Kim (2011) has introduced the q -Bernoulli numbers with weight α . In this paper, we consider the q -Bernoulli numbers and polynomials with weight $\alpha = 0$ and give p -adic q -integral representation of Bernstein polynomials associated with q -Bernoulli numbers and polynomials with weight 0. From these integral representation on $\mathbb{Z}_p$, we derive some interesting identities on the q -Bernoulli numbers and polynomials with weight 0.

1. Introduction

Let p be a fixed prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will denote the ring of p -adic integers, the field of p -adic rational numbers, and the completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$.

Let $|\cdot|_p$ be a p -adic norm with $|x|_p = p^{-r}$, where $x = p^r s/t$ and $(p, s) = (p, t) = (s, t) = 1$, $r \in \mathbb{Q}$. In this paper, we assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < p^{-1/(p-1)}$ so that $q^x = \exp(x \log q)$, and $[x]_q = (1 - q^x)/(1 - q)$.

Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the p -adic q -integral on $\mathbb{Z}_p$ is defined by Kim as follows:

$$\begin{aligned} I_q(f) &= \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \mu_q(x + p^N \mathbb{Z}_p) \\ &= \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \end{aligned} \quad (1.1)$$

(see [1–5]). For $n \in \mathbb{N}$, let $f_n(x) = f(x + n)$. From (1.1), we note that

$$q^n I_q(f_n) - I_q(f) = (q - 1) \sum_{l=0}^{n-1} q^l f(l) + \frac{q - 1}{\log q} \sum_{l=0}^{n-1} q^l f'(l), \quad (1.2)$$

where $f'(l) = df(x)/dx|_{x=l}$, (see [3, 6, 7]). In the special case, $n = 1$, we get

$$q \int_{\mathbb{Z}_p} f(x+1) d\mu_q(x) - \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = (q-1)f(0) + \frac{q-1}{\log q} f'(0). \quad (1.3)$$

Throughout this paper, we assume that $\alpha \in \mathbb{Q}$.

The q -Bernoulli numbers with weight α are defined by Kim [8] as follows:

$$\tilde{\beta}_{0,q}^{(\alpha)} = 1, \quad q \left(q^\alpha \tilde{\beta}_q^{(\alpha)} + 1 \right)^n - \tilde{\beta}_{n,q}^{(\alpha)} = \begin{cases} \frac{\alpha}{[\alpha]_q} & \text{if } n = 1, \\ 0 & \text{if } n > 1, \end{cases} \quad (1.4)$$

with the usual convention about replacing $(\tilde{\beta}_q^{(\alpha)})^n$ with $\tilde{\beta}_{n,q}^{(\alpha)}$. From (1.4), we can derive the following equation:

$$\begin{aligned} \tilde{\beta}_{n,q}^{(\alpha)} &= \frac{1}{(1-q)^n [\alpha]_q^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{\alpha l + 1}{[\alpha l + 1]_q} \\ &= -\frac{n\alpha}{[\alpha]_q} \sum_{m=0}^{\infty} q^{m\alpha+m} [m]_{q^\alpha}^{n-1} + (1-q) \sum_{m=0}^{\infty} q^m [m]_{q^\alpha}^n. \end{aligned} \quad (1.5)$$

By (1.1), (1.4), and (1.5), we get

$$\tilde{\beta}_{n,q}^{(\alpha)} = \int_{\mathbb{Z}_p} [x]_{q^\alpha}^n d\mu_q(x) = \frac{1}{(1-q)^n [\alpha]_q^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{\alpha l + 1}{[\alpha l + 1]_q}. \quad (1.6)$$

The q -Bernoulli polynomials with weight α are defined by

$$\tilde{\beta}_{n,q}^{(\alpha)}(x) = \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^n d\mu_q(y) = \sum_{l=0}^n \binom{n}{l} q^{\alpha l x} [x]_{q^\alpha}^{n-l} \tilde{\beta}_{l,q}^{(\alpha)}. \quad (1.7)$$

By (1.6) and (1.7), we easily see that

$$\tilde{\beta}_{n,q}^{(\alpha)}(x) = \frac{1}{(1-q)^n [\alpha]_q^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{\alpha l x} \frac{\alpha l + 1}{[\alpha l + 1]_q}. \quad (1.8)$$

Let $C(\mathbb{Z}_p)$ be the set of continuous functions on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the p -adic analogue of Bernstein operator of order n for f is given by

$$\mathbb{B}_{n,q}(f | x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}, \quad (1.9)$$

where $n, k \in \mathbb{Z}_+$ (see [1, 9, 10]). For $n, k \in \mathbb{Z}_+$, the p -adic Bernstein polynomials of degree n are defined by $B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}$ for $x \in \mathbb{Z}_p$, (see [1, 10, 11]).

In this paper, we consider Bernstein polynomials to express the p -adic q -integral on $\mathbb{Z}_p$ and investigate some interesting identities of Bernstein polynomials associated with the q -Bernoulli numbers and polynomials with weight 0 by using the expression of p -adic q -integral on $\mathbb{Z}_p$ of these polynomials.

2. q -Bernoulli Numbers with Weight 0 and Bernstein Polynomials

In the special case, $\alpha = 0$, the q -Bernoulli numbers with weight 0 will be denoted by $\tilde{\beta}_{n,q}^{(0)} = \tilde{\beta}_{n,q}$. From (1.4), (1.5), and (1.6), we note that

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{\beta}_{n,q} \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} x^n d\mu_q(x) \frac{t^n}{n!} = \int_{\mathbb{Z}_p} e^{xt} d\mu_q(x) \\ &= \left(\frac{q-1}{\log q} \right) \left(\frac{t + \log q}{qe^t - 1} \right). \end{aligned} \quad (2.1)$$

It is easy to show that

$$\begin{aligned} \frac{t + \log q}{qe^t - 1} &= \frac{t}{q-1} \left(\frac{1 - q^{-1}}{e^t - q^{-1}} \right) + \frac{\log q}{q-1} \left(\frac{1 - q^{-1}}{e^t - q^{-1}} \right) \\ &= \frac{t}{q-1} \sum_{n=0}^{\infty} H_n(q^{-1}) \frac{t^n}{n!} + \frac{\log q}{q-1} \sum_{n=0}^{\infty} H_n(q^{-1}) \frac{t^n}{n!} \\ &= \frac{1}{q-1} \sum_{n=1}^{\infty} n H_{n-1}(q^{-1}) \frac{t^n}{n!} + \frac{\log q}{q-1} \sum_{n=0}^{\infty} H_n(q^{-1}) \frac{t^n}{n!}, \end{aligned} \quad (2.2)$$

where $H_n(q^{-1})$ are the n th Frobenius-Euler numbers.

By (2.1) and (2.2), we get

$$\tilde{\beta}_{n,q} = \begin{cases} 1 & \text{if } n = 0, \\ \frac{n}{\log q} H_{n-1}(q^{-1}) + H_n(q^{-1}) & \text{if } n > 0. \end{cases} \quad (2.3)$$

Therefore, we obtain the following theorem.

Theorem 2.1. For $n \in \mathbb{Z}_+$, we have

$$\tilde{\beta}_{n,q} = \begin{cases} 1 & \text{if } n = 0, \\ \frac{n}{\log q} H_{n-1}(q^{-1}) + H_n(q^{-1}) & \text{if } n > 0, \end{cases} \quad (2.4)$$

where $H_n(q^{-1})$ are the n th Frobenius-Euler numbers.

From (1.5), (1.6), and (1.7), we have

$$\tilde{\beta}_{0,q} = 1, \quad q(\tilde{\beta}_q + 1)^n - \tilde{\beta}_{n,q} = \begin{cases} \frac{q-1}{\log q} & \text{if } n = 1, \\ 0 & \text{if } n > 1, \end{cases} \quad (2.5)$$

with the usual convention about replacing $(\tilde{\beta}_q)^n$ with $\tilde{\beta}_{n,q}$. By (1.7), the n th q -Bernoulli polynomials with weight 0 are given by

$$\tilde{\beta}_{n,q}(x) = \int_{\mathbb{Z}_p} (x+y)^n d\mu_q(y) = \sum_{l=0}^n \binom{n}{l} x^{n-l} \tilde{\beta}_{l,q}. \quad (2.6)$$

From (2.6), we can derive the following function equation:

$$\left(\frac{q-1}{\log q}\right) \left(\frac{t + \log q}{qe^t - 1}\right) e^{xt} = \sum_{n=0}^{\infty} \tilde{\beta}_{n,q}(x) \frac{t^n}{n!}. \quad (2.7)$$

Thus, by (2.7), we get that

$$\tilde{\beta}_{n,q^{-1}}(1-x) = (-1)^n \tilde{\beta}_{n,q}(x), \quad \text{for } n \in \mathbb{Z}_+. \quad (2.8)$$

By the definition of p -adic q -integral on $\mathbb{Z}_p$, we see that

$$\int_{\mathbb{Z}_p} (1-x)^n d\mu_q(x) = (-1)^n \int_{\mathbb{Z}_p} (x-1)^n d\mu_q(x) = (-1)^n \tilde{\beta}_{n,q}(-1). \quad (2.9)$$

Therefore, by (2.8) and (2.9), we obtain the following theorem.

Theorem 2.2. For $n \in \mathbb{Z}_+$, we have

$$(-1)^n \tilde{\beta}_{n,q}(x) = \tilde{\beta}_{n,q^{-1}}(1-x). \quad (2.10)$$

In particular, $x = -1$, we get

$$\int_{\mathbb{Z}_p} (1-y)^n d\mu_q(y) = (-1)^n \tilde{\beta}_{n,q}(-1) = \tilde{\beta}_{n,q^{-1}}(2). \quad (2.11)$$

From (2.5), we can derive the following equation:

$$q^2 \tilde{\beta}_{n,q}(2) = q^2 + nq \frac{q-1}{\log q} - q + \tilde{\beta}_{n,q}, \quad \text{if } n > 1. \quad (2.12)$$

Therefore, by (2.12), we obtain the following theorem.

Theorem 2.3. For $n \in \mathbb{N}$ with $n > 1$, we have

$$\tilde{\beta}_{n,q}(2) = 1 + \frac{n}{q} \left(\frac{q-1}{\log q} \right) - \frac{1}{q} + \frac{1}{q^2} \tilde{\beta}_{n,q}. \quad (2.13)$$

Taking the p -adic q -integral on $\mathbb{Z}_p$ for one Bernstein polynomials in (1.9), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_q(x) &= \binom{n}{k} \int_{\mathbb{Z}_p} x^k (1-x)^{n-k} d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} x^{k+l} d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \tilde{\beta}_{k+l,q}. \end{aligned} \quad (2.14)$$

From the symmetry of Bernstein polynomials, we note that

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_q(x) &= \int_{\mathbb{Z}_p} B_{n-k,n}(1-x) d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \int_{\mathbb{Z}_p} (1-x)^{n-l} d\mu_q(x). \end{aligned} \quad (2.15)$$

Let $n > k + 1$. Then, by Theorem 2.3 and (2.15), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_q(x) &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \left(1 - \frac{n-l}{q^{-1}} \left(\frac{q^{-1}-1}{\log q} \right) - q + q^2 \tilde{\beta}_{n-l,q^{-1}} \right) \\ &= \begin{cases} 1 + n \left(\frac{q-1}{\log q} \right) - q + q^2 \tilde{\beta}_{n,q^{-1}} & \text{if } k = 0, \\ n \left(\frac{1-q}{\log q} \right) + nq^2 \tilde{\beta}_{n,q^{-1}} + nq^2 \tilde{\beta}_{n-1,q^{-1}} & \text{if } k = 1, \\ \binom{n}{k} q^2 \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \tilde{\beta}_{n-l,q^{-1}} & \text{if } k > 1. \end{cases} \end{aligned} \quad (2.16)$$

By comparing the coefficients on the both sides of (2.14) and (2.16), we obtain the following theorem.

Theorem 2.4. For $n, k \in \mathbb{Z}_+$ with $n > k + 1$, we have

$$\begin{aligned} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \tilde{\beta}_{1+l,q} &= \frac{1-q}{\log q} + q^2 \tilde{\beta}_{n,q^{-1}} + q^2 \tilde{\beta}_{n-1,q^{-1}}, \\ \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \tilde{\beta}_{k+l,q} &= q^2 \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \tilde{\beta}_{n-l,q^{-1}}, \quad \text{if } k > 1. \end{aligned} \quad (2.17)$$

In particular, when $k = 0$, we have

$$\sum_{l=0}^n \binom{n}{l} (-1)^l \tilde{\beta}_{l,q} = 1 + n \frac{q-1}{\log q} - q + q^2 \tilde{\beta}_{n,q^{-1}}. \quad (2.18)$$

Let $m, n, k \in \mathbb{Z}_+$ with $m + n > 2k + 1$. Then we see that

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) d\mu_q(x) &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} x^{2k} (1-x)^{n+m-2k} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \int_{\mathbb{Z}_p} (1-x)^{n+m-1} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \left(1 - (n+m-l) \left(\frac{1-q}{\log q} \right) - q + q^2 \tilde{\beta}_{n+m-l,q^{-1}} \right) \\ &= \begin{cases} 1 + (n+m) \left(\frac{q-1}{\log q} \right) - q + q^2 \tilde{\beta}_{n+m,q^{-1}} & \text{if } k = 0, \\ \binom{n}{k} \binom{m}{k} q^2 \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \tilde{\beta}_{n+m-l,q^{-1}} & \text{if } k > 0. \end{cases} \end{aligned} \quad (2.19)$$

For $m, n, k \in \mathbb{Z}_+$, we have

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) d\mu_q(x) &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} x^{2k} (1-x)^{n+m-2k} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \int_{\mathbb{Z}_p} x^{2k+l} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \tilde{\beta}_{l+2k,q}. \end{aligned} \quad (2.20)$$

By comparing the coefficients on the both sides of (2.19) and (2.20), we obtain the following theorem.

Theorem 2.5. For $m, n, k \in \mathbb{Z}_+$ with $m + n > 2k + 1$, we have

$$\sum_{l=0}^{n+m} \binom{n+m}{l} (-1)^l \tilde{\beta}_{l,q} = 1 + (n+m) \left(\frac{q-1}{\log q} \right) - q + q^2 \tilde{\beta}_{n+m,q^{-1}}. \quad (2.21)$$

In particular, when $k \neq 0$, we have

$$\sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \tilde{\beta}_{l+2k,q} = q^2 \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \tilde{\beta}_{n+m-l,q^{-1}}. \quad (2.22)$$

For $s \in \mathbb{N}$, let $k, n_1, \dots, n_s \in \mathbb{Z}_+$ with $n_1 + n_2 + \dots + n_s > sk + 1$. By the same method above, we get

$$\int_{\mathbb{Z}_p} \left(\prod_{i=1}^s B_{k,n_i}(x) \right) d\mu_q(x) = \begin{cases} 1 + \left(\sum_{i=1}^s n_i \right) \left(\frac{q-1}{\log q} \right) - q + q^2 \tilde{\beta}_{n_1+n_2+\dots+n_s,q^{-1}} & \text{if } k = 0, \\ \left(\prod_{i=1}^s \binom{n_i}{k} \right) q^2 \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{l+sk} \tilde{\beta}_{n_1+n_2+\dots+n_s-l,q^{-1}} & \text{if } k > 0. \end{cases} \quad (2.23)$$

From the binomial theorem, we note that

$$\int_{\mathbb{Z}_p} \left(\prod_{i=1}^s B_{k,n_i}(x) \right) d\mu_q(x) = \left(\prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \tilde{\beta}_{l+sk,q}. \quad (2.24)$$

By comparing the coefficients on the both sides of (2.23) and (2.24), we obtain the following theorem.

Theorem 2.6. For $s \in \mathbb{N}$, let $k, n_1, \dots, n_s \in \mathbb{Z}_+$ with $n_1 + n_2 + \dots + n_s > sk + 1$. Then, we have

$$\sum_{l=0}^{n_1+\dots+n_s} \binom{n_1+\dots+n_s}{l} (-1)^l \tilde{\beta}_{l,q} = 1 + \left(\sum_{i=1}^s n_i \right) \left(\frac{q-1}{\log q} \right) - q + q^2 \tilde{\beta}_{n_1+\dots+n_s,q^{-1}}. \quad (2.25)$$

In particular, when $k \neq 0$, we have

$$\sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \tilde{\beta}_{l+sk,q} = q^2 \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{l+sk} \tilde{\beta}_{n_1+\dots+n_s-l,q^{-1}}. \quad (2.26)$$

Acknowledgment

The present research has been conducted by the Research Grant of Kwangwoon University in 2011.

References

- [1] T. Kim, " q -Bernstein polynomials, q -Stirling numbers and q -Bernoulli polynomials," <http://arxiv.org/abs/1008.4547>.
- [2] T. Kim, " q -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [3] T. Kim, " q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients," *Russian Journal of Mathematical Physics*, vol. 15, no. 1, pp. 51–57, 2008.
- [4] L.-C. Jang, "A study on the distribution of twisted q -Genocchi polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 2, pp. 181–189, 2009.
- [5] Y. Simsek, "Special functions related to Dedekind-type DC-sums and their applications," *Russian Journal of Mathematical Physics*, vol. 17, no. 4, pp. 495–508, 2010.
- [6] S.-H. Rim, J.-H. Jin, E.-J. Moon, and S.-J. Lee, "Some identities on the q -Genocchi polynomials of higher-order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *International Journal of Mathematics and Mathematical Sciences*, vol. 2010, Article ID 860280, 14 pages, 2010.
- [7] C. S. Ryoo, "Some identities of the twisted q -Euler numbers and polynomials associated with q -Bernstein polynomials," *Proceedings of the Jangjeon Mathematical Society*, vol. 14, pp. 239–248, 2011.
- [8] T. Kim, "On the weighted q -Bernoulli numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, pp. 207–215, 2011.
- [9] S. Araci, D. Erdal, and J. J. Seo, "A study on the fermionic p -adic q -integral representataion on $\mathbb{Z}_p$ associatied with weighted q -Bernstein and q -Genocchi polynomials," *Abstract and Applied Analysis*, vol. 2011, Article ID 649248, 7 pages, 2011.
- [10] Y. Simsek, "Construction a new generating function of Bernstein type polynomials," *Applied Mathematics and Computation*. In press.
- [11] D. Erdal, J. J. Seo, and S. Araci, "New construction weighted (h, q) -Genocchi numbers and polynomials related to zeta type function," *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 487490, 6 pages, 2011.