Research Article

The Fixed Point Property in c_0 with an Equivalent Norm

Berta Gamboa de Buen¹ and Fernando Núñez-Medina²

¹ *Matemáticas Básicas, Centro de Investigación en Matemáticas (CIMAT), Apartado Postal 402, 36000 Guanajuato, GTO, Mexico*

² *Departamento de Matemáticas Aplicadas, Universidad del Papaloapan (UNPA), 68400 Loma Bonita, OAX, Mexico*

Correspondence should be addressed to Berta Gamboa de Buen, gamboa@cimat.mx

Received 7 June 2011; Accepted 27 August 2011

Academic Editor: Elena Litsyn

Copyright © 2011 B. Gamboa de Buen and F. Núñez-Medina. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the fixed point property (FPP) in the Banach space c_0 with the equivalent norm $\|\cdot\|_D$. The space c_0 with this norm has the weak fixed point property. We prove that every infinite-dimensional subspace of $(c_0, \|\cdot\|_D)$ contains a complemented asymptotically isometric copy of c_0 , and thus does not have the FPP, but there exist nonempty closed convex and bounded subsets of $(c_0, \|\cdot\|_D)$ which are not ω -compact and do not contain asymptotically isometric c_0 -summing basis sequences. Then we define a family of sequences which are asymptotically isometric to different bases equivalent to the summing basis in the space $(c_0, \|\cdot\|_D)$, and we give some of its properties. We also prove that the dual space of $(c_0, \|\cdot\|_D)$ over the reals is the Bynum space $l_{1\infty}$ and that every infinite-dimensional subspace of $l_{1\infty}$ does not have the fixed point property.

1. Introduction

We start with some notations and terminologies. Let K be a nonempty, convex, closed and bounded subset of a Banach space $(X, \|\cdot\|)$. A mapping $T : K \rightarrow K$ is said to be nonexpansive if

$$\|Tx - Ty\| \leq \|x - y\|, \quad x, y \in K. \quad (1.1)$$

We say that K has the fixed point property for nonexpansive mappings (FPP) if every nonexpansive mapping $T : K \rightarrow K$ has a fixed point, that is, a point $x \in K$ such that $Tx = x$. We say that a Banach space $(X, \|\cdot\|)$ has the fixed point property for nonexpansive mappings (FPP) if every nonempty, convex, closed, and bounded subset K of $(X, \|\cdot\|)$ has the FPP, and we say that the Banach space $(X, \|\cdot\|)$ has the weak fixed point property for nonexpansive

mappings (ω -FPP) if every nonempty, convex and weakly compact subset K of $(X, \|\cdot\|)$ has the FPP.

In this paper we study the FPP in the Banach space c_0 with the equivalent norm $\|\cdot\|_D$ defined by

$$\|x\|_D = \sup_{i,j \in \mathbb{N}} |x_i - x_j|, \quad x = \{x_i\} \in c_0. \quad (1.2)$$

The norm $\|\cdot\|_D$ was used by Hagler in [1] to construct a separable Banach space X with non-separable dual such that l_1 does not embed in X and every normalized weakly null sequence in X has a subsequence equivalent to the canonical basis of c_0 .

In [2], Dowling et al. gave a characterization of nonempty, convex, closed and bounded subsets of c_0 which are not ω -compact. Specifically, they proved that if K is a convex, closed and bounded subset of c_0 , then K is ω -compact if and only if every nonempty, convex, closed and convex subset of K has the FPP. To do that, the authors showed that every closed, convex and bounded subset of c_0 which is not ω -compact contains an asymptotically isometric c_0 -summing basic sequence, *aisbc* $_0$ sequence for short, that is, a sequence $\{y_n\}_n \subset c_0$ such that for all $\{t_n\}_n \in l_1$,

$$\sup_{n \in \mathbb{N}} (1 + \varepsilon_n)^{-1} \left| \sum_{i=n}^{\infty} t_i \right| \leq \left\| \sum_{n=1}^{\infty} t_n y_n \right\| \leq \sup_{n \in \mathbb{N}} (1 + \varepsilon_n) \left| \sum_{i=n}^{\infty} t_i \right|, \quad (1.3)$$

for some $\{\varepsilon_n\}_n \subset \mathbb{R}$ with $0 \leq \varepsilon_{n+1} \leq \varepsilon_n$ and $\lim_n \varepsilon_n = 0$. They proved that if a convex, closed and bounded subset K of a Banach space contains an *aisbc* $_0$ sequence, then there exists a nonempty, convex, closed and bounded subset of K without the FPP. The authors used this fact in [3] as a tool to prove that a nonempty, closed, convex and bounded subset of c_0 is ω -compact if and only if it has the FPP.

It is easy to see that $(c_0, \|\cdot\|_D)$ contains c_0 isometrically, and then it contains *aisbc* $_0$ sequences.

First we prove that every infinite-dimensional subspace Y of $(c_0, \|\cdot\|_D)$ has a complemented asymptotically isometric copy of c_0 and hence by a result proved by Dowling et al. in [4], Y does not have the FPP. Also, as an immediate consequence we obtain that Y has an *aisbc* $_0$ sequence. Nevertheless, we exhibit a nonempty closed, convex and bounded subset of $(c_0, \|\cdot\|_D)$, which is not ω -compact and does not contain *aisbc* $_0$ sequences.

Then for every selection of signs $\Theta = \{\theta_i\}$, we define the Θ -basis of c_0 which is equivalent to the summing basis and define the corresponding asymptotically isometric Θ -basic sequence, *ai Θ bc* $_{0D}$ sequence for short. We prove that if $\Theta_1 \neq \pm \Theta_2$, then the *ai Θ_1 bc* $_{0D}$ and *ai Θ_2 bc* $_{0D}$ sequences are different in the sense that there exists a nonempty, closed, convex, and bounded subset of $(c_0, \|\cdot\|_D)$, which is not ω -compact, contains an *ai Θ_1 bc* $_{0D}$ sequence, and does not contain *ai Θ_2 bc* $_{0D}$ sequences. We also show that the *aisbc* $_0$ and *ai Θ bc* $_{0D}$ sequences are different in the last sense for all Θ . Hence, to give a similar result of Theorem 4 of [2] about convex, closed and bounded sets in $(c_0, \|\cdot\|_D)$ without the FPP, it is necessary to consider the *ai Θ bc* $_{0D}$ sequences.

Next we prove that if a convex and closed subset K of a Banach space contains an asymptotically isometric c_{0D} -summing basic sequence, that is, an *ai Θ bc* $_{0D}$ sequence, where Θ is such that $\theta_i = 1$ for all i , then there exists a nonempty, convex, closed and bounded subset of K without the FPP.

Finally, we show that the dual space of $(c_0, \|\cdot\|_D)$, over the reals, is the Bynum [5] space $l_{1\infty}$. Then, by a result of Dowling et al. in [6], the space $l_{1\infty} = (c_0, \|\cdot\|_D)^*$ has “many” subspaces and contains an asymptotically isometric copy of l_1 and does not have the FPP. In fact, we prove that every infinite dimensional subspace of $l_{1\infty}$ contains an asymptotically isometric copy of l_1 and does not have the FPP.

2. The Space $(c_0, \|\cdot\|_D)$

In the sequel, we will denote by $\{e_n\}$ the canonical basis of c_0 and by $\{\xi_n\}$ the summing basis of c_0 , that is, $\xi_n = \sum_{i=1}^n e_i$, $n \in \mathbb{N}$.

García Falset proved in [7] that a Banach space with strongly bimonotone basis and with the weak Banach-Saks property has the ω -FPP. It is easy to see that the canonical basis of c_0 is strongly bimonotone in $(c_0, \|\cdot\|_D)$. On the other hand, since c_0 has the weak Banach-Saks property and $\|\cdot\|_D$ and $\|\cdot\|_\infty$ are equivalent, we get that $(c_0, \|\cdot\|_D)$ has the weak Banach-Saks property. Hence we have that $(c_0, \|\cdot\|_D)$ has the ω -FPP.

To study the FPP in the space $(c_0, \|\cdot\|_D)$ using *aisbc* c_0 sequences, we would expect that nonempty, convex, closed and bounded subsets K of $(c_0, \|\cdot\|_D)$, which are not ω -compact, contain an *aisbc* c_0 sequence. This fact is true for some ω -compact sets in $(c_0, \|\cdot\|_D)$, since the space c_0 embeds isometrically in $(c_0, \|\cdot\|_D)$. In fact we have the following proposition.

Proposition 1. *Let $\{u_k\}_k \subset (c_0, \|\cdot\|_D)$ be a block basis of $\{e_n\}$ with $u_k = \sum_{i=p_k}^{q_k} a_i e_i$, $1 \leq p_1 \leq q_1 < p_2 \leq q_2 < \dots$. If $\|u_k\|_\infty = 1 = a_{i^k}$, for some $p_k \leq i^k \leq q_k$, and $y_k = (1/2)(u_{2k} - u_{2k-1})$, then the space $\text{span}\{y_k\}$ is isometric to $(c_0, \|\cdot\|_\infty)$.*

Proof. Since $\|u_k\|_\infty = 1 = a_{i^k}$ for every $k \in \mathbb{N}$, then $|a_j| \leq 1$ for all $j \in \mathbb{N}$ and

$$\max_{p_{2k-1} \leq i \leq q_{2k-1}, p_{2k} \leq j \leq q_{2k}} |a_i + a_j| = a_{i^{2k-1}} + a_{i^{2k}} = 2. \quad (2.1)$$

Hence, it is straightforward to see that $\|\sum_{k=1}^n t_k y_k\|_D = \|\sum_{k=1}^n t_k e_k\|_\infty$. $\square$

In the following theorem, we will show, using some results proved by Dowling et al. [4, 8], that every infinite-dimensional subspace Y of c_{0D} fails to have the FPP.

Theorem 2. *Let Y be an infinite-dimensional subspace of c_{0D} . Then Y has a complemented asymptotically isometric copy of c_0 and thus Y does not have the FPP.*

Proof. Let $\{\varepsilon_k\}_k \subset (0, 1)$ be a sequence such that $\varepsilon_{k+1} < \varepsilon_k$, $k \in \mathbb{N}$ and $\varepsilon_k \rightarrow 0$. As in [9] we construct sequences $\{n_k\} \subset \mathbb{N}$ and $\{y_k\}_k \subset Y$ such that $n_k < n_{k+1}$, $y_k = \sum_{i=n_k}^\infty \alpha_i^k e_i$, $\|y_k\|_\infty = 1$, and

$$\sup_{i \geq n_{k+1}} |\alpha_i^j| < \frac{\varepsilon_{k+2}}{4k} \quad \forall j = 1, \dots, k, \text{ and every } k \in \mathbb{N}. \quad (2.2)$$

Since $\|y_k\|_\infty = 1$, taking $-y_k$ instead of y_k , if necessary, we can suppose that there exists $n_k \leq r^k < n_{k+1}$ such that

$$\alpha_{r^k}^k = 1. \quad (2.3)$$

Define $x_k = (y_{2k-1} - y_{2k})/2$. Then, by (2.3) and (2.2), we get that $1 - (\varepsilon_k/2) < \|x_k\|_D < 1 + (\varepsilon_k/2)$ and

$$\begin{aligned} \sum_{k=1}^{\infty} t_k x_k &= \frac{1}{2} \sum_{k=1}^{\infty} t_k \left(\sum_{i=n_{2k-1}}^{\infty} \alpha_i^{2k-1} e_i - \sum_{i=n_{2k}}^{\infty} \alpha_i^{2k} e_i \right) \\ &= \frac{1}{2} \sum_{k=1}^{\infty} t_k \left(\sum_{i=n_{2k-1}}^{\infty} (\alpha_i^{2k-1} - \alpha_i^{2k}) e_i \right) \\ &= \frac{1}{2} \sum_{k=1}^{\infty} \left(\sum_{i=n_{2k-1}}^{n_{2k+1}-1} \left(\sum_{j=1}^k t_j (\alpha_i^{2j-1} - \alpha_i^{2j}) e_i \right) \right), \end{aligned} \quad (2.4)$$

where $\alpha_i^{2k} = 0$ for $i = n_{2k-1}, \dots, n_{2k} - 1$, $k \in \mathbb{N}$. Then by (2.3) and (2.2), if $k > 1$, we get

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} t_n x_n \right\|_D &\geq \frac{1}{2} \max_{n_{2k-1} \leq r < n_{2k}, n_{2k} \leq s < n_{2k+1}} \left| \sum_{j=1}^k t_j (\alpha_r^{2j-1} - \alpha_r^{2j} - \alpha_s^{2j-1} + \alpha_s^{2j}) \right| \\ &\geq \frac{1}{2} \left| \sum_{j=1}^k t_j (\alpha_{r^{2k-1}}^{2j-1} - \alpha_{r^{2k-1}}^{2j} - \alpha_{r^{2k}}^{2j-1} + \alpha_{r^{2k}}^{2j}) \right| \\ &\geq \frac{1}{2} |t_k| \left| \alpha_{r^{2k-1}}^{2k-1} - \alpha_{r^{2k-1}}^{2k} + \alpha_{r^{2k}}^{2k} \right| - \frac{1}{2} \sum_{j=1}^{k-1} |t_j| \left| \alpha_{r^{2k-1}}^{2j-1} - \alpha_{r^{2k-1}}^{2j} - \alpha_{r^{2k}}^{2j-1} + \alpha_{r^{2k}}^{2j} \right| \\ &\geq \frac{1}{2} |t_k| \left(\left| \alpha_{r^{2k-1}}^{2k-1} + \alpha_{r^{2k}}^{2k} \right| - \left| \alpha_{r^{2k-1}}^{2k-1} \right| \right) \\ &\quad - \frac{1}{2} \sum_{j=1}^{k-1} |t_j| \left(\left| \alpha_{r^{2k-1}}^{2j-1} \right| + \left| \alpha_{r^{2k-1}}^{2j} \right| + \left| \alpha_{r^{2k}}^{2j-1} \right| + \left| \alpha_{r^{2k}}^{2j} \right| \right) \\ &\geq \frac{1}{2} |t_k| (2 - \varepsilon_k) - \frac{1}{2} \sum_{j=1}^{k-1} |t_j| \frac{\varepsilon_k}{k} \\ &\geq |t_k| \left(1 - \frac{\varepsilon_k}{2} \right) - \max_{1 \leq j \leq k} |t_j| \frac{\varepsilon_k}{2}. \end{aligned} \quad (2.5)$$

On the other hand, if $n_{2k-1} \leq r < n_{2k+1}$, $n_{2m-1} \leq s < n_{2m+1}$, $k \leq m$, using (2.2), we get

$$\begin{aligned} &\frac{1}{2} \left| \sum_{j=1}^k t_j (\alpha_r^{2j-1} - \alpha_r^{2j}) - \sum_{j=1}^m t_j (\alpha_s^{2j-1} - \alpha_s^{2j}) \right| \\ &\leq \frac{1}{2} \left[|t_k| \left(\left| \alpha_r^{2k-1} \right| + \left| \alpha_r^{2k} \right| \right) + |t_m| \left(\left| \alpha_s^{2m-1} \right| + \left| \alpha_s^{2m} \right| \right) \right] \\ &\quad + \frac{1}{2} \left[\sum_{j=1}^{k-1} |t_j| \left(\left| \alpha_r^{2j-1} \right| + \left| \alpha_r^{2j} \right| \right) + \sum_{j=1}^{m-1} |t_j| \left(\left| \alpha_s^{2j-1} \right| + \left| \alpha_s^{2j} \right| \right) \right] \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2} \left[|t_k|(1 + \varepsilon_k) + |t_m|(1 + \varepsilon_m) + \sum_{j=1}^{k-1} |t_j| \frac{\varepsilon_k}{k-1} + \sum_{j=1}^{m-1} |t_j| \frac{\varepsilon_m}{m-1} \right] \\
&\leq \frac{1}{2} \left[|t_k|(1 + \varepsilon_k) + |t_m|(1 + \varepsilon_m) + \max_{1 \leq j < k} |t_j| \varepsilon_k + \max_{1 \leq j < m} |t_j| \varepsilon_m \right] \\
&\leq \frac{1}{2} \left[\max_{1 \leq j \leq k} |t_j| (1 + \varepsilon_k) + \max_{1 \leq j \leq m} |t_j| (1 + \varepsilon_m) \right] \\
&\leq \sup_{n \in \mathbb{N}} \left((1 + \varepsilon_n) \max_{1 \leq j \leq n} |t_j| \right) \leq \sup_{n \in \mathbb{N}} (1 + \varepsilon_n) |t_n|.
\end{aligned} \tag{2.6}$$

Then we obtain

$$\sup_{n \in \mathbb{N}} \left(|t_n| \left(1 - \frac{\varepsilon_n}{2} \right) - \max_{1 \leq j \leq n} |t_j| \frac{\varepsilon_n}{2} \right) \leq \left\| \sum_{n=1}^{\infty} t_n x_n \right\|_D \leq \sup_{n \in \mathbb{N}} (1 + \varepsilon_n) |t_n|. \tag{2.7}$$

Now, define $z_n = x_n / (1 + \varepsilon_n)$ and $m = (1 - \varepsilon_1) / (1 + \varepsilon_1)$; then $(1 - \varepsilon_n) / (1 + \varepsilon_n) \leq \|z_n\|_D$ and $\lim_n \|z_n\|_D = 1$. On the other hand,

$$\begin{aligned}
(1 + \varepsilon_1) m \sup_{n \in \mathbb{N}} |t_n| &= (1 - \varepsilon_1) \sup_{n \in \mathbb{N}} |t_n| = \left(1 - \frac{\varepsilon_1}{2} \right) \sup_{n \in \mathbb{N}} |t_n| - \frac{\varepsilon_1}{2} \sup_{n \in \mathbb{N}} |t_n| \\
&\leq \sup_{n \in \mathbb{N}} \left(|t_n| \left(1 - \frac{\varepsilon_n}{2} \right) - \max_{1 \leq j \leq n} |t_j| \frac{\varepsilon_n}{2} \right).
\end{aligned} \tag{2.8}$$

Thus

$$m \sup_{n \in \mathbb{N}} |t_n| \leq \sup_{n \in \mathbb{N}} \left(|t_n| \left(1 - \frac{\varepsilon_n}{2} \right) - \max_{1 \leq j \leq n} |t_j| \frac{\varepsilon_n}{2} \right) \leq \left\| \sum_{n=1}^{\infty} t_n z_n \right\|_D \leq \sup_{n \in \mathbb{N}} |t_n|. \tag{2.9}$$

Then by Theorem 2 of [8] Y contains an asymptotically isometric copy of c_0 and since Y does not contain a copy of l_1 , by Corollary 11 of [8] it contains a complemented asymptotically isometric copy of c_0 . Finally by Proposition 11 of [4], Y does not have the FPP. $\square$

As a consequence of the last theorem, we get that every infinite-dimensional subspace of $(c_0, \|\cdot\|_D)$ contains an *aisbc*₀ sequence. Nevertheless, the following result gives an example of a nonempty, convex, closed and bounded subset of $(c_0, \|\cdot\|_D)$ which is not weakly compact and without *aisbc*₀ sequences.

Proposition 3. *Let $\{\xi_n\}$ be the c_0 summing basis. Then*

$$C = \left\{ \sum_{n=1}^{\infty} \lambda_n \xi_n : \lambda_n \geq 0, \sum_{n=1}^{\infty} \lambda_n = 1 \right\} \tag{2.10}$$

*does not have *aisbc*₀ sequences with the norm $\|\cdot\|_D$.*

Proof. Suppose that $\{y_n\}$ is an $aisbc_0$ sequence in C with $\|\cdot\|_D$ for some sequence $\{\varepsilon_n\}$. Then $y_n = \sum_{i=1}^{\infty} \lambda_i^n \xi_i$ for some sequence $\{\lambda_i^n\}$ such that $\lambda_i^n \geq 0$ and $\sum_{i=1}^{\infty} \lambda_i^n = 1$. Fix $0 < \varepsilon < 1/4$. Passing to a subsequence we can suppose that $\varepsilon_{n+1} \leq \varepsilon_n < (1/2) - 2\varepsilon$ and $1/(1+\varepsilon_n) > 1-\varepsilon$, $n \in \mathbb{N}$.

Assume first that there exists $M \in \mathbb{N}$ such that for every $n \geq M$, $\sum_{i=M+1}^{\infty} \lambda_i^n \leq (1/2) - \varepsilon$. Let $u_n = \sum_{i=1}^M \lambda_i^n \xi_i$ and $v_n = \sum_{i=M+1}^{\infty} \lambda_i^n \xi_i$; then $y_n = u_n + v_n$. Since $\{u_n\} \subset [\xi_i]_{i=1}^M$ is bounded and $\dim [\xi_i]_{i=1}^M = M$, passing to another subsequence we can suppose that $u_n \rightarrow u$ for some $u \in C$. Then, there exist $n_1, n_2 \in \mathbb{N}$ with $M \leq n_1 < n_2$ such that

$$\|u_{n_1} - u_{n_2}\|_D < \varepsilon. \quad (2.11)$$

Since $\sum_{i=M+1}^{\infty} \lambda_i^n \leq (1/2) - \varepsilon$, $n \geq M$, we also get

$$\begin{aligned} \|v_{n_1} - v_{n_2}\|_D &= \max_{M+1 \leq r \leq k < \infty} \left| \sum_{i=r}^k \lambda_i^{n_1} - \sum_{i=r}^k \lambda_i^{n_2} \right| \\ &\leq \sum_{i=M+1}^{\infty} \lambda_i^{n_1} + \sum_{i=M+1}^{\infty} \lambda_i^{n_2} \leq 1 - 2\varepsilon. \end{aligned} \quad (2.12)$$

Hence $\|y_{n_1} - y_{n_2}\|_D \leq 1 - \varepsilon$. On the other hand, since $\{y_n\}$ is an $aisbc_0$ sequence, we have that $\|y_{n_1} - y_{n_2}\|_D \geq 1/(1 + \varepsilon_{n_2})$, which contradicts the fact that $1/(1 + \varepsilon_{n_2}) > 1 - \varepsilon$.

Assume now that for all $M \in \mathbb{N}$, there exist $n \geq M$ such that $\sum_{i=M+1}^{\infty} \lambda_i^n > (1/2) - \varepsilon$. Denote each y_n by $\{\alpha_i^n\} = \sum_{i=1}^{\infty} \alpha_i^n e_i$, where $\{e_n\}$ is the canonical basis of c_0 . Then $\alpha_i^n = \sum_{j=i}^{\infty} \lambda_j^n$. Since $y_1, y_2 \in c_0$, there exists $M \in \mathbb{N}$ such that

$$\alpha_i^1, \alpha_i^2 < \frac{\varepsilon}{2}, \quad i \geq M. \quad (2.13)$$

By hypothesis, there exists $n_0 \in \mathbb{N}$ such that $\sum_{i=M+1}^{\infty} \lambda_i^{n_0} > (1/2) - \varepsilon$. Then

$$\frac{3}{2} - 2\varepsilon \leq \|y_1 + y_2 - y_{n_0}\|_D. \quad (2.14)$$

On the other hand, since $\{y_n\}$ is an $aisbc_0$ sequence, we have that $\|y_1 + y_2 - y_{n_0}\|_D \leq 1 + \varepsilon_1$, which contradicts the fact that $\varepsilon_1 < (1/2) - 2\varepsilon$. $\square$

In view of the last proposition and motivated by the behavior of the c_0 summing basic sequence with the norm $\|\cdot\|_D$, we will define the asymptotically isometric c_{0D} -summing basic sequence. First we consider the following definition.

Definition 4. Let $\{x_n\}$ be a bounded basic sequence in a Banach space X . We say that $\{x_n\}$ is a convexly closed sequence if the set

$$C = \left\{ \sum_{n=1}^{\infty} t_n x_n : t_n \geq 0, \sum_{n=1}^{\infty} t_n = 1 \right\} \quad (2.15)$$

is closed, that is, if $\overline{\text{conv}}\{x_n\} = C$.

Note that subsequences of convexly closed sequences are again convexly closed and that every basic sequence equivalent to a convexly closed sequence is convexly closed.

It is easy to see that the c_0 summing basis, the canonical basis of l_1 , and $aisbc_0$ sequences are convexly closed. Moreover, a weakly null basic sequence in a Banach space is not a convexly closed sequence. Hence the canonical basis of c_0 and the canonical basis of l_p , $1 < p < \infty$, are not convexly closed.

Definition 5. Let $\{x_n\}$ be a sequence in a Banach space X . We say that $\{x_n\}$ is an asymptotically isometric c_{0D} -summing basic sequence, $aisbc_{0D}$ sequence for short, if $\{x_n\}$ is convexly closed and there exists $\{\varepsilon_n\} \subset (0, \infty)$ such that $\varepsilon_n \searrow 0$ and

$$\sup_{1 \leq n \leq m < \infty} (1 + \varepsilon_m)^{-1} \left| \sum_{k=n}^m t_k \right| \leq \left\| \sum_{n=1}^{\infty} t_n x_n \right\| \leq \sup_{1 \leq n \leq m < \infty} (1 + \varepsilon_m) \left| \sum_{k=n}^m t_k \right|, \quad \forall \{t_n\} \in l_1. \quad (2.16)$$

Now, we prove that the analogous of the operator defined in [2] is still contractive and then Banach spaces containing $aisbc_{0D}$ sequences does not have the FPP.

Proposition 6. Let K be a nonempty, convex, closed and bounded subset of a Banach space X . Let $\{\varepsilon_n\} \subset (0, \infty)$ be a sequence such that $\varepsilon_n \rightarrow 0$ and $\varepsilon_n < 2^{-1}4^{-n}$, $n \geq 2$. If K contains an $aisbc_{0D}$ sequence with this $\{\varepsilon_n\}$, then there exists a nonempty, convex and closed subset C of K and $T : C \rightarrow C$ affine, nonexpansive, and fixed-point-free. Moreover, T is contractive.

Proof. Let $\{x_n\}$ be an $aisbc_{0D}$ sequence in K with $\{\varepsilon_n\} \subset (0, \infty)$ such that $\varepsilon_n < 2^{-1}4^{-n}$, $n \geq 2$. Set

$$C = \overline{\text{conv}}\{x_n\} = \left\{ \sum_{n=1}^{\infty} t_n x_n : t_n \geq 0, n \in \mathbb{N}, \sum_{n=1}^{\infty} t_n = 1 \right\} \subset K. \quad (2.17)$$

Thus C is nonempty, convex, closed and bounded. Define $Tx_n = \sum_{j=1}^{\infty} ((x_{n+j})/2^j)$, $n \in \mathbb{N}$, and extend T linearly to C , that is, if $x = \sum_{n=1}^{\infty} t_n x_n \in C$ then define $T(\sum_{n=1}^{\infty} t_n x_n) = \sum_{n=1}^{\infty} t_n Tx_n$. It is easy to see that $T(C) \subset C$ and that T is affine and fixed-point-free, see [2]. We only need to show that T is a contractive mapping. Let $x, y \in C$, with $x \neq y$. Then $x = \sum_{n=1}^{\infty} t_n x_n$ and $y = \sum_{n=1}^{\infty} s_n x_n$, with $t_n, s_n \geq 0$, and $\sum_{n=1}^{\infty} t_n = \sum_{n=1}^{\infty} s_n = 1$. Let $\beta_n = t_n - s_n$, $n \in \mathbb{N}$, such that $\sum_{n=1}^{\infty} \beta_n = 0$. As in [2] we have

$$T(x) - T(y) = \sum_{n=1}^{\infty} B_n x_n, \quad (2.18)$$

where $B_1 = 0$ and $B_n = (\beta_1/2^{n-1}) + (\beta_2/2^{n-2}) + \cdots + (\beta_{n-1}/2)$, $n \geq 2$. Consequently,

$$\|T(x) - T(y)\| = \left\| \sum_{n=1}^{\infty} B_n x_n \right\| \leq \sup_{1 \leq n \leq m < \infty} (1 + \varepsilon_m) \left| \sum_{k=n}^m B_k \right|. \quad (2.19)$$

Take $n, m \in \mathbb{N}$ with $n \leq m$. Since

$$\begin{aligned}
 \sum_{k=n}^m B_k &= \frac{\beta_1}{2^{n-1}} + \frac{\beta_2}{2^{n-2}} + \cdots + \frac{\beta_{n-1}}{2} \\
 &\quad + \frac{\beta_1}{2^n} + \frac{\beta_2}{2^{n-1}} + \cdots + \frac{\beta_{n-1}}{2^2} + \frac{\beta_n}{2} + \cdots \\
 &\quad + \frac{\beta_1}{2^{m-1}} + \frac{\beta_2}{2^{m-2}} + \cdots + \frac{\beta_{n-1}}{2^{m-(n-1)}} + \frac{\beta_n}{2^{m-n}} + \frac{\beta_{n+1}}{2^{m-(n+1)}} + \cdots + \frac{\beta_{m-1}}{2} \\
 &= \frac{1}{2}(\beta_{n-1} + \beta_n + \cdots + \beta_{m-1}) \\
 &\quad + \frac{1}{2^2}(\beta_{n-2} + \beta_{n-1} + \cdots + \beta_{m-2}) + \cdots \\
 &\quad + \frac{1}{2^{n-1}}(\beta_1 + \beta_2 + \cdots + \beta_{m-(n-1)}) + \cdots \\
 &\quad + \frac{1}{2^{m-2}}(\beta_1 + \beta_2) + \frac{1}{2^{m-1}}(\beta_1),
 \end{aligned} \tag{2.20}$$

we have

$$\begin{aligned}
 (1 + \varepsilon_m) \left| \sum_{k=n}^m B_k \right| &\leq (1 + \varepsilon_m) \left(\frac{1 + 2\varepsilon_{m-1}}{2} \frac{1}{1 + 2\varepsilon_{m-1}} |\beta_{n-1} + \beta_n + \cdots + \beta_{m-1}| \right. \\
 &\quad + \frac{1 + 2\varepsilon_{m-2}}{2^2} \frac{1}{1 + 2\varepsilon_{m-2}} |\beta_{n-2} + \beta_{n-1} + \cdots + \beta_{m-2}| + \cdots \\
 &\quad + \frac{1 + 2\varepsilon_{m-(n-1)}}{2^{n-1}} \frac{1}{1 + 2\varepsilon_{m-(n-1)}} |\beta_1 + \beta_2 + \cdots + \beta_{m-(n-1)}| + \cdots \\
 &\quad \left. + \frac{1 + 2\varepsilon_2}{2^{m-2}} \frac{1}{1 + 2\varepsilon_2} |\beta_1 + \beta_2| + \frac{1 + 2\varepsilon_1}{2^{m-1}} \frac{1}{1 + 2\varepsilon_1} |\beta_1| \right) \\
 &\leq \left(\sup_{1 \leq i \leq j \leq m} (1 + 2\varepsilon_j)^{-1} \left| \sum_{k=i}^j \beta_k \right| \right) Q_{nm},
 \end{aligned} \tag{2.21}$$

where

$$\begin{aligned}
 Q_{nm} &= (1 + \varepsilon_m) \left(\frac{1 + 2\varepsilon_{m-1}}{2} + \frac{1 + 2\varepsilon_{m-2}}{2^2} + \cdots + \right. \\
 &\quad \left. + \frac{1 + 2\varepsilon_{m-(n-1)}}{2^{n-1}} + \frac{1 + 2\varepsilon_{m-n}}{2^n} + \cdots + \frac{1 + 2\varepsilon_2}{2^{m-2}} + \frac{1 + 2\varepsilon_1}{2^{m-1}} \right) \\
 &\leq \left(1 + \frac{1}{2 \cdot 4^m} \right) \left[\left(\frac{1}{2} + \frac{1}{2^2} + \cdots + \frac{1}{2^{m-1}} \right) + \left(\frac{1}{2 \cdot 4^{m-1}} + \cdots + \frac{1}{2^{m-1} \cdot 4^1} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
&= \left(1 + \frac{1}{2 \cdot 4^m}\right) \left[\left(1 - \frac{1}{2^{m-1}}\right) + \left(\frac{1}{2^{2m-1}} + \frac{1}{2^{2m-2}} + \cdots + \frac{1}{2^{m+1}}\right) \right] \\
&< \left(1 + \frac{1}{4^m}\right) \left[\left(1 - \frac{1}{2^{m-1}}\right) + \frac{1}{2^m} \right] < 1.
\end{aligned} \tag{2.22}$$

Then we get

$$\begin{aligned}
\sup_{1 \leq n \leq m < \infty} (1 + \varepsilon_m) \left| \sum_{k=n}^m B_k \right| &\leq \sup_{1 \leq n \leq m < \infty} (1 + 2\varepsilon_m)^{-1} \left| \sum_{k=n}^m \beta_k \right| < \sup_{1 \leq n \leq m < \infty} (1 + \varepsilon_m)^{-1} \left| \sum_{k=n}^m \beta_k \right| \\
&\leq \left\| \sum_{n=1}^{\infty} \beta_n x_n \right\| = \|x - y\|.
\end{aligned} \tag{2.23}$$

Thus T is contractive. $\square$

Next for any sequence of signs we will define a basis in c_0 equivalent to $\{\xi_n\}$, the summing basis of c_0 , and a sequence asymptotically isometric to it.

Let $\{e_n\}$ be the canonical basis of c_0 and for any selection of signs $\Theta = \{\theta_i\}_i$, that is, $\theta_i \in \{-1, 1\}$, $i \in \mathbb{N}$, let $\{\zeta_n^\Theta\}_n$ be the sequence defined by

$$\zeta_n^\Theta = \sum_{k=1}^n \theta_k e_k, \quad n \in \mathbb{N}. \tag{2.24}$$

Since $\|\sum_{n=1}^m t_n \xi_n\|_\infty = \|\sum_{n=1}^m t_n \zeta_n^\Theta\|_\infty$ for all $\{t_n\}_{n=1}^m \subset \mathbb{K}$, we get that $\{\zeta_n^\Theta\}$ is a basis of c_0 equivalent to the c_0 summing basis. The sequence $\{\zeta_n^\Theta\}$ is called the Θ -basis of c_0 . Let $\Theta_0 = \{\theta_i\}$, where $\theta_i = 1$, $i \in \mathbb{N}$. Then the Θ_0 -basis of c_0 is the c_0 summing basis. If we define $C = \{\sum_{n=1}^\infty t_n \zeta_n^\Theta : t_n \geq 0 \text{ and } \sum_{n=1}^\infty t_n = 1\}$, then C is nonempty, convex and bounded. Since $\|\cdot\|_\infty$ and $\|\cdot\|_D$ are equivalent, we have that $\{\zeta_n^\Theta\}$ is convexly closed in $(c_0, \|\cdot\|_D)$.

The set $C = \{\sum_{n=1}^\infty t_n \zeta_n^\Theta : t_n \geq 0 \text{ and } \sum_{n=1}^\infty t_n = 1\}$ is not ω -compact. The following result shows that the set C contains neither $aisbc_{0D}$ sequences nor $aisbc_0$ sequences with the norm $\|\cdot\|_D$ if $\Theta \neq \pm\Theta_0$.

Proposition 7. For $\Theta \neq \pm\Theta_0$, let $\{\zeta_n^\Theta\}$ be the Θ -basis of c_0 considered in $(c_0, \|\cdot\|_D)$. If

$$C = \left\{ \sum_{n=1}^\infty t_n \zeta_n^\Theta : t_n \geq 0, \sum_{n=1}^\infty t_n = 1 \right\}, \tag{2.25}$$

then the set C contains neither $aisbc_{0D}$ sequences nor $aisbc_0$ sequences with the norm $\|\cdot\|_D$.

Proof. Let $\{y_k\} \subset C$. Then $y_k = \sum_{n=1}^\infty \lambda_n^k \zeta_n^\Theta$ for some $\lambda_n^k \geq 0$ and $\sum_{n=1}^\infty \lambda_n^k = 1$. Suppose that $\{y_k\}$ is an $aisbc_{0D}$ sequence (resp. an $aisbc_0$ sequence) with the norm $\|\cdot\|_D$. Let $n_0 = \min\{n : \theta_n \neq \theta_1\}$. If there exists $0 < \rho < 1$ such that $\sum_{i=1}^{n_0-1} \lambda_i^k \leq 1 - \rho$ for all $k \geq 1$, then for all $k \geq 1$,

$$\|y_k\|_D \geq \sum_{n=1}^\infty \lambda_n^k + \sum_{n=n_0}^\infty \lambda_n^k \geq 1 + \rho. \tag{2.26}$$

Since $\{y_k\}$ is an $aisbc_{0D}$ sequence (resp. an $aisbc_0$ sequence) with the norm $\|\cdot\|_D$, then $\|y_k\|_D \leq 1 + \varepsilon_k \rightarrow 1$ and this is impossible. Now, if $\limsup_k \sum_{i=1}^{n_0-1} \lambda_i^k = 1$, as in the proof of Proposition 3, we obtain a subsequence $\{y_{k_i}\}$ of $\{y_k\}$ with $\|y_{k_i} - y_{k_{i+1}}\|_D \rightarrow 0$. Since $\{y_n\}$ is an $aisbc_{0D}$ with the norm $\|\cdot\|_D$, then $(1 + \varepsilon_{k_i})^{-1} \leq \|y_{k_i} - y_{k_{i+1}}\|_D$ (resp. $(1 + \varepsilon_{k_{i+1}})^{-1} \leq \|y_{k_i} - y_{k_{i+1}}\|_D$) and making $i \rightarrow \infty$ we get that $1 \leq 0$. This contradiction proves the result. $\square$

Although the set C of the last proposition has neither $aisbc_{0D}$ sequences nor $aisbc_0$ sequences, for some Θ it does not have the FPP.

For $\Theta = \{\theta_i\}$, let F_n be the set such that if $i, j \in F_n$, then $\theta_i = \theta_j$, and if $i \in F_{n+1}$ and $j \in F_n$, then $\theta_i \neq \theta_j$. Denote by r_n the cardinality of F_n . If $r_n < \infty$, define $p_0 = 0$, $p_{n-1} = \min F_n - 1$, and $p_n = \max F_n$.

Proposition 8. *Let $\Theta \neq \pm\Theta_0$. Then*

$$C = \left\{ \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta} : t_n \geq 0, \sum_{n=1}^{\infty} t_n = 1 \right\} \quad (2.27)$$

does not have the FPP in the following cases.

- (1) There exists $k \geq 1$ such that $r_n \leq r_{n+k} < \infty$, $n \in \mathbb{N}$.
- (2) $r_1 = 1$ and $r_2 = \infty$.
- (3) There exists $\{i_n\}$, with $i_1 > 1$, such that for any $k, l \in \mathbb{N}$ with $i_{k-1} < l < i_k$ we have $\theta_l = \theta_{i_k}$ and also $\theta_k \neq \theta_{i_k}$ for all $k \geq 2$ or $\theta_k = \theta_{i_k}$ for all $k \geq 2$.

Proof. Let $\Theta \neq \pm\Theta_0$.

(1) If there exists $k \geq 1$ such that $r_n \leq r_{n+k} < \infty$, $n \in \mathbb{N}$, define $q_n = \sum_{j=n-(k-2)}^{n+1} r_j$, $n \geq k$ and $T : C \rightarrow C$ by

$$T \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta} = T \sum_{n=1}^{\infty} \sum_{i=p_{n-1}+1}^{p_n} t_i \zeta_i^{\Theta} = \sum_{n=k}^{\infty} \sum_{i=w_n}^{p_{n+1}} t_{i-q_n} \zeta_i^{\Theta}, \quad (2.28)$$

where $w_n = p_n + r_{n+1} - r_{n+1-k} + 1$. The idea is to translate the coefficients of $\sum_{n=1}^{\infty} t_n \zeta_n^{\Theta}$ in the block F_n into the last r_n terms of the block F_{n+k} . Then it is easy to see that T does not have fixed points. To prove that T is nonexpansive first observe that if k is even the signs of the θ_i and θ_j with $i \in F_n$ and $j \in F_{n+k}$ are the same and are different if k is odd. Now let $x = \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta}$, $y = \sum_{n=1}^{\infty} s_n \zeta_n^{\Theta}$, and $x - y = \sum_{n=1}^{\infty} \alpha_n \zeta_n^{\Theta}$. Then $\alpha_n = t_n - s_n$ and $\sum_{n=1}^{\infty} \alpha_n = 0$. Hence

$$x - y = \sum_{n=1}^{\infty} \sum_{i=p_{n-1}+2}^{p_n} \theta_{p_n} \left(\sum_{n=i}^{\infty} \alpha_n \right) e_i, \quad (2.29)$$

$$T(x - y) = \sum_{n=k}^{\infty} \left(\theta_{p_{n+1}} \sum_{n=i-q_n}^{\infty} \alpha_n \right) \left(\sum_{i=p_n+1}^{w_n} e_i \right) \sum_{i=w_n+1}^{p_{n+1}} \theta_{p_{n+1}} \left(\sum_{n=i-q_n}^{\infty} \alpha_n \right) e_i \quad (2.30)$$

are the expressions of $x - y$ and $T(x - y)$ with respect to the canonical basis. Since the coefficients in (2.29) and (2.30) are the same, or the same with opposite signs, with perhaps some repetitions in (2.30), T is an isometry.

(2) Suppose now that $r_1 = 1$ and $r_2 = \infty$. In this case, define $T \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta} = \sum_{n=1}^{\infty} t_n \zeta_{n+1}^{\Theta}$. Clearly T is nonexpansive and fixed-point-free.

(3) In this case it is straightforward to see that the operator $T : C \rightarrow C$ defined by $T \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta} = \sum_{n=1}^{\infty} t_n \zeta_{i_n}^{\Theta}$ is nonexpansive and does not have fixed points. $\square$

Proposition 9. *Let $\Theta \neq \pm \Theta_0$. Suppose Θ does not satisfy the hypotheses of the above proposition, and let $\{i_n\}$ be a sequence with $i_1 > 1$. Then the operator $T : C \rightarrow C$ defined by $T \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta} = \sum_{n=1}^{\infty} t_n \zeta_{i_n}^{\Theta}$ is expansive.*

Proof. Since Θ does not satisfy the hypotheses (1) and (2) of the above proposition, there are three possibilities.

- (I) $r_n < \infty$ for every $n \geq 2$; then for every k there exists n such that $r_{n+k} < r_n$.
- (II) $r_2 = \infty$; then $r_1 > 1$.
- (III) There exists $k > 2$ such that $r_k = \infty$.

Let $\{i_n\}$ be fixed with $i_1 > 1$ and denote $i_0 = 0$. Since Θ does not satisfy the hypotheses (3) of the above proposition, there exist k and l with $i_{k-1} < l < i_k$ such that $\theta_l \neq \theta_{i_k}$ or there exists $k_1 \geq 2$ with $\theta_{k_1} = \theta_{i_{k_1}}$ and there exists $k_2 \geq 2$ with $\theta_{k_2} \neq \theta_{i_{k_2}}$.

Case 1. For every k there exists n such that $r_{n+k} < r_n$.

There are two subcases.

Subcase 1.1. There are k and l with $i_{k-1} < l < i_k$ such that $\theta_l \neq \theta_{i_k}$.

Let $x = (1/8)\zeta_1^{\Theta} + (3/8)\zeta_{k-1}^{\Theta} + (1/2)\zeta_k^{\Theta}$ and $y = (1/16)\zeta_1^{\Theta} + (3/16)\zeta_{k-1}^{\Theta} + (3/4)\zeta_k^{\Theta}$. Then $x - y = (1/16)\zeta_1^{\Theta} + (3/16)\zeta_{k-1}^{\Theta} - (1/4)\zeta_k^{\Theta} = -(1/16)\sum_{i=2}^{k-1}\theta_i e_i - (1/4)\theta_k e_k$ and $\|x - y\|_D \leq 5/16$. On the other hand, $Tx - Ty = (1/16)\zeta_{i_1}^{\Theta} + (3/16)\zeta_{i_{k-1}}^{\Theta} - (1/4)\zeta_{i_k}^{\Theta} = -(1/16)\sum_{j=i_1+1}^{i_{k-1}}\theta_j e_j - (1/4)\sum_{j=i_{k-1}+1}^{i_k}\theta_j e_j$ and $\|Tx - Ty\|_D = 1/2$.

Subcase 1.2. For any $k \in \mathbb{N}$ and l with $i_{k-1} < l < i_k$, we have $\theta_l = \theta_{i_k}$.

There are two subsubcases. (1) $\theta_1 = \theta_{i_1}$ and (2) $\theta_1 \neq \theta_{i_1}$.

(1) $\theta_1 = \theta_{i_1}$

If $\theta_k = \theta_{i_k}$ for every k ; then we would have $F_1 = \mathbb{N}$, which implies $\Theta = \pm \Theta_0$. Then there is k such that $\theta_k \neq \theta_{i_k}$. Let $s = \min\{l : \theta_l \neq \theta_{i_l}\}$. Then $s > 1$.

There are two possibilities: (A) there exists $r > s$ such that $\theta_r = \theta_{i_r}$ and (B) $\theta_k \neq \theta_{i_k}$ for all $k \geq s$.

(A) Let $k + 1 = \min\{r > s : \theta_r = \theta_{i_r}\}$. We need to consider the following cases.

(a) $\theta_k = \theta_{k+1}$.

Let $x = (1/2)\zeta_{k-1}^{\Theta} + (1/2)\zeta_{k+1}^{\Theta}$ and $y = (3/4)\zeta_{k-1}^{\Theta} + (1/4)\zeta_{k+1}^{\Theta}$. Then $x - y = -(1/4)\zeta_{k-1}^{\Theta} + (1/4)\zeta_{k+1}^{\Theta} = \theta_{k+1}((1/4)e_k + (1/4)e_{k+1})$ and $\|x - y\|_D = 1/4$. On the other hand, $Tx - Ty = -(1/4)\zeta_{i_{k-1}}^{\Theta} + (1/4)\zeta_{i_{k+1}}^{\Theta} = (1/4)\sum_{j=i_{k-1}+1}^{i_k}\theta_j e_j + (1/4)\sum_{j=i_{k+1}+1}^{i_{k+1}}\theta_j e_j = -(1/4)\theta_{k+1}\sum_{j=i_{k-1}+1}^{i_k}e_j + (1/4)\theta_{k+1}\sum_{j=i_{k+1}+1}^{i_{k+1}}e_j$ and $\|Tx - Ty\|_D = 1/2$.

(b) $\theta_k \neq \theta_{k+1}$.

Let $x = (1/2)\zeta_{k-1}^\Theta + (1/2)\zeta_{k+1}^\Theta$ and $y = (3/4)\zeta_k^\Theta + (1/4)\zeta_{k+1}^\Theta$. Then $x - y = (1/2)\zeta_{k-1}^\Theta - (3/4)\zeta_k^\Theta + (1/4)\zeta_{k+1}^\Theta = -(1/2)\theta_k e_k + (1/4)\theta_{k+1} e_{k+1} = \theta_{k+1}((1/2)e_k + (1/4)e_{k+1})$ and $\|x - y\|_D = 1/2$. On the other hand, $Tx - Ty = (1/2)\zeta_{i_{k-1}}^\Theta - (3/4)\zeta_{i_k}^\Theta + (1/4)\zeta_{i_{k+1}}^\Theta = -(1/2)\sum_{j=i_{k-1}+1}^{i_k} \theta_j e_j + (1/4)\sum_{j=i_k+1}^{i_{k+1}} \theta_j e_j = -(1/2)\theta_k \sum_{j=i_{k-1}+1}^{i_k} e_j + (1/4)\theta_{k+1} \sum_{j=i_k+1}^{i_{k+1}} e_j$ and $\|Tx - Ty\|_D = 3/4$.

(B) $\theta_k \neq \theta_{i_k}$ for all $k \geq s$. By hypothesis we have that $s > 2$. There are two cases.

(a) $\theta_{s-1} = \theta_s$. Then $\theta_{i_{s-1}} \neq \theta_{i_s}$. Let $x = (1/4)\zeta_{s-2}^\Theta + (1/2)\zeta_{s-1}^\Theta + (1/4)\zeta_s^\Theta$ and $y = (1/4)\zeta_{s-1}^\Theta + (3/4)\zeta_s^\Theta$. Then $x - y = (1/4)\zeta_{s-2}^\Theta + (1/4)\zeta_{s-1}^\Theta - (1/2)\zeta_s^\Theta = -\theta_s((1/4)e_{s-1} + (1/2)e_s)$ and $\|x - y\|_D = 1/2$. On the other hand, $Tx - Ty = (1/4)\zeta_{i_{s-2}}^\Theta + (1/4)\zeta_{i_{s-1}}^\Theta - (1/2)\zeta_{i_s}^\Theta = -(1/4)\theta_s \sum_{j=i_{s-2}+1}^{i_{s-1}} e_j + (1/2)\theta_s \sum_{j=i_{s-1}+1}^{i_s} e_j$ and $\|Tx - Ty\|_D = 3/4$.

(b) $\theta_{s-1} \neq \theta_s$. Then $\theta_{i_{s-1}} = \theta_{i_s}$. Let $x = (1/4)\zeta_{s-2}^\Theta + (1/4)\zeta_{s-1}^\Theta + (1/2)\zeta_s^\Theta$ and $y = (3/4)\zeta_{s-1}^\Theta + (1/4)\zeta_s^\Theta$. Then $x - y = (1/4)\zeta_{s-2}^\Theta - (1/2)\zeta_{s-1}^\Theta + (1/4)\zeta_s^\Theta = -(1/4)\theta_{s-1} e_{s-1} + (1/4)\theta_s e_s = \theta_s((1/4)e_{s-1} + (1/4)e_s)$ and $\|x - y\|_D = 1/4$. On the other hand, $Tx - Ty = (1/4)\zeta_{i_{s-2}}^\Theta - (1/2)\zeta_{i_{s-1}}^\Theta + (1/4)\zeta_{i_s}^\Theta = -(1/4)\theta_{i_{s-1}} \sum_{j=i_{s-2}+1}^{i_{s-1}} e_j + (1/4)\theta_{i_s} \sum_{j=i_{s-1}+1}^{i_s} e_j = \theta_{s-1}(-(1/4)\sum_{j=i_{s-2}+1}^{i_{s-1}} e_j + (1/4)\sum_{j=i_{s-1}+1}^{i_s} e_j)$ and $\|Tx - Ty\|_D = 1/2$.

(2) $\theta_1 \neq \theta_{i_1}$

In this case there exists k such that $\theta_k = \theta_{i_k}$. If $s = \min\{l : \theta_l = \theta_{i_l}\}$, then $s > 1$.

Hence consider the cases: (A) there exists $r > s$ such that $\theta_r \neq \theta_{i_r}$ and (B) $\theta_k = \theta_{i_k}$ for all $k \geq s$ and proceed as in the Case (1).

Case 2. $r_2 = \infty$ and $r_1 > 1$.

Then $\theta_{p_1} \neq \theta_{i_{p_1}}$ with $1 < p_1$. Hence we can proceed as in Subcase 1.2(1)(A) above taking $k = p_1$.

Case 3. There is $s > 1$ such that $r_{s+1} = \infty$.

Then $\theta_{p_s} \neq \theta_{i_{p_s}}$ with $1 < p_s$. Hence we can proceed as in Subcase 1.2(1)(A) above taking $k = p_s$. $\square$

Next, for every selection of signs $\Theta \neq \pm\Theta_0$, we will define the asymptotically isometric c_{0D} - Θ -basic sequences. To this end, let us consider the following notation.

Let

$$\begin{aligned}\mathfrak{S}_\Theta &= \{(n, m) : \theta_n = \theta_m\}, \\ \mathfrak{D}_\Theta &= \{(n, m) : \theta_n \neq \theta_m\}.\end{aligned}\tag{2.31}$$

Definition 10. Let $\{x_n\}$ be a sequence in a Banach space X . We say that $\{x_n\}$ is an asymptotically isometric c_{0D} - Θ -basic sequence (*aiOb c_{0D}* sequence for short) if $\{x_n\}$ is convexly closed

and there exists $\{\varepsilon_n^\Theta\} \subset (0, (1/2))$ such that $\varepsilon_n^\Theta \searrow 0$, and

$$\begin{aligned} L(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{S}_\Theta) \vee L(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{D}_\Theta) &\leq \left\| \sum_{n=1}^{\infty} t_n x_n \right\| \\ &\leq R(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{S}_\Theta) \vee R(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{D}_\Theta) \end{aligned} \quad (2.32)$$

holds for all $\{t_n\} \in l_1$, where

$$\begin{aligned} L(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{S}_\Theta) &= \left(\sup_{n < l, (n, l) \in \mathfrak{S}_\Theta} (1 + \varepsilon_{l-1}^\Theta)^{-1} \left| \sum_{k=n}^{l-1} t_k \right| \right), \\ L(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{D}_\Theta) &= \left(\sup_{n < l, (n, l) \in \mathfrak{D}_\Theta} (1 + \varepsilon_{l-1}^\Theta)^{-1} \left| \sum_{k=n}^{l-1} t_k + 2 \sum_{k=l}^{\infty} t_k \right| \right), \\ R(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{S}_\Theta) &= \left(\sup_{n < l, (n, l) \in \mathfrak{S}_\Theta} (1 + \varepsilon_{l-1}^\Theta) \left| \sum_{k=n}^{l-1} t_k \right| \right), \\ R(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{D}_\Theta) &= \left(\sup_{n < l, (n, l) \in \mathfrak{D}_\Theta} (1 + \varepsilon_{l-1}^\Theta) \left| \sum_{k=n}^{l-1} t_k + 2 \sum_{k=l}^{\infty} t_k \right| \right). \end{aligned} \quad (2.33)$$

We are interested in $ai\Theta b_{C_0D}$ sequences for which the numbers ε_n^Θ of Definition 10 are small. We are taking $\{\varepsilon_n^\Theta\} \subset (0, (1/2))$.

We know that the set C of Proposition 3 does not have $aisbc_0$ sequences. Now we also prove that C does not contain $ai\Theta b_{C_0D}$ sequences with the norm $\|\cdot\|_D$ if $\Theta \neq \pm\Theta_0$.

Proposition 11. *Let $\Theta \neq \pm\Theta_0$. The set $C = \{\sum_{n=1}^{\infty} t_n \xi_n : t_n \geq 0 \text{ and } \sum_{n=1}^{\infty} t_n = 1\}$ does not contain $ai\Theta b_{C_0D}$ sequences with the norm $\|\cdot\|_D$.*

Proof. Let $\{y_k\} \subset C$. Then $y_k = \sum_{n=1}^{\infty} \lambda_n^k \xi_n$ for some $\lambda_n^k \geq 0$ with $\sum_{n=1}^{\infty} \lambda_n^k = 1$. Suppose that $\{y_k\}$ is an $ai\Theta b_{C_0D}$ with $\|\cdot\|_D$. Since $\Theta \neq \pm\Theta_0$, there exist $m \in \mathbb{N}$ and $\{n_k\} \subset \mathbb{N}$ with $n_1 < n_2 < \dots$, such that for all $k \in \mathbb{N}$, $m < n_k$ and $\theta_{n_k} \neq \theta_m$. Let $t_n = 0$ for $n \neq m, n_k$ and $t_m = t_{n_k} = 1$. Thus

$$\begin{aligned} (1 + \varepsilon_{n_k-1}^\Theta)^{-1} 3 &\leq L(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{S}_\Theta) \vee L(\{\varepsilon_n^\Theta\}, \{t_n\}, \mathfrak{D}_\Theta) \\ &\leq \left\| \sum_{n=1}^{\infty} t_n y_n \right\|_D = \|y_m + y_{n_k}\|_D = 2. \end{aligned} \quad (2.34)$$

Since (2.34) holds for all $k \in \mathbb{N}$, making $k \rightarrow \infty$ in (2.34), we get that $3 \leq 2$, which is a contradiction. $\square$

Proposition 12. Let $\Theta_1 = \{\theta_i^1\}_i$ and $\Theta_2 = \{\theta_i^2\}_i$ such that $\Theta_1 \neq \pm\Theta_2$ and $\Theta_1, \Theta_2 \neq \pm\Theta_0$. Let $\{\zeta_n^{\Theta_1}\}$ be the Θ_1 -basis of c_0 considered in $(c_0, \|\cdot\|_D)$ and let

$$C(\Theta_1) = \left\{ \sum_{n=1}^{\infty} t_n \zeta_n^{\Theta_1} : t_n \geq 0, \sum_{n=1}^{\infty} t_n = 1 \right\}. \quad (2.35)$$

The set $C(\Theta_1)$ does not contain $ai\Theta_2bc_{0D}$ sequences with the norm $\|\cdot\|_D$.

Proof. Let $\{y_k\} \subset C$. Then $y_k = \sum_{n=1}^{\infty} \lambda_n^k \zeta_n^{\Theta_1}$ for some $\lambda_n^k \geq 0$ with $\sum_{n=1}^{\infty} \lambda_n^k = 1$. Suppose that $\{y_k\}$ is an $ai\Theta_2bc_{0D}$ with the norm $\|\cdot\|_D$.

Suppose first $\theta_1^1 = \theta_1^2$; since $\Theta_1 \neq \Theta_2$, there exists $m > 1$ such that $\theta_m^1 \neq \theta_m^2$.

There are two cases.

Case 1. $(1, m) \in \mathfrak{S}_{\Theta_1}$. In this case $(1, m) \in \mathfrak{D}_{\Theta_2}$. Let $t_n = 0$ for $n \neq 1, m$ and $t_1 = t_m = 1$. Thus

$$\begin{aligned} \left(1 + \varepsilon_{m-1}^{\Theta_2}\right)^{-1} 3 &\leq L\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{S}_{\Theta_2}\right) \vee L\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{D}_{\Theta_2}\right) \\ &\leq \left\| \sum_{n=1}^{\infty} t_n y_n \right\|_D = \|y_1 + y_m\|_D \leq 2. \end{aligned} \quad (2.36)$$

Since $\varepsilon_{m-1}^{\Theta_2} < 1/2$, we get a contradiction.

Case 2. $(1, m) \in \mathfrak{D}_{\Theta_1}$. In this case $(1, m) \in \mathfrak{S}_{\Theta_2}$. Let $t_n = 0$ for $n \neq 1, m$ and $t_1 = t_m = 1$. Thus

$$\begin{aligned} 2 &\leq \sum_{n=1}^{\infty} \lambda_n^1 + \sum_{n=1}^{\infty} \lambda_n^m + \sum_{n=m}^{\infty} \lambda_n^1 + \sum_{n=m}^{\infty} \lambda_n^m \leq \left\| \sum_{n=1}^{\infty} t_n y_n \right\|_D = \|y_1 + y_m\|_D \\ &\leq R\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{S}_{\Theta_2}\right) \vee R\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{D}_{\Theta_2}\right) \leq \left(1 + \varepsilon_{m-1}^{\Theta_2}\right). \end{aligned} \quad (2.37)$$

Since $\varepsilon_{m-1}^{\Theta_2} < 1/2$, we get a contradiction.

Suppose now $\theta_1^1 \neq \theta_1^2$; since $\Theta_1 \neq -\Theta_2$, there exists m such that $\theta_m^1 = \theta_m^2$.

There are two cases.

Case 1. $(1, m) \in \mathfrak{S}_{\Theta_1}$; in this case $(1, m) \in \mathfrak{D}_{\Theta_2}$. Let $t_n = 0$ for $n \neq 1, m$ and $t_1 = t_m = 1$. Thus

$$\begin{aligned} \left(1 + \varepsilon_{m-1}^{\Theta_2}\right)^{-1} 3 &\leq L\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{S}_{\Theta_2}\right) \vee L\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{D}_{\Theta_2}\right) \\ &\leq \left\| \sum_{n=1}^{\infty} t_n y_n \right\|_D = \|y_1 + y_m\|_D \leq 2. \end{aligned} \quad (2.38)$$

Since $\varepsilon_{m-1}^{\Theta_2} < 1/2$, we get a contradiction.

Case 2. $(1, m) \in \mathfrak{D}_{\Theta_1}$. In this case $(1, m) \in \mathfrak{S}_{\Theta_2}$. Let $t_n = 0$ for $n \neq 1, m$ and $t_1 = t_m = 1$. Thus

$$\begin{aligned} 2 &\leq \sum_{n=1}^{\infty} \lambda_n^1 + \sum_{n=1}^{\infty} \lambda_n^m + \sum_{n=m}^{\infty} \lambda_n^1 + \sum_{n=m}^{\infty} \lambda_n^m \leq \left\| \sum_{n=1}^{\infty} t_n y_n \right\|_D = \|y_1 + y_m\|_D \\ &\leq R\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{S}_{\Theta_2}\right) \vee R\left(\left\{\varepsilon_n^{\Theta_2}\right\}, \{t_n\}, \mathfrak{D}_{\Theta_2}\right) \leq \left(1 + \varepsilon_{m-1}^{\Theta_2}\right). \end{aligned} \quad (2.39)$$

Since $\varepsilon_{m-1}^{\Theta_2} < 1/2$, we get a contradiction. $\square$

Propositions 3, 7, and 11 show that, in contrast with Theorem 4 of the Dowling et al. paper [2] for *aisbc*₀ sequences in c_0 , in the space $(c_0, \|\cdot\|_D)$ we need an infinite number of sequences (at least *aisbc*₀ and *aiObc*_{0D} sequences) to have a similar result.

3. The Space $(c_0, \|\cdot\|_D)^*$

It is known that the dual of the Bynum space c_{01} is the Bynum space $l_{1\infty}$. Below we prove that the dual space of $(c_0, \|\cdot\|_D)$ when the scalar field is the set of real numbers is also the Bynum space $l_{1\infty}$. Let us suppose then that $\mathbb{K} = \mathbb{R}$. First we calculate the extreme points of the unit ball of $(c_0, \|\cdot\|_D)$.

Lemma 13. *Let $X = (c_0, \|\cdot\|_D)$. Then we have*

$$\mathcal{E}(B_X) = \{\{x_n\} \in S_X : x_n \in \{1, 0\}, n \in \mathbb{N}\} \cup \{\{x_n\} \in S_X : x_n \in \{-1, 0\}, n \in \mathbb{N}\}. \quad (3.1)$$

Proof. First note that if $\{x_n\} \in S_X$ then $|x_n - x_m| \leq 1$, $n, m \in \mathbb{N}$ and $|x_n| \leq 1$, $n \in \mathbb{N}$. Consequently, if $\{x_n\} \in S_X$ with $x_{n_0} = 1$ for some $n_0 \in \mathbb{N}$, then $0 \leq x_n \leq 1$, for all $n \in \mathbb{N}$. Analogously if $\{x_n\} \in S_X$ with $x_{n_0} = -1$ for some $n_0 \in \mathbb{N}$, then $-1 \leq x_n \leq 0$, for all $n \in \mathbb{N}$. Let $A = \{\{x_n\} \in S_X : x_n \in \{1, 0\}, n \in \mathbb{N}\}$ and $B = \{\{x_n\} \in S_X : x_n \in \{-1, 0\}, n \in \mathbb{N}\}$. Thus $A, B \subset S_X$.

Take $x = \{x_n\} \in A$ and suppose that $x = (y + z)/2$ with $y, z \in S_X$. Also suppose that $y = \{y_n\}$ and $z = \{z_n\}$. Since $x \in A$, there exists $n_0 \in \mathbb{N}$ such that $x_{n_0} = 1$. Since $x_n = (y_n + z_n)/2$ and $y_n, z_n \leq 1$, if $x_n = 1$ for some $n \in \mathbb{N}$, we have that $x_n = y_n = z_n$. Thus $x_{n_0} = y_{n_0} = z_{n_0} = 1$. On the other hand, if $x_n = 0$ for some $n \in \mathbb{N}$, we also have that $x_n = y_n = z_n$, because if $y_n < 0$ we get that $|y_n - y_{n_0}| > 1$, which contradicts that $|y_n - y_m| \leq 1$, $n, m \in \mathbb{N}$ and if $y_n > 0$ then $z_n < 0$ and we also have a contradiction. Therefore, $x = y = z$. Hence $x \in \mathcal{E}(B_X)$. Thus $A \subset \mathcal{E}(B_X)$. Analogously $B \subset \mathcal{E}(B_X)$.

Take now $x = \{x_n\} \in S_X \setminus \{A \cup B\}$. Then there exists $n_0 \in \mathbb{N}$ such that $0 < |x_{n_0}| < 1$. Let $a = \inf_n x_n$ and $b = \sup_n x_n$. If $x_{n_0} \in (a, b)$, define $c = \min(|x_{n_0} - a|, |x_{n_0} - b|)$, $y_n = z_n = x_n$, $n \neq n_0$, $y_{n_0} = x_{n_0} - c$, and $z_{n_0} = x_{n_0} + c$. Thus, $x_n = (y_n + z_n)/2$ with $y, z \in S_X$ and $x \neq y$, $x \neq z$. Therefore, $x \in S_X \setminus \mathcal{E}(B_X)$. Suppose now that $x_{n_0} = a$ or $x_{n_0} = b$. Since $0 < |x_{n_0}| < 1$ and $\sup_{n, m \in \mathbb{N}} |x_n - x_m| = 1$, we have that $0 \in (a, b)$. Since $x_n \rightarrow 0$, there exists $n_1 > n_0$ such that $x_{n_1} \in (a, b)$, which implies that $x \in S_X \setminus \mathcal{E}(B_X)$. Consequently, $\mathcal{E}(B_X) \subset A \cup B$. $\square$

Theorem 14. Let $f \in (c_0, \|\cdot\|_D)^*$. There exists a unique sequence $\{c_n\} \in l_1$ such that $f = \sum_{n=1}^{\infty} c_n e_n^*$ and

$$\|f\|_D = \max\left(\sum_{n=1}^{\infty} c_n^+, \sum_{n=1}^{\infty} c_n^-\right), \quad (3.2)$$

where $c_n^+ = \max(c_n, 0)$ and $c_n^- = -\min(c_n, 0)$.

Proof. Let $f \in (c_0, \|\cdot\|_D)^*$. Since $\{e_n\}$ is a shrinking basis of $(c_0, \|\cdot\|_D)$, there exists a unique sequence $\{c_n\} \subset \mathbb{K}$ such that $f = \sum_{n=1}^{\infty} c_n e_n^*$. As sets $(c_0, \|\cdot\|_D)^* = (c_0)^*$ and hence $f \in (c_0)^*$. Thus $f = R\{a_n\}$ where $R : l_1 \rightarrow c_0^*$ is the Riesz representation. Consequently,

$$c_n = f(e_n) = R\{a_n\}(e_n) = a_n. \quad (3.3)$$

Therefore, $\{c_n\} = \{a_n\} \in l_1$. Thus

$$\begin{aligned} \|f\|_D &= \sup_{x \in B_X} |f(x)| = \sup_{x \in \mathcal{E}(B_X)} |f(x)| \\ &= \sup \left\{ \left| \sum_{n \in F} c_n \right| : F \subset \mathbb{N}, F \text{ finite} \right\} \\ &= \max \left(\sum_{n=1}^{\infty} c_n^+, \sum_{n=1}^{\infty} c_n^- \right), \end{aligned} \quad (3.4)$$

where $c_n^+ = \max(c_n, 0)$ and $c_n^- = -\min(c_n, 0)$. □

Corollary 15. $(c_0, \|\cdot\|_D)^*$ is the Bynum space $l_{1\infty}$ and it has the ω -FPP.

Remark 16. It is well known that $l_1(c_0)^*$ has the ω^* fixed point property for left reversible semigroups, that is, whenever S is a semigroup such that $aS \cap bS \neq \emptyset$ for any $a, b \in S$, and $\mathcal{S} = \{T_s : s \in S\}$ is a representation of S as nonexpansive mappings on a nonempty ω^* -compact convex subset K of l_1 , there is a common fixed point in K for $\mathcal{S}$. (see [10–12]). In particular, l_1 has the ω^* fixed point property. Is this the case for $(c_0, \|\cdot\|_D)^*$?

Next we will see that every infinite-dimensional subspace of $l_{1\infty}$ contains an asymptotically isometric copy of l_1 and then, by a result of Dowling and Lennard [13], it does not have the FPP.

First recall that a Banach space $(X, \|\cdot\|)$ contains an asymptotically isometric copy of l_1 if there exists $\{x_n\}_n \subset X$ and $\{\varepsilon_n\} \subset (0, 1)$, $\varepsilon_n \rightarrow 0$ such that for every $k \in \mathbb{N}$ and every scalars $b_1, \dots, b_k$,

$$\sum_{i=1}^k (1 - \varepsilon_i) |b_i| \leq \left\| \sum_{i=1}^k b_i x_i \right\| \leq \sum_{i=1}^k (1 + \varepsilon_i) |b_i|. \quad (3.5)$$

In this case we say that $\{x_n\}_n$ is an asymptotically isometric l_1 -sequence (*ail*-sequence for short).

Observe that if $\{y_n\}_n$ is another sequence in X such that $\|y_n - x_n\| < \delta_n$ for all n , where $\{\varepsilon_n + \delta_n\} \subset (0, 1)$ and $\delta_n \rightarrow 0$, then for every k and every scalars $b_1, \dots, b_k$,

$$\sum_{i=1}^k (1 - \varepsilon_i - \delta_i) |b_i| \leq \left\| \sum_{i=1}^k b_i y_i \right\| \leq \sum_{i=1}^k (1 + \varepsilon_i + \delta_i) |b_i| \quad (3.6)$$

and $\{y_n\}$ is also an ail_1 -sequence.

Proposition 17. Let $\{u_i\}_i \subset l_{1\infty}$, and let $\{n_i\}$ be a strictly increasing sequence in $\mathbb{N}$ such that $u_i = \sum_{j=n_i+1}^{n_{i+1}} a_j^i e_j$. If $\sum_{j=n_i+1}^{n_{i+1}} (a_j^i)^+ = \sum_{j=n_i+1}^{n_{i+1}} (a_j^i)^-$, then $\{u_i\}_i$ is isometrically equivalent to the canonical basis in l_1 , that is, for every $k \in \mathbb{N}$ and every scalars $b_1, \dots, b_k$, we have that $\|\sum_{i=1}^k b_i u_i\| = \sum_{i=1}^k |b_i|$.

Proof. Let $b_1, \dots, b_k$ be scalars; then

$$\begin{aligned} \sum_{i=1}^k |b_i| &\leq \sum_{i=1}^k b_i^+ \sum_{j=n_i+1}^{n_{i+1}} (a_j^i)^+ + \sum_{i=1}^k b_i^- \sum_{j=n_i+1}^{n_{i+1}} (a_j^i)^- \\ &= \sum_{i=1}^k \sum_{j=n_i+1}^{n_{i+1}} (b_i a_j^i)^+ \\ &\leq \left\| \sum_{i=1}^k b_i u_i \right\|_{1\infty} \leq \sum_{i=1}^k |b_i|. \end{aligned} \quad (3.7)$$

□

Theorem 18. Every infinite-dimensional subspace of $l_{1\infty}$ contains an asymptotically isometric copy of l_1 and hence it does not have the FPP.

Proof. Let Y be an infinite-dimensional subspace of $l_{1\infty}$, $\{\varepsilon_n\} \subset (0, (1/2))$, $\varepsilon_n \searrow 0$ and $\{x_n\}$ a sequence in S_Y such that $x_i = \sum_{j=m_i+1}^{\infty} a_j^i e_j$, where $0 = m_0 < m_1 < \dots$ and $\sum_{j=m_{i+1}+1}^{\infty} |a_j^i| < \varepsilon_i/8$. Define

$$\begin{aligned} w_i &= \sum_{j=m_i+1}^{m_{i+1}} a_j^i e_j, \\ c_i^+ &= \frac{1}{\|w_i\|_{1\infty}} \sum_{j=m_i+1}^{m_{i+1}} (a_j^i)^+ \leq 1, \\ c_i^- &= \frac{1}{\|w_i\|_{1\infty}} \sum_{j=m_i+1}^{m_{i+1}} (a_j^i)^- \leq 1. \end{aligned} \quad (3.8)$$

Changing w_i by $-w_i$, if necessary, we can assume that $c_i^+ = 1$, $n \in \mathbb{N}$. If there is a sequence $\{k_i\}$ such that $c_{k_i}^- = 1$, then by Proposition 17, $\{w_{k_i}/\|w_{k_i}\|_{1\infty}\}$ is isometrically equivalent to the canonical basis of l_1 . It is straightforward to see that $\|x_{k_i} - (w_{k_i}/\|w_{k_i}\|_{1\infty})\|_{1\infty} < (1/4)\varepsilon_{k_i}$. Then by the above remark, $\{x_{k_i}\}$ is an ail_1 -sequence.

Suppose that $c_i^- \neq 1$ for all i and let

$$\alpha_i = \frac{1 - c_{2i}^-}{1 - c_{2i}^- c_{2i-1}^-}, \quad \beta_i = \frac{1 - c_{2i-1}^-}{1 - c_{2i-1}^- c_{2i}^-}. \quad (3.9)$$

Then $0 \leq \alpha_i < 1$, $0 \leq \beta_i < 1$ and

$$\alpha_i c_{2i-1}^+ + \beta_i c_{2i}^- = \alpha_i c_{2i-1}^- + \beta_i c_{2i}^+ = 1. \quad (3.10)$$

Now let

$$v_i = \alpha_i \frac{w_{2i-1}}{\|w_{2i-1}\|_{1\infty}} - \beta_i \frac{w_{2i}}{\|w_{2i}\|_{1\infty}}. \quad (3.11)$$

Suppose that $v_i = \sum_{j=m_{2i-1}+1}^{m_{2i+1}} b_j^i e_j$. It is easy to check, using (3.10), that

$$\sum_{j=m_{2i-1}+1}^{m_{2i+1}} (b_j^i)^+ = \sum_{j=m_{2i-1}+1}^{m_{2i+1}} (b_j^i)^- = 1. \quad (3.12)$$

Hence, by Proposition 17, $\{v_i\}$ is isometrically equivalent to the canonical basis of l_1 .

Now, if we define $y_n = \alpha_n x_{2n-1} - \beta_n x_{2n} \in Y$, it is straightforward to see that $\|y_n - v_n\|_{1\infty} < \varepsilon_n$ and by the above remark, $\{y_n\}$ is an ail_1 -sequence.

Finally in [13] Dowling and Lennard proved that if a Banach space contains an ail_1 -sequence, then it does not have the FPP. Hence Y does not have the FPP. $\square$

Acknowledgments

This work was partially funded by Grant SEP CONACYT 102380 and partially supported by Conacyt Scholarship 165342 and CIMAT Scholarship.

References

- [1] J. Hagler, "A counterexample to several questions about Banach spaces," *Studia Mathematica*, vol. 60, no. 3, pp. 289–308, 1977.
- [2] P. N. Dowling, C. J. Lennard, and B. Turett, "Characterizations of weakly compact sets and new fixed point free maps in c_0 ," *Studia Mathematica*, vol. 154, no. 3, pp. 277–293, 2003.
- [3] P. N. Dowling, C. J. Lennard, and B. Turett, "Weak compactness is equivalent to the fixed point property in c_0 ," *Proceedings of the American Mathematical Society*, vol. 132, no. 6, pp. 1659–1666, 2004.
- [4] P. N. Dowling, C. J. Lennard, and B. Turett, "Reflexivity and the fixed-point property for nonexpansive maps," *Journal of Mathematical Analysis and Applications*, vol. 200, no. 3, pp. 653–662, 1996.
- [5] W. L. Bynum, "A class of spaces lacking normal structure," *Compositio Mathematica*, vol. 25, pp. 233–236, 1972.
- [6] P. N. Dowling, W. B. Johnson, C. J. Lennard, and B. Turett, "The optimality of James's distortion theorems," *Proceedings of the American Mathematical Society*, vol. 125, no. 1, pp. 167–174, 1997.
- [7] J. García Falset, "Basis and fixed points for nonexpansive mappings," *Radovi Matematički*, vol. 8, no. 1, pp. 67–75, 1992/96.
- [8] P. N. Dowling, C. J. Lennard, and B. Turett, "Asymptotically isometric copies of c_0 in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 219, no. 2, pp. 377–391, 1998.

- [9] P. N. Dowling, C. J. Lennard, and B. Turett, "Failure of the FPP inside an asymptotically isometric-free copy of c_0 ," *Nonlinear Analysis, Theory, Methods and Applications*, vol. 73, no. 5, pp. 1175–1179, 2010.
- [10] R. D. Holmes and A. T. Lau, "Non-expansive actions of topological semigroups and fixed points," *Journal of the London Mathematical Society Second Series*, vol. 5, pp. 330–336, 1972.
- [11] A. T.-M. Lau and P. F. Mah, "Fixed point property for Banach algebras associated to locally compact groups," *Journal of Functional Analysis*, vol. 258, no. 2, pp. 357–372, 2010.
- [12] T. C. Lim, "Asymptotic centers and nonexpansive mappings in conjugate Banach spaces," *Pacific Journal of Mathematics*, vol. 90, no. 1, pp. 135–143, 1980.
- [13] P. N. Dowling and C. J. Lennard, "Every nonreflexive subspace of $L_1[0,1]$ fails the fixed point property," *Proceedings of the American Mathematical Society*, vol. 125, no. 2, pp. 443–446, 1997.