Research Article

Analogue of Lebesgue-Radon-Nikodym Theorem with respect to p -adic q -Measure on $\mathbb{Z}_p$

T. Kim,¹ D. V. Dolgy,² S. H. Lee,³ and C. S. Ryoo⁴

¹ Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

² Hanrimwon, Kwangwoon University, Seoul 139-701, Republic of Korea

³ Division of General Education, Kwangwoon University, Seoul 139-701, Republic of Korea

⁴ Department of Mathematics, Hannam University, Daejeon 306-791, Republic of Korea

Correspondence should be addressed to T. Kim, tkkim@kw.ac.kr

Received 3 July 2011; Accepted 12 August 2011

Academic Editor: H. B. Thompson

Copyright © 2011 T. Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently, Lebesgue-Radon-Nikodym theorem with respect to fermionic p -adic invariant measure on $\mathbb{Z}_p$ was studied in Kim. In this paper we will give the analogue of the Lebesgue-Radon-Nikodym theorem with respect to p -adic q -measure on $\mathbb{Z}_p$. In special case, $q = 1$, we can derive the same results in Kim.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, the symbols $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ denote the ring of p -adic integers, the field of p -adic rational numbers, and the p -adic completion of the algebraic closure of $\mathbb{Q}_p$, respectively. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p| = p^{-v_p(p)} = 1/p$ and $v_p(0) = \infty$.

When one speaks of q -extension, q can be regarded as an indeterminate, a complex $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. In this paper we assume that $q \in \mathbb{C}_p$ with $|1 - q| < 1$, and we use the notations of q -numbers as follows:

$$[x]_q = [x : q] = \frac{1 - q^x}{1 - q}, \quad [x]_{-q} = \frac{1 - (-q)^x}{1 + q}. \quad (1.1)$$

For any positive integer N , let

$$a + p^N \mathbb{Z}_p = \left\{ x \in \mathbb{Z}_p \mid x \equiv a \pmod{p^N} \right\}, \quad (1.2)$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < p^N$ (see [1–8]).

It is known that the fermionic p -adic q -measure on $\mathbb{Z}_p$ is given by Kim as follows:

$$\mu_{-q}(a + p^n \mathbb{Z}_p) = \frac{(-q)^a}{[p^N]_{-q}} = \frac{1+q}{1+q^{p^N}} (-q)^a, \quad (1.3)$$

(see [6, 9–12]). Let $C(\mathbb{Z}_p)$ be the space of continuous functions on $\mathbb{Z}_p$. From (1.3), the fermionic p -adic q -integral on $\mathbb{Z}_p$ is defined by Kim as follows:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \quad (1.4)$$

where $f \in C(\mathbb{Z}_p)$ (see [1, 6, 9–12]).

Various proofs of the Radon-Nikodym theorem can be found in many books on measure theory, analysis, or probability theory. Usually they use the Hahn decomposition theorem for signed measure, the Riesz representation theorem for functionals on Hilbert space, or a martingale theory (see [13, 14]). In the previous paper [3], the author has studied the analogue of the Lebesgue-Radon-Nikodym theorem with respect to fermionic p -adic invariant measure on $\mathbb{Z}_p$. The purpose of this paper is to derive the analogue of the Lebesgue-Radon-Nikodym theorem with respect to p -adic q -measure on $\mathbb{Z}_p$ in the sense of fermionic.

2. Lebesgue-Radon-Nikodym's Type Theorem with respect to p -Adic q -Measure on $\mathbb{Z}_p$

For any positive integer a and n with $a < p^n$ and $f \in C(\mathbb{Z}_p)$, let us define

$$\mu_{f,-q}(a + p^n \mathbb{Z}_p) = \int_{a+p^n \mathbb{Z}_p} f(x) d\mu_{-q}(x), \quad (2.1)$$

where the integral is the fermionic p -adic q -integral on $\mathbb{Z}_p$.

From (1.3), (1.4), and (2.1), we note that

$$\begin{aligned} \mu_{f,-q}(a + p^n \mathbb{Z}_p) &= \lim_{m \rightarrow \infty} \frac{1}{[p^{m+n}]_{-q}} \sum_{x=0}^{p^{m+n}-1} f(a + p^n x) (-q)^{a+p^n x} \\ &= \lim_{m \rightarrow \infty} \frac{1}{[p^m]_{-q}} \sum_{x=0}^{p^{m-n}-1} f(a + p^n x) (-q)^a q^{p^n x} (-1)^x \\ &= \frac{[2]_q}{[2]_{q^{p^n}}} (-q)^a \lim_{m \rightarrow \infty} \frac{1}{[p^{m-n}]_{-q^{p^n}}} \sum_{x=0}^{p^{m-n}-1} f(a + p^n x) (-q^{p^n})^x \\ &= \frac{[2]_q}{[2]_{q^{p^n}}} (-q)^a \int_{\mathbb{Z}_p} f(a + p^n x) d\mu_{-q^{p^n}}(x). \end{aligned} \quad (2.2)$$

By (2.2), we get

$$\mu_{f,-q}(a + p^n \mathbb{Z}_p) = \frac{[2]_q}{[2]_{q^{p^n}}} (-q)^a \int_{\mathbb{Z}_p} f(a + p^n x) d\mu_{-q^{p^n}}(x). \quad (2.3)$$

Therefore, by (2.3), we obtain the following theorem.

Theorem 2.1. For $f, g \in C(\mathbb{Z}_p)$, one has

$$\mu_{\alpha f + \beta g, -q} = \alpha \mu_{f, -q} + \beta \mu_{g, -q}, \quad (2.4)$$

where α, β are constants.

From (2.2) and (2.4), we note that

$$|\mu_{f, -q}(a + p^n \mathbb{Z}_p)| \leq M \|f\|_\infty, \quad (2.5)$$

where $\|f\|_\infty = \sup_{x \in \mathbb{Z}_p} |f(x)|$ and M is some positive constant.

Now, we recall the definition of the strongly fermionic p -adic q -measure on $\mathbb{Z}_p$. If μ_{-q} is satisfied the following equation:

$$\left| \mu_{-q}(a + p^n \mathbb{Z}_p) - \mu_{-q}(a + p^{n+1} \mathbb{Z}_p) \right| \leq \delta_{n,q}, \quad (2.6)$$

where $\delta_{n,q} \rightarrow 0$ and $n \rightarrow \infty$ and $\delta_{n,q}$ is independent of a , then μ_{-q} is called the weakly fermionic p -adic q -measure on $\mathbb{Z}_p$.

If $\delta_{n,q}$ is replaced by $Cp^{-v_p(1-q^n)}$ (C is some constant), then μ_{-q} is called strongly fermionic p -adic q -measure on $\mathbb{Z}_p$.

Let $P(x) \in C_p[[x]]_q$ be an arbitrary q -polynomial with $\sum a_i [x]_q^i$. Then we see that $\mu_{P,-q}$ is strongly fermionic p -adic q -measure on $\mathbb{Z}_p$. Without a loss of generality, it is enough to prove the statement for $P(x) = [x]_q^k$.

Let a be an integer with $0 \leq a < p^n$. Then we get

$$\begin{aligned} \mu_{P,-q}(a + p^n \mathbb{Z}_p) &= \frac{[2]_q}{[2]_{q^{p^n}}} (-q)^a \lim_{m \rightarrow \infty} \frac{1}{[p^{m-n}]_{-q^{p^n}}} \sum_{i=0}^{p^{m-n}} [a + ip^n]_q^k (-1)^i q^{p^n i}, \\ q^{p^n i} &= \sum_{l=0}^i \binom{i}{l} [p^n]_q^l (q-1)^l. \end{aligned} \quad (2.7)$$

By (2.7), we easily get

$$\begin{aligned} \mu_{P,-q}(a + p^n \mathbb{Z}_p) &\equiv \frac{[2]_q}{[2]_{q^{p^n}}} (-q)^a [a]_q^k \pmod{[p^n]_q} \\ &\equiv \frac{[2]_q}{[2]_{q^{p^n}}} (-q)^a P(a) \pmod{[p^n]_q}. \end{aligned} \quad (2.8)$$

Let x be an arbitrary in $\mathbb{Z}_p$ with $x \equiv x_n \pmod{p^n}$ and $x \equiv x_{n+1} \pmod{p^{n+1}}$, where x_n and x_{n+1} are positive integers such that $0 \leq x_n < p^n$ and $0 \leq x_{n+1} < p^{n+1}$. Thus, by (2.8), we have

$$\left| \mu_{P,-q}(a + p^n \mathbb{Z}_p) - \mu_{P,-q}(a + p^{n+1} \mathbb{Z}_p) \right| \leq Cp^{-v_p(1-q^{p^n})}, \quad (2.9)$$

where C is a positive some constant and $n \gg 0$.

Let

$$f_{\mu_{P,-q}}(a) = \lim_{n \rightarrow \infty} \mu_{P,-q}(a + p^n \mathbb{Z}_p). \quad (2.10)$$

Then, (2.5), (2.7), and (2.8), we get

$$f_{\mu_{P,-q}}(a) = \frac{[2]_q}{2} (-q)^a [a]_q^k = \frac{[2]_q}{2} (-q)^a P(a). \quad (2.11)$$

Since $f_{\mu_{P,-q}}(x)$ is continuous on $\mathbb{Z}_p$, it follows for all $x \in \mathbb{Z}_p$

$$f_{\mu_{P,-q}}(x) = \frac{[2]_q}{2} (-q)^x P(x). \quad (2.12)$$

Let $g \in C(\mathbb{Z}_p)$. By (2.10), (2.11), and (2.12), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} g(x) d\mu_{P,-q}(x) &= \lim_{m \rightarrow \infty} \sum_{i=0}^{p^m-1} g(i) \mu_{P,-q}(i + p^m \mathbb{Z}_p) \\ &= \frac{[2]_q}{2} \sum_{i=0}^{p^m-1} g(i) (-q)^i [i]_q^k \\ &= \int_{\mathbb{Z}_p} g(x) [x]_q^k d\mu_{-q}(x). \end{aligned} \quad (2.13)$$

Therefore, by (2.13), we obtain the following theorem.

Theorem 2.2. Let $P(x) \in C_p[[x]]_q$ be an arbitrary q -polynomial with $\sum a_i [x]_q^i$. Then $\mu_{P,-q}$ is a strongly fermionic p -adic q -measure on $\mathbb{Z}_p$ and for all $x \in \mathbb{Z}_p$

$$f_{\mu_{P,-q}} = (-1)^x \frac{[2]_q}{2} q^x P(x). \quad (2.14)$$

Furthermore, for all $g \in C(\mathbb{Z}_p)$, one has

$$\int_{\mathbb{Z}_p} g(x) d\mu_{P,-q}(x) = \int_{\mathbb{Z}_p} g(x) P(x) d\mu_{-q}(x), \quad (2.15)$$

where the second integral is fermionic p -adic q -integral on $\mathbb{Z}_p$.

Let $f(x) = \sum_{n=0}^{\infty} a_{n,q} \binom{x}{n}_q$ be the q -Mahler expansion of continuous function on $\mathbb{Z}_p$, where

$$\binom{x}{n}_q = \frac{[x]_q [x-1]_q \cdots [x-n+1]_q}{[n]_q!} \quad (2.16)$$

(see [4]). Then we note that $\lim_{m \rightarrow \infty} |a_{n,q}| = 0$.

Let

$$f_m(x) = \sum_{i=0}^m a_{i,q} \binom{x}{i}_q \in C_p[[x]]_q. \quad (2.17)$$

Then

$$\|f - f_m\|_{\infty} \leq \sup_{m \leq n} |a_{n,q}|. \quad (2.18)$$

Writing $f = f_m + f - f_m$, we easily get

$$\begin{aligned} & \left| \mu_{f,-q}(a + p^n \mathbb{Z}_p) - \mu_{f,-q}(a + p^{n+1} \mathbb{Z}_p) \right| \\ & \leq \max \left\{ \left| \mu_{f_m,-q}(a + p^n \mathbb{Z}_p) - \mu_{f_m,-q}(a + p^{n+1} \mathbb{Z}_p) \right|, \right. \\ & \quad \left. \left| \mu_{f-f_m,-q}(a + p^n \mathbb{Z}_p) - \mu_{f-f_m,-q}(a + p^{n+1} \mathbb{Z}_p) \right| \right\}. \end{aligned} \quad (2.19)$$

From Theorem 2.2, we note that

$$\left| \mu_{f-f_m,-q}(a + p^n \mathbb{Z}_p) \right| \leq \|f - f_m\|_{\infty} \leq C_1 p^{-v_p(1-q^{p^n})}, \quad (2.20)$$

where C_1 is some positive constant.

For $m \gg 0$, we have $\|f\|_{\infty} = \|f_m\|_{\infty}$.

So

$$\left| \mu_{f_m,-q}(a + p^n \mathbb{Z}_p) - \mu_{f_m,-q}(a + p^{n+1} \mathbb{Z}_p) \right| \leq C_2 p^{-v_p(1-q^{p^n})}, \quad (2.21)$$

where C_2 is also some positive constant.

By (2.20) and (2.21), we see that

$$\begin{aligned} & |f(a) - \mu_{f,-q}(a + p^n \mathbb{Z}_p)| \\ & \leq \max \{ |f(a) - f_m(a)|, |f_m(a) - \mu_{f_m,-q}(a + p^n \mathbb{Z}_p)|, |\mu_{f-f_m,-q}(a + p^n \mathbb{Z}_p)| \} \\ & \leq \max \{ |f(a) - f_m(a)|, |f_m(a) - \mu_{f_m,-q}(a + p^n \mathbb{Z}_p)|, \|f - f_m\|_{\infty} \}. \end{aligned} \quad (2.22)$$

If we fix $\epsilon > 0$ and fix m such that $\|f - f_m\| \leq \epsilon$, then, for $n \gg 0$, we have

$$|f(a) - \mu_{f,-q}(a + p^n \mathbb{Z}_p)| \leq \epsilon. \quad (2.23)$$

Hence, we have

$$f_{\mu_{f,-q}}(a) = \lim_{n \rightarrow \infty} \mu_{f,-q}(a + p^n \mathbb{Z}_p) = \frac{[2]_q}{2} (-q)^a f(a). \quad (2.24)$$

Let m be the sufficiently large number such that $\|f - f_m\|_\infty \leq p^{-n}$.

Then we get

$$\begin{aligned} \mu_{f,-q}(a + p^n \mathbb{Z}_p) &= \mu_{f_m,-q}(a + p^n \mathbb{Z}_p) + \mu_{f-f_m,-q}(a + p^n \mathbb{Z}_p) \\ &= \mu_{f_m,-q}(a + p^n \mathbb{Z}_p) \\ &= (-1)^a q^a \frac{[2]_q}{2} f(a) \pmod{[p^n]_q}. \end{aligned} \quad (2.25)$$

For all $g \in C(\mathbb{Z}_p)$, we have

$$\int_{\mathbb{Z}_p} g(x) d\mu_{f,-q}(x) = \int_{\mathbb{Z}_p} f(x) g(x) d\mu_{-q}(x). \quad (2.26)$$

Assume that f is the function from $C(\mathbb{Z}_p, C_p)$ to $Lip(\mathbb{Z}_p, C_p)$. By the definition of μ_{-q} , we easily see that μ_{-q} is a strongly p -adic q -measure on $\mathbb{Z}_p$ and for $n \gg 0$

$$|f_{\mu_{-q}}(a) - \mu_{-q}(a + p^n \mathbb{Z}_p)| \leq C_3 p^{-v_p(1-q^{p^n})}, \quad (2.27)$$

where C_3 is some positive constant.

If $\mu_{1,-q}$ is associated strongly fermionic p -adic q -measure on $\mathbb{Z}_p$, then we have

$$|\mu_{1,-q}(a + p^n \mathbb{Z}_p) - f_{\mu_{-q}}(a)| \leq C_4 p^{-v_p(1-q^{p^n})}, \quad (2.28)$$

where $n \gg 0$ and C_4 is some positive constant.

From (2.28), we get

$$|\mu_{-q}(a + p^n \mathbb{Z}_p) - \mu_{1,-q}(a + p^n \mathbb{Z}_p)| \leq |\mu_{-q}(a + p^n \mathbb{Z}_p) - f_{\mu_{-q}}(a)| + |f_{\mu_{-q}}(a) - \mu_{1,-q}(a + p^n \mathbb{Z}_p)| \leq K, \quad (2.29)$$

where K is some positive constant.

Therefore, $\mu_{-q} - \mu_{1,-q}$ is a q -measure on $\mathbb{Z}_p$. Hence, we obtain the following theorem.

Theorem 2.3. Let μ_{-q} be a strongly fermionic p -adic q -measure on $\mathbb{Z}_p$, and assume that the fermionic Radon-Nikodym derivative $f_{\mu_{-q}}$ on $\mathbb{Z}_p$ is continuous function on $\mathbb{Z}_p$. Suppose that $\mu_{1,-q}$ is the strongly fermionic p -adic q -measure associated to $f_{\mu_{-q}}$. Then there exists a q -measure $\mu_{2,-q}$ on $\mathbb{Z}_p$ such that

$$\mu_{-q} = \mu_{1,-q} + \mu_{2,-q}. \quad (2.30)$$

References

- [1] A. Bayad and T. Kim, "Identities involving values of Bernstein, q -Bernoulli, and q -Euler polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 2, pp. 133–143, 2011.
- [2] M. Cenkci, V. Kurt, S. H. Rim, and Y. Simsek, "On (i, q) Bernoulli and Euler numbers," *Applied Mathematics Letters*, vol. 21, no. 7, pp. 706–711, 2008.
- [3] T. Kim, "Lebesgue-Radon-Nikodym theorem with respect to fermionic p -adic invariant measure on $\mathbb{Z}_p$," to appear in *Russian Journal of Mathematical Physics*.
- [4] T. Kim, S. D. Kim, and D.-W. Park, "On uniform differentiability and q -Mahler expansions," *Advanced Studies in Contemporary Mathematics*, vol. 4, no. 1, pp. 35–41, 2001.
- [5] S.-H. Rim, K. H. Park, and E. J. Moon, "On Genocchi numbers and polynomials," *Abstract and Applied Analysis*, vol. 2008, Article ID 898471, 7 pages, 2008.
- [6] S.-H. Rim, J.-H. Jin, E.-J. Moon, and S.-J. Lee, "Some identities on the q -Genocchi polynomials of higher-order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *International Journal of Mathematics and Mathematical Sciences*, vol. 2010, Article ID 860280, 14 pages, 2010.
- [7] S.-H. Rim, J.-H. Jin, E.-J. Moon, and S.-J. Lee, "On multiple interpolation functions of the q -Genocchi polynomials," *Journal of Inequalities and Applications*, vol. 2010, Article ID 351419, 13 pages, 2010.
- [8] Y. Simsek, "Special functions related to Dedekind-type DC-sums and their applications," *Russian Journal of Mathematical Physics*, vol. 17, no. 4, pp. 495–508, 2010.
- [9] T. Kim, "Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [10] T. Kim, "New approach to q -Euler polynomials of higher order," *Russian Journal of Mathematical Physics*, vol. 17, no. 2, pp. 218–225, 2010.
- [11] T. Kim, "Symmetry of power sum polynomials and multivariate fermionic p -adic invariant integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 1, pp. 93–96, 2009.
- [12] T. Kim, "On the multiple q -Genocchi and Euler numbers," *Russian Journal of Mathematical Physics*, vol. 15, no. 4, pp. 481–486, 2008.
- [13] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, New York, NY, USA, 3rd edition, 1987.
- [14] H. L. Royden, *Real Analysis*, Prentice-Hall, Upper Saddle River, NJ, USA, 1998.