Research Article

Integral-Type Operators from Bloch-Type Spaces to Q_K Spaces

Stevo Stević¹ and Ajay K. Sharma²

¹ Mathematical Institute of the Serbian Academy of Sciences, Knez Mihailova 36/III, 11000 Beograd, Serbia

² School of Mathematics, Shri Mata Vaishno Devi University, Kakryal, Katra 182320, India

Correspondence should be addressed to Stevo Stević, sstevic@ptt.rs

Received 16 May 2011; Accepted 29 June 2011

Academic Editor: Narcisa C. Apreutesei

Copyright © 2011 S. Stević and A. K. Sharma. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The boundedness and compactness of the integral-type operator $I_{\varphi, g}^{(n)} f(z) = \int_0^z f^{(n)}(\varphi(\zeta))g(\zeta)d\zeta$, where $n \in \mathbb{N}_0$, φ is a holomorphic self-map of the unit disk $\mathbb{D}$, and g is a holomorphic function on $\mathbb{D}$, from α -Bloch spaces to Q_K spaces are characterized.

1. Introduction

Let $\mathbb{D}$ be the open unit disk in the complex plane, $\partial\mathbb{D}$ be its boundary, $D(w, r)$ be disk centered at w of radius r , and let $H(\mathbb{D})$ be the class of all holomorphic functions on $\mathbb{D}$. Let

$$\eta_a(z) = \frac{a - z}{1 - \bar{a}z}, \quad a \in \mathbb{D}, \quad (1.1)$$

be the involutive Möbius transformation which interchanges points 0 and a . If X is a Banach space, then by B_X we will denote the closed unit ball in X .

The α -Bloch space, $\mathcal{B}^\alpha(\mathbb{D}) = \mathcal{B}^\alpha$, $\alpha > 0$, consists of all $f \in H(\mathbb{D})$ such that

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^\alpha |f'(z)| < \infty. \quad (1.2)$$

The little α -Bloch space $\mathcal{B}_0^\alpha(\mathbb{D}) = \mathcal{B}_0^\alpha$ consists of all functions f holomorphic on $\mathbb{D}$ such that $\lim_{|z| \rightarrow 1} (1 - |z|^2)^\alpha |f'(z)| = 0$. The norm on $\mathcal{B}^\alpha$ is defined by

$$\|f\|_{\mathcal{B}^\alpha} = |f(0)| + \sup_{z \in \mathbb{D}} (1 - |z|^2)^\alpha |f'(z)|. \quad (1.3)$$

With this norm, $\mathcal{B}^\alpha$ is a Banach space, and the little α -Bloch space $\mathcal{B}_0^\alpha$ is a closed subspace of the α -Bloch space. Note that $\mathcal{B}^1 = \mathcal{B}$ is the usual Bloch space.

Given a nonnegative Lebesgue measurable function K on $(0, 1]$ the space Q_K consists of those $f \in H(\mathbb{D})$ for which

$$b_{Q_K}^2(f) = \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 K(1 - |\eta_a(z)|^2) dm(z) < \infty, \quad (1.4)$$

where $dm(z) = (1/\pi) dx dy = (1/\pi) r dr d\theta$ is the normalized area measure on $\mathbb{D}$ [1]. It is known that b_{Q_K} is a seminorm on Q_K which is Möbius invariant, that is,

$$b_{Q_K}(f \circ \eta) = b_{Q_K}(f), \quad \eta \in \text{Aut}(\mathbb{D}), \quad (1.5)$$

where $\text{Aut}(\mathbb{D})$ is the group of all automorphisms of the unit disk $\mathbb{D}$. It is a Banach space with the norm defined by

$$\|f\|_{Q_K} = |f(0)| + b_{Q_K}(f). \quad (1.6)$$

The space $Q_{K,0}$ consists of all $f \in H(\mathbb{D})$ such that

$$\lim_{|a| \rightarrow 1} \int_{\mathbb{D}} |f'(z)|^2 K(1 - |\eta_a(z)|^2) dm(z) = 0. \quad (1.7)$$

It is known that $Q_{K,0}$ is a closed subspace of Q_K . For classical Q spaces, see [2].

It is clear that each Q_K contains all constant functions. If Q_K consists of just constant functions, we say that it is trivial. Q_K is nontrivial if and only if

$$\sup_{t \in (0, 1)} \int_0^1 K(1 - r) \frac{(1 - t)^2}{(1 - tr^2)^3} r dr < \infty. \quad (1.8)$$

Throughout this paper, we assume that condition (1.8) is satisfied, so that the space Q_K is nontrivial. An important tool in the study of Q_K spaces is the auxiliary function λ_K defined by

$$\lambda_K(s) = \sup_{0 < t \leq 1} \frac{K(st)}{K(t)}, \quad 0 < s < \infty, \quad (1.9)$$

where the domain of K is extended to $(0, \infty)$ by setting $K(t) = K(1)$ when $t > 1$. The next two conditions play important role in the study of Q_K spaces.

(a) There is a constant $C > 0$ such that for all $t > 0$

$$K(2t) \leq CK(t). \quad (1.10)$$

(b) The auxiliary function λ_K satisfies the following condition:

$$\int_0^1 \frac{\lambda_K(s)}{s} ds < \infty. \quad (1.11)$$

Let $\Omega(0, \infty)$ denote the class of all nondecreasing continuous functions on $(0, \infty)$ satisfying conditions (1.8), (1.10), and (1.11).

A positive Borel measure μ on $\mathbb{D}$ is called a K -Carleson measure [3] if

$$\sup_I \int_{S(I)} K\left(\frac{1-|z|}{|I|}\right) d\mu(z) < \infty, \quad (1.12)$$

where the supremum is taken over all subarcs $I \subset \partial\mathbb{D}$, $|I|$ is the length of I , and $S(I)$ is the Carleson box defined by

$$S(I) = \left\{ z : 1 - |I| < |z| < 1, \frac{z}{|z|} \in I \right\}. \quad (1.13)$$

A positive Borel measure μ is called a vanishing K -Carleson measure if

$$\lim_{|I| \rightarrow 0} \int_{S(I)} K\left(\frac{1-|z|}{|I|}\right) d\mu(z) = 0. \quad (1.14)$$

We also need the following results of Wulan and Zhu in [3], in which Q_K spaces are characterized in terms of K -Carleson measures.

Theorem 1.1. *Let $K \in \Omega(0, \infty)$. Then a positive Borel measure μ on $\mathbb{D}$ is a K -Carleson measure if and only if*

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K\left(1 - |\eta_a(z)|^2\right) d\mu(z) < \infty. \quad (1.15)$$

Also, μ is a vanishing K -Carleson measure if and only if

$$\lim_{|a| \rightarrow 1} \int_{\mathbb{D}} K\left(1 - |\eta_a(z)|^2\right) d\mu(z) = 0. \quad (1.16)$$

From Theorem 1.1 and the definition of the spaces Q_K and $Q_{K,0}$, we see that when $K \in \Omega(0, \infty)$, then $f \in Q_K$ if and only if the measure $d\mu_f = |f'(z)|^2 dm(z)$ is a K -Carleson measure, while $f \in Q_{K,0}$ if and only if this measure is a vanishing K -Carleson measure.

Let $\varphi \in S(\mathbb{D})$ be the family of all holomorphic self-maps of $\mathbb{D}$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}_0$. We define an integral-type operator as follows:

$$I_{\varphi, g}^{(n)} f(z) = \int_0^z f^{(n)}(\varphi(\zeta)) g(\zeta) d\zeta, \quad z \in \mathbb{D}. \quad (1.17)$$

Operator (1.17) extends several operators which has been introduced and studied recently (see, e.g., [4–9]). For related operators in n -dimensional case, see, for example, [10–19]. For some classical operators see, for example, [20, 21] and the related references therein. For other product-type operators, see [22] and the references therein.

Motivated by [23, 24] (see also [25–29]), we characterize when φ and g induce bounded and/or compact operators in (1.17) from α -Bloch to Q_K spaces.

Throughout this paper, constants are denoted by C ; they are positive and not necessarily the same at each occurrence. The notation $A \asymp B$ means that there is a positive constant C such that $B/C \leq A \leq CB$.

2. Auxiliary Results

Here, we quote several lemmas which will be used in the proofs of the main results in this paper. The following lemma is folklore (see, e.g., [30]).

Lemma 2.1. *For any $f \in H(\mathbb{D})$ and $z \in \mathbb{D}$, the following inequalities hold*

$$|f(z)| \leq C \begin{cases} \|f\|_{\mathcal{B}^\alpha}, & \text{if } 0 < \alpha < 1, \\ \|f\|_{\mathcal{B}^\alpha} \ln \frac{e}{1-|z|^2}, & \text{if } \alpha = 1, \\ \frac{\|f\|_{\mathcal{B}^\alpha}}{(1-|z|^2)^{\alpha-1}}, & \text{if } \alpha > 1, \end{cases} \quad (2.1)$$

$$\begin{aligned} |f^{(n)}(z)| &\leq C \frac{\sup_{w \in D(z, (1-|z|)/2)} (1-|w|^2)^\alpha |f'(w)|}{(1-|z|^2)^{\alpha+n-1}} \\ &\leq C \frac{\|f\|_{\mathcal{B}^\alpha}}{(1-|z|^2)^{\alpha+n-1}}, \quad \text{if } n \in \mathbb{N}. \end{aligned} \quad (2.2)$$

The next lemma is obtained in [31, 32].

Lemma 2.2. *Let $\alpha > 0$. Then there are two functions $f_1, f_2 \in \mathcal{B}^\alpha$ such that*

$$|f_1'(z)| + |f_2'(z)| \geq \frac{C}{(1-|z|^2)^\alpha}, \quad z \in \mathbb{D}. \quad (2.3)$$

Also, if $\alpha \neq 1$, then there are two functions $f_3, f_4 \in \mathcal{B}^\alpha$ and $C > 0$, such that

$$|f_3(z)| + |f_4(z)| \geq \frac{C}{(1 - |z|^2)^{\alpha-1}}, \quad z \in \mathbb{D}. \quad (2.4)$$

The next Schwartz-type lemma [33] is proved in a standard way, so we omit the proof.

Lemma 2.3. *Let $\alpha > 0, K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}_0$. Then $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha$ (or $\mathcal{B}_0^\alpha$) $\rightarrow Q_K$ is compact if and only if for any bounded sequence $(f_m)_{m \in \mathbb{N}}$ in $\mathcal{B}^\alpha$ converging to zero on compacts of $\mathbb{D}$, we have $\lim_{m \rightarrow \infty} \|I_{\varphi,g}^{(n)} f_m\|_{Q_K} = 0$.*

Lemma 2.4. *Let $\alpha > 0$, $K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}_0$. Then $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ (or $Q_{K,0}$) is weakly compact if and only if it is compact.*

Proof. By a known theorem $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ (or $Q_{K,0}$) is weakly compact if and only if $(I_{\varphi,g}^{(n)})^* : Q_K^*$ (or $Q_{K,0}^*$) $\rightarrow (\mathcal{B}_0^\alpha)^*$ is weakly compact. Since $(\mathcal{B}_0^\alpha)^* \cong A^1$ (the Bergman space) and A^1 has the Schur property, it follows that it is equivalent to $(I_{\varphi,g}^{(n)})^* : Q_K^*$ (or $Q_{K,0}^*$) $\rightarrow (\mathcal{B}_0^\alpha)^*$, is compact, which is equivalent to $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ (or $Q_{K,0}$), is compact, as claimed. $\square$

Lemma 2.5. *Let $\alpha > 0$, $K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}_0$. Then $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is compact if and only if $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is bounded.*

Proof. By Lemma 2.4, $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is compact if and only if it is weakly compact, which, by Gantmacher's theorem ([34]), is equivalent to $(I_{\varphi,g}^{(n)})^{**} ((\mathcal{B}_0^\alpha)^{**}) \subseteq Q_{K,0}$. Since $(\mathcal{B}_0^\alpha)^{**} = \mathcal{B}^\alpha$ and by a standard duality argument $(I_{\varphi,g}^{(n)})^{**} = I_{\varphi,g}^{(n)}$ on $\mathcal{B}^\alpha$, this can be written as $I_{\varphi,g}^{(n)}(\mathcal{B}^\alpha) \subseteq Q_{K,0}$, which by the closed graph theorem is equivalent to $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is bounded. $\square$

For $a \in \mathbb{D}$, set

$$\Phi_{\varphi,g,K}(a) = \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z). \quad (2.5)$$

Lemma 2.6. *Let $\alpha > 0, K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}_0$. If $\Phi_{\varphi,g,K}$ is finite at some point $a \in \mathbb{D}$, then it is continuous on $\mathbb{D}$.*

Proof. We follow the lines of Lemma 2.3 in [24]. From the elementary inequality

$$\frac{(1 - |a|)(1 - |a_1|)}{4} \leq \frac{1 - |\eta_a(z)|^2}{1 - |\eta_{a_1}(z)|^2} \leq \frac{4}{(1 - |a|)(1 - |a_1|)}, \quad a, a_1, z \in \mathbb{D}, \quad (2.6)$$

and since K is nondecreasing and satisfies (1.10), we easily get

$$K(1 - |\eta_{a_1}(z)|^2) \leq C^{\lceil \log_2(4/(1-|a|)(1-|a_1|)) \rceil + 1} K(1 - |\eta_a(z)|^2). \quad (2.7)$$

From (2.7) and since $\Phi_{\varphi,g,K}(a)$ is finite, it follows that $\Phi_{\varphi,g,K}$ is finite at each point $a_1 \in \mathbb{D}$. Let $a \in \mathbb{D}$ be fixed, and let $(a_l)_{l \in \mathbb{N}} \subset \mathbb{D}$ be a sequence converging to a .

We have

$$|\Phi_{\varphi,g,K}(a) - \Phi_{\varphi,g,K}(a_l)| \leq \int_{\mathbb{D}} \frac{|g(z)|^2 |K(1 - |\eta_a(z)|^2) - K(1 - |\eta_{a_l}(z)|^2)|}{(1 - |\varphi(z)|^2)^{2(\alpha+n-1)}} dm(z). \quad (2.8)$$

From (2.6), we have that for l such that $1 - |a_l| \geq (1 - |a|)/2$, say $l \geq l_0$, holds

$$1 - |\eta_{a_l}(z)|^2 \leq \frac{8}{(1 - |a|)^2} (1 - |\eta_a(z)|^2), \quad (2.9)$$

and consequently for $l \geq l_0$, it holds

$$|K(1 - |\eta_a(z)|^2) - K(1 - |\eta_{a_l}(z)|^2)| \leq (1 + C^{\lceil \log_2(8/(1-|a|)^2) \rceil + 1}) K(1 - |\eta_a(z)|^2). \quad (2.10)$$

From this and since $\Phi_{\varphi,g,K}$ is finite at a , by the Lebesgue dominated convergence theorem, we get that the integral in (2.8) converges to zero as $l \rightarrow \infty$ which implies that $\Phi_{\varphi,g,K}(a_l) \rightarrow \Phi_{\varphi,g,K}(a)$ as $l \rightarrow \infty$, from which the lemma follows. $\square$

3. Boundedness and Compactness of $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha$ (or $\mathcal{B}_0^\alpha$) $\rightarrow \mathcal{Q}_K$ (or $\mathcal{Q}_{K,0}$)

In this section, we characterize the boundedness and compactness of the operators $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha$ (or $\mathcal{B}_0^\alpha$) $\rightarrow \mathcal{Q}_K$ (or $\mathcal{Q}_{K,0}$). Let

$$d\mu_{\varphi,g,n,\alpha}(z) = |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z). \quad (3.1)$$

Theorem 3.1. *Let $\alpha > 0$, $K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}$, or $n = 0$ and $\alpha > 1$. Then the following statements are equivalent.*

- (a) $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow \mathcal{Q}_K$ is bounded.
- (b) $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow \mathcal{Q}_K$ is bounded.
- (c) $M := \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) < \infty$.
- (d) $d\mu_{\varphi,g,n,\alpha}(z)$ is a K -Carleson measure.

Moreover, if $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow \mathcal{Q}_K$ is bounded, then the next asymptotic relations hold

$$\|I_{\varphi,g}^{(n)}\|_{\mathcal{B}^\alpha \rightarrow \mathcal{Q}_K} \asymp \|I_{\varphi,g}^{(n)}\|_{\mathcal{B}_0^\alpha \rightarrow \mathcal{Q}_K} \asymp M^{1/2}. \quad (3.2)$$

Proof. By Theorem 1.1, it is clear that (c) and (d) are equivalent.

(c) $\Rightarrow$ (a). Let $f \in \mathcal{B}^\alpha$. First note that $I_{\varphi,g}^{(n)}f(0) = 0$ for each $f \in H(\mathbb{B})$ and $n \in \mathbb{N}_0$. From this and by Lemma 2.1, we have

$$\begin{aligned} \|I_{\varphi,g}^{(n)}f\|_{Q_K}^2 &= \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left| (I_{\varphi,g}^{(n)}f)'(z) \right|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ &= \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left| f^{(n)}(\varphi(z)) \right|^2 |g(z)|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ &\leq C \|f\|_{\mathcal{B}^\alpha}^2 \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z), \end{aligned} \quad (3.3)$$

from which the boundedness of $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_K$ follows, and moreover

$$\|I_{\varphi,g}^{(n)}\|_{\mathcal{B}^\alpha \rightarrow Q_K} \leq CM^{1/2}. \quad (3.4)$$

(a) $\Rightarrow$ (b). This implication is obvious.

(b) $\Rightarrow$ (c). By Lemma 2.2, if $n \in \mathbb{N}$, there are two functions $f_1, f_2 \in \mathcal{B}^\alpha$ such that (2.3) holds, and if $n = 0$ and $\alpha > 1$ such that (2.4) holds. Let

$$h_1(z) = f_1(z) - \sum_{k=1}^{n-1} \frac{f_1^{(k)}(0)}{k!} z^k, \quad h_2(z) = f_2(z) - \sum_{k=1}^{n-1} \frac{f_2^{(k)}(0)}{k!} z^k. \quad (3.5)$$

It is known (see [30]) that for each $f \in H(\mathbb{D})$ and $n \in \mathbb{N}$, we have

$$(1 - |z|^2)^{\alpha+n-1} |f^{(n)}(z)| + \sum_{k=1}^{n-1} |f^{(k)}(0)| \asymp (1 - |z|^2)^\alpha |f'(z)|. \quad (3.6)$$

From this, Lemma 2.2, and since $h_1^{(k)}(0) = h_2^{(k)}(0) = 0$, $k = 0, 1, \dots, n-1$, we have that there is a $\delta > 0$ such that

$$C(1 - |z|^2)^{-(\alpha+n-1)} \leq |h_1^{(n)}(z)| + |h_2^{(n)}(z)|, \quad \text{for } |z| > \delta. \quad (3.7)$$

Now note that for any $f \in \mathcal{B}^\alpha$, the functions $f_r(z) = f(rz)$, $r \in (0, 1)$ belong to $\mathcal{B}^\alpha$, and moreover, $\sup_{0 < r < 1} \|f_r\|_{\mathcal{B}^\alpha} \leq \|f\|_{\mathcal{B}^\alpha}$.

Applying (3.7), using an elementary inequality, the boundedness of $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$, and the last inequality, we obtain

$$\begin{aligned}
& \int_{|r\varphi(z)|>\delta} r^{2n} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - (r|\varphi(z)|)^2)^{2(1-\alpha-n)} dm(z) \\
& \leq C \int_{\mathbb{D}} r^{2n} K(1 - |\eta_a(z)|^2) |g(z)|^2 \left(|h_1^{(n)}(r\varphi(z))|^2 + |h_2^{(n)}(r\varphi(z))|^2 \right) dm(z) \\
& = C \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) \left| \left(I_{\varphi,g}^{(n)}(h_1)_r \right)'(z) \right|^2 dm(z) \\
& \quad + C \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) \left| \left(I_{\varphi,g}^{(n)}(h_2)_r \right)'(z) \right|^2 dm(z) \\
& \leq \|I_{\varphi,g}^{(n)}\|_{\mathcal{B}_0^\alpha \rightarrow Q_K}^2 \left(\|h_1\|_{\mathcal{B}^\alpha}^2 + \|h_2\|_{\mathcal{B}^\alpha}^2 \right).
\end{aligned} \tag{3.8}$$

Letting $r \rightarrow 1$ in (3.8) and using the monotone convergence theorem, we get

$$\int_{|\varphi(z)|>\delta} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) \leq C \|I_{\varphi,g}^{(n)}\|_{\mathcal{B}_0^\alpha \rightarrow Q_K}^2. \tag{3.9}$$

On the other hand, for $f_0(z) = z^n/n! \in \mathcal{B}_0^\alpha$, we get $I_{\varphi,g}^{(n)} f_0 \in Q_K$ which implies

$$\sup_{\alpha \in \mathbb{D}} \int_{|\varphi(z)| \leq \delta} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) \leq \frac{\|I_{\varphi,g}^{(n)}\|_{\mathcal{B}_0^\alpha \rightarrow Q_K}^2 \|f_0\|_{\mathcal{B}^\alpha}^2}{(1 - \delta^2)^{2(\alpha+n-1)}}. \tag{3.10}$$

From (3.9) and (3.10), (c) follows. Moreover we get $M^{1/2} \leq C \|I_{\varphi,g}^{(n)}\|_{\mathcal{B}_0^\alpha \rightarrow Q_K}$. From this, (3.4) and since $\|I_{\varphi,g}^{(n)}\|_{\mathcal{B}_0^\alpha \rightarrow Q_K} \leq \|I_{\varphi,g}^{(n)}\|_{\mathcal{B}^\alpha \rightarrow Q_K}$ the asymptotic relations in (3.2) follow, finishing the proof of the theorem. $\square$

Theorem 3.2. Let $\alpha > 0$, $K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}$, or $n = 0$ and $\alpha > 1$. Let $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_K$ be bounded. Then the following statements are equivalent.

- (a) $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_K$ is compact.
- (b) $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is compact.
- (c) $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is weakly compact.
- (d) $\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \infty$, and

$$\lim_{r \rightarrow 1} \sup_{a \in \mathbb{D}} \int_{|\varphi(z)|>r} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) = 0. \tag{3.11}$$

Proof. By Lemma 2.4, we have that (b) is equivalent to (c).

(d) $\Rightarrow$ (a). Let $(f_l)_{l \in \mathbb{N}}$ be a bounded sequence in $\mathcal{B}^\alpha$, say by L , converging to zero uniformly on compacts of $\mathbb{D}$. Then $f_l^{(n)}$ also converges to zero uniformly on compacts of $\mathbb{D}$. From (3.11) we have that for every $\varepsilon > 0$ there is an $r_1 \in (0, 1)$ such that for $r \in (r_1, 1)$

$$\sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) < \varepsilon. \quad (3.12)$$

Therefore, by Lemma 2.1 and (3.12), we have that for $r \in (r_1, 1)$

$$\begin{aligned} \|I_{\varphi, g}^{(n)} f_l\|_{Q_K}^2 &= \left(\int_{|\varphi(z)| \leq r} + \int_{|\varphi(z)| > r} \right) |f_l^{(n)}(\varphi(z))|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) \\ &< \sup_{|\varphi(z)| \leq r} |f_l^{(n)}(\varphi(z))|^2 \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) + CL^2 \varepsilon. \end{aligned} \quad (3.13)$$

Letting $l \rightarrow \infty$ in (3.13), using the first condition in (d) and $\sup_{|w| \leq r} |f_l^{(n)}(w)| \rightarrow 0$ as $l \rightarrow \infty$, it follows that $\lim_{l \rightarrow \infty} \|I_{\varphi, g}^{(n)} f_l\|_{Q_K} = 0$. Thus, by Lemma 2.3, $I_{\varphi, g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_K$ is compact.

(a) $\Rightarrow$ (b). The implication is trivial since $\mathcal{B}_0^\alpha \subset \mathcal{B}^\alpha$.

(b) $\Rightarrow$ (d). By choosing $f(z) = z^n/n! \in \mathcal{B}_0^\alpha$, $n \in \mathbb{N}_0$, we have that the first condition in (d) holds. Let $f_l(z) = z^l/l$, $l \in \mathbb{N}$. It is easy to see that $(f_l)_{l \in \mathbb{N}}$ is a bounded sequence in $\mathcal{B}_0^\alpha$ converging to zero uniformly on compacts of $\mathbb{D}$. Hence, by Lemma 2.3, it follows that $\|I_{\varphi, g}^{(n)}(f_l)\|_{Q_K} \rightarrow 0$ as $l \rightarrow \infty$. Thus, for every $\varepsilon > 0$, there is an $l_0 \in \mathbb{N}$, $l_0 > n$ such that for $l \geq l_0$

$$\left(\prod_{j=1}^{n-1} (l-j) \right)^2 \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |\varphi(z)|^{2(l-n)} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.14)$$

From (3.14) we have that for each $r \in (0, 1)$ and $l \geq l_0$

$$r^{2(l-n)} \left(\prod_{j=1}^{n-1} (l-j) \right)^2 \sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.15)$$

Hence, for $r \in [(\prod_{j=1}^{n-1} (l_0 - j))^{-1/(l_0-n)}, 1)$, we have that

$$\sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.16)$$

Let $f \in B_{\mathcal{B}_0^\alpha}$, and let $f_t(z) = f(tz)$, $0 < t < 1$. Then $\sup_{0 < t < 1} \|f_t\|_{\mathcal{B}^\alpha} \leq \|f\|_{\mathcal{B}^\alpha}$, $f_t \in \mathcal{B}_0^\alpha$, $t \in (0, 1)$, and $f_t \rightarrow f$ uniformly on compact subsets of $\mathbb{D}$ as $t \rightarrow 1$. The compactness of $I_{\varphi, g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ implies

$$\lim_{t \rightarrow 1} \|I_{\varphi, g}^{(n)} f_t - I_{\varphi, g}^{(n)} f\|_{Q_K} = 0. \quad (3.17)$$

Hence, for every $\varepsilon > 0$, there is a $t \in (0, 1)$ such that

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left| f_t^{(n)}(\varphi(z)) - f^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.18)$$

From this and (3.16), we have that for such t and each $r \in [(\prod_{j=1}^{n-1} (l_0 - j))^{-1/(l_0-n)}, 1)$

$$\begin{aligned} & \sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| f^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) \\ & \leq 2 \sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| f_t^{(n)}(\varphi(z)) - f^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) \\ & \quad + 2 \sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| f_t^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) \\ & < 2\varepsilon \left(1 + \|f_t^{(n)}\|_\infty^2 \right). \end{aligned} \quad (3.19)$$

From (3.19) we conclude that for every $f \in B_{\mathcal{B}_0^\alpha}$, there is a $\delta_0 \in (0, 1)$ and $\delta_0 = \delta_0(f, \varepsilon)$ such that for $r \in (\delta_0, 1)$

$$\sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| f^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.20)$$

Since $I_{\varphi, g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is compact, we have that for every $\varepsilon > 0$ there is a finite collection of functions $f_1, f_2, \dots, f_k \in B_{\mathcal{B}_0^\alpha}$ such that, for each $f \in B_{\mathcal{B}_0^\alpha}$, there is a $j \in \{1, \dots, k\}$, such that

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} \left| f^{(n)}(\varphi(z)) - f_j^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.21)$$

On the other hand, from (3.20), it follows that if $\hat{\delta} := \max_{1 \leq j \leq k} \delta_j(f_j, \varepsilon)$, then for $r \in (\hat{\delta}, 1)$ and all $j \in \{1, \dots, k\}$, we have

$$\sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| f_j^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \varepsilon. \quad (3.22)$$

From (3.21) and (3.22), we have that for $r \in (\widehat{\delta}, 1)$ and every $f \in B_{\mathcal{B}_0^\alpha}$

$$\sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| f^{(n)}(\varphi(z)) \right|^2 K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < 4\varepsilon. \quad (3.23)$$

If we apply (3.23) to the delays of the functions in (3.5) which are normalized so that they belong to $B_{\mathcal{B}^\alpha}$ and then use the monotone convergence theorem, we easily get that for $r > \max\{\delta, \widehat{\delta}\}$ where δ is chosen as in (3.7)

$$\sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) < C\varepsilon, \quad (3.24)$$

from which (3.11) follows, as desired. $\square$

Theorem 3.3. Let $\alpha > 0$, $K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$ and $n \in \mathbb{N}$, or $n = 0$ and $\alpha > 1$. Then the next statements are equivalent.

- (a) $I_{\varphi, g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is bounded.
- (b) $I_{\varphi, g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is compact.
- (c) $I_{\varphi, g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is compact.
- (d) $I_{\varphi, g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is weakly compact.
- (e) $\lim_{|a| \rightarrow 1} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) = 0$.
- (f) $d\mu_{\varphi, g, n, \alpha}(z)$ is a vanishing K -Carleson measure.

Proof. By Theorem 1.1, (e) and (f) are equivalent; by Lemma 2.4, (c) is equivalent to (d), while, by Lemma 2.5, (a) is equivalent to (c). Also (b) obviously implies (a).

(a) $\Rightarrow$ (e) Let h_1 and h_2 be as in (3.5). Then from (3.7) and an elementary inequality, we get

$$\begin{aligned} & \int_{|\varphi(z)| > \delta} K(1 - |\eta_a(z)|^2) (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} |g(z)|^2 dm(z) \\ & \leq C \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) \left| \left(I_{\varphi, g}^{(n)} h_1 \right)'(z) \right|^2 dm(z) \\ & \quad + C \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) \left| \left(I_{\varphi, g}^{(n)} h_2 \right)'(z) \right|^2 dm(z). \end{aligned} \quad (3.25)$$

For $f_0(z) = z^n/n! \in \mathcal{B}^\alpha$, we get $I_{\varphi, g}^{(n)} f_0 \in Q_{K,0}$. From this and since $I_{\varphi, g}^{(n)}(h_j) \in Q_{K,0}$, $j = 1, 2$, by letting $|a| \rightarrow 1$, we get that (e) holds.

(e) $\Rightarrow$ (b). We have that for every $\varepsilon > 0$ there is a $\delta \in (0, 1)$ so that for $|a| > \delta$

$$\Phi_{\varphi, g, K}(a) < \varepsilon. \quad (3.26)$$

On the other hand, by Lemma 2.6, $\Phi_{\varphi,g,K}$ is continuous on $|a| \leq \delta$, so uniformly bounded therein, which along with (3.26) gives the boundedness of $\Phi_{\varphi,g,K}$ on $\mathbb{D}$. Hence, by Theorem 3.1, $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_K$ is bounded. By Lemma 2.1, we have

$$\begin{aligned} & \lim_{|a| \rightarrow 1} \sup_{\|f\|_{\mathcal{B}^\alpha} \leq 1} \int_{\mathbb{D}} \left| \left(I_{\varphi,g}^{(n)} f \right)'(z) \right|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ & \leq C \sup_{\|f\|_{\mathcal{B}^\alpha} \leq 1} \|f\|_{\mathcal{B}^\alpha}^2 \lim_{|a| \rightarrow 1} \Phi_{\varphi,g,K}(a) = C \lim_{|a| \rightarrow 1} \Phi_{\varphi,g,K}(a) = 0, \end{aligned} \quad (3.27)$$

so $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is bounded.

Now assume that $(f_l)_{l \in \mathbb{N}}$ is a bounded sequence in $\mathcal{B}^\alpha$, say by L , converging to zero uniformly on compacta of $\mathbb{D}$ as $l \rightarrow \infty$. To show that the operator $I_{\varphi,g}^{(n)} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is compact, it is enough to prove that there is a subsequence $(f_{l_k})_{k \in \mathbb{N}}$ of $(f_l)_{l \in \mathbb{N}}$ such that $I_{\varphi,g}^{(n)} f_{l_k}$ converges in $Q_{K,0}$ as $k \rightarrow \infty$. By Lemma 2.1 and Montel's theorem, it follows that there is a subsequence, which we may denote again by $(f_l)_{l \in \mathbb{N}}$ converging uniformly on compacta of $\mathbb{D}$ to an $f \in \mathcal{B}^\alpha$, such that $\|f\|_{\mathcal{B}^\alpha} \leq L$. Since $I_{\varphi,g}^{(n)}(\mathcal{B}^\alpha) \subseteq Q_{K,0}$, then clearly $I_{\varphi,g}^{(n)} f \in Q_{K,0}$. We show that

$$\lim_{l \rightarrow \infty} \left\| I_{\varphi,g}^{(n)} f_l - I_{\varphi,g}^{(n)} f \right\|_{Q_K} = 0. \quad (3.28)$$

From (3.26), Lemma 2.1, and some simple calculation, we obtain

$$\sup_{\delta < |a| < 1} \int_{\mathbb{D}} \left| \left(I_{\varphi,g}^{(n)} f_l(z) - I_{\varphi,g}^{(n)} f(z) \right)' \right|^2 K(1 - |\eta_a(z)|^2) dm(z) < 4CL^2 \varepsilon. \quad (3.29)$$

For $a \in \mathbb{D}$ and $t \in (0, 1)$, let

$$\Psi_t(a) = \int_{\mathbb{D} \setminus t\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z). \quad (3.30)$$

Lemma 2.6 essentially shows that Ψ_t is continuous on $\mathbb{D}$. Hence, for each $a \in \mathbb{D}$, there is a $t(a) \in (r, 1)$ such that $\Psi_{t(a)}(a) < \varepsilon/2$. Moreover, there is a neighborhood $\mathcal{O}(a)$ of a such that, for every $b \in \mathcal{O}(a)$, $\Psi_{t(a)}(b) < \varepsilon$. From this and since the set $|a| \leq \delta$ is compact, it follows that there is a $t_0 \in (0, 1)$ such that $\Psi_{t_0}(a) < \varepsilon$ when $|a| \leq \delta$. This along with Lemma 2.1 implies that

$$\begin{aligned} & \sup_{|a| \leq \delta} \int_{\mathbb{D} \setminus t_0\mathbb{D}} \left| \left(I_{\varphi,g}^{(n)} f_l(z) - I_{\varphi,g}^{(n)} f(z) \right)' \right|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ & \leq C \|f_l - f\|_{\mathcal{B}^\alpha}^2 \sup_{|a| \leq \delta} \Psi_{t_0}(a) < 4CL^2 \varepsilon. \end{aligned} \quad (3.31)$$

By the Weierstrass theorem $f_l^{(n)} \rightarrow f^{(n)}$ uniformly on compacta as $l \rightarrow \infty$, from which along with (2.2) and since $\varphi(t_0\mathbb{D})$ is compact, for $r = \sup_{w \in \varphi(t_0\mathbb{D})} |w|$, it follows that

$$\begin{aligned} & \sup_{|a| \leq \delta} \int_{t_0\mathbb{D}} \left| \left(I_{\varphi,g}^{(n)} f_l(z) - I_{\varphi,g}^{(n)} f(z) \right)' \right|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ & \leq C \sup_{|z| \leq r} \left| (f_l - f)^{(n)}(z) \right|^2 \sup_{|a| \leq \delta} \Phi_{\varphi,g,K}(a) \rightarrow 0, \quad \text{as } l \rightarrow \infty. \end{aligned} \quad (3.32)$$

From (3.29)–(3.32) and since $I_{\varphi,g}^{(n)} f(0) = 0$ for each $f \in H(\mathbb{D})$, we easily get (3.28), from which (b) follows, finishing the proof of this theorem. $\square$

Theorem 3.4. *Let $\alpha > 0$, $K \in \Omega(0, \infty)$, $\varphi \in S(\mathbb{D})$, $g \in H(\mathbb{D})$, and $n \in \mathbb{N}$, or $n = 0$ and $\alpha > 1$. Then the following statements are equivalent.*

- (a) $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is bounded,
- (b) $\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |g(z)|^2 K(1 - |\eta_a(z)|^2) (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} dm(z) < \infty$, and

$$\lim_{|a| \rightarrow 1} \int_{\mathbb{D}} |g(z)|^2 K(1 - |\eta_a(z)|^2) dm(z) = 0. \quad (3.33)$$

Proof. Suppose (b) holds and $f \in \mathcal{B}_0^\alpha$. Then by Theorem 3.1, $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is bounded. We show $I_{\varphi,g}^{(n)} f \in Q_{K,0}$, for every $f \in \mathcal{B}_0^\alpha$. Since $f \in \mathcal{B}_0^\alpha$, we have that, for every $\varepsilon > 0$, there is an $r \in (0, 1)$ such that (see, e.g., the idea in [35, Lemma 2.4])

$$\left| f^{(n)}(\varphi(z)) \right|^2 (1 - |\varphi(z)|^2)^{2(\alpha+n-1)} < \varepsilon \quad \text{for } |\varphi(z)| > r. \quad (3.34)$$

Thus,

$$\begin{aligned} & \sup_{a \in \mathbb{D}} \int_{|\varphi(z)| > r} \left| \left(I_{\varphi,g}^{(n)} f(z) \right)' \right|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ & < \varepsilon \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) (1 - |\varphi(z)|^2)^{2(1-\alpha-n)} |g(z)|^2 dm(z). \end{aligned} \quad (3.35)$$

We also have

$$\begin{aligned} & \lim_{|a| \rightarrow 1} \int_{|\varphi(z)| \leq r} \left| \left(I_{\varphi,g}^{(n)} f(z) \right)' \right|^2 K(1 - |\eta_a(z)|^2) dm(z) \\ & \leq C \frac{\|f\|_{\mathcal{B}^\alpha}^2}{(1 - r^2)^{2(\alpha+n-1)}} \lim_{|a| \rightarrow 1} \int_{|\varphi(z)| \leq r} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) \\ & \leq C \frac{\|f\|_{\mathcal{B}^\alpha}^2}{(1 - r^2)^{2(\alpha+n-1)}} \lim_{|a| \rightarrow 1} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) = 0. \end{aligned} \quad (3.36)$$

Combining (3.35) and (3.36), we get $I_{\varphi,g}^{(n)} f \in Q_{K,0}$. Hence, $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is bounded.

Conversely, if $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is bounded, then $I_{\varphi,g}^{(n)} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is bounded too. Thus, by Theorem 3.1, we get the first condition in (b). For $f_0(z) = z^n/n! \in \mathcal{B}_0^\alpha$, we get $I_{\varphi,g}^{(n)} f_0 \in Q_{K,0}$, which is equivalent to (3.33), finishing the proof of the theorem. $\square$

If $n = 0$, we simply denote the operator $I_{\varphi,g}^{(0)}$ by $I_{\varphi,g}$.

Theorem 3.5. *Let $\alpha \in (0,1)$, $K \in \Omega(0,\infty)$, $\varphi \in S(\mathbb{D})$, and $g \in H(\mathbb{D})$. Then the following statements are equivalent.*

- (a) $I_{\varphi,g} : \mathcal{B}^\alpha \rightarrow Q_K$ is bounded.
- (b) $I_{\varphi,g} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is bounded.
- (c) $M_1 := \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) < \infty$.
- (d) $d\mu_1(z) = |g(z)|^2 dm(z)$ is a K -Carleson measure.
- (e) $I_{\varphi,g} : \mathcal{B}^\alpha \rightarrow Q_K$ is compact.
- (f) $I_{\varphi,g} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is compact.
- (g) $I_{\varphi,g} : \mathcal{B}_0^\alpha \rightarrow Q_K$ is weakly compact.

Moreover, if $I_{\varphi,g} : \mathcal{B}^\alpha \rightarrow Q_K$ is bounded, then the next asymptotic relations hold

$$\|I_{\varphi,g}\|_{\mathcal{B}^\alpha \rightarrow Q_K} \asymp \|I_{\varphi,g}\|_{\mathcal{B}_0^\alpha \rightarrow Q_K} \asymp M_1^{1/2}. \quad (3.37)$$

Proof. The proof of the equivalence of statements (a)–(d) of this theorem is similar to the proof of Theorem 3.1; moreover, the implication (b) $\Rightarrow$ (c) is much simpler since it follows by using the test function $f_0(z) \equiv 1$. That (c) is equivalent to (e)–(g) is proved similarly as in Theorem 3.2, by using the well-known fact that if a bounded sequence $(f_l)_{l \in \mathbb{N}}$ in $\mathcal{B}^\alpha$, $\alpha \in (0,1)$ converges to zero uniformly on compacts of $\mathbb{D}$, then it converges to zero uniformly on the whole $\mathbb{D}$. The details are omitted. $\square$

The proof of the next theorem is similar to the proofs of Theorems 3.3 and 3.4 and will be omitted.

Theorem 3.6. *Let $\alpha \in (0,1)$, $K \in \Omega(0,\infty)$, $\varphi \in S(\mathbb{D})$, and $g \in H(\mathbb{D})$. Then the following statements are equivalent.*

- (a) $I_{\varphi,g} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is bounded.
- (b) $I_{\varphi,g} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is bounded.
- (c) $I_{\varphi,g} : \mathcal{B}^\alpha \rightarrow Q_{K,0}$ is compact.
- (d) $I_{\varphi,g} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is compact.
- (e) $I_{\varphi,g} : \mathcal{B}_0^\alpha \rightarrow Q_{K,0}$ is weakly compact.
- (f) $\lim_{|a| \rightarrow 1} \int_{\mathbb{D}} K(1 - |\eta_a(z)|^2) |g(z)|^2 dm(z) = 0$.
- (g) $d\mu_1(z) = |g(z)|^2 dm(z)$ is a vanishing K -Carleson measure.

Acknowledgment

This work is partially supported by the National Board of Higher Mathematics (NBHM)/DAE, India (Grant no. 48/4/2009/R&D-II/426) and by the Serbian Ministry of Science (Projects III41025 and III44006).

References

- [1] H. Wulan and K. Zhu, " Q_K spaces via higher order derivatives," *The Rocky Mountain Journal of Mathematics*, vol. 38, no. 1, pp. 329–350, 2008.
- [2] J. Xiao, *Holomorphic Q Classes*, vol. 1767 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 2001.
- [3] H. Wulan and K. Zhu, "Derivative-free characterizations of Q_K spaces," *Journal of the Australian Mathematical Society*, vol. 82, no. 2, pp. 283–295, 2007.
- [4] S. Li and S. Stević, "Generalized composition operators on Zygmund spaces and Bloch type spaces," *Journal of Mathematical Analysis and Applications*, vol. 338, no. 2, pp. 1282–1295, 2008.
- [5] S. Li and S. Stević, "Products of composition and integral type operators from H^∞ to the Bloch space," *Complex Variables and Elliptic Equations*, vol. 53, no. 5, pp. 463–474, 2008.
- [6] S. Li and S. Stević, "Products of Volterra type operator and composition operator from H^∞ and Bloch spaces to Zygmund spaces," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 1, pp. 40–52, 2008.
- [7] S. Li and S. Stević, "Products of integral-type operators and composition operators between Bloch-type spaces," *Journal of Mathematical Analysis and Applications*, vol. 349, no. 2, pp. 596–610, 2009.
- [8] A. Sharma and A. K. Sharma, "Carleson measures and a class of generalized integration operators on the Bergman space," *Rocky Mountain Journal of Mathematics*, vol. 41, no. 5, pp. 1711–1724, 2011.
- [9] S. Stević, "Generalized composition operators from logarithmic Bloch spaces to mixed-norm spaces," *Utilitas Mathematica*, vol. 77, pp. 167–172, 2008.
- [10] S. Stević, "On a new integral-type operator from the Bloch space to Bloch-type spaces on the unit ball," *Journal of Mathematical Analysis and Applications*, vol. 354, no. 2, pp. 426–434, 2009.
- [11] S. Stević, "On an integral-type operator from logarithmic Bloch-type and mixed-norm spaces to Bloch-type spaces," *Nonlinear Analysis*, vol. 71, no. 12, pp. 6323–6342, 2009.
- [12] S. Stević, "Products of integral-type operators and composition operators from the mixed norm space to Bloch-type spaces," *Siberian Mathematical Journal*, vol. 50, no. 4, pp. 726–736, 2009.
- [13] S. Stević, "Norm and essential norm of an integral-type operator from the Dirichlet space to the Bloch-type space on the unit ball," *Abstract and Applied Analysis*, vol. 2010, Article ID 134969, 9 pages, 2010.
- [14] S. Stević, "On an integral operator between Bloch-type spaces on the unit ball," *Bulletin des Sciences Mathématiques*, vol. 134, no. 4, pp. 329–339, 2010.
- [15] S. Stević, "On an integral-type operator from Zygmund-type spaces to mixed-norm spaces on the unit ball," *Abstract and Applied Analysis*, vol. 2010, Article ID 198608, 7 pages, 2010.
- [16] S. Stević, "On operator P_ϕ^S from the logarithmic Bloch-type space to the mixed-norm space on the unit ball," *Applied Mathematics and Computation*, vol. 215, no. 12, pp. 4248–4255, 2010.
- [17] S. Stević and S. I. Ueki, "On an integral-type operator between weighted-type spaces and Bloch-type spaces on the unit ball," *Applied Mathematics and Computation*, vol. 217, no. 7, pp. 3127–3136, 2010.
- [18] W. Yang and X. Meng, "Generalized composition operators from $F(p, q, s)$ spaces to Bloch-type spaces," *Applied Mathematics and Computation*, vol. 217, no. 6, pp. 2513–2519, 2010.
- [19] X. Zhu, "Integral-type operators from iterated logarithmic Bloch spaces to Zygmund-type spaces," *Applied Mathematics and Computation*, vol. 215, no. 3, pp. 1170–1175, 2009.
- [20] A. G. Siskakis and R. Zhao, "A Volterra type operator on spaces of analytic functions," *Contemporary Mathematics*, vol. 232, pp. 299–311, 1999.
- [21] R. Yoneda, "Pointwise multipliers from $BMOA^\alpha$ to $BMOA^\beta$," *Complex Variables*, vol. 49, no. 14, pp. 1045–1061, 2004.
- [22] S. Stević, A. K. Sharma, and A. Bhat, "Products of multiplication composition and differentiation operators on weighted Bergman spaces," *Applied Mathematics and Computation*, vol. 217, pp. 8115–8125, 2011.
- [23] S. Li and H. Wulan, "Composition operators on Q_K spaces," *Journal of Mathematical Analysis and Applications*, vol. 327, no. 2, pp. 948–958, 2007.

- [24] W. Smith and R. Zhao, "Composition operators mapping into the Q_p spaces," *Analysis*, vol. 17, no. 2-3, pp. 239–263, 1997.
- [25] L. Jiang and Y. He, "Composition operators from β^a to $F(p, q, s)$," *Acta Mathematica Scientia—Series B*, vol. 23, no. 2, pp. 252–260, 2003.
- [26] S. Li, "Composition operators on Q_p spaces," *Georgian Mathematical Journal*, vol. 12, no. 3, pp. 505–514, 2005.
- [27] Z. Lou, "Composition operators on Q^p spaces," *Journal of the Australian Mathematical Society*, vol. 70, no. 2, pp. 161–188, 2001.
- [28] P. Wu and H. Wulan, "Composition operators from the Bloch space into the spaces Q_K ," *International Journal of Mathematics and Mathematical Science*, vol. 31, pp. 1973–1979, 2003.
- [29] H. Wulan, "Compactness of composition operators from the Bloch space β to Q_K spaces," *Acta Mathematica Sinica*, vol. 21, no. 6, pp. 1415–1424, 2005.
- [30] K. Zhu, "Bloch type spaces of analytic functions," *The Rocky Mountain Journal of Mathematics*, vol. 23, no. 3, pp. 1143–1177, 1993.
- [31] Z. Lou, *Bloch type spaces of analytic functions*, Ph.D. thesis, 1997.
- [32] J. Xiao, "Riemann-Stieltjes operators on weighted Bloch and Bergman spaces of the unit ball," *Journal of the London Mathematical Society*, vol. 70, no. 1, pp. 199–214, 2004.
- [33] H. J. Schwartz, *Composition operators on H^p* , M.S. thesis, University of Toledo, 1969.
- [34] N. Dunford and J. Schwartz, *Linear Operators*, vol. 1, Interscience, New York, NY, USA, 1958.
- [35] S. Stević and S. I. Ueki, "Integral-type operators acting between weighted-type spaces on the unit ball," *Applied Mathematics and Computation*, vol. 215, no. 7, pp. 2464–2471, 2009.