Research Article

The Existence of Positive Solutions for Fractional Differential Equations with Sign Changing Nonlinearities

Weihua Jiang, Jiqing Qiu, and Weiwei Guo

College of Sciences, Hebei University of Science and Technology, Hebei, Shijiazhuang 050018, China

Correspondence should be addressed to Jiqing Qiu, qiujiqing@263.net

Received 23 March 2012; Revised 11 June 2012; Accepted 12 June 2012

Academic Editor: Bashir Ahmad

Copyright © 2012 Weihua Jiang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the existence of at least two positive solutions to eigenvalue problems of fractional differential equations with sign changing nonlinearities in more generalized boundary conditions. Our analysis relies on the Avery-Peterson fixed point theorem in a cone. Some examples are given for the illustration of main results.

1. Introduction

The theory of fractional differential equations has become an important aspect of differential equations (see [1–8]). Boundary value problems of fractional differential equations have been investigated in many papers (see [9–46]). The existence of positive solutions to boundary value problems of fractional differential equations has been studied by many authors when nonlinearities are positive (see [9–24]). There are a few papers to study the existence of positive solutions of semipositone fractional differential equations. For example, using the Krasnoselskii fixed point theorem, Yuan et al. [9] discussed the existence of positive solutions for the singular positone and semipositone two-point boundary value problems

$$\begin{aligned} D_{0^+}^\alpha u(t) &= \mu a(t) f(t, u(t)), \\ u(0) = u'(0) &= u(1) = u'(1) = 0, \quad 3 < \alpha \leq 4, \end{aligned} \tag{1.1}$$

where $\mu > 0$, a and f are continuous. In [10], Wang et al. studied the existence of positive solutions for the singular semipositone two-point boundary value problems

$$D_{0+}^{\alpha} u(t) + \lambda f(t, u(t)) = 0, \quad (1.2)$$

$$u(0) = u'(0) = u(1) = 0, \quad 2 < \alpha \leq 3, \quad (1.3)$$

where $\lambda > 0$, f is continuous.

In [11], using Krasnoselskii fixed point theorem, Goodrich discussed the existence of at least one positive solutions for the system of fractional boundary value problems

$$\begin{aligned} -D_{0+}^{\nu_1} y_1(t) &= \lambda_1 a_1(t) f(y_1(t), y_2(t)), & -D_{0+}^{\nu_2} y_2(t) &= \lambda_2 a_2(t) g(y_1(t), y_2(t)), \\ y_1^{(i)}(0) &= y_2^{(i)}(0) = 0, \quad 0 \leq i \leq n-2, & D_{0+}^{\alpha} y_1(t)|_{t=1} &= \phi_1(y), \quad D_{0+}^{\alpha} y_2(t)|_{t=1} = \phi_2(y), \end{aligned} \quad (1.4)$$

where a_1, a_2, f , and g are nonnegative for $t \in [0, 1]$.

Motivated by the excellent results mentioned above, in this paper, we investigate the existence of at least two positive solutions for the problem

$$\begin{aligned} D_{0+}^{\alpha} y(t) + \lambda f(t, y(t)) &= 0, \quad t \in [0, 1], \\ y^{(i)}(0) &= 0, \quad 0 \leq i \leq n-2, \\ D_{0+}^{\beta} y(t)|_{t=1} &= h(y), \end{aligned} \quad (1.5)$$

where $\lambda > 0$, $\alpha \in (n-1, n]$, $n \geq 3$, $1 \leq \beta \leq n-2 < \alpha-1$, $f \in C([0, 1] \times \mathbb{R}^+, [-M, \infty))$, $M > 0$, $h \in C(C(\mathbb{R}^+), \mathbb{R}^+)$, $\mathbb{R}^+ = [0, \infty)$. The main tool is the Avery-Peterson theorem. To the best of our knowledge, this is the first paper dealing with eigenvalue problems of fractional differential equations with sign changing nonlinearities involving more general boundary conditions. Our results improve some of the earlier work presented in [10, 17, 46].

2. Preliminaries

For the convenience of the readers, we present here some necessary definitions and lemmas from fractional calculus theory. For more details see [1, 2].

Definition 2.1. The Riemann-Liouville fractional integral of order $\alpha > 0$ of a function $y : (0, \infty) \rightarrow \mathbb{R}$ is given by

$$I_{0+}^{\alpha} y(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds, \quad (2.1)$$

provided that the right-hand side is pointwise defined on $(0, \infty)$.

Definition 2.2. The Riemann-Liouville fractional derivative of order $\alpha > 0$ of a function $y : (0, \infty) \rightarrow \mathbb{R}$ is given by

$$D_{0+}^{\alpha} y(t) = \frac{1}{\Gamma(n-\alpha)} \frac{d^n}{dt^n} \int_0^t (t-s)^{n-\alpha-1} y(s) ds, \quad (2.2)$$

provided that the right-hand side is pointwise defined on $(0, \infty)$, where $n = [\alpha] + 1$.

Lemma 2.3. Assume $f \in C[0, 1]$, $q \geq p \geq 0$, then

$$D_{0+}^p I_{0+}^q f(t) = I_{0+}^{q-p} f(t). \quad (2.3)$$

Lemma 2.4. Assume $\alpha > 0$, then

(1) If $\lambda > -1$, $\lambda \neq \alpha - i$, $i = 1, 2, \dots, [\alpha] + 1$, $t > 0$, then

$$D_{0+}^{\alpha} t^{\lambda} = \frac{\Gamma(\lambda + 1)}{\Gamma(\lambda - \alpha + 1)} t^{\lambda - \alpha}. \quad (2.4)$$

(2) $D_{0+}^{\alpha} t^{\alpha-i} = 0$, $i = 1, 2, \dots, n$.

(3) $D_{0+}^{\alpha} I_{0+}^{\alpha} u(t) = u(t)$, for a.e. $t \in [0, 1]$, where $u \in L^1[0, 1]$.

(4) $D_{0+}^{\alpha} u(t) = 0$ if and only if

$$u(t) = c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \dots + c_n t^{\alpha-n}, \quad (2.5)$$

for some $c_i \in \mathbb{R}$, $i = 1, 2, \dots, n$, where n is the least integer greater than or equal to α .

Lemma 2.5 (see [11]). Given $g \in C[0, 1]$, y is a solution of the problem

$$\begin{aligned} D_{0+}^{\alpha} y(t) + g(t) &= 0, \quad t \in [0, 1], \\ y^{(i)}(0) &= 0, \quad 0 \leq i \leq n-2, \\ D_{0+}^{\beta} y(t)|_{t=1} &= h(y), \end{aligned} \quad (2.6)$$

if and only if it satisfies

$$y(t) = \frac{\Gamma(\alpha - \beta)}{\Gamma(\alpha)} t^{\alpha-1} h(y) + \int_0^1 G(t, s) g(s) ds, \quad (2.7)$$

where

$$G(t, s) = \begin{cases} \frac{t^{\alpha-1}(1-s)^{\alpha-\beta-1} - (t-s)^{\alpha-1}}{\Gamma(\alpha)}, & 0 \leq s \leq t \leq 1, \\ \frac{t^{\alpha-1}(1-s)^{\alpha-\beta-1}}{\Gamma(\alpha)}, & 0 \leq t \leq s \leq 1. \end{cases} \quad (2.8)$$

Lemma 2.6 (see[11]). $G(t, s)$ is continuous on $[0, 1] \times [0, 1]$ and

$$0 \leq G(t, s) \leq G(1, s), \quad t, s \in [0, 1]. \quad (2.9)$$

Lemma 2.7. $G(t, s) \geq t^{\alpha-1}G(1, s)$, $t, s \in [0, 1]$.

Proof. For $s \leq t$,

$$\begin{aligned} G(t, s) &= \frac{t^{\alpha-1}(1-s)^{\alpha-\beta-1} - (t-s)^{\alpha-1}}{\Gamma(\alpha)} \\ &= t^{\alpha-1} \frac{(1-s)^{\alpha-\beta-1} - (1-s/t)^{\alpha-1}}{\Gamma(\alpha)} \\ &\geq t^{\alpha-1} \frac{(1-s)^{\alpha-\beta-1} - (1-s)^{\alpha-1}}{\Gamma(\alpha)} = t^{\alpha-1}G(1, s). \end{aligned} \quad (2.10)$$

For $s > t$,

$$G(t, s) = \frac{t^{\alpha-1}(1-s)^{\alpha-\beta-1}}{\Gamma(\alpha)} \geq t^{\alpha-1} \frac{(1-s)^{\alpha-\beta-1} - (t-s)^{\alpha-1}}{\Gamma(\alpha)} = t^{\alpha-1}G(1, s). \quad (2.11)$$

□

By simple calculation, we can get

$$\int_0^1 G(1, s)ds = \frac{\beta}{(\alpha - \beta)\Gamma(\alpha + 1)}, \quad \int_{1/2}^1 G(1, s)ds = \frac{2^\beta \alpha - \alpha + \beta}{2^\alpha (\alpha - \beta)\Gamma(\alpha + 1)}. \quad (2.12)$$

By Lemma 2.5, we can easily get the following lemma.

Lemma 2.8. *The boundary value problem*

$$\begin{aligned} D_{0+}^\alpha u(t) + 1 &= 0, \quad 0 < t < 1, \\ u^{(i)}(0) &= D_{0+}^\beta u(t)|_{t=1} = 0, \quad 0 \leq i \leq n-2, \end{aligned} \quad (2.13)$$

has a unique solution

$$u(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)} \left(\frac{1}{\alpha-\beta} - \frac{t}{\alpha} \right). \quad (2.14)$$

Obviously, u satisfies

$$\frac{\beta t^{\alpha-1}}{(\alpha-\beta)\Gamma(\alpha+1)} \leq u(t) \leq \frac{t^{\alpha-1}}{(\alpha-\beta)\Gamma(\alpha)}, \quad t \in [0, 1]. \quad (2.15)$$

Lemma 2.9. $\tilde{y} \geq \lambda Mu$ is a solution of the following problem:

$$\begin{aligned} D_{0+}^{\alpha} \tilde{y} + \lambda [f(t, \tilde{y} - \lambda Mu) + M] &= 0, \quad t \in [0, 1], \\ \tilde{y}^{(i)}(0) &= 0, \quad 0 \leq i \leq n-2, \\ D_{0+}^{\beta} \tilde{y}(t)|_{t=1} &= h(\tilde{y} - \lambda Mu), \end{aligned} \quad (2.16)$$

if and only if $y = \tilde{y} - \lambda Mu$ is a positive solution of (1.5).

Proof. In fact, if y is a positive solution of the problem (1.5), by Lemma 2.8, we get that y satisfies

$$\begin{aligned} D_{0+}^{\alpha} (y + \lambda Mu) + \lambda [f(t, y) + M] &= 0, \quad t \in [0, 1], \\ (y + \lambda Mu)^{(i)}(0) &= 0, \quad 0 \leq i \leq n-2, \\ D_{0+}^{\beta} (y + \lambda Mu)|_{t=1} &= h(y). \end{aligned} \quad (2.17)$$

Take $\tilde{y} = y + \lambda Mu$. Then $\tilde{y}$ satisfies (2.16) and $\tilde{y} \geq \lambda Mu$.

On the other hand, if $\tilde{y}$ is a solution of (2.16) and $\tilde{y} \geq \lambda Mu$. Take $y = \tilde{y} - \lambda Mu$. By Lemma 2.8, we can easily get that y satisfies (1.5). Clearly, $y \geq 0$. $\square$

Define functions $\tilde{h}, \tilde{f}$ and an operator $T : C[0, 1] \rightarrow C[0, 1]$ by

$$\begin{aligned} \tilde{h}(y) &= h(\max\{y - \lambda Mu, 0\}), \quad \tilde{f}(t, y) = f(t, \max\{y - \lambda Mu, 0\}) + M, \\ Ty(t) &= \frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha)} t^{\alpha-1} \tilde{h}(y) + \lambda \int_0^1 G(t, s) \tilde{f}(s, y(s)) ds. \end{aligned} \quad (2.18)$$

Obviously, $y \geq \lambda Mu$ is a fixed point of the operator T if and only if $y - \lambda Mu$ is a positive solution of the problem (1.5).

Take $X = C[0, 1]$ with norm $\|x\| = \max_{t \in [0, 1]} |x(t)|$. Define a cone P by

$$P = \left\{ y \in C[0, 1] \mid y(t) \geq t^{\alpha-1} \|y\|, \quad t \in [0, 1] \right\}. \quad (2.19)$$

Lemma 2.10. $T : P \rightarrow P$ is a completely continuous operator.

Proof. Take $y \in P$. By Lemmas 2.6 and 2.7, we get

$$Ty(t) \geq t^{\alpha-1} \left[\frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha)} \tilde{h}(y) + \lambda \int_0^1 G(1,s) \tilde{f}(s, y(s)) ds \right] \geq t^{\alpha-1} \|Ty\|. \quad (2.20)$$

So, $T : P \rightarrow P$. Let $\Omega \subset P$ be bounded. It follows from the continuity of h, f that there exist constants M_1 and M_2 such that $\tilde{h}(y) \leq M_1$ and $\tilde{f}(t, y) \leq M_2$ for $t \in [0, 1]$, $y \in \Omega$. Thus,

$$\|Ty\| \leq \frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha)} M_1 + \lambda M_2 \int_0^1 G(1,s) ds. \quad (2.21)$$

That is $T(\Omega)$ is bounded. For $y \in \Omega$, $t_1, t_2 \in [0, 1]$,

$$|Ty(t_1) - Ty(t_2)| \leq M_1 \frac{\Gamma(\alpha-\beta)}{\Gamma(\alpha)} |t_1^{\alpha-1} - t_2^{\alpha-1}| + \lambda M_2 \int_0^1 |G(t_1, s) - G(t_2, s)| ds. \quad (2.22)$$

By the uniform continuity of $t^{\alpha-1}$ and $G(t, s)$, we get that $T(\Omega)$ is equicontinuous. Obviously, $T : P \rightarrow P$ is continuous. By the Arzela-Ascoli theorem, we get that $T : P \rightarrow P$ is completely continuous. $\square$

Definition 2.11. A map ϕ is said to be a nonnegative, continuous, and concave functional on a cone P of a real Banach space E if and only if $\phi : P \rightarrow \mathbb{R}^+$ is continuous and

$$\phi(tx + (1-t)y) \geq t\phi(x) + (1-t)\phi(y), \quad (2.23)$$

for all $x, y \in P$ and $t \in [0, 1]$.

Definition 2.12. A map Φ is said to be a nonnegative, continuous, and convex functional on a cone P of a real Banach space E iff $\Phi : P \rightarrow \mathbb{R}^+$ is continuous and

$$\Phi(tx + (1-t)y) \leq t\Phi(x) + (1-t)\Phi(y), \quad (2.24)$$

for all $x, y \in P$ and $t \in [0, 1]$.

Let φ and Θ be nonnegative, continuous, and convex functional on P , Φ a nonnegative, continuous, and concave functional on P , and Ψ a nonnegative continuous functional on P . Then, for positive numbers a, b, c , and d , we define the following sets:

$$\begin{aligned} P(\varphi, d) &= \{x \in P : \varphi(x) < d\}, \\ P(\varphi, \Phi, b, d) &= \{x \in P : b \leq \Phi(x), \varphi(x) \leq d\}, \\ P(\varphi, \Theta, \Phi, b, c, d) &= \{x \in P : b \leq \Phi(x), \Theta(x) \leq c, \varphi(x) \leq d\}, \\ R(\varphi, \Psi, a, d) &= \{x \in P : a \leq \Psi(x), \varphi(x) \leq d\}. \end{aligned} \quad (2.25)$$

We will use the following fixed point theorem of Avery and Peterson to study the problem (1.5).

Theorem 2.13 (see [47]). *Let P be a cone in a real Banach space E . Let φ and Θ be nonnegative, continuous, and convex functionals on P , Φ a nonnegative, continuous, and concave functional on P , and Ψ a nonnegative continuous functional on P satisfying $\Psi(kx) \leq k\Psi(x)$ for $0 \leq k \leq 1$, such that for some positive numbers M and d ,*

$$\Phi(x) \leq \Psi(x), \quad \|x\| \leq M\varphi(x) \quad (2.26)$$

for all $x \in \overline{P(\varphi, d)}$. Suppose that

$$T : \overline{P(\varphi, d)} \longrightarrow \overline{P(\varphi, d)} \quad (2.27)$$

is completely continuous and there exist positive numbers a, b, c with $a < b$, such that the following conditions are satisfied:

- (S1) $\{x \in P(\varphi, \Theta, \Phi, b, c, d) : \Phi(x) > b\} \neq \emptyset$ and $\Phi(Tx) > b$ for $x \in P(\varphi, \Theta, \Phi, b, c, d)$;
- (S2) $\Phi(Tx) > b$ for $x \in P(\varphi, \Phi, b, d)$ with $\Theta(Tx) > c$;
- (S3) $0 \notin R(\varphi, \Psi, a, d)$ and $\Psi(Tx) < a$ for $x \in R(\varphi, \Psi, a, d)$ with $\Psi(x) = a$.

Then T has at least three fixed points $x_1, x_2, x_3 \in \overline{P(\varphi, d)}$, such that

$$\begin{aligned} \varphi(x_i) &\leq d, \quad \text{for } i = 1, 2, 3, \\ b &< \Phi(x_1), \quad a < \Psi(x_2), \quad \Phi(x_2) < b, \quad \Psi(x_3) < a. \end{aligned} \quad (2.28)$$

3. Main Results

We define a concave function $\Phi(x) = \min_{t \in [1/2, 1]} |x(t)|$ and convex functions $\Psi(x) = \Theta(x) = \varphi(x) = \|x\|$.

Theorem 3.1. *Assume that there exists a constant $0 < l < \Gamma(\alpha)/\Gamma(\alpha - \beta)$, such that $h(y) \leq l\|y\|$ for $y \in P$. In addition, suppose that there exist constants k, a, b, c, d with $k > 2^{2\alpha-1}\beta\Gamma(\alpha)/(\Gamma(\alpha) - \Gamma(\alpha - \beta)l)$, $[\Gamma(\alpha) - \Gamma(\alpha - \beta)l]\alpha M/\beta\Gamma(\alpha) < a < b - [\Gamma(\alpha) - \Gamma(\alpha - \beta)l]\alpha M/\beta\Gamma(\alpha) < 2^{\alpha-1}b < c < d$, such that the following conditions hold:*

- (C1) $f(t, y) \leq d - M$, for $(t, y) \in [0, 1] \times [0, d]$;
- (C2) $f(t, y) \geq kb - M$, for $(t, y) \in [1/2, 1] \times [b - [\Gamma(\alpha) - \Gamma(\alpha - \beta)l]\alpha M/\beta\Gamma(\alpha), c]$;
- (C3) $f(t, y) \leq a - M$, for $(t, y) \in [0, 1] \times [0, a]$.

Then the problem (1.5) has at least two positive solutions for

$$\frac{2^{2\alpha-1}(\alpha - \beta)\Gamma(\alpha + 1)}{k[\alpha(2^\beta - 1) + \beta]} < \lambda < \frac{\alpha(\alpha - \beta)[\Gamma(\alpha) - \Gamma(\alpha - \beta)l]}{\beta}. \quad (3.1)$$

Proof. Take $y \in \overline{P(\varphi, d)}$. By $\|\max\{y - \lambda Mu, 0\}\| \leq d$, (C1), Lemma 2.6, (2.12), and (3.1), we have

$$\begin{aligned} \|Ty\| &\leq \frac{\Gamma(\alpha - \beta)}{\Gamma(\alpha)} ld + \lambda d \int_0^1 G(1, s) ds \\ &= \left[\frac{\Gamma(\alpha - \beta)}{\Gamma(\alpha)} l + \lambda \int_0^1 G(1, s) ds \right] d \leq d. \end{aligned} \quad (3.2)$$

This means that $T : \overline{P(\varphi, d)} \rightarrow \overline{P(\varphi, d)}$.

It is easy to see that $\{y \in P(\varphi, \Theta, \Phi, b, c, d) : \Phi(y) > b\} \neq \emptyset$. $y \in P(\varphi, \Theta, \Phi, b, c, d)$ implies $\min_{t \in [1/2, 1]} y(t) \geq b$, $\|y\| \leq c$. It follows from (2.15) and (3.1) that $\min_{t \in [1/2, 1]} (y - \lambda Mu) \geq b - \alpha[\Gamma(\alpha) - \Gamma(\alpha - \beta)l]M/\beta\Gamma(\alpha)$. By (C2), (2.12), (3.1), and Lemma 2.7, we get

$$\Phi(Ty) = \min_{t \in [1/2, 1]} Ty(t) \geq \lambda \min_{t \in [1/2, 1]} \int_0^1 G(t, s) \tilde{f}(s, y(s)) ds \geq \left(\frac{1}{2}\right)^{\alpha-1} \lambda kb \int_{1/2}^1 G(1, s) ds > b. \quad (3.3)$$

So, the condition (S1) of Theorem 2.13 holds.

Take $y \in P(\varphi, \Phi, b, d)$ with $\Theta(Ty) > c$. By $Ty \in P$, we get

$$\min_{t \in [1/2, 1]} Ty(t) \geq \min_{t \in [1/2, 1]} t^{\alpha-1} \|Ty\| \geq \frac{1}{2^{\alpha-1}} \|Ty\| > \frac{1}{2^{\alpha-1}} c > b. \quad (3.4)$$

Thus, (S2) holds.

By $a > 0$, we have $0 \notin R(\varphi, \Psi, a, d)$. Take $y \in R(\varphi, \Psi, a, d)$ with $\Psi(y) = a$. By (C3), we get

$$\begin{aligned} \Psi(Ty) = \|Ty\| &\leq \frac{\Gamma(\alpha - \beta)}{\Gamma(\alpha)} la + \lambda a \int_0^1 G(1, s) ds \\ &= \left[\frac{\Gamma(\alpha - \beta)}{\Gamma(\alpha)} l + \lambda \int_0^1 G(1, s) ds \right] a \leq a. \end{aligned} \quad (3.5)$$

By Theorem 2.13, we get that T has at least three fixed points $y_1, y_2, y_3 \in \overline{P(\varphi, d)}$ such that $\|y_i\| \leq d$, $i = 1, 2, 3$, and

$$b < \Phi(y_1), \quad a < \Psi(y_2), \quad \Phi(y_2) < b, \quad \Psi(y_3) < a. \quad (3.6)$$

If $y \in P$ and $\|y\| > a$, by (2.15) and (3.1), we have

$$y(t) \geq t^{\alpha-1} \|y\| > at^{\alpha-1} > \lambda Mu(t). \quad (3.7)$$

Obviously, $\|y_1\| > b > a$ and $\|y_2\| > a$. So, $y_1 - \lambda Mu$, $y_2 - \lambda Mu$ are two positive solutions of (1.5). The proof is completed. $\square$

4. Example

For convenience, we define the following notations:

$$[a, b] := \{x : x \in \mathbb{R}, a \leq x \leq b\}, \quad (a, b] := \{x : x \in \mathbb{R}, a < x \leq b\}, \quad \text{for } a, b \in \mathbb{R}, \quad a < b. \quad (4.1)$$

Example 4.1. Consider the following boundary value problem:

$$\begin{aligned} D_{0+}^{5/2} y(t) + \lambda f(t, y(t)) &= 0, \quad t \in [0, 1], \\ y(0) = y'(0) &= 0, \\ y'(1) &= h(y), \end{aligned} \quad (4.2)$$

where $h(y) = \int_0^1 y(s) dg(s)$, $g(s)$ is a bounded variation function on $[0, 1]$ with $0 < V_0^1(g) \leq 1 < 3/2$,

$$f(t, y) = \begin{cases} \sin\left(t - \frac{1}{2}\right)\pi - 1 - \sqrt{y}, & (t, y) \in [0, 1] \times [0, 6], \\ \sin\left(t - \frac{1}{2}\right)\pi - 1 + 601(y - 6) - \sqrt{y}, & (t, y) \in [0, 1] \times (6, 7], \\ \sin\left(t - \frac{1}{2}\right)\pi + 600 - \sqrt{y}, & (t, y) \in [0, 1] \times (7, 900], \\ \sin\left(t - \frac{1}{2}\right)\pi + 570, & (t, y) \in [0, 1] \times (900, +\infty). \end{cases} \quad (4.3)$$

Corresponding to the problem (1.5), we get that $\alpha = 5/2$, $\beta = 1$, $n = 3$, $h(y) \leq \|y\|V_0^1(g)$. Take $l = 1$, $k = 50$, $M = a = 6$, $b = 12$, $c = 36$, $d = 620$.

By simple calculation, we can get that the conditions of Theorem 3.1 are satisfied. So, when $(9/35)\sqrt{\pi} < \lambda < (15/16)\sqrt{\pi}$, the problem (4.2) has at least two positive solutions.

Example 4.2. Consider the following boundary value problem:

$$\begin{aligned} D_{0+}^{7/2} y(t) + \lambda f(t, y(t)) &= 0, \quad t \in [0, 1], \\ y(0) = y'(0) = y''(0) &= 0, \\ D_{0+}^{3/2} y(t)|_{t=1} &= h(y), \end{aligned} \quad (4.4)$$

where $h(y) = \int_0^1 y(s)dg(s)$, $g(s)$ is a bounded variation function on $[0, 1]$ with $0 < V_0^1(g) \leq (15/8)\sqrt{\pi} - 1$,

$$f(t, y) = \begin{cases} -\frac{3}{4} \cos \frac{\pi}{2}t - \frac{1}{4}\sqrt{y}, & (t, y) \in [0, 1] \times [0, 1], \\ -\frac{3}{4} \cos \frac{\pi}{2}t + 3205(y - 1) - \frac{1}{4}\sqrt{y}, & (t, y) \in [0, 1] \times (1, 1.2], \\ -\frac{3}{4} \cos \frac{\pi}{2}t + 641 - \frac{1}{4}\sqrt{y}, & (t, y) \in [0, 1] \times (1.2, 12], \\ -\frac{3}{4} \cos \frac{\pi}{2}t + 641 - \frac{1}{2}\sqrt{3}, & (t, y) \in [0, 1] \times (12, +\infty). \end{cases} \quad (4.5)$$

Corresponding to the problem (1.5), we get that $\alpha = 7/2$, $\beta = 3/2$, $n = 4$, $h(y) \leq \|y\|V_0^1(g)$. Take $l = (15/8)\sqrt{\pi} - 1$, $k = 320$, $M = a = 1$, $b = 2$, $c = 12$, $d = 642$.

Obviously, $f \in C([0, 1] \times \mathbb{R}^+, [-1, \infty))$, $h(y) \leq l\|y\|$. By simple calculation, we can get that $k, l, a, b, c, d, \alpha, \beta, M$ satisfy $0 < l < \Gamma(\alpha)/\Gamma(\alpha - \beta)$, $k > 2^{2\alpha-1}\beta\Gamma(\alpha)/(\Gamma(\alpha) - \Gamma(\alpha - \beta)l)$, and

$$\frac{[\Gamma(\alpha) - \Gamma(\alpha - \beta)l]\alpha M}{\beta\Gamma(\alpha)} < a < b - \frac{[\Gamma(\alpha) - \Gamma(\alpha - \beta)l]\alpha M}{\beta\Gamma(\alpha)} < 2^{\alpha-1}b < c < d. \quad (4.6)$$

It is easy to see that $f(t, y) + 1 \leq 642$, for $(t, y) \in [0, 1] \times [0, 642]$, and $f(t, y) + 1 \leq 1$, for $(t, y) \in [0, 1] \times [0, 1]$. So, conditions (C1) and (C3) of Theorem 3.1 hold.

For $(t, y) \in [1/2, 1] \times [2 - 56/45\sqrt{\pi}, 12]$, $f(t, y) + 1 \geq kb = 640$. Therefore, condition (C2) of Theorem 3.1 holds. So, for $(21/8(7\sqrt{2} - 2))\sqrt{\pi} < \lambda < 14/3$, the problem (4.4) has at least two positive solutions.

Specially, in Example 4.2, we take

$$g(t) = \begin{cases} 0, & t < \xi, \\ 1, & t \geq \xi, \end{cases} \quad (4.7)$$

where $0 < \xi < 1$ and all other conditions remain unchanged. Then $h(y) = y(\xi)$. Clearly, $V_0^1(g) = 1 < (15/8)\sqrt{\pi} - 1$. The problem (4.4) has at least two positive solutions for $(21/8(7\sqrt{2} - 2))\sqrt{\pi} < \lambda < 14/3$.

Acknowledgments

The authors are grateful to Editor Bashir Ahmad and anonymous referees for their constructive comments and suggestions which led to improvement of the paper. This work is supported by the Natural Science Foundation of China (11171088), the Doctoral Program Foundation of Hebei University of Science and Technology (QD201020), and the Foundation of Hebei University of Science and Technology (XL201236).

References

- [1] I. Podlubny, *Fractional Differential Equations*, vol. 198 of *Mathematics in Science and Engineering*, Academic Press, San Diego, Calif, USA, 1999.
- [2] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, New York, NY, USA, 1993.
- [3] V. Lakshmikantham and A. S. Vatsala, "Basic theory of fractional differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 69, no. 8, pp. 2677–2682, 2008.
- [4] R. P. Agarwal, V. Lakshmikantham, and J. J. Nieto, "On the concept of solution for fractional differential equations with uncertainty," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 6, pp. 2859–2862, 2010.
- [5] V. Lakshmikantham, "Theory of fractional functional differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 69, no. 10, pp. 3337–3343, 2008.
- [6] V. Lakshmikantham and S. Leela, "Nagumo-type uniqueness result for fractional differential equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 7-8, pp. 2886–2889, 2009.
- [7] T. G. Bhaskar, V. Lakshmikantham, and S. Leela, "Fractional differential equations with a Krasnoselskii-Krein type condition," *Nonlinear Analysis. Hybrid Systems*, vol. 3, no. 4, pp. 734–737, 2009.
- [8] V. Lakshmikantham and A. S. Vatsala, "General uniqueness and monotone iterative technique for fractional differential equations," *Applied Mathematics Letters*, vol. 21, no. 8, pp. 828–834, 2008.
- [9] C. Yuan, D. Jiang, and X. Xu, "Singular positive and semipositone boundary value problems of nonlinear fractional differential equations," *Mathematical Problems in Engineering*, vol. 2009, Article ID 535209, 17 pages, 2009.
- [10] Y. Wang, L. Liu, and Y. Wu, "Positive solutions for a class of fractional boundary value problem with changing sign nonlinearity," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 17, pp. 6434–6441, 2011.
- [11] C. S. Goodrich, "Existence of a positive solution to systems of differential equations of fractional order," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1251–1268, 2011.
- [12] X. Xu, D. Jiang, and C. Yuan, "Multiple positive solutions for the boundary value problem of a nonlinear fractional differential equation," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 10, pp. 4676–4688, 2009.
- [13] X. Li, S. Liu, and W. Jiang, "Positive solutions for boundary value problem of nonlinear fractional functional differential equations," *Applied Mathematics and Computation*, vol. 217, no. 22, pp. 9278–9285, 2011.
- [14] S. Liang and J. Zhang, "Existence and uniqueness of strictly nondecreasing and positive solution for a fractional three-point boundary value problem," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1333–1340, 2011.
- [15] H. Jafari and V. Daftardar-Gejji, "Positive solutions of nonlinear fractional boundary value problems using Adomian decomposition method," *Applied Mathematics and Computation*, vol. 180, no. 2, pp. 700–706, 2006.
- [16] Y. Wang, L. Liu, and Y. Wu, "Positive solutions for a nonlocal fractional differential equation," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 11, pp. 3599–3605, 2011.
- [17] S. Liang and J. Zhang, "Positive solutions for boundary value problems of nonlinear fractional differential equation," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 11, pp. 5545–5550, 2009.
- [18] Z. Bai and H. Lü, "Positive solutions for boundary value problem of nonlinear fractional differential equation," *Journal of Mathematical Analysis and Applications*, vol. 311, no. 2, pp. 495–505, 2005.
- [19] J. Caballero, J. Harjani, and K. Sadarangani, "Positive solutions for a class of singular fractional boundary value problems," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1325–1332, 2011.
- [20] R. P. Agarwal, D. O'Regan, and S. Staněk, "Positive solutions for Dirichlet problems of singular nonlinear fractional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 371, no. 1, pp. 57–68, 2010.
- [21] D. Jiang and C. Yuan, "The positive properties of the Green function for Dirichlet-type boundary value problems of nonlinear fractional differential equations and its application," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 2, pp. 710–719, 2010.

- [22] E. R. Kaufmann and E. Mboumi, "Positive solutions of a boundary value problem for a nonlinear fractional differential equation," *Electronic Journal of Qualitative Theory of Differential Equations*, no. 3, pp. 1–11, 2008.
- [23] S. Staněk, "The existence of positive solutions of singular fractional boundary value problems," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1379–1388, 2011.
- [24] T. Jankowski, "Positive solutions for second order impulsive differential equations involving Stieltjes integral conditions," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 11, pp. 3775–3785, 2011.
- [25] R. P. Agarwal, B. de Andrade, and G. Siracusa, "On fractional integro-differential equations with state-dependent delay," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1143–1149, 2011.
- [26] Z.-W. Lv, J. Liang, and T.-J. Xiao, "Solutions to the Cauchy problem for differential equations in Banach spaces with fractional order," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1303–1311, 2011.
- [27] C. Kou, H. Zhou, and Y. Yan, "Existence of solutions of initial value problems for nonlinear fractional differential equations on the half-axis," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 17, pp. 5975–5986, 2011.
- [28] M. Benchohra, S. Hamani, and S. K. Ntouyas, "Boundary value problems for differential equations with fractional order and nonlocal conditions," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 7–8, pp. 2391–2396, 2009.
- [29] Y. Zhou, F. Jiao, and J. Li, "Existence and uniqueness for fractional neutral differential equations with infinite delay," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 7–8, pp. 3249–3256, 2009.
- [30] G. M. N'Guérékata, "A Cauchy problem for some fractional abstract differential equation with non local conditions," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 70, no. 5, pp. 1873–1876, 2009.
- [31] Z. Bai, "On positive solutions of a nonlocal fractional boundary value problem," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 2, pp. 916–924, 2010.
- [32] R. P. Agarwal, Y. Zhou, and Y. He, "Existence of fractional neutral functional differential equations," *Computers & Mathematics with Applications*, vol. 59, no. 3, pp. 1095–1100, 2010.
- [33] S. Zhang, "Positive solutions for boundary-value problems of nonlinear fractional differential equations," *Electronic Journal of Differential Equations*, vol. 2006, pp. 1–12, 2006.
- [34] H. A. H. Salem, "On the fractional order m -point boundary value problem in reflexive Banach spaces and weak topologies," *Journal of Computational and Applied Mathematics*, vol. 224, no. 2, pp. 565–572, 2009.
- [35] W. Zhong and W. Lin, "Nonlocal and multiple-point boundary value problem for fractional differential equations," *Computers & Mathematics with Applications*, vol. 59, no. 3, pp. 1345–1351, 2010.
- [36] C. F. Li, X. N. Luo, and Y. Zhou, "Existence of positive solutions of the boundary value problem for nonlinear fractional differential equations," *Computers & Mathematics with Applications*, vol. 59, no. 3, pp. 1363–1375, 2010.
- [37] M. Al-Refai and M. Ali Hajji, "Monotone iterative sequences for nonlinear boundary value problems of fractional order," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 11, pp. 3531–3539, 2011.
- [38] X. Su, "Solutions to boundary value problem of fractional order on unbounded domains in a Banach space," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 8, pp. 2844–2852, 2011.
- [39] H. A. H. Salem, "Fractional order boundary value problem with integral boundary conditions involving Pettis integral," *Acta Mathematica Scientia B*, vol. 31, no. 2, pp. 661–672, 2011.
- [40] Z. Bai, "Solvability for a class of fractional m -point boundary value problem at resonance," *Computers & Mathematics with Applications*, vol. 62, no. 3, pp. 1292–1302, 2011.
- [41] M. U. Rehman, R. A. Khan, and N. A. Asif, "Three point boundary value problems for nonlinear fractional differential equations," *Acta Mathematica Scientia B*, vol. 31, no. 4, pp. 1337–1346, 2011.
- [42] B. Ahmad, J. J. Nieto, and A. Alsaedi, "Existence and uniqueness of solutions for nonlinear fractional differential equations with non-separated type integral boundary conditions," *Acta Mathematica Scientia B*, vol. 31, pp. 2122–2130, 2011.
- [43] G. Wang, B. Ahmad, and L. Zhang, "Impulsive anti-periodic boundary value problem for nonlinear differential equations of fractional order," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 74, no. 3, pp. 792–804, 2011.
- [44] B. Ahmad and S. K. Ntouyas, "A four-point nonlocal integral boundary value problem for fractional differential equations of arbitrary order," *Electronic Journal of Qualitative Theory of Differential Equations*, no. 22, pp. 1–15, 2011.

- [45] B. Ahmad and J. J. Nieto, "Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions," *Computers & Mathematics with Applications*, vol. 58, no. 9, pp. 1838–1843, 2009.
- [46] M. El-Shahed, "Positive solutions for boundary value problem of nonlinear fractional differential equation," *Abstract and Applied Analysis*, vol. 2007, Article ID 10368, 8 pages, 2007.
- [47] R. I. Avery and A. C. Peterson, "Three positive fixed points of nonlinear operators on ordered Banach spaces," *Computers & Mathematics with Applications*, vol. 42, no. 3–5, pp. 313–322, 2001.