Research Article

Common Fixed Point Theorems for Six Mappings in Generalized Metric Spaces

Feng Gu

*Institute of Applied Mathematics and Department of Mathematics, Hangzhou Normal University,
Hangzhou, Zhejiang 310036, China*

Correspondence should be addressed to Feng Gu, gufeng99@sohu.com

Received 23 September 2012; Accepted 6 October 2012

Academic Editor: RuDong Chen

Copyright © 2012 Feng Gu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By using the weakly commutative and weakly compatible conditions of self-mapping pairs, we prove some new common fixed point theorems for six self-mappings in the framework of generalized metric spaces. An example is provided to support our result. The results presented in this paper generalize the well-known comparable results in the literature due to Abbas, Nazir, Saadati, Mustafa, and Sims.

1. Introduction and Preliminaries

The study of fixed points of mappings satisfying certain conditions has been at the center of vigorous research activity. In 2006, Mustafa and Sims [1] introduced a new structure of generalized metric spaces, which are called G -metric spaces as follows.

Definition 1.1 (see [1]). Let X be a nonempty set and let $G : X \times X \times X \rightarrow R^+$ be a function satisfying the following properties:

- (G₁) $G(x, y, z) = 0$ if $x = y = z$;
- (G₂) $0 < G(x, x, y)$, for all $x, y \in X$ with $x \neq y$;
- (G₃) $G(x, x, y) \leq G(x, y, z)$ for all $x, y, z \in X$ with $z \neq y$;
- (G₄) $G(x, y, z) = G(x, z, y) = G(y, z, x) = \dots$, symmetry in all three variables;
- (G₅) $G(x, y, z) \leq G(x, a, a) + G(a, y, z)$ for all $x, y, z, a \in X$ (rectangle inequality).

Then the function G is called a generalized metric, or, more specifically, a G -metric on X , and the pair (X, G) is called a G -metric space.

Since then the fixed point theory in G -metric spaces has been studied and developed by authors (see [2–43]). Fixed point problems have also been considered in partially ordered G -metric spaces (see [44–56]).

The purpose of this paper is to use the concept of weakly commuting mappings and weakly compatible mappings to discuss some new common fixed point problem for six self-mappings in G -metric spaces. The results presented in this paper extend and improve the corresponding results of Abbas et al. [2] and Mustafa and Sims [3].

We now recall some of the basic concepts and results in G -metric spaces.

Proposition 1.2 (see [1]). *Let (X, G) be a G -metric space, then the function $G(x, y, z)$ is jointly continuous in three of its variables.*

Definition 1.3 (see [1]). Let (X, G) be a G -metric space, and let (x_n) be a sequence of points of X . A point $x \in X$ is said to be the limit of the sequence (x_n) , if $\lim_{n,m \rightarrow +\infty} G(x, x_n, x_m) = 0$, and we say that the sequence (x_n) is G -convergent to x or (x_n) G -convergent to x , that is, for any $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that $G(x, x_n, x_m) < \epsilon$ for all $m, n \geq N$ (throughout this paper we mean by $\mathbb{N}$ the set of all natural numbers).

Proposition 1.4 (see [1]). *Let (X, G) be a G -metric space, then the following are equivalent:*

- (i) (x_n) is G -convergent to x ;
- (ii) $G(x_n, x_n, x) \rightarrow 0$ as $n \rightarrow +\infty$;
- (iii) $G(x_n, x, x) \rightarrow 0$ as $n \rightarrow +\infty$;
- (iv) $G(x_n, x_m, x) \rightarrow 0$ as $n, m \rightarrow +\infty$.

Definition 1.5 (see [1]). Let (X, G) be a G -metric space. A sequence (x_n) is called G -cauchy if for every $\epsilon > 0$, there is $N \in \mathbb{N}$ such that $G(x_n, x_m, x_l) < \epsilon$ for all $m, n, l \geq N$, that is $G(x_n, x_m, x_l) \rightarrow 0$ as $n, m, l \rightarrow +\infty$.

Proposition 1.6 (see [1]). *Let (X, G) be a G -metric space, then the following are equivalent:*

- (i) the sequence (x_n) is G -cauchy;
- (ii) for every $\epsilon > 0$, there is $N \in \mathbb{N}$ such that $G(x_n, x_m, x_m) < \epsilon$ for all $m, n \geq N$.

Definition 1.7 (see [1]). A G -metric space (X, G) is G -complete if every G -cauchy sequence in (X, G) is G -convergent in X .

Definition 1.8 (see [1]). Let (X, G) and (X', G') be G -metric spaces, and let $f : (X, G) \rightarrow (X', G')$ be a function. Then f is said to be G -continuous at a point $a \in X$ if and only if for every $\epsilon > 0$, there is $\delta > 0$ such that $x, y \in X$, and $G(a, x, y) < \delta$ implies $G'(f(a), f(x), f(y)) < \epsilon$. A function f is G -continuous at X if and only if it is G -continuous at all $a \in X$.

Proposition 1.9 (see [1]). *Let (X, G) and (X', G') be G -metric spaces, then a function $f : X \rightarrow X'$ is G -continuous at a point $x \in X$ if and only if it is G -sequentially continuous at x , that is, whenever (x_n) is G -convergent to x , $(f(x_n))$ is G -convergent to $f(x)$.*

Definition 1.10 (see [4]). Two self-mappings f and g of a G -metric space (X, G) are said to be weakly commuting if $G(fgx, gfx, gfx) \leq G(fx, gx, gx)$ for all x in X .

Definition 1.11 (see [4]). Let f and g be two self-mappings from a G -metric space (X, G) into itself. Then the mappings f and g are said to be weakly compatible if $G(fgx, gfx, gfx) = 0$ whenever $G(fx, gx, gx) = 0$.

Proposition 1.12 (see [1]). Let (X, G) be a G -metric space. Then, for all x, y, z, a in X , it follows that

- (i) if $G(x, x, y) = 0$, then $x = y = z$;
- (ii) $G(x, y, z) \leq G(x, x, y) + G(x, x, z)$;
- (iii) $G(x, y, y) \leq 2G(y, x, x)$;
- (iv) $G(x, y, z) \leq G(x, a, z) + G(a, y, z)$;
- (v) $G(x, y, z) \leq (2/3)(G(x, y, a) + G(x, a, z) + G(a, y, z))$;
- (vi) $G(x, y, z) \leq G(x, a, a) + G(y, a, a) + G(z, a, a)$.

2. Common Fixed Point Theorems

Theorem 2.1. Let (X, G) be a complete G -metric space, and let f, g, h, A, B , and C be six mappings of X into itself satisfying the following conditions:

- (i) $f(X) \subset B(X)$, $g(X) \subset C(X)$, $h(X) \subset A(X)$;
- (ii) for all $x, y, z \in X$,

$$G(fx, gy, hz) \leq k \max \left\{ \begin{array}{l} G(Ax, gy, gy) + G(By, fx, fx), \\ G(By, hz, hz) + G(Cz, gy, gy), \\ G(Cz, fx, fx) + G(Ax, hz, hz) \end{array} \right\} \quad (2.1)$$

or

$$G(fx, gy, hz) \leq k \max \left\{ \begin{array}{l} G(Ax, Ax, gy) + G(By, By, fx), \\ G(By, By, hz) + G(Cz, Cz, gy), \\ G(Cz, Cz, fx) + G(Ax, Ax, hz) \end{array} \right\}, \quad (2.2)$$

where $k \in [0, 1/3)$. Then one of the pairs (f, A) , (g, B) , and (h, C) has a coincidence point in X . Further, if one of the following conditions is satisfied, then the mappings f, g, h, A, B , and C have a unique common fixed point in X .

- (a) Either f or A is G -continuous, the pair (f, A) is weakly commutative, the pairs (g, B) and (h, C) are weakly compatible;
- (b) Either g or B is G -continuous, the pair (g, B) is weakly commutative, the pairs (f, A) and (h, C) are weakly compatible;
- (c) Either h or C is G -continuous, the pair (h, C) is weakly commutative, the pairs (f, A) and (g, B) are weakly compatible.

Proof. Suppose that mappings f, g, h, A, B , and C satisfy condition (2.1).

Let x_0 in X be arbitrary point, since $f(X) \subset B(X)$, $g(X) \subset C(X)$, $h(X) \subset A(X)$, there exist the sequences $\{x_n\}$ and $\{y_n\}$ in X such that

$$y_{3n} = fx_{3n} = Bx_{3n+1}, \quad y_{3n+1} = gx_{3n+1} = Cx_{3n+2}, \quad y_{3n+2} = hx_{3n+2} = Ax_{3n+3}, \quad (2.3)$$

for $n = 0, 1, 2, \dots$

If $y_n = y_{n+1}$ for some n , with $n = 3m$, then $p = x_{3m+1}$ is a coincidence point of the pair (g, B) ; if $y_{n+1} = y_{n+2}$ for some n , with $n = 3m$, then $p = x_{3m+2}$ is a coincidence point of the pair (h, C) ; if $y_{n+2} = y_{n+3}$ for some n , with $n = 3m$, then $p = x_{3m+3}$ is a coincidence point of the pair (f, A) . Without loss of generality, we can assume that $y_n \neq y_{n+1}$ for all $n = 0, 1, 2, \dots$

Now we prove that $\{y_n\}$ is a G -cauchy sequence in X .

Actually, using the condition (2.1) and (G_3) , we have

$$\begin{aligned} & G(y_{3n-1}, y_{3n}, y_{3n+1}) \\ &= G(fx_{3n}, gx_{3n+1}, hx_{3n-1}) \\ &\leq k \max \left\{ \begin{array}{l} G(Ax_{3n}, gx_{3n+1}, gx_{3n+1}) + G(Bx_{3n+1}, fx_{3n}, fx_{3n}), \\ G(Bx_{3n+1}, hx_{3n-1}, hx_{3n-1}) + G(Cx_{3n-1}, gx_{3n+1}, gx_{3n+1}), \\ G(Cx_{3n-1}, fx_{3n}, fx_{3n}) + G(Ax_{3n}, hx_{3n-1}, hx_{3n-1}) \end{array} \right\} \\ &= k \max \left\{ \begin{array}{l} G(y_{3n-1}, y_{3n+1}, y_{3n+1}) + G(y_{3n}, y_{3n}, y_{3n}), \\ G(y_{3n}, y_{3n-1}, y_{3n-1}) + G(y_{3n-2}, y_{3n+1}, y_{3n+1}), \\ G(y_{3n-2}, y_{3n}, y_{3n}) + G(y_{3n-1}, y_{3n-1}, y_{3n-1}) \end{array} \right\} \\ &\leq k \max \left\{ \begin{array}{l} G(y_{3n-1}, y_{3n}, y_{3n+1}), G(y_{3n-2}, y_{3n-1}, y_{3n}), \\ G(y_{3n-1}, y_{3n}, y_{3n+1}) + [G(y_{3n-2}, y_{3n-1}, y_{3n-1}) + G(y_{3n-1}, y_{3n+1}, y_{3n+1})] \end{array} \right\} \\ &\leq k \max \left\{ \begin{array}{l} G(y_{3n-1}, y_{3n}, y_{3n+1}), G(y_{3n-2}, y_{3n-1}, y_{3n}), \\ 2G(y_{3n-1}, y_{3n}, y_{3n+1}) + G(y_{3n-2}, y_{3n-1}, y_{3n}) \end{array} \right\} \\ &= k [2G(y_{3n-1}, y_{3n}, y_{3n+1}) + G(y_{3n-2}, y_{3n-1}, y_{3n})], \end{aligned} \quad (2.4)$$

which further implies that

$$(1 - 2k)G(y_{3n-1}, y_{3n}, y_{3n+1}) \leq kG(y_{3n-2}, y_{3n-1}, y_{3n}). \quad (2.5)$$

Thus

$$G(y_{3n-1}, y_{3n}, y_{3n+1}) \leq \lambda G(y_{3n-2}, y_{3n-1}, y_{3n}), \quad (2.6)$$

where $\lambda = k/(1 - 2k)$. Obviously $0 \leq \lambda < 1$.

Similarly it can be shown that

$$\begin{aligned} G(y_{3n}, y_{3n+1}, y_{3n+2}) &\leq \lambda G(y_{3n-1}, y_{3n}, y_{3n+1}), \\ G(y_{3n+1}, y_{3n+2}, y_{3n+3}) &\leq \lambda G(y_{3n}, y_{3n+1}, y_{3n+2}). \end{aligned} \quad (2.7)$$

It follows from (2.6) and (2.7) that, for all $n \in \mathbb{N}$,

$$G(y_n, y_{n+1}, y_{n+2}) \leq \lambda G(y_{n-1}, y_n, y_{n+1}) \leq \lambda^2 G(y_{n-2}, y_{n-1}, y_n) \leq \cdots \leq \lambda^n G(y_0, y_1, y_2). \quad (2.8)$$

Therefore, for all $n, m \in \mathbb{N}$, $n < m$, by (G_3) and (G_5) we have

$$\begin{aligned} G(y_n, y_m, y_m) &\leq G(y_n, y_{n+1}, y_{n+1}) + G(y_{n+1}, y_{n+2}, y_{n+2}) + G(y_{n+2}, y_{n+3}, y_{n+3}) \\ &\quad + \cdots + G(y_{m-1}, y_m, y_m) \\ &\leq G(y_n, y_{n+1}, y_{n+2}) + G(y_{n+1}, y_{n+2}, y_{n+3}) + \cdots + G(y_{m-1}, y_m, y_{m+1}) \\ &\leq (\lambda^n + \lambda^{n+1} + \lambda^{n+2} + \cdots + \lambda^{m-1}) G(y_0, y_1, y_2) \\ &\leq \frac{\lambda^n}{1-\lambda} G(y_0, y_1, y_2) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (2.9)$$

Hence $\{y_n\}$ is a G -Cauchy sequence in X , since X is complete G -metric space, there exists a point $u \in X$ such that $y_n \rightarrow u$ ($n \rightarrow \infty$).

Since the sequences $\{fx_{3n}\} = \{Bx_{3n+1}\}$, $\{gx_{3n+1}\} = \{Cx_{3n+2}\}$, and $\{hx_{3n-1}\} = \{Ax_{3n}\}$ are all subsequences of $\{y_n\}$, then they all converge to u , that is,

$$\begin{aligned} y_{3n} = fx_{3n} = Bx_{3n+1} &\rightarrow u, & y_{3n+1} = gx_{3n+1} = Cx_{3n+2} &\rightarrow u, \\ y_{3n-1} = hx_{3n-1} = Ax_{3n} &\rightarrow u \quad (n \rightarrow \infty). \end{aligned} \quad (2.10)$$

Now we prove that u is a common fixed point of f , g , h , A , B , and C under the condition (a).

First, we suppose that A is continuous, the pair (f, A) is weakly commutative, the pairs (g, B) and (h, C) are weakly compatible.

Step 1. We prove that $u = fu = Au$.

By (2.10) and weakly commutativity of mapping pair (f, A) , we have

$$G(fAx_{3n}, Af x_{3n}, Af x_{3n}) \leq G(fx_{3n}, Ax_{3n}, Ax_{3n}) \rightarrow 0 \quad (n \rightarrow \infty). \quad (2.11)$$

Since A is continuous, then $A^2x_{3n} \rightarrow Au$ ($n \rightarrow \infty$), $Afx_{3n} \rightarrow Au$ ($n \rightarrow \infty$). By (2.11), we know that $fAx_{3n} \rightarrow Au$ ($n \rightarrow \infty$).

From the condition (2.1), we get

$$\begin{aligned} &G(fAx_{3n}, gx_{3n+1}, hx_{3n+2}) \\ &\leq k \max \left\{ \begin{aligned} &G(A^2x_{3n}, gx_{3n+1}, gx_{3n+1}) + G(Bx_{3n+1}, fAx_{3n}, fAx_{3n}), \\ &G(Bx_{3n+1}, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gx_{3n+1}, gx_{3n+1}), \\ &G(Cx_{3n+2}, fAx_{3n}, fAx_{3n}) + G(A^2x_{3n}, hx_{3n+2}, hx_{3n+2}) \end{aligned} \right\}. \end{aligned} \quad (2.12)$$

Letting $n \rightarrow \infty$ and using the Proposition 1.12(iii), we have

$$\begin{aligned} G(Au, u, u) &\leq k \max \left\{ \begin{array}{l} G(Au, u, u) + G(u, Au, Au), \\ G(u, u, u) + G(u, u, u), \\ G(u, Au, Au) + G(Au, u, u) \end{array} \right\} \\ &= k[G(Au, u, u) + G(u, Au, Au)] \\ &\leq 3kG(Au, u, u). \end{aligned} \quad (2.13)$$

Hence, $G(Au, u, u) = 0$ and $Au = u$, since $0 \leq k < 1/3$.

Again by using condition (2.1), we have

$$\begin{aligned} &G(fu, gx_{3n+1}, hx_{3n+2}) \\ &\leq k \max \left\{ \begin{array}{l} G(Au, gx_{3n+1}, gx_{3n+1}) + G(Bx_{3n+1}, fu, fu), \\ G(Bx_{3n+1}, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gx_{3n+1}, gx_{3n+1}), \\ G(Cx_{3n+2}, fu, fu) + G(Au, hx_{3n+2}, hx_{3n+2}) \end{array} \right\}. \end{aligned} \quad (2.14)$$

Letting $n \rightarrow \infty$, we have

$$G(fu, u, u) \leq kG(u, fu, fu). \quad (2.15)$$

From the Proposition 1.12(iii), we get

$$G(fu, u, u) \leq kG(u, fu, fu) \leq 2kG(fu, u, u). \quad (2.16)$$

Hence, $G(fu, u, u) = 0$ and $fu = u$, since $0 \leq k < 1/3$.

So we have $u = Au = fu$.

Step 2. We prove that $u = gu = Bu$.

Since $f(X) \subset B(X)$ and $u = fu \in f(X)$, there is a point $v \in X$ such that $u = fu = Bv$.

Again by using condition (2.1), we have

$$G(fu, gv, hx_{3n+2}) \leq k \max \left\{ \begin{array}{l} G(Au, gv, gv) + G(Bv, fu, fu), \\ G(Bv, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gv, gv), \\ G(Cx_{3n+2}, fu, fu) + G(Au, hx_{3n+2}, hx_{3n+2}) \end{array} \right\}. \quad (2.17)$$

Letting $n \rightarrow \infty$, using $u = Au = fu$ and the Proposition 1.12(iii), we obtain

$$G(u, gv, u) \leq kG(u, gv, gv) \leq 2kG(u, gv, u). \quad (2.18)$$

Hence, $G(u, gv, u) = 0$ and so $gv = u = Bv$, since $0 \leq k < 1/3$.

Since the pair (g, B) is weakly compatible, we have

$$gu = gBv = Bgv = Bu. \quad (2.19)$$

Again by using condition (2.1), we have

$$G(fu, gu, hx_{3n+2}) \leq k \max \left\{ \begin{array}{l} G(Au, gu, gu) + G(Bu, fu, fu), \\ G(Bu, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gu, gu), \\ G(Cx_{3n+2}, fu, fu) + G(Au, hx_{3n+2}, hx_{3n+2}) \end{array} \right\}. \quad (2.20)$$

Letting $n \rightarrow \infty$, using $u = Au = fu$, $gu = Bu$ and the Proposition 1.12(iii), we have

$$G(u, gu, u) \leq k[G(u, gu, gu) + G(u, gu, u)] \leq 3kG(u, gu, u). \quad (2.21)$$

Hence, $G(u, gu, u) = 0$ and so $u = gu = Bu$, since $0 \leq k < 1/3$.

So we have $u = gu = Bu$.

Step 3. We prove that $u = hu = Cu$.

Since $g(X) \subset C(X)$ and $u = gu \in g(X)$, there is a point $w \in X$ such that $u = gu = Cw$. Again by using condition (2.1), we have

$$G(fu, gu, hw) \leq k \max \left\{ \begin{array}{l} G(Au, gu, gu) + G(Bu, fu, fu), \\ G(Bu, hw, hw) + G(Cw, gu, gu), \\ G(Cw, fu, fu) + G(Au, hw, hw) \end{array} \right\}. \quad (2.22)$$

Using $u = Au = fu$, $u = gu = Bu = Cw$ and the Proposition 1.12(iii), we obtain

$$G(u, u, hw) \leq kG(u, hw, hw) \leq 2kG(u, u, hw). \quad (2.23)$$

Hence, $G(u, u, hw) = 0$ and so $hw = u = Cw$, since $0 \leq k < 1/3$.

Since the pair (h, C) is weakly compatible, we have

$$hu = hCw = Chw = Cu. \quad (2.24)$$

Again by using condition (2.1), we have

$$G(fu, gu, hu) \leq k \max \left\{ \begin{array}{l} G(Au, gu, gu) + G(Bu, fu, fu), \\ G(Bu, hu, hu) + G(Cu, gu, gu), \\ G(Cu, fu, fu) + G(Au, hu, hu) \end{array} \right\}. \quad (2.25)$$

Using $u = Au = fu$, $u = gu = Bu$, $Cu = hu$ and the Proposition 1.12(iii), we have

$$G(u, u, hu) \leq k[G(u, hu, hu) + G(hu, u, u)] \leq 3kG(u, u, hu). \quad (2.26)$$

Hence, $G(u, u, hu) = 0$ and so $u = hu = Cu$, since $0 \leq k < 1/3$.

Therefore, u is the common fixed point of f, g, h, A, B , and C when A is continuous and the pair (f, A) is weakly commutative, the pairs (g, B) and (h, C) are weakly compatible.

Next, we suppose that f is continuous, the pair (f, A) is weakly commutative, the pairs (g, B) and (h, C) are weakly compatible.

Step 1. We prove that $u = fu$.

By (2.10) and weak commutativity of mapping pair (f, A) , we have

$$G(fAx_{3n}, Af x_{3n}, Af x_{3n}) \leq G(fx_{3n}, Ax_{3n}, Ax_{3n}) \longrightarrow 0 \quad (n \longrightarrow \infty). \quad (2.27)$$

Since f is continuous, then $f^2x_{3n} \rightarrow fu$ ($n \rightarrow \infty$), $fAx_{3n} \rightarrow fu$ ($n \rightarrow \infty$). By (2.10), we know that $Afx_{3n} \rightarrow fu$ ($n \rightarrow \infty$).

From the condition (2.1), we have

$$G(f^2x_{3n}, gx_{3n+1}, hx_{3n+2}) \leq k \max \left\{ \begin{array}{l} G(Afx_{3n}, gx_{3n+1}, gx_{3n+1}) + G(Bx_{3n+1}, f^2x_{3n}, f^2x_{3n}), \\ G(Bx_{3n+1}, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gx_{3n+1}, gx_{3n+1}), \\ G(Cx_{3n+2}, f^2x_{3n}, f^2x_{3n}) + G(Afx_{3n}, hx_{3n+2}, hx_{3n+2}) \end{array} \right\}. \quad (2.28)$$

Letting $n \rightarrow \infty$ and noting the Proposition 1.12(iii), we have

$$\begin{aligned} G(fu, u, u) &\leq k \max \left\{ \begin{array}{l} G(fu, u, u) + G(u, fu, fu), \\ G(u, u, u) + G(u, u, u), \\ G(u, fu, fu) + G(fu, u, u) \end{array} \right\} \\ &= k[G(fu, u, u) + G(u, fu, fu)] \\ &\leq 3kG(fu, u, u). \end{aligned} \quad (2.29)$$

Hence, $G(fu, u, u) = 0$ and so $fu = u$, since $0 \leq k < 1/3$.

Step 2. We prove that $u = gu = Bu$.

Since $f(X) \subset B(X)$ and $u = fu \in f(X)$, there is a point $z \in X$ such that $u = fu = Bz$. Again by using condition (2.1), we have

$$G(f^2x_{3n}, gz, hx_{3n+2}) \leq k \max \left\{ \begin{array}{l} G(Afx_{3n}, gz, gz) + G(Bz, f^2x_{3n}, f^2x_{3n}), \\ G(Bz, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gz, gz), \\ G(Cx_{3n+2}, f^2x_{3n}, f^2x_{3n}) + G(Afx_{3n}, hx_{3n+2}, hx_{3n+2}) \end{array} \right\}. \quad (2.30)$$

Letting $n \rightarrow \infty$ and using $u = fu$ and the Proposition 1.12(iii), we have

$$G(u, gz, u) \leq kG(u, gz, gz) \leq 2kG(u, gz, u). \quad (2.31)$$

Hence $G(u, gz, u) = 0$ and so $gz = u = Bz$, since $0 \leq k < 1/3$.

Since the pair (g, B) is weakly compatible, we have

$$gu = gBz = Bgz = Bu. \quad (2.32)$$

Again by using condition (2.1), we have

$$G(fx_{3n}, gu, hx_{3n+2}) \leq k \max \left\{ \begin{array}{l} G(Ax_{3n}, gu, gu) + G(Bu, fx_{3n}, fx_{3n}), \\ G(Bu, hx_{3n+2}, hx_{3n+2}) + G(Cx_{3n+2}, gu, gu), \\ G(Cx_{3n+2}, fx_{3n}, fx_{3n}) + G(Ax_{3n}, hx_{3n+2}, hx_{3n+2}) \end{array} \right\}. \quad (2.33)$$

Letting $n \rightarrow \infty$ and using $u = fu$, $gu = Bu$ and the Proposition 1.12(iii), we have

$$G(u, gu, u) \leq k[G(u, gu, gu) + G(gu, u, u)] \leq 3kG(u, gu, u). \quad (2.34)$$

Hence, $G(u, gu, u) = 0$ and so $gu = u = Bu$, since $0 \leq k < 1/3$.

So we have $u = gu = Bu$.

Step 3. We prove that $u = hu = Cu$.

Since $g(X) \subset C(X)$ and $u = gu \in g(X)$, there is a point $t \in X$ such that $u = gu = Ct$. Again by using condition (2.1), we have

$$G(fx_{3n}, gu, ht) \leq k \max \left\{ \begin{array}{l} G(Ax_{3n}, gu, gu) + G(Bu, fx_{3n}, fx_{3n}), \\ G(Bu, ht, ht) + G(Ct, gu, gu), \\ G(Ct, fx_{3n}, fx_{3n}) + G(Ax_{3n}, ht, ht) \end{array} \right\}. \quad (2.35)$$

Letting $n \rightarrow \infty$ and using $u = gu = Bu$ and the Proposition 1.12(iii), we obtain

$$G(u, u, ht) \leq kG(u, ht, ht) \leq 2kG(u, u, ht). \quad (2.36)$$

Hence, $G(u, u, ht) = 0$ and so $ht = u = Ct$, since $0 \leq k < 1/3$.

Since the pair (h, C) is weakly compatible, we have

$$hu = hCt = Cht = Cu. \quad (2.37)$$

Again by using condition (2.1), we have

$$G(fx_{3n}, gu, hu) \leq k \max \left\{ \begin{array}{l} G(Ax_{3n}, gu, gu) + G(Bu, fx_{3n}, fx_{3n}), \\ G(Bu, hu, hu) + G(Cu, gu, gu), \\ G(Cu, fx_{3n}, fx_{3n}) + G(Ax_{3n}, hu, hu) \end{array} \right\}. \quad (2.38)$$

Letting $n \rightarrow \infty$ and using $u = gu = Bu$ and the Proposition 1.12(iii), we have

$$G(u, u, hu) \leq k[G(u, hu, hu) + G(u, u, hu)] \leq 3kG(u, u, hu). \quad (2.39)$$

Hence, $G(u, u, hu) = 0$ and so $hu = u = Cu$, since $0 \leq k < 1/3$.

Step 4. We prove that $u = Au$.

Since $h(X) \subset A(X)$ and $u = hu \in h(X)$, there is a point $p \in X$ such that $u = hu = Ap$. Again by using condition (2.1), we have

$$G(fp, gu, hu) \leq k \max \left\{ \begin{array}{l} G(Ap, gu, gu) + G(Bu, fp, fp), \\ G(Bu, hu, hu) + G(Cu, gu, gu), \\ G(Cu, fp, fp) + G(Ap, hu, hu) \end{array} \right\}. \quad (2.40)$$

Using $u = gu = Bu$, $u = hu = Cu$, and the Proposition 1.12(iii), we obtain

$$G(fp, u, u) \leq kG(u, fp, fp) \leq 2kG(fp, u, u). \quad (2.41)$$

Hence $G(fp, u, u) = 0$ and $fp = u = Ap$, since $0 \leq k < 1/3$.

Since the pair (f, A) is weakly compatible, we have

$$fu = fAp = Afp = Au = u. \quad (2.42)$$

Therefore, u is the common fixed point of f, g, h, A, B , and C when f is continuous and the pair (f, A) is weakly commutative, the pairs (g, B) and (h, C) are weakly compatible.

Similarly, we can prove the result that u is a common fixed point of f, g, h, A, B , and C when under the condition of (b) or (c).

Finally, we prove uniqueness of common fixed point u .

Let u and q be two common fixed points of f, g, h, A, B , and C , by using condition (2.1), we have

$$\begin{aligned} G(q, u, u) &= G(fq, gu, hu) \\ &\leq k \max \left\{ \begin{array}{l} G(Aq, gu, gu) + G(Bu, fq, fq), \\ G(Bu, hu, hu) + G(Cu, gu, gu), \\ G(Cu, fq, fq) + G(Aq, hu, hu) \end{array} \right\} \\ &= k[G(q, u, u) + G(u, q, q)] \\ &\leq 3kG(q, u, u). \end{aligned} \quad (2.43)$$

Hence, $G(q, u, u) = 0$ and so $q = u$, since $0 \leq k < 1/3$. Thus common fixed point is unique.

The proof using (2.2) is similar. This completes the proof. $\square$

Now we introduce an example to support Theorem 2.1.

Example 2.2. Let $X = [0, 1]$, and let (X, G) be a G -metric space defined by $G(x, y, z) = |x - y| + |y - z| + |z - x|$ for all x, y , and z in X . Let f, g, h, A, B , and C be self-mappings defined by

$$fx = \begin{cases} 1, & x \in \left[0, \frac{1}{2}\right], \\ \frac{7}{8}, & x \in \left(\frac{1}{2}, 1\right], \end{cases} \quad gx = \begin{cases} \frac{10}{11}, & x \in \left[0, \frac{1}{2}\right], \\ \frac{7}{8}, & x \in \left(\frac{1}{2}, 1\right], \end{cases} \quad hx = \begin{cases} \frac{9}{10}, & x \in \left[0, \frac{1}{2}\right], \\ \frac{7}{8}, & x \in \left(\frac{1}{2}, 1\right], \end{cases}$$

$$Ax = x, \quad Bx = \begin{cases} 1, & x \in \left[0, \frac{1}{2}\right], \\ \frac{7}{8}, & x \in \left(\frac{1}{2}, 1\right), \\ 0, & x = 1, \end{cases} \quad Cx = \begin{cases} 1, & x \in \left[0, \frac{1}{2}\right], \\ \frac{7}{8}, & x \in \left(\frac{1}{2}, 1\right), \\ \frac{10}{11}, & x = 1. \end{cases} \quad (2.44)$$

Note that A is G -continuous in X , and f, g, h, B , and C are not G -continuous in X .

Clearly we can get $f(X) \subset B(X)$, $g(X) \subset C(X)$, and $h(X) \subset A(X)$.

Actually, since $fX = \{7/8, 1\}$, $BX = \{0, 7/8, 1\}$, $gX = \{7/8, 10/11\}$, $CX = \{7/8, 10/11, 1\}$, $hX = \{7/8, 9/10\}$, and $AX = X = [0, 1]$, so we know $f(X) \subset B(X)$, $g(X) \subset C(X)$, and $h(X) \subset A(X)$.

By the definition of the mappings of f and A , for all $x \in [0, 1]$, $G(fAx, Afx, Afx) = G(fx, fx, fx) = 0 \leq G(fx, Ax, Ax)$, so we can get the pair (f, A) is weakly commuting.

By the definition of the mappings of g and B , only for $x \in (1/2, 1)$, $gx = Bx = 7/8$, at this time $gBx = g(7/8) = 7/8 = B(7/8) = Bgx$, so $gBx = Bgx$, so we can obtain that the pair (g, B) is weakly compatible. Similarly, we can show that the pair (h, C) is also weakly compatible.

Now we proof that the mappings f, g, h, A, B , and C are satisfying the condition (2.1) of Theorem 2.1 with $k = 5/16 \in [0, 1/3)$. Let

$$M(x, y, z) = \max \left\{ \begin{array}{l} G(Ax, gy, gy) + G(By, fx, fx), \\ G(By, hz, hz) + G(Cz, gy, gy), \\ G(Cz, fx, fx) + G(Ax, hz, hz) \end{array} \right\}. \quad (2.45)$$

Case 1. If $x, y, z \in [0, 1/2]$, then

$$G(fx, gy, hz) = G\left(1, \frac{10}{11}, \frac{9}{10}\right) = \frac{1}{5},$$

$$G(Ax, gy, gy) + G(By, fx, fx) = G\left(x, \frac{10}{11}, \frac{10}{11}\right) + G(1, 1, 1) = 2\left|x - \frac{10}{11}\right| \geq \frac{9}{11}. \quad (2.46)$$

Thus, we have

$$G(fx, gy, hz) = \frac{1}{5} < \frac{5}{16} \cdot \frac{9}{11} \leq k(G(Ax, gy, gy) + G(By, fx, fx)) \leq kM(x, y, z). \quad (2.47)$$

Case 2. If $x, y \in [0, 1/2]$, $z \in (1/2, 1]$, then

$$\begin{aligned} G(fx, gy, hz) &= G\left(1, \frac{10}{11}, \frac{7}{8}\right) = \frac{1}{4}, \\ G(Ax, gy, gy) + G(By, fx, fx) &= G\left(x, \frac{10}{11}, \frac{10}{11}\right) + G(1, 1, 1) = 2\left|x - \frac{10}{11}\right| \geq \frac{9}{11}. \end{aligned} \quad (2.48)$$

Therefore, we get

$$G(fx, gy, hz) = \frac{1}{4} < \frac{5}{16} \cdot \frac{9}{11} \leq k(G(Ax, gy, gy) + G(By, fx, fx)) \leq kM(x, y, z). \quad (2.49)$$

Case 3. If $x, z \in [0, 1/2]$, $y \in (1/2, 1]$, then

$$\begin{aligned} G(fx, gy, hz) &= G\left(1, \frac{7}{8}, \frac{9}{10}\right) = \frac{1}{4}, \\ G(Cz, fx, fx) + G(Ax, hz, hz) &= G(1, 1, 1) + G\left(x, \frac{9}{10}, \frac{9}{10}\right) = 2\left|x - \frac{9}{10}\right| \geq \frac{4}{5}. \end{aligned} \quad (2.50)$$

Hence, we have

$$G(fx, gy, hz) = \frac{1}{4} = \frac{5}{16} \cdot \frac{4}{5} \leq k(G(Cz, fx, fx) + G(Ax, hz, hz)) \leq kM(x, y, z). \quad (2.51)$$

Case 4. If $y, z \in [0, 1/2]$, $x \in (1/2, 1]$, then

$$\begin{aligned} G(fx, gy, hz) &= G\left(\frac{7}{8}, \frac{10}{11}, \frac{9}{10}\right) = \frac{3}{44}, \\ G(By, hz, hz) + G(Cz, gy, gy) &= G\left(1, \frac{9}{10}, \frac{9}{10}\right) + G\left(1, \frac{10}{11}, \frac{10}{11}\right) = \frac{21}{55}. \end{aligned} \quad (2.52)$$

So we get

$$G(fx, gy, hz) = \frac{3}{44} < \frac{5}{16} \cdot \frac{21}{55} \leq k(G(By, hz, hz) + G(Cz, gy, gy)) \leq kM(x, y, z). \quad (2.53)$$

Case 5. If $x \in [0, 1/2]$, $y, z \in (1/2, 1]$, then

$$G(fx, gy, hz) = G\left(1, \frac{7}{8}, \frac{7}{8}\right) = \frac{1}{4}. \quad (2.54)$$

If $y \in (1/2, 1]$, then

$$G(Ax, gy, gy) + G(By, fx, fx) = G\left(x, \frac{7}{8}, \frac{7}{8}\right) + G\left(\frac{7}{8}, 1, 1\right) = 2\left|x - \frac{7}{8}\right| + \frac{1}{4} \geq \frac{3}{4} + \frac{1}{4} = 1. \quad (2.55)$$

If $y = 1$, then

$$G(Ax, gy, gy) + G(By, fx, fx) = G\left(x, \frac{7}{8}, \frac{7}{8}\right) + G(0, 1, 1) = 2\left|x - \frac{7}{8}\right| + 2 \geq \frac{3}{4} + 2 = \frac{11}{4}. \quad (2.56)$$

And so we have

$$G(Ax, gy, gy) + G(By, fx, fx) \geq 1 \quad (2.57)$$

for all $y \in (1/2, 1]$. Hence we have

$$G(fx, gy, hz) = \frac{1}{4} < \frac{5}{16} \cdot 1 \leq k(G(Ax, gy, gy) + G(By, fx, fx)) \leq kM(x, y, z). \quad (2.58)$$

Case 6. If $y \in [0, 1/2]$, $x, z \in (1/2, 1]$, then

$$\begin{aligned} G(fx, gy, hz) &= G\left(\frac{7}{8}, \frac{10}{11}, \frac{7}{8}\right) = \frac{3}{44}, \\ G(Ax, gy, gy) + G(By, fx, fx) &= G\left(x, \frac{10}{11}, \frac{10}{11}\right) + G\left(1, \frac{7}{8}, \frac{7}{8}\right) = 2\left|x - \frac{10}{11}\right| + \frac{1}{4} \geq \frac{1}{4}. \end{aligned} \quad (2.59)$$

Thus, we have

$$G(fx, gy, hz) = \frac{3}{44} < \frac{5}{16} \cdot \frac{1}{4} \leq k(G(Ax, gy, gy) + G(By, fx, fx)) \leq kM(x, y, z). \quad (2.60)$$

Case 7. If $z \in [0, 1/2]$, $x, y \in (1/2, 1]$, then

$$G(fx, gy, hz) = G\left(\frac{7}{8}, \frac{7}{8}, \frac{9}{10}\right) = \frac{1}{20}. \quad (2.61)$$

If $y \in (1/2, 1)$, then

$$G(By, hz, hz) + G(Cz, gy, gy) = G\left(\frac{7}{8}, \frac{9}{10}, \frac{9}{10}\right) + G\left(1, \frac{7}{8}, \frac{7}{8}\right) = \frac{3}{10}. \quad (2.62)$$

If $y = 1$, then

$$G(By, hz, hz) + G(Cz, gy, gy) = G\left(0, \frac{9}{10}, \frac{9}{10}\right) + G\left(1, \frac{7}{8}, \frac{7}{8}\right) = \frac{41}{20}. \quad (2.63)$$

And so we have

$$G(By, hz, hz) + G(Cz, gy, gy) \geq \frac{3}{10} \quad (2.64)$$

for all $y \in (1/2, 1]$. Hence we have

$$G(fx, gy, hz) = \frac{1}{20} < \frac{5}{16} \cdot \frac{3}{10} \leq k(G(By, hz, hz) + G(Cz, gy, gy)) \leq kM(x, y, z). \quad (2.65)$$

Case 8. If $x, y, z \in (1/2, 1]$, then

$$G(fx, gy, hz) = G\left(\frac{7}{8}, \frac{7}{8}, \frac{7}{8}\right) = 0 \leq \frac{5}{16}M(x, y, z) = kM(x, y, z). \quad (2.66)$$

Then in all the above cases, the mappings f, g, h, A, B , and C are satisfying the condition (2.1) of Theorem 2.1 with $k = 5/16$ so that all the conditions of Theorem 2.1 are satisfied. Moreover, $7/8$ is the unique common fixed point for all of the mappings f, g, h, A, B , and C .

In Theorem 2.1, if we take $A = B = C = I$ (I is identity mapping, the same as below), then we have the following corollary.

Corollary 2.3 (see [2, Theorem 2.4]). *Let (X, G) be a complete G -metric space, and let f, g , and h be three mappings of X into itself satisfying the following conditions:*

$$G(fx, gy, hz) \leq k \max \left\{ \begin{array}{l} G(x, gy, gy) + G(y, fx, fx), \\ G(y, hz, hz) + G(z, gy, gy), \\ G(z, fx, fx) + G(x, hz, hz) \end{array} \right\} \quad (2.67)$$

or

$$G(fx, gy, hz) \leq k \max \left\{ \begin{array}{l} G(x, x, gy) + G(y, y, fx), \\ G(y, y, hz) + G(z, z, gy), \\ G(z, z, fx) + G(x, x, hz) \end{array} \right\} \quad (2.68)$$

for all $x, y, z \in X$, where $k \in [0, 1/3)$. Then f, g , and h have a unique common fixed point in X .

Also, if we take $f = g = h$ and $A = B = C = I$ in Theorem 2.1, then we get the following.

Corollary 2.4 (see [3, Theorem 2.4]). *Let (X, G) be a complete G -metric space, and let f be a mapping of X into itself satisfying the following conditions:*

$$G(fx, fy, fz) \leq k \max \left\{ \begin{array}{l} G(x, fy, fy) + G(y, fx, fx), \\ G(y, fz, fz) + G(z, fy, fy), \\ G(z, fx, fx) + G(x, fz, fz) \end{array} \right\} \quad (2.69)$$

or

$$G(fx, fy, fz) \leq k \max \left\{ \begin{array}{l} G(x, x, fy) + G(y, y, fx), \\ G(y, y, fz) + G(z, z, fy), \\ G(z, z, fx) + G(x, x, fz) \end{array} \right\}, \quad (2.70)$$

for all $x, y, z \in X$, where $k \in [0, 1/3)$. Then, f has a unique fixed point in X .

Remark 2.5. Theorem 2.1 and Corollaries 2.3 and 2.4 generalize and extend the corresponding results of Abbas and Rhoades [5] and Mustafa et al. [6].

Remark 2.6. In Theorem 2.1, if we take: (1) $f = g = h$; (2) $A = B = C$; (3) $g = h$ and $B = C$; (4) $g = h, B = C = I$, several new results can be obtained.

Theorem 2.7. Let (X, G) be a complete G -metric space, and let f, g, h, A, B , and C be six mappings of X into itself satisfying the following conditions:

- (i) $f(X) \subset B(X), g(X) \subset C(X), h(X) \subset A(X)$;
- (ii) the pairs $(f, A), (g, B)$, and (h, C) are commutative mappings;
- (iii) for all $x, y, z \in X$,

$$G(f^m x, g^m y, h^m z) \leq k \max \left\{ \begin{array}{l} G(Ax, g^m y, g^m y) + G(By, f^m x, f^m x), \\ G(By, h^m z, h^m z) + G(Cz, g^m y, g^m y), \\ G(Cz, f^m x, f^m x) + G(Ax, h^m z, h^m z) \end{array} \right\} \quad (2.71)$$

or

$$G(f^m x, g^m y, h^m z) \leq k \max \left\{ \begin{array}{l} G(Ax, Ax, g^m y) + G(By, By, f^m x), \\ G(By, By, h^m z) + G(Cz, Cz, g^m y), \\ G(Cz, Cz, f^m x) + G(Ax, Ax, h^m z) \end{array} \right\}, \quad (2.72)$$

where $k \in [0, 1/2), m \in \mathbb{N}$, then f, g, h, A, B , and C have a unique common fixed point in X .

Proof. Suppose that mappings f, g, h, A, B , and C satisfy the condition (2.71). Since $f^m X \subset f^{m-1} X \subset \dots \subset fX, fX \subset BX$ so that $f^m X \subset BX$. Similarly, we can show that $g^m X \subset CX$ and $h^m X \subset AX$. From the Theorem 2.1, we see that f^m, g^m, h^m, A, B , and C have a unique common fixed point u .

Since $fu = f(f^m u) = f^{m+1} u = f^m(fu)$, so that

$$G(f^m fu, g^m u, h^m u) \leq k \max \left\{ \begin{array}{l} G(Afu, g^m u, g^m u) + G(Bu, f^m fu, f^m fu), \\ G(Bu, h^m u, h^m u) + G(Cu, g^m u, g^m u), \\ G(Cu, f^m fu, f^m fu) + G(Afu, h^m u, h^m u) \end{array} \right\}, \quad (2.73)$$

note that $Afu = fAu = fu$ and the Proposition 1.12(iii), we obtain

$$\begin{aligned} G(fu, u, u) &\leq k \max \left\{ \begin{array}{l} G(fu, u, u) + G(u, fu, fu), \\ G(u, u, u) + G(u, u, u), \\ G(u, fu, fu) + G(fu, u, u) \end{array} \right\} \\ &= k[G(u, fu, fu) + G(fu, u, u)] \\ &\leq 3kG(fu, u, u). \end{aligned} \quad (2.74)$$

Since $k \in [0, 1/3)$, hence $G(fu, u, u) = 0$ and so $fu = u$.

By the same argument, we can prove that $gu = u$ and $hu = u$. Thus, we have $u = fu = gu = hu = Au = Bu = Cu$ so that f, g, h, A, B , and C have a common fixed point u in X . Let v be any other common fixed point of f, g, h, A, B , and C , then by using condition (2.71), we have

$$\begin{aligned} G(u, u, v) &= G(f^m u, g^m u, h^m v) \\ &\leq k \max \left\{ \begin{array}{l} G(Au, g^m u, g^m u) + G(Bu, f^m u, f^m u), \\ G(Bu, h^m v, h^m v) + G(Cv, g^m u, g^m u), \\ G(Cv, f^m u, f^m u) + G(Au, h^m v, h^m v) \end{array} \right\} \\ &= k \max \left\{ \begin{array}{l} G(u, u, u) + G(u, u, u), \\ G(u, v, v) + G(v, u, u), \\ G(v, u, u) + G(u, v, v) \end{array} \right\} \\ &= k[G(u, v, v) + G(v, u, u)] \\ &\leq 3kG(u, u, v). \end{aligned} \quad (2.75)$$

Hence, $G(u, u, v) = 0$ and so $u = v$, since $0 \leq k < 1/3$. Thus, common fixed point is unique.

The proof using (2.72) is similar. This completes the proof. $\square$

In Theorem 2.7, if we take $A = B = C = I$, then we have the following corollary.

Corollary 2.8 (see [2, Corollary 2.5]). *Let (X, G) be a complete G -metric space, and let f, g , and h be three mappings of X into itself satisfying the following conditions:*

$$G(f^m x, g^m y, h^m z) \leq k \max \left\{ \begin{array}{l} G(x, g^m y, g^m y) + G(y, f^m x, f^m x), \\ G(y, h^m z, h^m z) + G(z, g^m y, g^m y), \\ G(z, f^m x, f^m x) + G(x, h^m z, h^m z) \end{array} \right\} \quad (2.76)$$

or

$$G(f^m x, g^m y, h^m z) \leq k \max \left\{ \begin{array}{l} G(x, x, g^m y) + G(y, y, f^m x), \\ G(y, y, h^m z) + G(z, z, g^m y), \\ G(z, z, f^m x) + G(x, x, h^m z) \end{array} \right\} \quad (2.77)$$

for all $x, y, z \in X$, where $k \in [0, 1/3)$, $m \in \mathbb{N}$; then f , g , and h have a unique common fixed point in X .

Also, if we take $f = g = h$ and $A = B = C = I$ in Theorem 2.7, then we get the following.

Corollary 2.9 (see [3, Corollary 2.5]). *Let (X, G) be a complete G -metric space, and let f be a mapping of X into itself satisfying the following conditions:*

$$G(f^m x, f^m y, f^m z) \leq k \max \left\{ \begin{array}{l} G(x, f^m y, f^m y) + G(y, f^m x, f^m x), \\ G(y, f^m z, f^m z) + G(z, f^m y, f^m y), \\ G(z, f^m x, f^m x) + G(x, f^m z, f^m z) \end{array} \right\} \quad (2.78)$$

or

$$G(f^m x, f^m y, f^m z) \leq k \max \left\{ \begin{array}{l} G(x, x, f^m y) + G(y, y, f^m x), \\ G(y, y, f^m z) + G(z, z, f^m y), \\ G(z, z, f^m x) + G(x, x, f^m z) \end{array} \right\} \quad (2.79)$$

for all $x, y, z \in X$, where $k \in [0, 1/3)$, $m \in \mathbb{N}$; then f has a unique fixed point in X .

Remark 2.10. In Theorem 2.7, if we take: (1) $f = g = h$; (2) $g = h$ and $B = C$; (3) $g = h$, $B = C = I$, several new results can be obtained.

Corollary 2.11. *Let (X, G) be a complete G -metric space, and let f, g, h, A, B , and C be six mappings of X into itself satisfying the following conditions:*

- (i) $f(X) \subset B(X)$, $g(X) \subset C(X)$, $h(X) \subset A(X)$;
- (ii) for all $x, y, z \in X$,

$$\begin{aligned} G(fx, gy, hz) &\leq a\{G(Ax, gy, gy) + G(By, fx, fx)\} \\ &\quad + b\{G(By, hz, hz) + G(Cz, gy, gy)\} \\ &\quad + c\{G(Cz, fx, fx) + G(Ax, hz, hz)\} \end{aligned} \quad (2.80)$$

or

$$\begin{aligned} G(fx, gy, hz) &\leq a\{G(Ax, Ax, gy) + G(By, By, fx)\} \\ &\quad + b\{G(By, By, hz) + G(Cz, Cz, gy)\} \\ &\quad + c\{G(Cz, Cz, fx) + G(Ax, Ax, hz)\}, \end{aligned} \quad (2.81)$$

where $0 \leq a+b+c < 1/3$. Then one of the pairs (f, A) , (g, B) , and (h, C) has a coincidence point in X . Further, if one of the following conditions is satisfied, then the mappings f, g, h, A, B , and C have a unique common fixed point in X .

- (a) Either f or A is G -continuous, the pair (f, A) is weakly commutative, the pairs (g, B) and (h, C) are weakly compatible;

- (b) Either g or B is G -continuous, the pair (g, B) is weakly commutative, the pairs (f, A) and (h, C) are weakly compatible;
- (c) Either h or C is G -continuous, the pair (h, C) is weakly commutative, the pairs (f, A) and (g, B) are weakly compatible.

Proof. Suppose that mappings f, g, h, A, B , and C satisfy the condition (2.80). For $x, y, z \in X$, let

$$M(x, y, z) = \max \left\{ \begin{array}{l} G(Ax, gy, gy) + G(By, fx, fx), \\ G(By, hz, hz) + G(Cz, gy, gy), \\ G(Cz, fx, fx) + G(Ax, hz, hz) \end{array} \right\}. \quad (2.82)$$

Then

$$\begin{aligned} & a\{G(Ax, gy, gy) + G(By, fx, fx)\} + b\{G(By, hz, hz) + G(Cz, gy, gy)\} \\ & + c\{G(Cz, fx, fx) + G(Ax, hz, hz)\} \\ & \leq (a + b + c)M(x, y, z). \end{aligned} \quad (2.83)$$

So, if

$$\begin{aligned} G(fx, gy, hz) & \leq a\{G(Ax, gy, gy) + G(By, fx, fx)\} \\ & + b\{G(By, hz, hz) + G(Cz, gy, gy)\} \\ & + c\{G(Cz, fx, fx) + G(Ax, hz, hz)\}, \end{aligned} \quad (2.84)$$

then $G(fx, gy, hz) \leq (a + b + c)M(x, y, z)$. Taking $k = a + b + c$ in Theorem 2.1, the conclusion of Corollary 2.11 can be obtained from Theorem 2.1 immediately.

The proof using (2.81) is similar. This completes the proof. $\square$

Corollary 2.12. Let (X, G) be a complete G -metric space, and let f, g, h, A, B , and C be six mappings of X into itself satisfying the following conditions:

- (i) $f(X) \subset B(X)$, $g(X) \subset C(X)$, $h(X) \subset A(X)$;
- (ii) the pairs (f, A) , (g, B) and (h, C) are commutative mappings;
- (iii) for all $x, y, z \in X$,

$$\begin{aligned} G(f^m x, g^m y, h^m z) & \leq a\{G(Ax, g^m y, g^m y) + G(By, f^m x, f^m x)\} \\ & + b\{G(By, h^m z, h^m z) + G(Cz, g^m y, g^m y)\} \\ & + c\{G(Cz, f^m x, f^m x) + G(Ax, h^m z, h^m z)\} \end{aligned} \quad (2.85)$$

or

$$\begin{aligned} G(f^m x, g^m y, h^m z) \leq & a\{G(Ax, Ax, g^m y) + G(By, By, f^m x)\} \\ & + b\{G(By, By, h^m z) + G(Cz, Cz, g^m y)\} \\ & + c\{G(Cz, Cz, f^m x) + G(Ax, Ax, h^m z)\}, \end{aligned} \quad (2.86)$$

where $0 \leq a + b + c < 1/3$, $m \in \mathbb{N}$; then f, g, h, A, B , and C have a unique common fixed point in X .

Proof. The proof follows from Corollary 2.11, and from an argument similar to that used in Theorem 2.7. $\square$

Remark 2.13. In Corollaries 2.11 and 2.12, if we take: (1) $A = B = C = I$; (2) $f = g = h$; (3) $A = B = C$; (4) $g = h$ and $B = C$; (5) $g = h, B = C = I$, several new results can be obtained.

Acknowledgment

This work was supported by the National Natural Science Foundation of China Grant (nos. 11071169, 11271105) and the Natural Science Foundation of Zhejiang Province (nos. Y6110287, LY12A01030).

References

- [1] Z. Mustafa and B. Sims, "A new approach to generalized metric spaces," *Journal of Nonlinear and Convex Analysis*, vol. 7, no. 2, pp. 289–297, 2006.
- [2] M. Abbas, T. Nazir, and R. Saadati, "Common fixed point results for three maps in generalized metric space," *Advances in Difference Equations*, vol. 49, pp. 1–20, 2011.
- [3] Z. Mustafa and B. Sims, "Fixed point theorems for contractive mappings in complete G-metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 917175, 10 pages, 2009.
- [4] S. Manro, S. S. Bhatia, and S. Kumar, "Expansion mappings theorems in G-metric spaces," *International Journal of Contemporary Mathematical Sciences*, vol. 5, no. 49–52, pp. 2529–2535, 2010.
- [5] M. Abbas and B. E. Rhoades, "Common fixed point results for noncommuting mappings without continuity in generalized metric spaces," *Applied Mathematics and Computation*, vol. 215, no. 1, pp. 262–269, 2009.
- [6] Z. Mustafa, H. Obiedat, and F. Awawdeh, "Some fixed point theorem for mapping on complete G-metric spaces," *Fixed Point Theory and Applications*, vol. 2008, Article ID 189870, 12 pages, 2008.
- [7] Z. Mustafa, W. Shatanawi, and M. Bataineh, "Existence of fixed point results in G-metric spaces," *International Journal of Mathematics and Mathematical Sciences*, vol. 2009, Article ID 283028, 10 pages, 2009.
- [8] W. Shatanawi, "Fixed point theory for contractive mappings satisfying Φ -maps in G-metric spaces," *Fixed Point Theory and Applications*, vol. 2010, Article ID 181650, 9 pages, 2010.
- [9] R. Chugh, T. Kadian, A. Rani, and B. E. Rhoades, "Property P in G-metric spaces," *Fixed Point Theory and Applications*, vol. 2010, Article ID 401684, 12 pages, 2010.
- [10] M. Abbas, T. Nazir, and S. Radenović, "Some periodic point results in generalized metric spaces," *Applied Mathematics and Computation*, vol. 217, no. 8, pp. 4094–4099, 2010.
- [11] Z. Mustafa, F. Awawdeh, and W. Shatanawi, "Fixed point theorem for expansive mappings in G-metric spaces," *International Journal of Contemporary Mathematical Sciences*, vol. 5, no. 49–52, pp. 2463–2472, 2010.
- [12] H. Obiedat and Z. Mustafa, "Fixed point results on a nonsymmetric G-metric spaces," *Jordan Journal of Mathematics and Statistics*, vol. 3, no. 2, pp. 65–79, 2010.

- [13] Z. Mustafa and H. Obiedat, "A fixed point theorem of Reich in G -metric spaces," *CUBO Mathematics Journal*, vol. 12, no. 1, pp. 83–93, 2010.
- [14] M. Abbas, S. H. Khan, and T. Nazir, "Common fixed points of R -weakly commuting maps in generalized metric spaces," *Fixed Point Theory and Applications*, vol. 2011, article 41, no. 1, 11 pages, 2011.
- [15] H. Aydi, W. Shatanawi, and C. Vetro, "On generalized weak G -contraction mapping in G -metric spaces," *Computers & Mathematics with Applications*, vol. 62, no. 11, pp. 4223–4229, 2011.
- [16] M. Abbas, A. R. Khan, and T. Nazir, "Coupled common fixed point results in two generalized metric spaces," *Applied Mathematics and Computation*, vol. 217, no. 13, pp. 6328–6336, 2011.
- [17] W. Shatanawi, M. Abbas, and T. Nazir, "Common coupled coincidence and coupled fixed point results in two generalized metric spaces," *Fixed Point Theory and Applications*, vol. 2011, article 80, 2011.
- [18] R. K. Vats, S. Kumar, and V. Sihag, "Some common fixed point theorems for compatible mappings of type (A) in complete G -metric space," *Advances in Fuzzy Mathematics*, vol. 6, no. 1, pp. 27–38, 2011.
- [19] H. Aydi, "A fixed point result involving a generalized weakly contractive condition in G -metric spaces," *Bulletin of Mathematical Analysis and Applications*, vol. 3, no. 4, pp. 180–188, 2011.
- [20] Z. Mustafa, M. Khandagji, and W. Shatanawi, "Fixed point results on complete G -metric spaces," *Studia Scientiarum Mathematicarum Hungarica*, vol. 48, no. 3, pp. 304–319, 2011.
- [21] W. Shatanawi, "Coupled fixed point theorems in generalized metric spaces," *Haceteppe Journal of Mathematics and Statistics*, vol. 40, no. 3, pp. 441–447, 2011.
- [22] M. Abbas, T. Nazir, and P. Vetro, "Common fixed point results for three maps in G -metric spaces," *Filomat*, vol. 25, no. 4, pp. 1–17, 2011.
- [23] L. Gajić and Z. Lozanov-Crvenković, "A fixed point result for mappings with contractive iterate at a point in G -metric spaces," *Filomat*, vol. 25, no. 2, pp. 53–58, 2011.
- [24] S. Manro, S. Kumar, and S. S. Bhatia, " R -weakly commuting maps in G -metric spaces," *Fasciculi Mathematici*, no. 47, pp. 11–17, 2011.
- [25] R. K. Vats, S. Kumar, and V. Sihag, "Fixed point theorems in complete G -metric space," *Fasciculi Mathematici*, no. 47, pp. 127–139, 2011.
- [26] M. Abbas, T. Nazir, and D. Dorić, "Common fixed point of mappings satisfying $(E.A)$ property in generalized metric spaces," *Applied Mathematics and Computation*, vol. 218, no. 14, pp. 7665–7670, 2012.
- [27] Z. Mustafa, H. Aydi, and E. Karapinar, "On common fixed points in G -metric spaces using $(E.A)$ property," *Computers & Mathematics with Applications*, vol. 64, no. 6, pp. 1944–1956, 2012.
- [28] L. Gajić and M. Stojaković, "On Ćirić generalization of mappings with a contractive iterate at a point in G -metric spaces," *Applied Mathematics and Computation*, vol. 219, no. 1, pp. 435–441, 2012.
- [29] H. Ye and F. Gu, "Common fixed point theorems for a class of twice power type contraction maps in G -metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 736214, 19 pages, 2012.
- [30] W. Long, M. Abbas, T. Nazir, and S. Radenović, "Common fixed point for two pairs of mappings satisfying $(E.A)$ property in generalized metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 394830, 15 pages, 2012.
- [31] A. Kaewcharoen, "Common fixed point theorems for contractive mappings satisfying Φ -maps in G -metric spaces," *Banach Journal of Mathematical Analysis*, vol. 6, no. 1, pp. 101–111, 2012.
- [32] N. Tahat, H. Aydi, E. Karapinar, and W. Shatanawi, "Common fixed points for single-valued and multi-valued maps satisfying a generalized contraction in G -metric spaces," *Fixed Point Theory and Applications*, vol. 2012, article 48, 2012.
- [33] Z. Mustafa, "Some new common fixed point theorems under strict contractive conditions in G -metric spaces," *Journal of Applied Mathematics*, vol. 2012, Article ID 248937, 21 pages, 2012.
- [34] A. Kaewcharoen, "Common fixed points for four mappings in G -metric spaces," *International Journal of Mathematics Analysis*, vol. 6, no. 47, pp. 2345–2356, 2012.
- [35] V. Popa and A.-M. Patriciu, "A general fixed point theorem for pairs of weakly compatible mappings in G -metric spaces," *Journal of Nonlinear Science and Its Applications*, vol. 5, no. 2, pp. 151–160, 2012.
- [36] M. Khandagji, S. Al-Sharif, and M. Al-Khaleel, "Property P and some fixed point results on (ψ, ϕ) -weakly contractive G -metric spaces," *International Journal of Mathematics and Mathematical Sciences*, vol. 2012, Article ID 675094, 11 pages, 2012.
- [37] Z. Mustafa, "Common fixed points of weakly compatible mappings in G -metric spaces," *Applied Mathematical Sciences*, vol. 6, no. 92, pp. 4589–4600, 2012.
- [38] M. Aggarwal, R. Chugh, and R. Kamal, "Suzuki-type fixed point results in G -metric spaces and applications," *International Journal of Computer Applications*, vol. 47, no. 12, pp. 14–17, 2012.
- [39] M. Gugnani, M. Aggarwal, and R. Chugh, "Common fixed point results in G -metric spaces and applications," *International Journal of Computer Applications*, vol. 43, no. 11, pp. 38–42, 2012.

- [40] S. K. Mohanta, "Some fixed point theorems in G -metric spaces," *Analele Stiintifice ale Universitatii Ovidius Constanta*, vol. 20, no. 1, pp. 285–306, 2012.
- [41] H. Aydi, E. Karapinar, and W. Shatanawi, "Tripled fixed point results in generalized metric spaces," *Journal of Applied Mathematics*, vol. 2012, Article ID 314279, 10 pages, 2012.
- [42] H. Aydi, "A common fixed point of integral type contraction in generalized metric spaces," *Journal of Advanced Mathematical Studies*, vol. 5, no. 1, pp. 111–117, 2012.
- [43] K. P. R. Rao, K. B. Lakshmi, Z. Mustafa, and V. C. C. Raju, "Fixed and related fixed point theorems for three maps in G -metric spaces," *Journal of Advanced Studies in Topology*, vol. 3, no. 4, pp. 12–19, 2012.
- [44] R. Saadati, S. M. Vaezpour, P. Vetro, and B. E. Rhoades, "Fixed point theorems in generalized partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 52, no. 5-6, pp. 797–801, 2010.
- [45] B. S. Choudhury and P. Maity, "Coupled fixed point results in generalized partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 54, no. 1-2, pp. 73–79, 2011.
- [46] H. Aydi, B. Damjanović, B. Samet, and W. Shatanawi, "Coupled fixed point theorems for nonlinear contractions in partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 54, no. 9-10, pp. 2443–2450, 2011.
- [47] W. Shatanawi, "Some fixed point theorems in ordered G -metric spaces and applications," *Abstract and Applied Analysis*, vol. 2011, Article ID 126205, 11 pages, 2011.
- [48] N. V. Luong and N. X. Thuan, "Coupled fixed point theorems in partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 55, no. 3-4, pp. 1601–1609, 2012.
- [49] H. Aydi, M. Postolache, and W. Shatanawi, "Coupled fixed point results for (ψ, φ) -weakly contractive mappings in ordered G -metric spaces," *Computers & Mathematics with Applications*, vol. 63, no. 1, pp. 298–309, 2012.
- [50] M. Abbas, T. Nazir, and S. Radenović, "Common fixed point of generalized weakly contractive maps in partially ordered G -metric spaces," *Applied Mathematics and Computation*, vol. 218, no. 18, pp. 9383–9395, 2012.
- [51] Y. J. Cho, B. E. Rhoades, R. Saadati, B. Samet, and W. Shatanawi, "Nonlinear coupled fixed point theorems in ordered generalized metric spaces with integral type," *Fixed Point Theory and Applications*, vol. 2012, article 8, 2012.
- [52] H. Aydi, E. Karapinar, and W. Shatanawi, "Tripled common fixed point results for generalized contractions in ordered generalized metric spaces," *Fixed Point Theory and Applications*, vol. 2012, article 101, 2012.
- [53] M. Abbas, W. Sintunavarat, and P. Kumam, "Coupled fixed point of generalized contractive mappings on partially ordered G -metric spaces," *Fixed Point Theory and Applications*, vol. 2012, article 31, 2012.
- [54] H. Kumar Nashine, "Coupled common fixed point results in ordered G -metric spaces," *Journal of Nonlinear Science and Its Applications*, vol. 5, no. 1, pp. 1–13, 2012.
- [55] Z. Mustafa, H. Aydi, and E. Karapinar, "Mixed g -monotone property and quadruple fixed point theorems in partially ordered metric space," *Fixed Point Theory and Applications*, vol. 2012, article 71, 2012.
- [56] W. Shatanawi, M. Abbas, H. Aydi, and N. Tahat, "Common coupled coincidence and coupled fixed points in G -metric spaces," *Nonlinear Analysis and Application*, vol. 2012, Article ID jnaa-00162, 16 pages, 2012.