Research Article

Identities Involving q -Bernoulli and q -Euler Numbers

D. S. Kim,¹ T. Kim,² J. Choi,³ and Y. H. Kim³

¹ Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

² Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

³ Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to T. Kim, tkkim@kw.ac.kr

Received 18 January 2012; Revised 24 February 2012; Accepted 24 February 2012

Academic Editor: Toka Diagana

Copyright © 2012 D. S. Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We give some identities on the q -Bernoulli and q -Euler numbers by using p -adic integral equations on $\mathbb{Z}_p$.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will denote the ring of p -adic integers, the field of p -adic rational numbers, and the completion of algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. The p -adic norm $|\cdot|_p$ is normally defined by $|p|_p = 1/p$.

As it is well known, the Euler polynomials are defined by

$$\frac{2}{e^t + 1} e^{xt} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (1.1)$$

with the usual convention about replacing $E^n(x)$ by $E_n(x)$ (see [1–14]). In the special case, $x = 0$, $E_n(0) = E_n$ is called the n th Euler number.

The ordinary Bernoulli polynomials are also defined by

$$\frac{t}{e^t - 1} e^{xt} = e^{B(x)t} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}, \quad (1.2)$$

with the usual convention about replacing $B^n(x)$ by $B_n(x)$ (see [1–14]). In the special case, $x = 0$, $B_n(0) = B_n$ is called the n th Bernoulli number.

Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the bosonic p -adic integral on $\mathbb{Z}_p$ is defined by

$$I(f) = \int_{\mathbb{Z}_p} f(x) d\mu(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \mu(x + p^N \mathbb{Z}_p) = \lim_{N \rightarrow \infty} \frac{1}{p^N} \sum_{x=0}^{p^N-1} f(x), \quad (1.3)$$

(see [1, 7]). Let f_1 be the translation of f with $f_1(x) = f(x+1)$. From (1.3) we have

$$I(f_1) - I(f) = f'(0), \quad (1.4)$$

(see [1, 7]).

The fermionic p -adic integral on $\mathbb{Z}_p$ is also defined by T. Kim as follows:

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \mu_{-1}(x + p^N \mathbb{Z}_p) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x, \quad (1.5)$$

(see [6, 15, 16]). By (1.5), we get

$$I_{-1}(f_1) + I_{-1}(f) = 2f(0), \quad (1.6)$$

(see [6, 8]).

Let $f(x) = e^{xt} \in UD(\mathbb{Z}_p)$ with $|t|_p < |p|_p^{1/(p-1)}$ and $|x|_p \leq 1$. From (1.4), we have

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu(y) = \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}. \quad (1.7)$$

Thus, by (1.7), we see that

$$\int_{\mathbb{Z}_p} (x+y)^n d\mu(y) = B_n(x), \quad n \in \mathbb{Z}_+. \quad (1.8)$$

By (1.8), we get

$$B_n(x) = \sum_{l=0}^n \binom{n}{l} x^{n-l} \int_{\mathbb{Z}_p} y^l d\mu(y) = \sum_{l=0}^n \binom{n}{l} x^{n-l} B_l. \quad (1.9)$$

As an indeterminate, let us assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$.

From (1.1) and (1.6), we note that the q -Euler polynomials are given by

$$\int_{\mathbb{Z}_p} q^y e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{qe^t + 1} e^{xt} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}, \quad (1.10)$$

where $E_{n,q}(x)$ are called the n th q -Euler polynomials (see [1, 3, 6, 8]).

Thus, by (1.10), we get

$$E_{n,q}(x) = \int_{\mathbb{Z}_p} q^y (x+y)^n d\mu_{-1}(y), \quad n \in \mathbb{Z}_+. \quad (1.11)$$

In the special case, $x = 0$, $E_{n,q}(0) = E_{n,q}$ is called the n th q -Euler number (see [8, 9]). By (1.10) and (1.11), we get the recurrence formula for the q -Euler numbers as follows:

$$E_{0,q} = \frac{2}{[2]_q}, \quad q(E_q + 1)^n + E_{n,q} = 2\delta_{0,n}, \quad (1.12)$$

with the usual convention about replacing E_q^n by $E_{n,q}$. Here $[x]_q = (1 - q^x)/(1 - q)$ is the q -number of x and $\delta_{k,n}$ is the Kronecker symbol (see [10, 11]).

From (1.2), (1.7), and (1.8), we have

$$B_0 = 1, \quad (B + 1)^n - B_n = \delta_{1,n}, \quad (1.13)$$

with the usual convention about replacing B^n by B_n (see [1, 3, 14]).

From (1.11), we easily see that

$$E_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} x^{n-l} E_{l,q}, \quad (1.14)$$

(see [14]).

In this paper we give some interesting properties of p -adic integrals on $\mathbb{Z}_p$ associated with the q -Bernoulli and the q -Euler numbers. From those properties, we derive new identities involving the q -Bernoulli and the q -Euler numbers arising from p -adic integrals of polynomial identities.

2. Identities on q -Bernoulli and q -Euler Numbers

Let C_{p^n} be the cyclic group of order p^n with $C_{p^n} = \{\zeta \in \mathbb{C}_p \mid \zeta^{p^n} = 1\}$. Then T_p is defined by the direct limit as $T_p = \lim_{n \rightarrow \infty} C_{p^n} = C_{p^\infty}$. In this section, we assume that $q (\neq 1) \in T_p$, then $|q - 1|_p < 1$. From (1.4), we can derive the following equation (2.1):

$$\int_{\mathbb{Z}_p} q^y e^{(x+y)t} d\mu(y) = \frac{t}{qe^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_{n,q}(x) \frac{t^n}{n!}, \quad (2.1)$$

where $B_{n,q}(x)$ is called the n th q -Bernoulli polynomial (see [7]). In the special case, $x = 0$, $B_{n,q}(0) = B_{n,q}$ is called the n th q -Bernoulli number.

By (2.1), we get

$$B_{0,q} = 0, \quad q(B_q + 1)^n - B_{n,q} = \delta_{1,n}, \quad (2.2)$$

with the usual convention about replacing B_q^n by $B_{n,q}$ (see [7, 14]).

From (1.3), we have

$$\int_{\mathbb{Z}_p} f(-y) d\mu(y) = \int_{\mathbb{Z}_p} f(y+1) d\mu(y). \quad (2.3)$$

By (2.3), we get

$$q \int_{\mathbb{Z}_p} (1-x+y)^n q^y d\mu(y) = (-1)^n \int_{\mathbb{Z}_p} q^{-y} (x+y)^n d\mu(y). \quad (2.4)$$

From (2.1) and (2.4), we have

$$qB_{n,q}(1-x) = (-1)^n B_{n,q^{-1}}(x), \quad n \in \mathbb{Z}_+. \quad (2.5)$$

By using (1.4), we see that

$$q \int_{\mathbb{Z}_p} (1+x+y)^n q^y d\mu(y) = \int_{\mathbb{Z}_p} q^y (x+y)^n d\mu(y) + nx^{n-1}. \quad (2.6)$$

Thus, by (2.1) and (2.6), we get

$$qB_{n,q}(1+x) = B_{n,q}(x) + nx^{n-1}, \quad n \in \mathbb{Z}_+. \quad (2.7)$$

Therefore, by (2.5) and (2.7), we obtain the following theorem.

Theorem 2.1. *For $n \in \mathbb{Z}_+$, one has*

$$(-1)^n B_{n,q^{-1}}(-x) = qB_{n,q}(1+x) = B_{n,q}(x) + nx^{n-1}. \quad (2.8)$$

From (1.5) and (1.6), we note that

$$\int_{\mathbb{Z}_p} (1-x+y)^n q^y d\mu_{-1}(y) = (-1)^n \int_{\mathbb{Z}_p} (x+y)^n q^{-(y+1)} d\mu_{-1}(y). \quad (2.9)$$

Therefore, by (1.11) and (2.9), we obtain the following theorem.

Theorem 2.2. *For $n \in \mathbb{Z}_+$, one has*

$$qE_{n,q}(1-x) = (-1)^n E_{n,q^{-1}}(x). \quad (2.10)$$

By (1.6), we get

$$q \int_{\mathbb{Z}_p} q^y (x+1+y)^n d\mu_{-1}(y) + \int_{\mathbb{Z}_p} q^y (x+y)^n d\mu_{-1}(y) = 2x^n. \quad (2.11)$$

Thus, by (1.11) and (2.11), we have

$$qE_{n,q}(x+1) = -E_{n,q}(x) + 2x^n, \quad n \in \mathbb{Z}_+. \quad (2.12)$$

Therefore, by Theorem 2.2 and (2.12), we obtain the following theorem.

Theorem 2.3. *For $n \in \mathbb{Z}_+$, one has*

$$(-1)^n E_{n,q^{-1}}(-x) = qE_{n,q}(1+x) = -E_{n,q}(x) + 2x^n. \quad (2.13)$$

By using the p -adic integrals on $\mathbb{Z}_p$, we have the following equation (2.14):

$$\begin{aligned} \int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} q^{(x+y)} e^{(x+y)t} d\mu(x) d\mu_{-1}(y) &= \int_{\mathbb{Z}_p} q^x e^{xt} d\mu(x) \int_{\mathbb{Z}_p} q^y e^{yt} d\mu_{-1}(y) \\ &= \left(\frac{t}{qe^t - 1} \right) \left(\frac{2}{qe^t + 1} \right) = \frac{2t}{q^2 e^{2t} - 1} \\ &= \int_{\mathbb{Z}_p} q^{2x} e^{2xt} d\mu(x). \end{aligned} \quad (2.14)$$

By (2.14), we get

$$\int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} q^{(x+y)} (x+y)^n d\mu(x) d\mu_{-1}(y) = 2^n \int_{\mathbb{Z}_p} q^{2x} x^n d\mu(x). \quad (2.15)$$

It is not difficult to show that

$$\int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} q^{(x+y)} (x+y)^n d\mu(x) d\mu_{-1}(y) = \sum_{l=0}^n \binom{n}{l} \int_{\mathbb{Z}_p} \int_{\mathbb{Z}_p} q^x x^{n-l} q^y y^l d\mu(x) d\mu_{-1}(y). \quad (2.16)$$

Therefore, by (2.15) and (2.16), we obtain the following theorem.

Theorem 2.4. *For $n \in \mathbb{Z}_+$, one has*

$$\sum_{l=0}^n \binom{n}{l} B_{n-l,q} E_{l,q} = 2^n B_{n,q^2}. \quad (2.17)$$

By (2.5), (2.7), (2.12), Theorems 2.1, and 2.3, we get

$$qB_{n,q}(x) = (-1)^n B_{n,q^{-1}}(1-x) = B_{n,q}(x-1) + n(x-1)^{n-1}, \quad (2.18)$$

$$qE_{n,q}(x) = (-1)^n E_{n,q^{-1}}(1-x) = -E_{n,q}(x-1) + 2(x-1)^n. \quad (2.19)$$

From (2.1), we have

$$B_{n,q}(x) = \int_{\mathbb{Z}_p} q^y (x+y)^n d\mu(y) = \sum_{l=0}^n \binom{n}{l} B_{l,q} x^{n-l}, \quad n \in \mathbb{Z}_+. \quad (2.20)$$

Let

$$I_1 = q \int_{\mathbb{Z}_p} q^x B_{n,q}(x) d\mu(x) = q \sum_{l=0}^n \binom{n}{l} B_{n-l,q} B_{l,q}. \quad (2.21)$$

From (2.18), (2.20), and (2.21), we note that

$$\begin{aligned} I_1 &= (-1)^n \sum_{l=0}^n \binom{n}{l} B_{n-l,q} \int_{\mathbb{Z}_p} q^x (x-1)^l d\mu(x) + n \int_{\mathbb{Z}_p} q^x (x-1)^{n-1} d\mu(x) \\ &= (-1)^n \sum_{l=0}^n \binom{n}{l} B_{n-l,q} B_{l,q}(-1) + n B_{n-1,q}(-1) \\ &= (-1)^n \sum_{l=0}^n \binom{n}{l} B_{l,q} B_{n-l,q}(-1) + n B_{n-1,q}(-1). \end{aligned} \quad (2.22)$$

By (2.5), we get

$$q B_{n,q}(-1) = (-1)^n B_{n,q^{-1}}(2), \quad n \in \mathbb{Z}_+. \quad (2.23)$$

By (2.3), we easily see that

$$q^2 B_{n,q}(2) = nq + q B_{n,q}(1) = nq + B_{n,q} + \delta_{1,n}, \quad (2.24)$$

where $\delta_{1,n}$ is the Kronecker symbol.

Thus, by (2.23) and (2.24), we get

$$B_{n,q}(-1) = (-1)^n (n + q B_{n,q^{-1}} + q \delta_{1,n}). \quad (2.25)$$

By (2.22) and (2.25), we get

$$\begin{aligned} I_1 &= (-1)^n \sum_{l=0}^n \binom{n}{l} B_{l,q}(-1)^{n-l} (n-l + q B_{n-l,q^{-1}} + q \delta_{1,n-l}) + n B_{n-1,q}(-1) \\ &= n \sum_{l=0}^{n-1} \binom{n-1}{l} B_{l,q}(-1)^l + q \sum_{l=0}^n \binom{n}{l} (-1)^l B_{l,q} B_{n-l,q^{-1}} + q(-1)^{n-1} n B_{n-1,q} + n B_{n-1,q}(-1) \\ &= q \sum_{l=0}^n \binom{n}{l} (-1)^l B_{l,q} B_{n-l,q^{-1}} + q(-1)^{n-1} n B_{n-1,q} + (1 + (-1)^{n-1}) n B_{n-1,q}(-1). \end{aligned} \quad (2.26)$$

Therefore, by (2.21) and (2.26), we obtain the following theorem.

Theorem 2.5. For $n \in \mathbb{Z}_+$, one has

$$\sum_{l=0}^{2n} \binom{2n}{l} B_{2n-l,q} B_{l,q} - \sum_{l=0}^{2n} \binom{2n}{l} (-1)^l B_{l,q} B_{2n-l,q^{-1}} = -2n B_{2n-1,q}. \quad (2.27)$$

Let us consider the following integral:

$$I_2 = \int_{\mathbb{Z}_p} q^{x+1} E_{n,q}(x) d\mu_{-1}(x) = q \sum_{l=0}^n \binom{n}{l} E_{n-l,q} E_{l,q}. \quad (2.28)$$

By (2.19), we get

$$\begin{aligned} I_2 &= - \int_{\mathbb{Z}_p} q^x E_{n,q}(x-1) d\mu_{-1}(x) + 2 \int_{\mathbb{Z}_p} q^x (x-1)^n d\mu_{-1}(x) \\ &= - \sum_{l=0}^n \binom{n}{l} E_{l,q} E_{n-l,q}(-1) + 2E_{n,q}(-1) \\ &= - \sum_{l=0}^n \binom{n}{l} E_{n-l,q} E_{l,q}(-1) + 2E_{n,q}(-1). \end{aligned} \quad (2.29)$$

From Theorem 2.2, we note that

$$qE_{n,q}(-1) = (-1)^n E_{n,q^{-1}}(2), \quad n \in \mathbb{Z}_+. \quad (2.30)$$

By (1.12), we get

$$q^2 E_{n,q}(2) = 2q + E_{n,q} - 2\delta_{0,n}. \quad (2.31)$$

Thus, by (2.30) and (2.31), we get

$$E_{n,q}(-1) = (-1)^n (2 + qE_{n,q^{-1}} - 2q\delta_{0,n}). \quad (2.32)$$

From (2.29) and (2.32), we note that

$$\begin{aligned} I_2 &= - \sum_{l=0}^n \binom{n}{l} E_{l,q} (-1)^{n-l} (2 + qE_{n-l,q^{-1}} - 2q\delta_{0,n-l}) + 2(-1)^n (2 + qE_{n,q^{-1}} - 2q\delta_{0,n}) \\ &= -q \sum_{l=0}^n \binom{n}{l} (-1)^{n-l} E_{l,q} E_{n-l,q^{-1}} + 2qE_{n,q}. \end{aligned} \quad (2.33)$$

Therefore, by (2.28) and (2.33), we obtain the following theorem.

Theorem 2.6. For $n \in \mathbb{Z}_+$, one has

$$\sum_{l=0}^n \binom{n}{l} E_{n-l,q} E_{l,q} + \sum_{l=0}^n \binom{n}{l} (-1)^{n-l} E_{l,q} E_{n-l,q^{-1}} = 2E_{n,q}. \quad (2.34)$$

Now we consider the fermionic p -adic integral on $\mathbb{Z}_p$ for the n th q -Euler polynomials as follows:

$$\begin{aligned} I_3 &= \int_{\mathbb{Z}_p} q^x E_{n,q}(x) d\mu_{-1}(x) = \sum_{l=0}^n \binom{n}{l} E_{l,q} \int_{\mathbb{Z}_p} q^x x^{n-l} d\mu_{-1}(x) \\ &= \sum_{l=0}^n \binom{n}{l} E_{l,q} E_{n-l,q}, \quad n \in \mathbb{Z}_+. \end{aligned} \quad (2.35)$$

On the other hand, by Theorem 2.2, we get

$$\begin{aligned} I_3 &= (-1)^n q^{-1} \int_{\mathbb{Z}_p} E_{n,q^{-1}}(1-x) q^x d\mu_{-1}(x) \\ &= (-1)^n q^{-1} \sum_{l=0}^n \binom{n}{l} E_{n-l,q^{-1}} \int_{\mathbb{Z}_p} q^x (1-x)^l d\mu_{-1}(x) \\ &= q^{-1} \sum_{l=0}^n \binom{n}{l} E_{n-l,q^{-1}} (-1)^{n-l} E_{l,q}(-1). \end{aligned} \quad (2.36)$$

From (2.32) and (2.36), we note that

$$\begin{aligned} I_3 &= (-1)^n q^{-1} \sum_{l=0}^n \binom{n}{l} E_{n-l,q^{-1}} (2 + qE_{l,q^{-1}} - 2q\delta_{0,l}) \\ &= 2(-1)^n q^{-1} E_{n,q^{-1}}(1) + (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l,q^{-1}} E_{l,q^{-1}} - 2(-1)^n E_{n,q^{-1}} \\ &= -2(-1)^n E_{n,q^{-1}} + 4(-1)^n \delta_{0,n} + (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l,q^{-1}} E_{l,q^{-1}} - 2(-1)^n E_{n,q^{-1}}. \end{aligned} \quad (2.37)$$

Therefore, by (2.35) and (2.37), we obtain the following theorem.

Theorem 2.7. *For $n \in \mathbb{Z}_+$, one has*

$$\sum_{l=0}^{2n+1} \binom{2n+1}{l} E_{l,q} E_{2n+1-l,q} + \sum_{l=0}^{2n+1} \binom{2n+1}{l} E_{l,q^{-1}} E_{2n+1-l,q^{-1}} = 4E_{2n+1,q^{-1}}. \quad (2.38)$$

From (2.1) and (2.7), we note that

$$x^n = \frac{q}{n+1} \sum_{l=0}^n \binom{n+1}{l} B_{l,q}(x) + \frac{q-1}{n+1} B_{n+1,q}(x). \quad (2.39)$$

Let us consider the following fermionic p -adic integral on $\mathbb{Z}_p$:

$$\begin{aligned} \int_{\mathbb{Z}_p} x^n q^x d\mu_{-1}(x) &= \frac{q}{n+1} \sum_{l=0}^n \binom{n+1}{l} \int_{\mathbb{Z}_p} q^x B_{l,q}(x) d\mu_{-1}(x) + \frac{q-1}{n+1} \int_{\mathbb{Z}_p} q^x B_{n+1,q}(x) d\mu_{-1}(x) \\ &= \frac{q}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k,q} E_{k,q} + \frac{q-1}{n+1} \sum_{l=0}^{n+1} \binom{n+1}{l} B_{n+1-l,q} E_{l,q}. \end{aligned} \quad (2.40)$$

Therefore, by (2.40), we obtain the following theorem.

Theorem 2.8. For $n \in \mathbb{Z}_+$, one has

$$E_{n,q} = \frac{q}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k,q} E_{k,q} + \frac{q-1}{n+1} \sum_{l=0}^{n+1} \binom{n+1}{l} B_{n+1-l,q} E_{l,q}. \quad (2.41)$$

From (1.10) and (2.12), we note that

$$x^n = \frac{[2]_q}{2} E_{n,q}(x) + \frac{q}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_{l,q}(x). \quad (2.42)$$

Thus, by (2.42), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} q^x x^n d\mu_{-1}(x) &= \frac{[2]_q}{2} \int_{\mathbb{Z}_p} E_{n,q}(x) q^x d\mu_{-1}(x) + \frac{q}{2} \sum_{l=0}^{n-1} \binom{n}{l} \int_{\mathbb{Z}_p} q^x E_{l,q}(x) d\mu_{-1}(x) \\ &= \frac{[2]_q}{2} \sum_{l=0}^n \binom{n}{l} E_{n-l,q} E_{l,q} + \frac{q}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k,q} E_{k,q}. \end{aligned} \quad (2.43)$$

Thus, by (2.43), we have

$$E_{n,q} = \frac{[2]_q}{2} \sum_{l=0}^n \binom{n}{l} E_{n-l,q} E_{l,q} + \frac{q}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k,q} E_{k,q}. \quad (2.44)$$

Acknowledgments

The authors would like to express their sincere gratitude to referees for their valuable suggestions and comments. The first author was supported by National Research Foundation of Korea Grant funded by the Korean Government 2011-0002486.

References

- [1] A. Bayad and T. Kim, "Identities involving values of Bernstein, q -Bernoulli, and q -Euler polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 2, pp. 133–143, 2011.

- [2] A. Bayad, "Modular properties of elliptic Bernoulli and Euler functions," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 389–401, 2010.
- [3] A. Bayad, T. Kim, B. Lee, and S.-H. Rim, "Some identities on Bernstein polynomials associated with q -Euler polynomials," *Abstract and Applied Analysis*, vol. 2011, Article ID 294715, 10 pages, 2011.
- [4] L. Carlitz, "The product of two Eulerian polynomials," *Mathematics Magazine*, vol. 36, no. 1, pp. 37–41, 1963.
- [5] L. C. Jang, "A note on Nörlund-type twisted q -Euler polynomials and numbers of higher order associated with fermionic invariant q -integrals," *Journal of Inequalities and Applications*, vol. 2010, Article ID 417452, 12 pages, 2010.
- [6] T. Kim, "Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [7] T. Kim, "An analogue of Bernoulli numbers and their congruences," *Reports of the Faculty of Science and Engineering. Saga University. Mathematics*, vol. 22, no. 2, pp. 21–26, 1994.
- [8] T. Kim, "New approach to q -Euler polynomials of higher order," *Russian Journal of Mathematical Physics*, vol. 17, no. 2, pp. 218–225, 2010.
- [9] Y.-H. Kim, K.-W. Hwang, and T. Kim, "Interpolation functions of the q -Genocchi and the q -Euler polynomials of higher order," *Journal of Computational Analysis and Applications*, vol. 12, no. 1-B, pp. 228–238, 2010.
- [10] H. Ozden and Y. Simsek, "A new extension of q -Euler numbers and polynomials related to their interpolation functions," *Applied Mathematics Letters*, vol. 21, no. 9, pp. 934–939, 2008.
- [11] H. Ozden, I. N. Cangul, and Y. Simsek, "Multivariate interpolation functions of higher-order q -Euler numbers and their applications," *Abstract and Applied Analysis*, vol. 2008, Article ID 390857, 16 pages, 2008.
- [12] K.-H. Park, Y.-H. Kim, and T. Kim, "A note on the generalized q -Euler numbers (2)," *Journal of Computational Analysis and Applications*, vol. 12, no. 3, pp. 630–636, 2010.
- [13] C. S. Ryoo, "Some identities of the twisted q -Euler numbers and polynomials associated with q -Bernstein polynomials," *Proceedings of the Jangjeon Mathematical Society*, vol. 14, no. 2, pp. 239–248, 2011.
- [14] Y. Simsek, "On p -adic twisted p - L -functions related to generalized twisted Bernoulli numbers," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 340–348, 2006.
- [15] T. Kim, "A note on q -Volkenborn integration," *Proceedings of the Jangjeon Mathematical Society*, vol. 8, no. 1, pp. 13–17, 2005.
- [16] T. Kim, " q -Euler numbers and polynomials associated with p -adic q -integrals," *Journal of Nonlinear Mathematical Physics*, vol. 14, no. 1, pp. 15–27, 2007.