Research Article

Eigenvalue Problem of Nonlinear Semipositone Higher Order Fractional Differential Equations

Jing Wu¹ and Xinguang Zhang²

¹ School of Economic Mathematics, Southwestern University of Finance and Economics,
 Chengdu 610074, Sichuan, China

² School of Mathematical and Informational Sciences, Yantai University, Yantai 264005, Shandong, China

Correspondence should be addressed to Jing Wu, wujing8119@163.com
 and Xinguang Zhang, zxcg123242@sina.com

Received 22 October 2012; Accepted 19 November 2012

Academic Editor: Dragoş-Pătru Covei

Copyright © 2012 J. Wu and X. Zhang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the eigenvalue interval for the existence of positive solutions to a semipositone higher order fractional differential equation $-\mathfrak{D}_t^\mu x(t) = \lambda f(t, x(t), \mathfrak{D}_t^{\mu_1} x(t), \mathfrak{D}_t^{\mu_2} x(t), \dots, \mathfrak{D}_t^{\mu_{n-1}} x(t))$, $\mathfrak{D}_t^{\mu_i} x(0) = 0, 1 \leq i \leq n-1, \mathfrak{D}_t^{\mu_{n-1}+1} x(0) = 0, \mathfrak{D}_t^{\mu_{n-1}} x(1) = \sum_{j=1}^{m-2} a_j \mathfrak{D}_t^{\mu_{n-1}} x(\xi_j)$, where $n-1 < \mu \leq n, n \geq 3, 0 < \mu_1 < \mu_2 < \dots < \mu_{n-2} < \mu_{n-1}, n-3 < \mu_{n-1} < \mu-2, a_j \in \mathbb{R}, 0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$ satisfying $0 < \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1} < 1$, $\mathfrak{D}_t^\mu$ is the standard Riemann-Liouville derivative, $f \in C((0,1) \times \mathbb{R}^n, (-\infty, +\infty))$, and f is allowed to be changing-sign. By using reducing order method, the eigenvalue interval of existence for positive solutions is obtained.

1. Introduction

In this paper, we consider the eigenvalue interval for existence of positive solutions to the following semipositone higher order fractional differential equation:

$$\begin{aligned} -\mathfrak{D}_t^\mu x(t) &= \lambda f(t, x(t), \mathfrak{D}_t^{\mu_1} x(t), \mathfrak{D}_t^{\mu_2} x(t), \dots, \mathfrak{D}_t^{\mu_{n-1}} x(t)), \\ \mathfrak{D}_t^{\mu_i} x(0) &= 0, \quad 1 \leq i \leq n-1, \quad \mathfrak{D}_t^{\mu_{n-1}+1} x(0) = 0, \\ \mathfrak{D}_t^{\mu_{n-1}} x(1) &= \sum_{j=1}^{m-2} a_j \mathfrak{D}_t^{\mu_{n-1}} x(\xi_j), \end{aligned} \tag{1.1}$$

where $n - 1 < \mu \leq n$, $n \geq 3$, $n \in \mathbb{N}$, $0 < \mu_1 < \mu_2 < \cdots < \mu_{n-2} < \mu_{n-1}$, and $n - 3 < \mu_{n-1} < \mu - 2$, $a_j \in \mathbb{R}$, $0 < \xi_1 < \xi_2 < \cdots < \xi_{m-2} < 1$ satisfying $0 < \sum_{j=1}^{m-2} a_j \xi_j^{\mu - \mu_{n-1} - 1} < 1$, $\mathfrak{D}_t^\mu$ is the standard Riemann-Liouville derivative, $f : (0, 1) \times \mathbb{R}^n \rightarrow (-\infty, +\infty)$ is continuous.

Recently, one has found that fractional models can sufficiently describe the operation of variety of computational, economic mathematics, physical, and biological processes and systems, see [1–9]. Accordingly, considerable attention has been paid to the solution of fractional differential equations, integral equations, and fractional partial differential equations of physical phenomena [10–24]. One of the most frequently used tools in the theory of fractional calculus is furnished by the Riemann-Liouville operators. It possesses advantages of fast convergence, higher stability and higher accuracy to derive the solution of different types of fractional equations.

In this work, we will deal with the eigenvalue interval for existence of positive solutions to the higher order fractional differential equation when f may be negative. This type of differential equation is called semipositone problem which arises in many interesting applications as pointed out by Lions in [25]. For example, the semipositone differential equation which can be derived from chemical reactor theory, design of suspension bridges, combustion, and management of natural resources, see [26–28]. To our knowledge, few results were established, especially for higher order multipoint boundary value problems with the fractional derivatives.

2. Preliminaries and Lemmas

We use the following assumptions in this paper:

(B1) $f : (0, 1) \times \mathbb{R}^n \rightarrow (-\infty, +\infty)$ is continuous, and there exist functions $\alpha, \beta \in L^1[(0, 1), (0, +\infty)]$ and continuous function $h : \mathbb{R}^n \rightarrow [0, +\infty)$ such that

$$-\alpha(t) \leq f(t, x_1, x_2, \dots, x_n) \leq \beta(t)h(x_1, x_2, \dots, x_n), (t, x_1, x_2, \dots, x_n) \in (0, 1) \times \mathbb{R}^n. \quad (2.1)$$

Now we begin this section with some preliminaries of fractional calculus. Let $\mu > 0$ and $n = [\mu] + 1 = N + 1$, where N is the smallest integer greater than or equal to μ . For a function $x : (0, 1) \rightarrow \mathbb{R}$, we define the fractional integral of order μ of x as

$$I^\mu x(t) = \frac{1}{\Gamma(\mu)} \int_0^t (t-s)^{\mu-1} x(s) ds \quad (2.2)$$

provided the integral exists. The fractional derivative of order μ of a continuous function x is defined by

$$\mathfrak{D}_t^\mu x(t) = \frac{1}{\Gamma(n-\mu)} \left(\frac{d}{dt} \right)^n \int_0^t (t-s)^{n-\mu-1} x(s) ds, \quad (2.3)$$

provided the right side is pointwise defined on $(0, +\infty)$. We recall the following properties [8, 9] which are useful for the sequel.

Lemma 2.1 (see [8, 9]).

(1) If $x \in L^1(0, 1)$, $\rho > \sigma > 0$, and $n \in \mathbb{N}$, then

$$I^\rho I^\sigma x(t) = I^{\rho+\sigma} x(t), \quad \mathfrak{D}_t^\sigma I^\rho x(t) = I^{\rho-\sigma} x(t), \quad (2.4)$$

$$\mathfrak{D}_t^\sigma I^\sigma x(t) = x(t), \quad \frac{d}{dt^n} (\mathfrak{D}_t^\sigma x(t)) = \mathfrak{D}_t^{n+\sigma} x(t). \quad (2.5)$$

(2) If $\nu > 0$, $\sigma > 0$, then

$$\mathfrak{D}_t^\nu t^{\sigma-1} = \frac{\Gamma(\sigma)}{\Gamma(\sigma-\nu)} t^{\sigma-\nu-1}. \quad (2.6)$$

Lemma 2.2 (see [8]). Assume that $x \in L^1(0, 1)$ and $\mu > 0$. Then

$$I^\mu \mathfrak{D}_t^\mu x(t) = x(t) + c_1 t^{\mu-1} + c_2 t^{\mu-2} + \dots + c_n t^{\mu-n}, \quad (2.7)$$

where $c_i \in \mathbb{R}$ ($i = 1, 2, \dots, n$), n is the smallest integer greater than or equal to μ .

Let $x(t) = I^{\mu_{n-1}} v(t)$, and consider the following modified integro-differential equation:

$$\begin{aligned} -\mathfrak{D}_t^{\mu-\mu_{n-1}} v(t) &= \lambda f(t, I^{\mu_{n-1}} v(t), I^{\mu_{n-1}-\mu_1} v(t), \dots, I^{\mu_{n-1}-\mu_{n-2}} v(t), v(t)), \\ v(0) = v'(0) &= 0 \quad v(1) = \sum_{j=1}^{m-2} a_j v(\xi_j). \end{aligned} \quad (2.8)$$

The following Lemmas 2.3–2.5 are obtained by Zhang et al. [10].

Lemma 2.3. The higher order multipoint boundary value problem (1.1) has a positive solution if and only if nonlinear integro-differential equation (2.8) has a positive solution. Moreover, if v is a positive solution of (2.8), then $x(t) = I^{\mu_{n-1}} v(t)$ is positive solution of the higher order multipoint boundary value problem (1.1).

Lemma 2.4. If $2 < \mu - \mu_{n-1} < 3$ and $\alpha \in L^1[0, 1]$, then the boundary value problem

$$\begin{aligned} \mathfrak{D}_t^{\mu-\mu_{n-1}} w(t) + \lambda \alpha(t) &= 0, \\ w(0) = w'(0) &= 0, \quad w(1) = \sum_{j=1}^{m-2} a_j w(\xi_j) \end{aligned} \quad (2.9)$$

has the unique solution

$$w(t) = \lambda \int_0^1 K(t, s) \alpha(s) ds, \quad (2.10)$$

where

$$K(t, s) = k(t, s) + \frac{t^{\mu-\mu_{n-1}-1}}{1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1}} \sum_{j=1}^{m-2} a_j k(\xi_j, s), \quad (2.11)$$

is the Green function of the boundary value problem (2.9), and

$$k(t, s) = \begin{cases} \frac{[t(1-s)]^{\mu-\mu_{n-1}-1} - (t-s)^{\mu-\mu_{n-1}-1}}{\Gamma(\mu-\mu_{n-1})}, & 0 \leq s \leq t \leq 1, \\ \frac{[t(1-s)]^{\mu-\mu_{n-1}-1}}{\Gamma(\mu-\mu_{n-1})}, & 0 \leq t \leq s \leq 1. \end{cases} \quad (2.12)$$

Lemma 2.5. *The Green function of the boundary value problem (2.9) satisfies*

$$K(t, s) \leq \tau, \quad (2.13)$$

where

$$\tau = \frac{1 + \sum_{j=1}^{m-2} a_j (1 - \xi_j^{\mu-\mu_{n-1}-1})}{\Gamma(\mu-\mu_{n-1}) (1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1})}. \quad (2.14)$$

Define a modified function $[\cdot]^*$ for any $z \in C[0, 1]$ by

$$[z(t)]^* = \begin{cases} z(t), & z(t) \geq 0, \\ 0, & z(t) < 0, \end{cases} \quad (2.15)$$

and consider the following boundary value problem

$$\begin{aligned} -\mathfrak{D}_t^{\mu-\mu_{n-1}} u(t) &= \lambda [f(t, I^{\mu_{n-1}}[u(t) - w(t)]^*, I^{\mu_{n-1}-\mu_1}[u(t) - w(t)]^* \cdots, \\ &\quad I^{\mu_{n-1}-\mu_{n-2}}[u(t) - w(t)]^*, [u(t) - w(t)]^*) + \alpha(t)], \\ u(0) = u'(0) &= 0, \quad u(1) = \sum_{j=1}^{m-2} a_j u(\xi_j). \end{aligned} \quad (2.16)$$

Lemma 2.6. *Suppose $u(t) \geq w(t)$, $t \in [0, 1]$ is a solution of the problem (2.16), then $u - w$ is a positive solution of the problem (2.8), consequently, $I^{\mu_{n-1}}[u(t) - w(t)]$ is also a positive solution of the semipositone higher differential equation (1.1).*

Proof. Since u is a solution of the BVP (2.16) and $u(t) \geq w(t)$ for any $t \in [0, 1]$, then we have

$$\begin{aligned} -\mathfrak{D}_t^{\mu-\mu_{n-1}}u(t) &= \lambda [f(t, I^{\mu_{n-1}}[u(t) - w(t)], I^{\mu_{n-1}-\mu_1}[u(t) - w(t)], \dots, \\ &\quad I^{\mu_{n-1}-\mu_{n-2}}[u(t) - w(t)], [u(t) - w(t)] + \alpha(t)], \\ u(0) = u'(0) &= 0, \quad u(1) = \sum_{j=1}^{m-2} a_j u(\xi_j). \end{aligned} \quad (2.17)$$

Let $v = u - w$, then we have

$$w(0) = w'(0) = 0, \quad w(1) = \sum_{j=1}^{m-2} a_j w(\xi_j), \quad (2.18)$$

and $\mathfrak{D}_t^{\mu-\mu_{n-1}}v(t) = \mathfrak{D}_t^{\mu-\mu_{n-1}}u(t) - \mathfrak{D}_t^{\mu-\mu_{n-1}}w(t)$, which implies that

$$-\mathfrak{D}_t^{\mu-\mu_{n-1}}u(t) = -\mathfrak{D}_t^{\mu-\mu_{n-1}}v(t) + \lambda \alpha(t). \quad (2.19)$$

Substituting the above into (2.17), then $v = u - w$ solves the (2.8), that is, $u - w$ is a positive solution of the semipositone differential equation (2.8). By Lemma 2.3, $I^{\mu_{n-1}}[u(t) - w(t)]$ is a positive solution of the singular semipositone differential equation (1.1). This completes the proof of Lemma 2.5. $\square$

Let

$$\gamma(t) = \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1}\right) t^{\mu-\mu_{n-1}-1} (1-t) + \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1} (1-\xi_j) t^{\mu-\mu_{n-1}-1}. \quad (2.20)$$

Lemma 2.7 (see [10]). *The solution $w(t)$ of (2.9) satisfies*

$$w(t) \leq \lambda \eta \gamma(t), \quad t \in [0, 1], \quad (2.21)$$

where

$$\eta = \frac{\int_0^1 \alpha(s) ds}{\Gamma(\mu - \mu_{n-1}) \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1}\right)}. \quad (2.22)$$

It is well known that the BVP (2.17) is equivalent to the fixed points for the mapping T by

$$\begin{aligned} (Tu)(t) &= \lambda \int_0^1 K(t, s) [f(s, I^{\mu_{n-1}}[u(s) - w(s)], I^{\mu_{n-1}-\mu_1}[u(s) - w(s)], \dots, \\ &\quad I^{\mu_{n-1}-\mu_{n-2}}[u(s) - w(s)], [u(s) - w(s)] + \alpha(s)] ds. \end{aligned} \quad (2.23)$$

The basic space used in this paper is $E = C([0, 1]; \mathbb{R})$, where $\mathbb{R}$ is the set of real numbers. Obviously, the space E is a Banach space if it is endowed with the norm as follows:

$$\|u\| = \max_{t \in [0, 1]} |u(t)|, \quad (2.24)$$

for any $u \in E$. Let

$$P = \left\{ u \in E : u(t) \geq \frac{1}{2} \gamma(t) \|u\| \right\}, \quad (2.25)$$

then P is a cone of E .

Lemma 2.8. *Assume that (B1) holds. Then $T : P \rightarrow P$ is a completely continuous operator.*

Proof. By using similar method to [10] and standard arguments, according to the Ascoli-Arzelà Theorem, one can show that $T : P \rightarrow P$ is a completely continuous operator. $\square$

Lemma 2.9 (see [29]). *Let E be a real Banach space, $P \subset E$ be a cone. Assume Ω_1, Ω_2 are two bounded open subsets of E with $\theta \in \Omega_1$, $\overline{\Omega_1} \subset \Omega_2$, and let $T : P \cap (\overline{\Omega_2} \setminus \Omega_1) \rightarrow P$ be a completely continuous operator such that either*

- (1) $\|Tx\| \leq \|x\|, x \in P \cap \partial\Omega_1$ and $\|Tx\| \geq \|x\|, x \in P \cap \partial\Omega_2$, or
- (2) $\|Tx\| \geq \|x\|, x \in P \cap \partial\Omega_1$ and $\|Tx\| \leq \|x\|, x \in P \cap \partial\Omega_2$.

Then T has a fixed point in $P \cap (\overline{\Omega_2} \setminus \Omega_1)$.

3. Main Results

Theorem 3.1. *Suppose that (B1) holds, and*

$$\lim_{\sum_{i=1}^n |x_i| \rightarrow \infty} \min_{t \in [1/4, 3/4]} \frac{f(t, x_1, x_2, \dots, x_n)}{\sum_{i=1}^n |x_i|} = +\infty. \quad (3.1)$$

Then there exists some constant $\lambda^ > 0$ such that the higher order multipoint boundary value problem (1.1) has at least one positive solution for any $\lambda \in (0, \lambda^*)$.*

Proof. By Lemma 2.8, we know T is a completely continuous operator. Take

$$r = \tau \int_0^1 [\beta(s) + \alpha(s)] ds, \quad (3.2)$$

where τ is defined by Lemma 2.5. Let $\Omega_1 = \{u \in P : \|u\| < r\}$. Then, for any $u \in \partial\Omega_1$, $s \in [0, 1]$, we have

$$\begin{aligned} 0 &\leq [u(s) - w(s)]^* \leq u(s) \leq \|u\| \leq r, \\ 0 &\leq I^{\mu_{n-1}}[u(s) - w(s)]^* \leq \frac{r}{\Gamma(\mu_{n-1})}, \\ 0 &\leq I^{\mu_{n-1}-\mu_i}[u(s) - w(s)]^* \leq \frac{r}{\Gamma(\mu_{n-1}-\mu_i)}, \quad i = 1, \dots, n-2. \end{aligned} \quad (3.3)$$

Choose

$$\lambda^* = \min \left\{ \frac{r}{4\eta}, \frac{1}{N+1} \right\}, \quad (3.4)$$

where

$$N = \max_{(u_1, u_2, \dots, u_n) \in [0,1] \times [0, r/\Gamma(\mu_{n-1})] \times \dots \times [0, r/\Gamma(\mu_{n-1}-\mu_{n-2})] \times [0, r]} h(u_1, u_2, \dots, u_n), \quad (3.5)$$

and η is defined by Lemma 2.7. Thus, for any $u \in \partial\Omega_1$, $s \in [0, 1]$, and $\lambda \in (0, \lambda^*)$, by (3.3), we have

$$\begin{aligned} \|Tu\| &= \max_{t \in [0,1]} (Tu)(t) \\ &= \lambda \max_{t \in [0,1]} \int_0^1 K(t, s) [f(s, I^{\mu_{n-1}}[u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1}[u(s) - w(s)]^*, \dots, \\ &\quad I^{\mu_{n-1}-\mu_{n-2}}[u(s) - w(s)]^*, [u(s) - w(s)]^*) + \alpha(s)] ds \\ &\leq \lambda \tau \int_0^1 [\beta(s) h(I^{\mu_{n-1}}[u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1}[u(s) - w(s)]^*, \dots, \\ &\quad I^{\mu_{n-1}-\mu_{n-2}}[u(s) - w(s)]^*, [u(s) - w(s)]^*) + \alpha(s)] ds \\ &\leq \lambda \tau (N+1) \int_0^1 [\beta(s) + \alpha(s)] ds \leq r = \|u\|. \end{aligned} \quad (3.6)$$

Therefore,

$$\|Tu\| \leq \|u\|, \quad u \in P \cap \partial\Omega_1. \quad (3.7)$$

On the other hand, choose a real number $L > 0$ such that

$$\lambda L \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) ds \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}}\right) \left(\frac{1}{4}\right)^{\mu-\mu_{n-1}+1} \geq 1. \quad (3.8)$$

By (3.1), for any $t \in [1/4, 3/4]$, there exists a constant $B > 0$ such that

$$\frac{f(t, x_1, x_2, \dots, x_n)}{\sum_{i=1}^n |x_i|} > L, \quad \text{for } \sum_{i=1}^n |x_i| \geq B. \quad (3.9)$$

Take

$$R = \max \left\{ 2r, 4\lambda\eta, \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}} \right)^{-1} 4^{\mu-\mu_{n-1}+1} B \right\}, \quad (3.10)$$

let $\Omega_2 = \{u \in P : \|u\| < R\}$ and $\partial\Omega_2 = \{u \in P : \|u\| = R\}$. Then for any $u \in \partial\Omega_2$, $s \in [0, 1]$, by Lemma 2.7, we have

$$u(s) - w(s) \geq u(s) - \lambda\eta\gamma(s) \geq \frac{1}{2}R\gamma(s) - \lambda\eta\gamma(s) \geq \frac{1}{4}R\gamma(s) \geq 0. \quad (3.11)$$

And then, for any $u \in \partial\Omega_2$, $s \in [1/4, 3/4]$, one gets

$$\begin{aligned} & \left(\sum_{i=1}^{n-2} |I^{\mu_{n-1}-\mu_i} [u(s) - w(s)]^*| \right) + |I^{\mu_{n-1}} [u(s) - w(s)]^*| + |[u(s) - w(s)]^*| \\ & \geq u(s) - w(s) \geq \frac{1}{4}R\gamma(t) \geq \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1} \right) \left(\frac{1}{4} \right)^{\mu-\mu_{n-1}+1} R \\ & \quad + \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1} (1 - \xi_j) \left(\frac{1}{4} \right)^{\mu-\mu_{n-1}} R \geq \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}} \right) \left(\frac{1}{4} \right)^{\mu-\mu_{n-1}+1} R \geq B. \end{aligned} \quad (3.12)$$

It follows from (3.12) that, for any $u \in \partial\Omega_2$,

$$\begin{aligned} \|Tu\| & \geq \lambda \int_0^1 K\left(\frac{1}{2}, s\right) [f(s, I^{\mu_{n-1}} [u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1} [u(s) - w(s)]^*, \dots, \\ & \quad I^{\mu_{n-1}-\mu_{n-2}} [u(s) - w(s)]^*, [u(s) - w(s)]^*) + \alpha(s)] ds \\ & \geq \lambda \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) f(s, I^{\mu_{n-1}} [u(s) - w(s)], I^{\mu_{n-1}-\mu_1} [u(s) - w(s)], \dots, \\ & \quad I^{\mu_{n-1}-\mu_{n-2}} [u(s) - w(s)], [u(s) - w(s)]) ds \end{aligned}$$

$$\begin{aligned}
&\geq \lambda \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) L(|I^{\mu_{n-1}}[u(s) - w(s)]| + |I^{\mu_{n-1}-\mu_1}[u(s) - w(s)]| \\
&\quad + \dots + |I^{\mu_{n-1}-\mu_{n-2}}[u(s) - w(s)]| + |[u(s) - w(s)]|) ds \\
&\geq \lambda L \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) |[u(s) - w(s)]| ds \\
&\geq \lambda L \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) ds \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}}\right) \left(\frac{1}{4}\right)^{\mu-\mu_{n-1}+1} R = R = \|u\|.
\end{aligned} \tag{3.13}$$

So, we have

$$\|Tu\| \geq \|u\|, \quad u \in P \cap \partial\Omega_2. \tag{3.14}$$

By Lemma 2.9, T has at least a fixed point $u \in (P \cap \overline{\Omega_2}) \setminus \Omega_1$ such that $r \leq \|u\| \leq R$.

It follows from $\lambda^* \leq r/4\eta$ and $\lambda \in (0, \lambda^*)$ that

$$u(t) - w(t) \geq u(t) - \lambda\eta\gamma(t) \geq \frac{1}{2}r\gamma(t) - \lambda\eta\gamma(t) \geq \frac{r}{4}\gamma(t) > 0, \quad t \in (0, 1). \tag{3.15}$$

Let $x(t) = I^{\mu_{n-1}}[u(t) - w(t)]$, then

$$x(t) > 0, \quad t \in (0, 1). \tag{3.16}$$

By Lemma 2.6, we know that the differential equation (1.1) has at least a positive solutions x . $\square$

Theorem 3.2. Suppose (B1) holds and

$$\liminf_{\sum_{i=1}^n |u_i| \rightarrow \infty} \min_{t \in [1/4, 3/4]} f(t, u_1, u_2, \dots, u_n) = +\infty, \quad \lim_{\sum_{i=1}^n |u_i| \rightarrow \infty} \frac{h(u_1, u_2, \dots, u_n)}{\sum_{i=1}^n |u_i|} = 0. \tag{3.17}$$

Then there exists $\lambda^* > 0$ such that the higher order multipoint boundary value problem (1.1) has at least one positive solution for any $\lambda \in (\lambda^*, +\infty)$.

Proof. By (3.17), there exists $M > 0$ such that for any $t \in [1/4, 3/4]$ we have

$$f(t, u_1, u_2, \dots, u_n) \geq \frac{4\eta}{\int_{1/4}^{3/4} K(1/2, s) ds}, \quad \text{if } \sum_{i=1}^n |u_i| \geq M. \tag{3.18}$$

Let

$$\lambda^* = \frac{M}{\eta \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu - \mu_{n-1}} \right) (1/4)^{\mu - \mu_{n-1} + 1}}. \quad (3.19)$$

In the following of the proof, we suppose $\lambda > \lambda^*$. Take

$$r = 4\lambda\eta, \quad (3.20)$$

and let $\Omega_1 = \{u \in P : \|u\| < r\}$ and $\partial\Omega_1 = \{u \in P : \|u\| = r\}$. Then for any $u \in \partial\Omega_1$, $s \in [0, 1]$, by Lemma 2.7, we have

$$u(s) - w(s) \geq u(s) - \lambda\eta\gamma(s) \geq \frac{1}{2}r\gamma(s) - \lambda\eta\gamma(s) = \lambda\eta\gamma(s) \geq 0. \quad (3.21)$$

So for any $u \in \partial\Omega_1$, $s \in [1/4, 3/4]$, one gets

$$\begin{aligned} & \left(\sum_{i=1}^{n-2} |I^{\mu_{n-1}-\mu_i} [u(s) - w(s)]^*| \right) + |I^{\mu_{n-1}} [u(s) - w(s)]^*| + |[u(s) - w(s)]^*| \\ & \geq u(s) - w(s) \geq \lambda\eta\gamma(s) \geq \lambda\eta \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu - \mu_{n-1}} \right) \left(\frac{1}{4} \right)^{\mu - \mu_{n-1} + 1} \\ & \geq \lambda^* \eta \left(1 - \sum_{j=1}^{m-2} a_j \xi_j^{\mu - \mu_{n-1}} \right) \left(\frac{1}{4} \right)^{\mu - \mu_{n-1} + 1} \geq M. \end{aligned} \quad (3.22)$$

Thus, by (3.22), for any $u \in \partial\Omega_1$, we have

$$\begin{aligned} \|Tu\| & \geq \lambda \int_0^1 K\left(\frac{1}{2}, s\right) [f(s, I^{\mu_{n-1}} [u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1} [u(s) - w(s)]^*, \dots, \\ & \quad I^{\mu_{n-1}-\mu_{n-2}} [u(s) - w(s)]^*, [u(s) - w(s)]^*) + \alpha(s)] ds \\ & \geq \lambda \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) f(s, I^{\mu_{n-1}} [u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1} [u(s) - w(s)]^*, \dots, \\ & \quad I^{\mu_{n-1}-\mu_{n-2}} [u(s) - w(s)]^*, [u(s) - w(s)]^*) ds \\ & \geq \lambda \int_{1/4}^{3/4} K\left(\frac{1}{2}, s\right) ds \times \frac{4\eta}{\int_{1/4}^{3/4} K(1/2, s) ds} = r = \|u\|. \end{aligned} \quad (3.23)$$

So, we have

$$\|Tu\| \geq \|u\|, \quad u \in P \cap \partial\Omega_1. \quad (3.24)$$

Next, take

$$\sigma = \sum_{i=1}^{n-2} \frac{1}{\Gamma(\mu_{n-1} - \mu_i)} + \frac{1}{\Gamma(\mu_{n-1})} + 1. \quad (3.25)$$

Let us choose $\varepsilon > 0$ such that

$$\lambda \tau \sigma \varepsilon \int_0^1 \beta(s) ds < 1. \quad (3.26)$$

Then for the above ε , by (3.17), there exists $N > r > 0$ such that, for any $t \in [0, 1]$,

$$h(u_1, u_2, \dots, u_n) \leq \varepsilon \sum_{i=1}^n |u_i|, \quad \text{if } \sum_{i=1}^n |u_i| > N. \quad (3.27)$$

Thus, by (3.3) and (3.27), if

$$\left(\sum_{i=1}^{n-2} |I^{\mu_{n-1}-\mu_i} [u(s) - w(s)]^*| \right) + |I^{\mu_{n-1}} [u(s) - w(s)]^*| + |[u(s) - w(s)]^*| > N, \quad (3.28)$$

we have

$$\begin{aligned} & h(I^{\mu_{n-1}} [u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1} [u(s) - w(s)]^*, \dots, \\ & I^{\mu_{n-1}-\mu_{n-2}} [u(s) - w(s)]^*, [u(s) - w(s)]^*) ds \\ & \leq |I^{\mu_{n-1}} [u(s) - w(s)]^*| + |I^{\mu_{n-1}-\mu_1} [u(s) - w(s)]^*| + \dots \\ & \quad + |I^{\mu_{n-1}-\mu_{n-2}} [u(s) - w(s)]^*| + |[u(s) - w(s)]^*| \varepsilon \\ & \leq \left(\sum_{i=1}^{n-2} \frac{1}{\Gamma(\mu_{n-1} - \mu_i)} + \frac{1}{\Gamma(\mu_{n-1})} + 1 \right) \|u\| \varepsilon = \sigma \|u\| \varepsilon. \end{aligned} \quad (3.29)$$

Take

$$R = \frac{\lambda \tau \theta \int_0^1 [\beta(s) + \alpha(s)] ds + \lambda \tau \int_0^1 \alpha(s) ds}{1 - \lambda \tau \sigma \varepsilon \int_0^1 \beta(s) ds} + N, \quad (3.30)$$

where

$$\theta = \max_{\sum_{i=1}^n |u_i| \leq N} h(u_1, u_2, \dots, u_n) + 1. \quad (3.31)$$

Then $R > N > r$.

Now let $\Omega_2 = \{u \in P : \|u\| < R\}$ and $\partial\Omega_2 = \{u \in P : \|u\| = R\}$. Then, for any $u \in P \cap \partial\Omega_2$, we have

$$\begin{aligned}
\|Tu\| &= \max_{t \in [0,1]} (Tu)(t) \\
&= \lambda \max_{t \in [0,1]} \int_0^1 K(t,s) [f(s, I^{\mu_{n-1}}[u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1}[u(s) - w(s)]^*, \dots, \\
&\quad I^{\mu_{n-1}-\mu_{n-2}}[u(s) - w(s)]^*, [u(s) - w(s)]^*) + \alpha(s)] ds \\
&\leq \lambda \tau \int_0^1 [\beta(s) h(I^{\mu_{n-1}}[u(s) - w(s)]^*, I^{\mu_{n-1}-\mu_1}[u(s) - w(s)]^*, \dots, \\
&\quad I^{\mu_{n-1}-\mu_{n-2}}[u(s) - w(s)]^*, [u(s) - w(s)]^*) + \alpha(s)] ds \\
&\leq \lambda \tau \left(\max_{\sum_{i=1}^n |u_i| \leq N} h(u_1, u_2, \dots, u_n) + 1 \right) \int_0^1 [\beta(s) + \alpha(s)] ds + \lambda \tau \int_0^1 [\beta(s) \sigma \varepsilon \|u\| + \alpha(s)] ds \\
&\leq \lambda \tau \theta \int_0^1 [\beta(s) + \alpha(s)] ds + \lambda \tau \int_0^1 \alpha(s) ds + \lambda \tau \sigma \varepsilon \int_0^1 \beta(s) ds R \leq R = \|u\|,
\end{aligned} \tag{3.32}$$

which implies that

$$\|Tu\| \leq \|u\|, \quad u \in P \cap \partial\Omega_2. \tag{3.33}$$

By Lemma 2.9, T has at least a fixed point $u \in (P \cap \overline{\Omega_2}) \setminus \Omega_1$ such that $r \leq \|u\| \leq R$.

It follows from $r = 4\lambda\eta$ that

$$u(t) - w(t) \geq \frac{1}{2} \|u\| \gamma(t) - \lambda \eta \gamma(t) = \lambda \eta \gamma(t) > 0, \quad t \in (0, 1). \tag{3.34}$$

Let $x(t) = I^{\mu_{n-1}}[u(t) - w(t)]$, then

$$x(t) > 0, \quad t \in (0, 1). \tag{3.35}$$

By Lemma 2.6, we know that the differential equation (1.1) has at least a positive solutions x . $\square$

Example 3.3. Consider the existence of positive solutions for the nonlinear higher order fractional differential equation with four-point boundary condition

$$\begin{aligned}
-\mathfrak{D}_t^{5/2} x(t) &= \left[|x(t)| + \left| \mathfrak{D}_t^{1/4} x(t) \right| + \left| \mathfrak{D}_t^{3/8} x(t) \right| \right]^{1/2} + \ln t, \\
\mathfrak{D}_t^{1/4} x(0) &= \mathfrak{D}_t^{3/8} x(0) = \mathfrak{D}_t^{11/8} x(0) = 0, \quad \mathfrak{D}_t^{3/8} x(1) = 2\mathfrak{D}_t^{3/8} x\left(\frac{1}{2}\right) - \mathfrak{D}_t^{3/8} x\left(\frac{3}{4}\right).
\end{aligned} \tag{3.36}$$

Then there exists $\lambda^* > 0$ such that the higher order four-point boundary value problem (1.1) has at least one positive solution for any $\lambda \in (\lambda^*, +\infty)$.

Proof. Let

$$f(t, x, y, z) = (|x| + |y| + |z|)^{1/2} + \ln t, \quad t \in (0, 1) \times \mathbb{R}^3, \quad (3.37)$$

then

$$\begin{aligned} -|\ln t| = \ln t \leq f(t, x, y, z) &= (|x| + |y| + |z|)^{1/2} + \ln t \leq (|x| + |y| + |z|)^{1/2}, \\ \alpha(t) = |\ln t|, \quad \beta(t) &= 1, \quad h(x, y, z) = (|x| + |y| + |z|)^{1/2}, \end{aligned} \quad (3.38)$$

$$\liminf_{|x|+|y|+|z| \rightarrow +\infty} \min_{t \in [1/4, 3/4]} f(t, x, y, z) = +\infty, \quad \lim_{|x|+|y|+|z| \rightarrow +\infty} \frac{h(x, y, z)}{|x| + |y| + |z|} = 0.$$

Clearly, $\alpha, \beta \in L^1[(0, 1), (0, +\infty)]$, and

$$0 < \sum_{j=1}^{m-2} a_j \xi_j^{\mu-\mu_{n-1}-1} = 1 - 2\left(\frac{1}{2}\right)^{9/8} + \left(\frac{3}{4}\right)^{9/8} = 0.8065 < 1. \quad (3.39)$$

By Theorem 3.2, there exists $\lambda^* > 0$ such that the higher order multipoint boundary value problem (3.36) has at least one positive solution for any $\lambda \in (\lambda^*, +\infty)$. $\square$

References

- [1] H. M. Srivastava and S. Owa, *Univalent Functions, Fractional Calculus, and Their Applications*, Halsted Press, New York, NY, USA, John Wiley and Sons, Chichester, UK, 1989.
- [2] L. Gaul, P. Klein, and S. Kemple, "Damping description involving fractional operators," *Mechanical Systems and Signal Processing*, vol. 5, no. 2, pp. 81–88, 1991.
- [3] W. G. Glockle and T. F. Nonnenmacher, "A fractional calculus approach to self-similar protein dynamics," *Biophysical Journal*, vol. 68, no. 1, pp. 46–53, 1995.
- [4] K. Diethelm, *The Analysis of Fractional Differential Equations*, Springer-Verlag, Berlin, Germany, 2010.
- [5] R. Metzler, W. Schick, H. G. Kilian, and T. F. Nonnenmacher, "Relaxation in filled polymers: a fractional calculus approach," *The Journal of Chemical Physics*, vol. 103, no. 16, pp. 7180–7186, 1995.
- [6] K. B. Oldham and J. Spanier, *The Fractional Calculus*, Academic Press, New York, NY, USA, 1974.
- [7] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, "Theory and applications of fractional differential equations," in *North-Holland Mathematics Studies*, vol. 204, Elsevier, Amsterdam, The Netherlands, 2006.
- [8] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, New York, NY, USA, 1993.
- [9] I. Podlubny, *Fractional Differential Equations*, *Mathematics in Science and Engineering*, Academic Press, New York, NY, USA, 1999.
- [10] X. Zhang, L. Liu, and Y. Wu, "Existence results for multiple positive solutions of nonlinear higher order perturbed fractional differential equations with derivatives," *Applied Mathematics and Computation*, vol. 219, no. 4, pp. 1420–1433, 2012.
- [11] B. Ahmad, A. Alsaedi, and B. S. Alghamdi, "Analytic approximation of solutions of the forced Duffing equation with integral boundary conditions," *Nonlinear Analysis: Real World Applications*, vol. 9, no. 4, pp. 1727–1740, 2008.

- [12] X. Zhang and Y. Han, "Existence and uniqueness of positive solutions for higher order nonlocal fractional differential equations," *Applied Mathematics Letters*, vol. 25, no. 3, pp. 555–560, 2012.
- [13] X. Zhang, L. Liu, and Y. Wu, "The eigenvalue problem for a singular higher order fractional differential equation involving fractional derivatives," *Applied Mathematics and Computation*, vol. 218, no. 17, pp. 8526–8536, 2012.
- [14] X. Zhang, L. Liu, B. Wiwatanapataphee, and Y. Wu, "Positive solutions of eigenvalue problems for a class of fractional differential equations with derivatives," *Abstract and Applied Analysis*, vol. 2012, Article ID 512127, 16 pages, 2012.
- [15] X. Zhang, L. Liu, and Y. Wu, "Multiple positive solutions of a singular fractional differential equation with negatively perturbed term," *Mathematical and Computer Modelling*, vol. 55, no. 3–4, pp. 1263–1274, 2012.
- [16] H. A. H. Salem, "On the fractional order m -point boundary value problem in reflexive Banach spaces and weak topologies," *Journal of Computational and Applied Mathematics*, vol. 224, no. 2, pp. 565–572, 2009.
- [17] C. S. Goodrich, "Existence of a positive solution to a class of fractional differential equations," *Applied Mathematics Letters*, vol. 23, no. 9, pp. 1050–1055, 2010.
- [18] M. Rehman and R. A. Khan, "Existence and uniqueness of solutions for multi-point boundary value problems for fractional differential equations," *Applied Mathematics Letters*, vol. 23, no. 9, pp. 1038–1044, 2010.
- [19] B. Ahmad and J. J. Nieto, "Riemann-Liouville fractional integro-differential equations with fractional nonlocal integral boundary conditions," *Boundary Value Problems*, vol. 2011, 2011.
- [20] C. S. Goodrich, "Existence of a positive solution to systems of differential equations of fractional order," *Computers and Mathematics with Applications*, vol. 62, no. 3, pp. 1251–1268, 2011.
- [21] C. S. Goodrich, "Existence and uniqueness of solutions to a fractional difference equation with nonlocal conditions," *Computers and Mathematics with Applications*, vol. 61, no. 2, pp. 191–202, 2011.
- [22] C. S. Goodrich, "Positive solutions to boundary value problems with nonlinear boundary conditions," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 75, no. 1, pp. 417–432, 2012.
- [23] M. Jia, X. Zhang, and X. Gu, "Nontrivial solutions for a higher fractional differential equation with fractional multi-point boundary conditions," *Boundary Value Problems*, vol. 2012, 70 pages, 2012.
- [24] M. Jia, X. Liu, and X. Gu, "Uniqueness and asymptotic behavior of positive solutions for a fractional-order integral boundary value problem," *Abstract and Applied Analysis*, vol. 2012, Article ID 294694, 21 pages, 2012.
- [25] P.-L. Lions, "On the existence of positive solutions of semilinear elliptic equations," *SIAM Review*, vol. 24, no. 4, pp. 441–467, 1982.
- [26] R. Aris, *Introduction to the Analysis of Chemical Reactors*, Prentice-Hall, Englewood Cliffs, NJ, USA, 1965.
- [27] A. Castro, C. Maya, and R. Shivaji, "Nonlinear eigenvalue problems with semipositone," *Electronic Journal of Differential Equations*, vol. 5, pp. 33–49, 2000.
- [28] V. Anuradha, D. D. Hai, and R. Shivaji, "Existence results for superlinear semipositone BVP'S," *Proceedings of the American Mathematical Society*, vol. 124, no. 3, pp. 757–763, 1996.
- [29] D. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cone*, Academic Press, New York, NY, USA, 1988.