Research Article

On the q -Euler Numbers and Polynomials with Weight 0

T. Kim and J. Choi

Division of General Education-Mathematics, Kwangjuon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to T. Kim, tkkim@kw.ac.kr

Received 13 October 2011; Accepted 29 November 2011

Academic Editor: Ibrahim Sadek

Copyright © 2012 T. Kim and J. Choi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of this paper is to investigate some properties of q -Euler numbers and polynomials with weight 0. From those q -Euler numbers with weight 0, we derive some identities on the q -Euler numbers and polynomials with weight 0.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will denote the ring of p -adic rational integers, the field of p -adic rational numbers, and the completion of algebraic closure of $\mathbb{Q}_p$. The p -adic absolute value is defined by $|x|_p = 1/p^r$ where $x = p^r s/t$ for $s, t \in \mathbb{Z}$ with $(p, t) = (p, s) = 1$ and $r \in \mathbb{Q}$. In this paper, we assume that $\alpha \in \mathbb{Q}$ and $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$. As well-known definition, the Euler polynomials are defined by

$$\frac{2}{e^t + 1} e^{xt} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (1.1)$$

with the usual convention about replacing $E^n(x)$ by $E_n(x)$ (see [1–15]).

In this special case, $x = 0$, $E_n(0) = E_n$ are called the n th Euler numbers (see [1]). Recently, the q -Euler numbers with weight α are defined by

$$\tilde{E}_{0,q}^{(\alpha)} = 1, \quad q \left(q^\alpha \tilde{E}_q^{(\alpha)} + 1 \right)^n + \tilde{E}_{n,q}^{(\alpha)} = 0 \quad \text{if } n > 0, \quad (1.2)$$

with the usual convention about replacing $(\tilde{E}_q^{(\alpha)})^n$ by $\tilde{E}_{n,q}^{(\alpha)}$ (see [3, 12]). The q -number of x is defined by $[x]_q = (1 - q^x)/(1 - q)$ (see [1–15]). Note that $\lim_{q \rightarrow 1} [x]_q = x$. Let us define

the notation of q -Euler numbers with weight 0 as $\tilde{E}_{n,q}^{(0)} = \tilde{E}_{n,q}$. The purpose of this paper is to investigate some interesting identities on the q -Euler numbers with weight 0.

2. On the Extended q -Euler Numbers of Higher-Order with Weight 0

Let $C(\mathbb{Z}_p)$ be the space of continuous functions on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, the fermionic p -adic q -integral on $\mathbb{Z}_p$ is defined by Kim as follows:

$$\begin{aligned} I_q(f) &= \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) \\ &= \lim_{N \rightarrow \infty} \frac{[2]_q}{1 + q^{p^N}} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \end{aligned} \quad (2.1)$$

(see [1–12]). By (2.1), we get

$$q^n I_q(f_n) + (-1)^{n-1} I_q(f) = [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l) q^l, \quad (2.2)$$

where $f_n(x) = f(x+n)$ and $n \in \mathbb{N}$ (see [4, 5]).

By (1.2), (2.1), and (2.2), we see that

$$\int_{\mathbb{Z}_p} [x]_{q^a}^n d\mu_{-q}(x) = \tilde{E}_{n,q}^{(\alpha)} = \frac{[2]_q}{(1-q)^n [\alpha]_q^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^{al+1}}. \quad (2.3)$$

In the special case, $n = 1$, we get

$$\int_{\mathbb{Z}_p} e^{xt} d\mu_{-q}(x) = \frac{[2]_q}{qe^t + 1} = \frac{1+q^{-1}}{e^t + q^{-1}} = \sum_{n=0}^{\infty} H_n(-q^{-1}) \frac{t^n}{n!}, \quad (2.4)$$

where $H_n(-q^{-1})$ are the n th Frobenius-Euler numbers. From (2.4), we note that the q -Euler numbers with weight 0 are given by

$$\tilde{E}_{n,q} = \int_{\mathbb{Z}_p} x^n d\mu_{-q}(x) = H_n(-q^{-1}), \quad \text{for } n \in \mathbb{Z}_+. \quad (2.5)$$

Therefore, by (2.5), we obtain the following theorem.

Theorem 2.1. For $n \in \mathbb{Z}_+$, one has

$$\tilde{E}_{n,q} = H_n(-q^{-1}), \quad (2.6)$$

where $H_n(-q^{-1})$ are called the n th Frobenius-Euler numbers.

Let us define the generating function of the q -Euler numbers with weight 0 as follows:

$$\tilde{F}_q(t) = \sum_{n=0}^{\infty} \tilde{E}_{n,q} \frac{t^n}{n!}. \quad (2.7)$$

Then, by (2.3) and (2.7), we get

$$\tilde{F}_q(t) = [2]_q \sum_{m=0}^{\infty} (-1)^m q^m e^{mt} = \frac{1+q}{qe^t + 1}. \quad (2.8)$$

Now we define the q -Euler polynomials with weight 0 as follows:

$$\sum_{n=0}^{\infty} \tilde{E}_{n,q}(x) \frac{t^n}{n!} = \frac{1+q}{qe^t + 1} e^{xt}. \quad (2.9)$$

Thus, (2.4) and (2.9), we get

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-q}(y) = \frac{1+q}{qe^t + 1} e^{xt} = \sum_{n=0}^{\infty} \tilde{E}_{n,q}(x) \frac{t^n}{n!}. \quad (2.10)$$

From (2.10), we have

$$\sum_{n=0}^{\infty} \tilde{E}_{n,q}(x) \frac{t^n}{n!} = \left(\frac{1+q^{-1}}{e^t + q^{-1}} \right) e^{xt} = \sum_{n=0}^{\infty} H_n(-q^{-1}, x) \frac{t^n}{n!}, \quad (2.11)$$

where $H_n(-q^{-1}, x)$ are called the n th Frobenius-Euler polynomials (see [9]).

Therefore, by (2.11), we obtain the following theorem.

Theorem 2.2. For $n \in \mathbb{Z}_+$, one has

$$\tilde{E}_{n,q}(x) = \int_{\mathbb{Z}_p} (x+y)^n d\mu_{-q}(x) = H_n(-q^{-1}, x), \quad (2.12)$$

where $H_n(-q^{-1}, x)$ are called the n th Frobenius-Euler polynomials.

From (2.2) and Theorem 2.2, we note that

$$q^n H_m(-q^{-1}, n) + H_m(-q^{-1}) = [2]_q \sum_{l=0}^{n-1} (-1)^l l^m q^l, \quad (2.13)$$

where $n \in \mathbb{N}$ with $n \equiv 1 \pmod{2}$.

Therefore, by (2.13), we obtain the following corollary.

Corollary 2.3. For $n \in \mathbb{N}$, with $n \equiv 1 \pmod{2}$ and $m \in \mathbb{Z}_+$, one has

$$q^n H_m(-q^{-1}, n) + H_m(-q^{-1}) = [2]_q \sum_{l=0}^{n-1} (-1)^l l^m q^l. \quad (2.14)$$

In particular, $q = 1$, we get $E_m(n) + E_m = 2 \sum_{l=0}^{n-1} (-1)^l l^m$, where E_m and $E_m(n)$ are called the m th Euler numbers and polynomials which are defined by

$$\frac{2}{e^t + 1} = \sum_{m=0}^{\infty} E_m \frac{t^m}{m!}, \quad \frac{2}{e^t + 1} e^{xt} = \sum_{m=0}^{\infty} E_m(x) \frac{t^m}{m!}. \quad (2.15)$$

By (2.2), we easily see that

$$q \int_{\mathbb{Z}_p} f(x+1) d\mu_{-q}(x) + \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = [2]_q f(0). \quad (2.16)$$

Thus, by (2.16), we get

$$\begin{aligned} [2]_q &= q \int_{\mathbb{Z}_p} e^{(x+1)t} d\mu_{-q}(x) + \int_{\mathbb{Z}_p} e^{xt} d\mu_{-q}(x) \\ &= \sum_{n=0}^{\infty} \left(q \int_{\mathbb{Z}_p} (x+1)^n d\mu_{-q}(x) + \int_{\mathbb{Z}_p} x^n d\mu_{-q}(x) \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(q H_n(-q^{-1}, 1) + H_n(-q^{-1}) \right) \frac{t^n}{n!}. \end{aligned} \quad (2.17)$$

Therefore, by (2.16), we obtain the following theorem.

Theorem 2.4. For $n \in \mathbb{Z}_+$, one has

$$q H_n(-q^{-1}, 1) + H_n(-q^{-1}) = \begin{cases} 1 + q, & \text{if } n = 0, \\ 0, & \text{if } n > 0, \end{cases} \quad (2.18)$$

where $H_n(-q^{-1}, x)$ are called the n th Frobenius-Euler polynomials and $H_n(-q^{-1})$ are called the n th Frobenius-Euler numbers. In particular, $q = 1$, we have

$$E_n(1) + E_n = \begin{cases} 2, & \text{if } n = 0, \\ 0, & \text{if } n > 0, \end{cases} \quad (2.19)$$

where E_n are called the n th Euler numbers.

From (2.5) and Theorem 2.2, we note that

$$\tilde{E}_{n,q}(x) = \int_{\mathbb{Z}_p} (x+y)^n d\mu_{-q}(y) = \sum_{l=0}^n \binom{n}{l} \int_{\mathbb{Z}_p} y^l d\mu_{-q}(y) x^{n-l} = \sum_{l=0}^n \binom{n}{l} \tilde{E}_{n,q} x^{n-l} = (x + \tilde{E}_q)^n, \quad (2.20)$$

where the usual convention about replacing $(\tilde{E}_q)^l$ by $\tilde{E}_{l,q}$. By Theorems 2.2 and 2.4, we get

$$q \tilde{E}_{n,q}(1) + \tilde{E}_{n,q} = \begin{cases} [2]_q, & \text{if } n = 0, \\ 0, & \text{if } n > 0. \end{cases} \quad (2.21)$$

From (2.20) and (2.21), we have

$$q\left(\tilde{E}_q + 1\right)^n + \tilde{E}_{n,q} = \begin{cases} [2]_q, & \text{if } n = 0, \\ 0, & \text{if } n > 0. \end{cases} \quad (2.22)$$

For $n \in \mathbb{N}$, by (2.20) and (2.22), we have

$$\begin{aligned} q^2 \tilde{E}_{n,q}(2) &= q^2 \left(\tilde{E}_q + 1 + 1\right)^n = q^2 \sum_{l=1}^n \binom{n}{l} \left(\tilde{E}_q + 1\right)^l + q \left(1 + q - \tilde{E}_{0,q}\right) = q + q^2 - q \sum_{l=0}^n \binom{n}{l} \tilde{E}_{l,q} \\ &= q + q^2 - q \left(\tilde{E}_q + 1\right)^n = q + q^2 + \tilde{E}_{n,q} - q[2]_q \delta_{0,n}. \end{aligned} \quad (2.23)$$

Therefore, by (2.23), we obtain the following theorem.

Theorem 2.5. *For $n \in \mathbb{N}$, one has*

$$q^2 \tilde{E}_{n,q}(2) = q + q^2 + \tilde{E}_{n,q}. \quad (2.24)$$

For $n \in \mathbb{Z}_+$, we have

$$\tilde{E}_{n,q^{-1}}(1-x) = \int_{\mathbb{Z}_p} (1-x+x_1)^n d\mu_{-q^{-1}}(x_1) = (-1)^n \int_{\mathbb{Z}_p} (x_1+x)^n d\mu_{-q}(x_1) = (-1)^n \tilde{E}_{n,q}(x). \quad (2.25)$$

Therefore, by (2.25), we obtain the following theorem.

Theorem 2.6. *For $n \in \mathbb{Z}_+$, one has*

$$\tilde{E}_{n,q^{-1}}(1-x) = (-1)^n \tilde{E}_{n,q}(x). \quad (2.26)$$

From (2.20), we have

$$\int_{\mathbb{Z}_p} (1-x)^n d\mu_{-q}(x) = (-1)^n \int_{\mathbb{Z}_p} (x-1)^n d\mu_{-q}(x) = (-1)^n \tilde{E}_{n,q}(-1). \quad (2.27)$$

By Theorem 2.6 and (2.27), we get

$$\int_{\mathbb{Z}_p} (1-x)^n d\mu_{-q}(x) = \tilde{E}_{n,q^{-1}}(2) = 1 + q + q^2 \tilde{E}_{n,q^{-1}} \quad \text{if } n > 0. \quad (2.28)$$

Therefore, by (2.28), we obtain the following theorem.

Theorem 2.7. *For $n \in \mathbb{N}$, one has*

$$\int_{\mathbb{Z}_p} (1-x)^n d\mu_{-q}(x) = 1 + q + q^2 \tilde{E}_{n,q^{-1}}. \quad (2.29)$$

Let $C(\mathbb{Z}_p)$ be the space of continuous functions on $\mathbb{Z}_p$. For $f \in C(\mathbb{Z}_p)$, p -adic analogue of Bernstein operator of order n for f is given by

$$\mathbb{B}_n(f | x) = \sum_{k=0}^n B_{k,n}(x) f\left(\frac{k}{n}\right) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k}, \quad (2.30)$$

where $n, k \in \mathbb{Z}_+$ (see [1, 6, 7]).

For $n, k \in \mathbb{Z}_+$, p -adic Bernstein polynomial of degree n is defined by

$$B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}, \quad x \in \mathbb{Z}_p \quad (2.31)$$

(see [1, 6, 7]).

Let us take the fermionic p -adic q -integral on $\mathbb{Z}_p$ for one Bernstein polynomials in (2.31) as follows:

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_{-q}(x) &= \binom{n}{k} \int_{\mathbb{Z}_p} x^k (1-x)^{n-k} d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} x^{k+l} d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \tilde{E}_{k+l,q}. \end{aligned} \quad (2.32)$$

By simple calculation, we easily get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) d\mu_{-q}(x) &= \int_{\mathbb{Z}_p} B_{n-k,n}(1-x) d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \int_{\mathbb{Z}_p} (1-x)^{n-l} d\mu_{-q}(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} (1+q+q^2 \tilde{E}_{n-l,q^{-1}}) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} q^2 \tilde{E}_{n-l,q^{-1}} + [2]_q \binom{n}{k} (-1)^k \delta_{0,k} \quad \text{if } n > k. \end{aligned} \quad (2.33)$$

Therefore, by (2.32) and (2.33), we obtain the following theorem.

Theorem 2.8. For $n \in \mathbb{Z}_+$ with $n > k > 0$, one has

$$\sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \tilde{E}_{k+l,q} = \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} q^2 \tilde{E}_{n-l,q^{-1}}. \quad (2.34)$$

In particular, $k = 0$, we get

$$\sum_{l=0}^n \binom{n}{l} (-1)^l \tilde{E}_{l,q} = q^2 \tilde{E}_{n,q^{-1}} + [2]_q. \quad (2.35)$$

By Theorems 2.1 and 2.2, we get

$$\sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l H_{k+l}(-q^{-1}) = \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} q^2 H_{n-l}(-q), \quad (2.36)$$

where $n, k \in \mathbb{Z}_+$ with $n > k > 0$.

References

- [1] S. Araci, D. Erdal, and J. J. Seo, "A Study on the fermionic p -adic q -integral on $\mathbb{Z}_p$ associated with weighted q -Bernstein and q -Genocchi polynomials," *Abstract and Applied Analysis*, vol. 2011, Article ID 649248, 10 pages, 2011.
- [2] D. Erdal, J. J. Seo, and S. Araci, "New construction weighted (h, g) -Genocchi numbers and polynomials related to Zeta type function," *Discrete Dynamics in Nature and Society*, vol. 2011, Article ID 487490, 7 pages, 2011.
- [3] T. Kim, B. Lee, J. Choi, Y. H. Kim, and S. H. Rim, "On the q -Euler numbers and weighted q -Bernstein polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, pp. 13–18, 2011.
- [4] T. Kim, "Some identities on the q -Euler polynomials of higher order and q -stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [5] T. Kim, J. Y. Choi, and J. Y. Sug, "Extended q -euler numbers and polynomials associated with fermionic p -adic q -integral on $\mathbb{Z}_p$," *Russian Journal of Mathematical Physics*, vol. 14, no. 2, pp. 160–163, 2007.
- [6] T. Kim, "A note on q -Bernstein polynomials," *Russian Journal of Mathematical Physics*, vol. 18, no. 4, pp. 73–82, 2011.
- [7] L. C. Jang, W.-J. Kim, and Y. Simsek, "A study on the p -adic integral representation on $\mathbb{Z}_p$ associated with Bernstein and Bernoulli polynomials," *Advances in Difference Equations*, vol. 2010, Article ID 163217, 6 pages, 2010.
- [8] L. C. Jang, K.-W. Hwang, and Y.-H. Kim, "A note on (h, q) -Genocchi polynomials and numbers of higher order," *Advances in Difference Equations*, vol. 2010, Article ID 309480, 6 pages, 2010.
- [9] M. Can, M. Cenkci, V. Kurt, and Y. Simsek, "Twisted Dedekind type sums associated with Barnes' type multiple Frobenius-Euler l -functions," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 2, pp. 135–160, 2009.
- [10] Y. Simsek, "Special functions related to Dedekind-type DC-sums and their applications," *Russian Journal of Mathematical Physics*, vol. 17, no. 4, pp. 495–508, 2010.
- [11] Y. Simsek, O. Yurekli, and V. Kurt, "On interpolation functions of the twisted generalized Frobenius-Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 15, pp. 187–194, 2007.
- [12] C. S. Ryoo, "A note on the weighted q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, pp. 47–54, 2011.
- [13] S. H. Rim, S. J. Lee, E. J. Moon, and J. H. Jin, "On the q -Genocchi numbers and polynomials associated with Q -zeta function," *Proceedings of the Jangjeon Mathematical Society*, vol. 12, no. 3, pp. 261–267, 2009.
- [14] C. S. Ryoo, "On the generalized Barnes type multiple q -Euler polynomials twisted by ramified roots of unity," *Proceedings of the Jangjeon Mathematical Society*, vol. 13, no. 2, pp. 255–263, 2010.
- [15] A. Bayad, "Modular properties of elliptic Bernoulli and Euler functions," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 389–401, 2010.