Research Article

Minimum-Norm Fixed Point of Pseudocontractive Mappings

Habtu Zegeye,¹ Naseer Shahzad,² and Mohammad Ali Alghamdi²

¹ Department of Mathematics, University of Botswana, Private Bag 00704, Gaborone, Botswana

² Department of Mathematics, King Abdulaziz University, P.O. Box. 80203, Jeddah 21589, Saudi Arabia

Correspondence should be addressed to Naseer Shahzad, nshahzad@kau.edu.sa

Received 7 May 2012; Accepted 14 June 2012

Academic Editor: Yonghong Yao

Copyright © 2012 Habtu Zegeye et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let K be a closed convex subset of a real Hilbert space H and let $T : K \rightarrow K$ be a continuous pseudocontractive mapping. Then for $\beta \in (0, 1)$ and each $t \in (0, 1)$, there exists a sequence $\{y_t\} \subset K$ satisfying $y_t = \beta P_K[(1-t)y_t] + (1-\beta)T(y_t)$ which converges strongly, as $t \rightarrow 0^+$, to the minimum-norm fixed point of T . Moreover, we provide an explicit iteration process which converges strongly to a minimum-norm fixed point of T provided that T is Lipschitz. Applications are also included. Our theorems improve several results in this direction.

1. Introduction

Let K be a nonempty subset of a real Hilbert space H . A mapping $T : K \rightarrow H$ is called *Lipschitz* if there exists $L \geq 0$ such that

$$\|Tx - Ty\| \leq L\|x - y\|, \quad \forall x, y \in K. \quad (1.1)$$

If $L \in [0, 1)$, then T is called a *contraction*; if $L = 1$ then T is called a *nonexpansive*. It is easy to see from (1.1) that every contraction mapping is nonexpansive, and every nonexpansive mapping is Lipschitz.

A mapping T is called *strongly pseudocontractive* if there exists $\alpha \in (0, 1)$ such that inequality

$$\langle Tx - Ty, x - y \rangle \leq \alpha \|x - y\|^2, \quad (1.2)$$

holds for all $x, y \in K$. T is called *pseudocontractive* if the inequality

$$\langle Tx - Ty, x - y \rangle \leq \|x - y\|^2, \quad (1.3)$$

holds for all $x, y \in K$. Note that inequality (1.3) can be equivalently written as

$$\|Tx - Ty\|^2 \leq \|x - y\|^2 + \|(I - T)x - (I - T)y\|^2, \quad \forall x, y \in K. \quad (1.4)$$

It is easy to see that nonexpansive and strongly pseudocontractive mappings are pseudocontractive mappings. However, the converse may not be true (see [1, 2] for details).

Interest in pseudocontractive mappings stems mainly from their firm connection with the important class of nonlinear *monotone* mappings, where a mapping A with domain $D(A)$ and range $R(A)$ in H is called *monotone* if the inequality

$$\langle Ax - Ay, x - y \rangle \geq 0, \quad (1.5)$$

holds for every $x, y \in D(A)$. We note that A is monotone if and only if $T := I - A$ is pseudocontractive, and hence a zero of A , $N(A) := \{x \in D(A) : Ax = 0\}$ is a fixed point of T , $F(T) := \{x \in D(T) : Tx = x\}$.

Let K be a nonempty closed convex subset of a real Hilbert space H and $T : K \rightarrow K$ a pseudocontractive mapping. Assume that the set of fixed points of T is nonempty. It is known from [3] that $F(T)$ is closed and convex.

Let the variational inequality (VI) be given as finding a point x^* with the property that

$$x^* \in F(T) \text{ such that } \langle x^*, x - x^* \rangle \geq 0, \quad \forall x \in F(T). \quad (1.6)$$

Then, x^* is the minimum-norm fixed point of T which exists uniquely and is exactly the (nearest point or metric) projection of the origin onto $F(T)$, that is, $x^* = P_{F(T)}(0)$. We also observe that the minimum-norm fixed point of pseudocontractive T is the minimum-norm solution of a monotone operator equation $Ax = 0$, where $A = (I - T)$.

It is quite often to seek the minimum-norm solution of a given nonlinear problem. In an abstract way, we may formulate such problems as finding a point x^* with the property

$$x^* \in K, \quad \|x^*\| = \min_{x \in K} \|x\|. \quad (1.7)$$

In other words, x^* is the projection of the origin onto K , that is,

$$x^* = P_K(0). \quad (1.8)$$

A typical example is the split feasibility problem (SFP), formulated as finding a point x^* with the property that

$$x^* \in K, \quad Ax^* \in Q, \quad (1.9)$$

where K and Q are nonempty closed convex subsets of the infinite-dimension real Hilbert spaces H_1 and H_2 , respectively, and A is bounded linear mapping from H_1 to H_2 . Equation (1.9) models many applied problems arising from image reconstructions and learning theory (see, e.g., [4]). Some works on the finite dimensional setting with relevant projection methods for solving image recovery problems can be found in [5–7]. Defining the proximity function f by

$$f(x) := \frac{1}{2} \|Ax - P_Q Ax\|^2, \quad (1.10)$$

we consider the convex optimization problem:

$$\min_{x \in K} f(x) := \min_{x \in K} \frac{1}{2} \|Ax - P_Q Ax\|^2. \quad (1.11)$$

It is clear that x^* is a solution to the split feasibility problem (1.9) if and only if $x^* \in K$ and $Ax^* - P_Q Ax^* = 0$ which is the minimum-norm solution of the minimization problem (1.11).

Motivated by the above split feasibility problem, we study the general case of finding the minimum-norm fixed point of a pseudocontractive mapping $T : K \rightarrow K$, that is, we find minimum norm fixed point of T which satisfies

$$x^* \in F(T) \quad \text{such that } \|x^*\| = \min\{\|x\| : x \in F(T)\}. \quad (1.12)$$

Let $T : K \rightarrow K$ be a nonexpansive self-mapping on *closed convex* subset K of a Banach space E . For a given $u \in K$ and for a given $t \in (0, 1)$ define a contraction $T_t : K \rightarrow K$ by

$$T_t x = (1 - t)u + tTx, \quad x \in K. \quad (1.13)$$

By Banach contraction principle, it yields a fixed point $z_t \in K$ of T_t , that is, z_t is the unique solution of the equation:

$$z_t = (1 - t)u + tTz_t. \quad (1.14)$$

Browder [8] proved that as $t \rightarrow 1$, z_t converges strongly to a fixed point of T which is closer to u , that is, the nearest point projection of u onto $F(T)$. In 1980, Reich [9] extended the result of Browder to a more general Banach spaces. Furthermore, Takahashi and Ueda [10] and Morales and Jung [11] improved results of Reich [9] to the class of continuous pseudocontractive mappings. For other results on pseudocontractive mappings, we refer to [12–15].

We note that the above methods can be used to find the minimum-norm fixed point x^* of T if $0 \in K$. However, if $0 \notin K$ neither Browder's, Reich's, Takahashi and Ueda's, nor Morales and Jung's method works to find minimum-norm fixed point of T .

Our concern is now the following: is it possible to construct a scheme, implicit or explicit, which converges strongly to the minimum-norm fixed point of T for any closed convex domain K of T ?

In this direction, Yang et al. [4] introduced an implicit and explicit iteration processes which converge strongly to the minimum-norm fixed point of nonexpansive self-mapping T , in real Hilbert spaces. In fact, they proved the following theorems.

Theorem YLY1 (see [4]). *Let K be a nonempty closed convex subset of a real Hilbert space H and $T : K \rightarrow K$ a nonexpansive mapping with $F(T) \neq \emptyset$. For $\beta \in (0, 1)$ and each $t \in (0, 1)$, let y_t be defined as the unique solution of fixed point equation:*

$$y_t = \beta T y_t + (1 - \beta) P_K [(1 - t) y_t], \quad t \in (0, 1). \quad (1.15)$$

Then the net $\{y_t\}$ converges strongly, as $t \rightarrow 0$, to the minimum-norm fixed point of T .

Theorem YLY2 (see [4]). *Let K be a nonempty closed convex subset of a real Hilbert space H , and let $T : K \rightarrow K$ be a nonexpansive mapping with $F(T) \neq \emptyset$. For a given $x_0 \in K$, define a sequence $\{x_n\}$ iteratively by*

$$x_{n+1} = \beta T x_n + (1 - \beta) P_K [(1 - \alpha_n) x_n], \quad n \geq 1, \quad (1.16)$$

where $\beta \in (0, 1)$ and $\alpha_n \in (0, 1)$, satisfying certain conditions. Then the sequence $\{x_n\}$ converges strongly to the minimum-norm fixed point of T .

A natural question arises whether the above theorems can be extended to a more general class of pseudocontractive mappings or not.

Let K be a closed convex subset a real Hilbert space H and let $T : K \rightarrow K$ be continuous pseudocontractive mapping.

It is our purpose in this paper to prove that for $\beta \in (0, 1)$ and each $t \in (0, 1)$, there exists a sequence $\{y_t\} \subset K$ satisfying $y_t = \beta P_K [(1 - t) y_t] + (1 - \beta) T(y_t)$ which converges strongly, as $t \rightarrow 0^+$, to the minimum-norm fixed point of T . Moreover, we provide an explicit iteration process which converges strongly to the minimum-norm fixed point of T provided that T is Lipschitz. Our theorems improve Theorem YLY1 and Theorem YLY2 of Yang et al. [4] and Theorems 3.1, and 3.2 of Cai et al. [16].

2. Preliminaries

In what follows, we shall make use of the following lemmas.

Lemma 2.1 (see [11]). *Let H be a real Hilbert space. Then, for any given $x, y \in H$, the following inequality holds:*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle. \quad (2.1)$$

Lemma 2.2 (see [17]). *Let K be a closed and convex subset of a real Hilbert space H . Let $x \in H$. Then $x_0 = P_K x$ if and only if*

$$\langle z - x_0, x - x_0 \rangle \leq 0, \quad \forall z \in K. \quad (2.2)$$

Lemma 2.3 (see [18]). Let $\{\lambda_n\}$, $\{\alpha_n\}$, and $\{\gamma_n\}$ be sequences of nonnegative numbers satisfying the conditions: $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\sum_{n=1}^{\infty} \alpha_n = \infty$, and $\gamma_n/\alpha_n \rightarrow 0$, as $n \rightarrow \infty$. Let the recursive inequality:

$$\lambda_{n+1} \leq \lambda_n - \alpha_n \psi(\lambda_{n+1}) + \gamma_n, \quad n = 1, 2, \dots, \quad (2.3)$$

be given where $\psi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing function such that it is positive on $(0, \infty)$ and $\psi(0) = 0$. Then $\lambda_n \rightarrow 0$, as $n \rightarrow \infty$.

Lemma 2.4 (see [3]). Let H be a real Hilbert space, K be a closed convex subset of H and $T : K \rightarrow K$ be a continuous pseudocontractive mapping, then

- (i) $F(T)$ is closed convex subset of K ;
- (ii) $(I - T)$ is demiclosed at zero, that is, if $\{x_n\}$ is a sequence in K such that $x_n \rightharpoonup x$ and $Tx_n - x_n \rightarrow 0$, as $n \rightarrow \infty$, then $x = T(x)$.

Lemma 2.5 (see [19]). Let H be a real Hilbert space. Then for all $x, y \in H$ and $\alpha \in [0, 1]$, the following equality holds:

$$\|\alpha x + (1 - \alpha)x\|^2 = \alpha\|x\|^2 + (1 - \alpha)\|y\|^2 - \alpha(1 - \alpha)\|x - y\|^2. \quad (2.4)$$

3. Main Results

Theorem 3.1. Let K be a nonempty closed and convex subset of a real Hilbert space H . Let $T : K \rightarrow K$ be a continuous pseudocontractive mapping with $F(T) \neq \emptyset$. Then for $\beta \in (0, 1)$ and each $t \in (0, 1)$, there exists a sequence $\{y_t\} \subset K$ satisfying the following condition:

$$y_t = \beta P_K[(1 - t)y_t] + (1 - \beta)T(y_t) \quad (3.1)$$

and the net $\{y_t\}$ converges strongly, as $t \rightarrow 0^+$, to the minimum-norm fixed point of T .

Proof. For $\beta \in (0, 1)$ and each $t \in (0, 1)$ let $T_t(y) := \beta P_K[(1 - t)y] + (1 - \beta)T(y)$. Then using nonexpansiveness of P_K and pseudocontractivity of T , for $x, y \in K$, we have that

$$\begin{aligned} \langle T_t x - T_t y, x - y \rangle &= \beta \langle P_K[(1 - t)x] - P_K[(1 - t)y], x - y \rangle \\ &\quad + (1 - \beta) \langle T(x) - T(y), x - y \rangle \\ &\leq \beta(1 - t) \|x - y\|^2 + (1 - \beta) \|x - y\|^2 \\ &\leq (1 - t\beta) \|x - y\|^2. \end{aligned} \quad (3.2)$$

This implies that T_t is strongly pseudocontractive on K . Thus, by Corollary 1 of [20] T_t has a unique fixed point, y_t , in K . This means that the equation:

$$y_t = \beta P_K[(1 - t)y_t] + (1 - \beta)T(y_t) \quad (3.3)$$

has a unique solution for each $t \in (0, 1)$. Furthermore, since $F(T) \neq \emptyset$, for $y^* \in F(T)$, we have that

$$\begin{aligned}
 \|y_t - y^*\|^2 &= \langle \beta P_K[(1-t)y_t] + (1-\beta)Ty_t - y^*, y_t - y^* \rangle \\
 &= \beta \langle P_K[(1-t)y_t] - P_K y^*, y_t - y^* \rangle + (1-\beta) \langle Ty_t - Ty^*, y_t - y^* \rangle \\
 &\leq \beta \|(1-t)y_t - y^*\| \cdot \|y_t - y^*\| + (1-\beta) \|y_t - y^*\|^2 \\
 &\leq \beta[(1-t)\|y_t - y^*\| + t\|y^*\|] \|y_t - y^*\| + (1-\beta) \|y_t - y^*\|^2,
 \end{aligned} \tag{3.4}$$

which implies that

$$\|y_t - y^*\| \leq \beta(1-t)\|y_t - y^*\| + \beta t\|y^*\| + (1-\beta)\|y_t - y^*\|, \tag{3.5}$$

and hence $\|y_t - y^*\| \leq \|y^*\|$. Therefore, $\{y_t\}$ and hence $\{Ty_t\}$ is bounded.

Furthermore, from (3.3) and using nonexpansiveness of P_K we get that

$$\begin{aligned}
 \|y_t - Ty_t\| &= \|\beta P_K[(1-t)y_t] + (1-\beta)Ty_t - Ty_t\| \\
 &= \beta \|P_K[(1-t)y_t] - P_K Ty_t\| \\
 &\leq \beta \|(1-t)y_t - Ty_t\| \\
 &\leq \beta \|y_t - Ty_t\| + \beta t\|y_t\|,
 \end{aligned} \tag{3.6}$$

which implies that

$$\|y_t - Ty_t\| \leq \frac{\beta}{(1-\beta)} t \|y_t\| \longrightarrow 0, \quad \text{as } t \longrightarrow 0. \tag{3.7}$$

Furthermore, from (3.3), convexity of $\|\cdot\|^2$, (1.4), and (3.7), we get that

$$\begin{aligned}
 \|y_t - y^*\|^2 &= \|(1-\beta)(Ty_t - y^*) + \beta(P_K[(1-t)y_t] - P_K y^*)\|^2 \\
 &= (1-\beta)\|Ty_t - y^*\|^2 + \beta\|P_K[(1-t)y_t] - P_K y^*\|^2 \\
 &\leq (1-\beta)[\|y_t - y^*\|^2 + \|Ty_t - y_t\|^2] + \beta\|(1-t)y_t - y^*\|^2 \\
 &\leq (1-\beta)\|y_t - y^*\|^2 + (1-\beta)\|Ty_t - y_t\|^2 + \beta\|(1-t)y_t - y^*\|^2 \\
 &\leq (1-\beta)\|y_t - y^*\|^2 + \frac{\beta^2}{(1-\beta)} t^2 \|y_t\|^2 \\
 &\quad + \beta[\|y_t - y^*\|^2 - 2t\|y_t - y^*\|^2 - 2t\langle y^*, y_t - y^* \rangle + t^2 \|y_t\|^2].
 \end{aligned} \tag{3.8}$$

This implies that

$$\|y_t - y^*\|^2 \leq \langle y^*, y^* - y_t \rangle + tM, \quad \text{for some } M > 0. \quad (3.9)$$

Now, for $t_n \rightarrow 0$, as $n \rightarrow \infty$, let $\{y_n := y_{t_n}\}$ be a subsequence of $\{y_t\}$ such that $y_n \rightarrow y'$. Then, we have from (3.7) and Lemma 2.4 that $y' \in F(T)$. Furthermore, replacing y^* by y' in (3.9) and the fact that $y_n \rightarrow y'$ imply that

$$\|y_n - y'\|^2 \leq \langle y', y' - y_n \rangle + t_n M \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (3.10)$$

which implies that

$$y_n \rightarrow y', \quad \text{as } n \rightarrow \infty. \quad (3.11)$$

Thus, from (3.9) and (3.11), we have that

$$\|y' - y^*\|^2 \leq \langle y^*, y^* - y' \rangle, \quad \text{as } n \rightarrow \infty, \quad (3.12)$$

which is equivalent to the inequality:

$$\langle y', y^* - y' \rangle \geq 0 \quad \text{and hence } y' = P_F 0. \quad (3.13)$$

If there is another subsequence $\{y_m\}$ of $\{y_t\}$ such that $y_m \rightarrow y''$, similar argument gives that $y'' = P_F 0$, which implies, by uniqueness of $P_F 0$, that $y'' = y'$. Therefore, the net $y_t \rightarrow y' = P_F 0$ which is the minimum-norm of fixed point of T . The proof is complete. $\square$

We now state and prove a convergence theorem for the minimum-norm zero of a monotone mapping A .

Theorem 3.2. *Let H be a real Hilbert space. Let $A : H \rightarrow H$ be a continuous monotone mapping with $N(A) \neq \emptyset$. Then for $\beta \in (0, 1)$ and each $t \in (0, 1)$, there exists a sequence $\{y_t\} \subset H$ satisfying the following condition:*

$$y_t = \beta(1 - t)y_t + (1 - \beta)(I - A)y_t, \quad (3.14)$$

and the net $\{y_t\}$ converges strongly, as $t \rightarrow 0^+$, to the minimum-norm zero of A .

Proof. Let $Tx := (I - A)x$. Then, we get that T is continuous pseudocontractive mapping with $F(T) = N(A) \neq \emptyset$. Moreover, since P_H is an identity mapping on H , when A is replaced with $(I - T)$ scheme (3.14) reduces to scheme (3.1), and hence the conclusion follows from Theorem 3.1. $\square$

If in Theorem 3.1, we consider $\{t_n\}, \{\beta_n\} \subset (0, 1)$ such that $t_n \rightarrow 0$, $\beta_n \rightarrow 0$ and $y_n := y_{t_n}$, the method of proof of Theorem 3.1 provides the following corollary.

Corollary 3.3. *Let K be a nonempty closed and convex subset of a real Hilbert space H . Let $T : K \rightarrow K$ be continuous pseudocontractive mapping with $F(T) \neq \emptyset$. Then the sequence $\{y_n\} \subset K$ defined by*

$$y_n = \beta_n P_K[(1 - t_n)y_n] + (1 - \beta_n)T(y_n), \quad (3.15)$$

where $\{t_n\}, \{\beta_n\} \subset (0, 1)$ such that $t_n \rightarrow 0, \beta_n \rightarrow 0$, as $n \rightarrow \infty$, converges strongly, as $n \rightarrow \infty$, to the minimum-norm fixed point of T .

The following proposition and lemma play an important role in proving the next theorem.

Proposition 3.4. *Let K be a nonempty closed and convex subset of a real Hilbert space H . Let $T : K \rightarrow K$ be continuous pseudocontractive mapping. Then the sequence $\{y_n\}$ in (3.15) satisfies the following inequality:*

$$\|y_n - y_{n-1}\| \leq \frac{|\theta_{n-1} - \theta_n|}{\theta_n t_n} [\|y_n\| + \|P_K[(1 - t_n)y_{n-1}]\|] + \frac{\theta_{n-1}}{\theta_n} \frac{|t_n - t_{n-1}|}{t_n} \|y_{n-1}\|, \quad (3.16)$$

where $\theta_n := \beta_n / (1 - \beta_n)$ for $\{\beta_n\}$ decreasing sequence.

Proof. If we put $\theta_n := \beta_n / (1 - \beta_n)$, (3.15) reduces to

$$y_n = Ty_n + \theta_n(P_K[(1 - t_n)y_n] - y_n). \quad (3.17)$$

Thus, using pseudocontractivity of T and nonexpansiveness of P_K we get that

$$\begin{aligned} \|y_n - y_{n-1}\|^2 &= \|Ty_n + \theta_n(P_K[(1 - t_n)y_n] - y_n) - Ty_{n-1} - \theta_{n-1}(P_K[(1 - t_{n-1})y_{n-1}] - y_{n-1})\|^2 \\ &= \|Ty_n - Ty_{n-1} + \theta_{n-1}y_{n-1} - \theta_n y_n + \theta_{n-1}y_n - \theta_{n-1}y_n \\ &\quad + \theta_n P_K[(1 - t_n)y_n] - \theta_{n-1} P_K[(1 - t_{n-1})y_{n-1}]\|^2 \\ &= \langle Ty_n - Ty_{n-1} + \theta_{n-1}(y_{n-1} - y_n) + (\theta_{n-1} - \theta_n) y_n, y_n - y_{n-1} \rangle \\ &\quad + \langle \theta_n P_K[(1 - t_n)y_n] - \theta_{n-1} P_K[(1 - t_{n-1})y_{n-1}], y_n - y_{n-1} \rangle \\ &\quad + \langle \theta_n P_K[(1 - t_n)y_{n-1}] - \theta_{n-1} P_K[(1 - t_n)y_{n-1}], y_n - y_{n-1} \rangle \\ &\quad + \langle \theta_{n-1} P_K[(1 - t_n)y_{n-1}] - \theta_{n-1} P_K[(1 - t_{n-1})y_{n-1}], y_n - y_{n-1} \rangle \\ &\leq \|y_n - y_{n-1}\|^2 - \theta_{n-1} \|y_n - y_{n-1}\|^2 + (\theta_{n-1} - \theta_n) \|y_n\| \\ &\quad \times \|y_n - y_{n-1}\| + \theta_n (1 - t_n) \|y_n - y_{n-1}\|^2 \\ &\quad + (\theta_n - \theta_{n-1}) \|P_K[(1 - t_n)y_{n-1}]\| \cdot \|y_{n-1} - y_n\| \\ &\quad + \theta_{n-1} |t_n - t_{n-1}| \cdot \|y_{n-1}\| \|y_n - y_{n-1}\|, \end{aligned} \quad (3.18)$$

which implies, using the fact that θ_n is decreasing, that

$$\begin{aligned}
\|y_n - y_{n-1}\| &\leq [1 - \theta_{n-1} + \theta_n(1 - t_n)]\|y_n - y_{n-1}\| + |\theta_{n-1} - \theta_n|[\|y_n\| + \|P_K[(1 - t_n)y_{n-1}]\|] \\
&\quad + \theta_{n-1}|t_n - t_{n-1}| \cdot \|y_{n-1}\| \\
&\leq (1 - t_n\theta_n)\|y_n - y_{n-1}\| + |\theta_{n-1} - \theta_n|[\|y_n\| + \|P_K[(1 - t_n)y_{n-1}]\|] \\
&\quad + \theta_{n-1}|t_n - t_{n-1}| \cdot \|y_{n-1}\|,
\end{aligned} \tag{3.19}$$

and hence

$$\|y_n - y_{n-1}\| \leq \frac{|\theta_{n-1} - \theta_n|}{\theta_n t_n} [\|y_n\| + \|P_K[(1 - t_n)y_{n-1}]\|] + \frac{\theta_{n-1}}{\theta_n} \frac{|t_n - t_{n-1}|}{t_n} \|y_{n-1}\|. \tag{3.20}$$

The proof is complete. $\square$

For the rest of this paper, let $\{\lambda_n\}$, $\{\theta_n\}$ (decreasing) and $\{t_n\}$ be real sequences in $(0, 1]$ satisfying the following conditions: (i) $\lim_{n \rightarrow \infty} \theta_n = 0 = \lim_{n \rightarrow \infty} t_n$; (ii) $\lambda_n(1 + \theta_n) \leq 1$, $\sum \lambda_n \theta_n t_n = \infty$, $\lim_{n \rightarrow \infty} \lambda_n / \theta_n t_n = 0$; (iii) $\lim_{n \rightarrow \infty} [\theta_{n-1} - \theta_n] / \lambda_n \theta_n^2 t_n^2 = 0$ and $\lim_{n \rightarrow \infty} [t_{n-1} - t_n] / \lambda_n \theta_n t_n^2 = 0$. Examples of real sequences which satisfy these conditions are $\lambda_n = 1/(n + 1)^{1/2}$, $\theta_n = 1/(n + 1)^{1/3}$ and $t_n = 1/(n + 1)^{1/4}$.

Lemma 3.5. *Let K be a nonempty closed convex subset of a real Hilbert space H . Let $T : K \rightarrow K$ be a Lipschitz pseudocontractive mapping with Lipschitz constant $L \geq 0$ and $F(T) \neq \emptyset$. Let a sequence $\{x_n\}$ be generated from arbitrary $x_1 \in K$ by*

$$x_{n+1} := (1 - \lambda_n)x_n + \lambda_n T x_n - \lambda_n \theta_n (x_n - P_K[(1 - t_n)x_n]), \tag{3.21}$$

for all positive integers $n \geq 1$. Then $\{x_n\}$ is bounded.

Proof. We follow the method of proof of Chidume and Zegeye [21]. Since $\lambda_n / (\theta_n t_n) \rightarrow 0$, there exists $N_0 > 0$ such that $\lambda_n / (\theta_n t_n) \leq d := 1/(2(3 + L)^2)$, for all $n \geq N_0$. Let $x^* \in F(T)$ and $r > 0$ be sufficiently large such that $x_{N_0} \in B_r(x^*)$ and $\|x^*\| \leq r/(2(4 + L))$. Now, we show by induction that $\{x_n\}$ belongs to $B := \overline{B_r(x^*)}$ for all integers $n \geq N_0$. By construction, we have $x_{N_0} \in B$. Assume that $x_n \in B$ for any $n > N_0$. Then, we prove that $x_{n+1} \in B$. Suppose x_{n+1} is

not in B . Then $\|x_{n+1} - x^*\| > r$, and thus from the recursion formula (1.2) and Lemma 2.1 we get that

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &= \|x_n - x^* - \lambda_n((x_n - Tx_n) + \theta_n(x_n - P_K[(1 - t_n)x_n]))\|^2 \\
 &= \|x_n - x^*\|^2 - 2\lambda_n\langle x_n - Tx_n \\
 &\quad + \theta_n(x_n - P_K[(1 - t_n)x_n]), j(x_{n+1} - x^*) \rangle \\
 &= \|x_n - x^*\|^2 - 2\lambda_n\theta_n\langle x_{n+1} - x^*, x_{n+1} - x^* \rangle \\
 &\quad + 2\lambda_n\langle \theta_n(x_{n+1} - x_n) - (x_n - Tx_n) + \theta_n(P_K[(1 - t_n)x_n] - x^*) \\
 &\quad + (x_{n+1} - Tx_{n+1}) - (x_{n+1} - Tx_{n+1}), j(x_{n+1} - x^*) \rangle.
 \end{aligned} \tag{3.22}$$

Since T is pseudocontractive we have $\langle x_{n+1} - Tx_{n+1}, j(x_{n+1} - x^*) \rangle \geq 0$. Thus, (3.22) gives

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq \|x_n - x^*\|^2 - 2\lambda_n\theta_n\|x_{n+1} - x^*\|^2 \\
 &\quad + 2\lambda_n(2 + L)\|x_{n+1} - x_n\| \cdot \|x_{n+1} - x^*\| \\
 &\quad + 2\lambda_n\theta_n\langle P_K[(1 - t_n)x_n] - x^*, j(x_{n+1} - x^*) \rangle \\
 &= \|x_n - x^*\|^2 - 2\lambda_n\theta_n\|x_{n+1} - x^*\|^2 \\
 &\quad + 2\lambda_n(2 + L)\|x_{n+1} - x_n\| \cdot \|x_{n+1} - x^*\| \\
 &\quad + 2\lambda_n\theta_n\langle P_K[(1 - t_n)x_n] - P_K[(1 - t_n)x_{n+1}] + P_K[(1 - t_n)x_{n+1}] \\
 &\quad - P_K[(1 - t_n)x^*] + P_K[(1 - t_n)x^*] - x^*, j(x_{n+1} - x^*) \rangle,
 \end{aligned} \tag{3.23}$$

which implies that

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq \|x_n - x^*\|^2 - 2\lambda_n\theta_nt_n\|x_{n+1} - x^*\|^2 \\
 &\quad + 2\lambda_n(2 + L + (1 - t_n))\|x_{n+1} - x_n\| \cdot \|x_{n+1} - x^*\| \\
 &\quad + 2\lambda_n\theta_n\|P_K[(1 - t_n)x^*] - x^*\| \cdot \|x_{n+1} - x^*\| \\
 &= \|x_n - x^*\|^2 - 2\lambda_n\theta_nt_n\|x_{n+1} - x^*\|^2 \\
 &\quad + 2\lambda_n(3 + L)[\lambda_n\theta_n(P_K[(1 - t_n)x_n] - P_K[(1 - t_n)x^*] \\
 &\quad + P_K[(1 - t_n)x^*] - x^* + x^* - x_n) + Tx_n - Tx^* + x^* - x_n] \\
 &\quad \times \|x_{n+1} - x^*\| + 2\lambda_n\theta_nt_n\|x^*\| \cdot \|x_{n+1} - x^*\| \\
 &\leq \|x_n - x^*\|^2 - 2\lambda_n\theta_nt_n\|x_{n+1} - x^*\|^2 \\
 &\quad + 2\lambda_n^2(3 + L)^2\|x_n - x^*\| \cdot \|x_{n+1} - x^*\| \\
 &\quad + 2\lambda_n\theta_nt_n(4 + L)\|x^*\|\|x_{n+1} - x^*\|.
 \end{aligned} \tag{3.24}$$

Since $\|x_{n+1} - x^*\| > \|x_n - x^*\|$, from (3.24) we get that

$$\|x_{n+1} - x^*\| \leq \frac{\lambda_n}{\theta_n t_n} (3 + L)^2 \|x_n - x^*\| + (4 + L) \|x^*\|, \quad (3.25)$$

and hence $\|x_{n+1} - x^*\| \leq r$, since $x_n \in B$, $\|x^*\| \leq r/(2(4 + L))$ and $\lambda_n/\theta_n t_n \leq 1/2(3 + L)^2$ for all $n \geq N_0$. But this is a contradiction. Therefore, $x_n \in B$ for all positive integers $n \geq N_0$, and hence the sequence $\{x_n\}$ is bounded. $\square$

For the next theorem, let $\{y_n\}$ denotes the sequence defined by $y_n := y_{s_n} = s_n T y_{s_n} + (1 - s_n) P_K[(1 - t_n) y_n]$, $s_n = 1/(1 + \theta_n)$, for all $n \geq 1$, guaranteed by Corollary 3.3 (which reduces to $\theta_n(P_K[(1 - t_n) y_n] - y_n) - (y_n - T y_n) = 0$).

Theorem 3.6. *Let K be a nonempty closed convex subset of a real Hilbert space H . Let $T : K \rightarrow K$ be a Lipschitz pseudocontractive mapping with Lipschitz constant $L \geq 0$ and $F(T) \neq \emptyset$. Let a sequence $\{x_n\}$ be generated from arbitrary $x_1 \in K$ by*

$$x_{n+1} := (1 - \lambda_n) x_n + \lambda_n T x_n - \lambda_n \theta_n (x_n - P_K[(1 - t_n) x_n]), \quad (3.26)$$

for all positive integers $n \geq 1$. Then $\{x_n\}$ converges strongly to the minimum-norm fixed point of T , as $n \rightarrow \infty$.

Proof. By Lemma 3.5, we have that the sequence $\{x_n\}$ is bounded. Now, we show that it converges strongly to a minimum-norm fixed point of T . But from (3.26) and Lemma 2.1, we have that

$$\begin{aligned} \|x_{n+1} - y_n\|^2 &\leq \|x_n - y_n\|^2 - 2\lambda_n \theta_n \langle x_{n+1} - y_n, j(x_{n+1} - y_n) \rangle \\ &\quad + 2\lambda_n \langle \theta_n (x_{n+1} - y_n) - (x_n - T x_n) \\ &\quad - \theta_n (x_n - P_K[(1 - t_n) x_n]), j(x_{n+1} - y_n) \rangle \\ &= \|x_n - y_n\|^2 - 2\lambda_n \theta_n \|x_{n+1} - y_n\|^2 + 2\lambda_n \langle \theta_n (x_{n+1} - x_n) \\ &\quad + [\theta_n (P_K[(1 - t_n) y_n] - y_n) - (y_n - T y_n)] - [(x_{n+1} - T x_{n+1}) \\ &\quad - (y_n - T y_n)] + \theta_n (P_K[(1 - t_n) x_n] - P_K[(1 - t_n) y_n]) \\ &\quad + [(x_{n+1} - T x_{n+1}) - (x_n - T x_n)], j(x_{n+1} - y_n) \rangle. \end{aligned} \quad (3.27)$$

Observe that by the property of y_n and pseudocontractivity of T we have $\theta_n(P_K[(1 - t_n) y_n] - y_n) - (y_n - T y_n) = 0$ (see (3.17)) and $\langle (x_{n+1} - T x_{n+1}) - (y_n - T y_n), j(x_{n+1} - y_n) \rangle \geq 0$ for all $n \geq 1$. Thus, we have from (3.27) that

$$\begin{aligned} \|x_{n+1} - y_n\|^2 &\leq \|x_n - y_n\|^2 - 2\lambda_n \theta_n \|x_{n+1} - y_n\|^2 + 2\lambda_n \langle \theta_n (x_{n+1} - x_n) \\ &\quad + \theta_n (P_K[(1 - t_n) x_n] - P_K[(1 - t_n) x_{n+1}]) \\ &\quad + P_K[(1 - t_n) x_{n+1}] - P_K[(1 - t_n) y_n] \rangle \\ &\quad + (x_{n+1} - T x_{n+1}) - (x_n - T x_n), j(x_{n+1} - y_n) \rangle \end{aligned}$$

$$\begin{aligned}
&\leq \|x_n - y_n\|^2 - 2\lambda_n\theta_n t_n \|x_{n+1} - y_n\|^2 \\
&\quad + 2\lambda_n(3+L)\|x_{n+1} - x_n\| \cdot \|x_{n+1} - y_n\|.
\end{aligned} \tag{3.28}$$

But by Corollary 3.3, we have that $\{y_n\}$ is bounded. Therefore, there exists $M_1 > 0$ such that $\max\{(3+L)\|x_{n+1} - y_n\| \cdot \|x_n - Tx_n + \theta_n(x_n - P_K[(1-t_n)x_n])\|\} \leq M_1$. Thus from (3.28), we get that

$$\|x_{n+1} - y_n\|^2 \leq \|x_n - y_n\|^2 - 2\lambda_n\theta_n t_n \|x_{n+1} - y_n\|^2 + 2\lambda_n^2 M_1. \tag{3.29}$$

But using triangle inequality and Proposition 3.4, we have that

$$\begin{aligned}
\|x_n - y_n\|^2 &\leq [\|x_n - y_{n-1}\| + \|y_{n-1} - y_n\|]^2 \\
&\leq \|x_n - y_{n-1}\|^2 + \|y_{n-1} - y_n\|^2 M_2 \\
&\leq \|x_n - y_{n-1}\|^2 + \frac{|\theta_{n-1} - \theta_n|}{\theta_n t_n} M_3 + \frac{|t_n - t_{n-1}|}{t_n} M_3,
\end{aligned} \tag{3.30}$$

for some $M_2, M_3 > 0$, and for all $n \geq N_0$. Now, substituting (3.30) in (3.29) we obtain that

$$\begin{aligned}
\|x_{n+1} - y_n\|^2 &\leq \|x_n - y_{n-1}\|^2 - 2\lambda_n\theta_n t_n \|x_{n+1} - y_n\|^2 \\
&\quad + 2\lambda_n^2 M_4 + \frac{\theta_{n-1} - \theta_n}{\theta_n t_n} M_4 + \frac{|t_n - t_{n-1}|}{t_n} M_4,
\end{aligned} \tag{3.31}$$

for some constant $M_4 > 0$. Now, by Lemma 2.3 and the conditions on $\{\lambda_n\}$, $\{\theta_n\}$, and $\{t_n\}$ we get $x_{n+1} - y_n \rightarrow 0$. Consequently, $\|x_n - y_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Therefore, since by Corollary 3.3 we have that $y_t \rightarrow y^* \in F(T)$, where y^* is with the minimum-norm in $F(T)$, we get that $\{x_n\}$ converges strongly to the minimum-norm of fixed point of T . $\square$

Corollary 3.7. *Let H be a real Hilbert space. Let $A : H \rightarrow H$ be a Lipschitz monotone mapping with Lipschitz constant $L \geq 0$ and $N(A) \neq \emptyset$. Let a sequence $\{x_n\}$ be generated from arbitrary $x_1 \in H$ by*

$$x_{n+1} = x_n - \lambda_n A x_n + \lambda_n \theta_n t_n x_n, \tag{3.32}$$

for all positive integers n . Then $\{x_n\}$ converges strongly to the minimum-norm solution of the equation $Ax = 0$.

Proof. Let $T := (I - A)$. Then T is a Lipschitz pseudocontractive mapping with Lipschitz constant $L' := (L + 1)$, and the minimum-norm solution of the equation $Ax = 0$ is the minimum-norm fixed point of T . Moreover, if we replace T by $(I - A)$ in (3.26), then the equation reduces to (3.32). Thus, the conclusion follows from Theorem 3.6. $\square$

4. Applications

For the rest of this paper, let H be a Hilbert space and $A : H \rightarrow H$ a bounded linear operator. Consider the convexly constrained linear inverse problem, which has extensively been discussed in the literature (see, e.g., [22]), given by:

$$x \in K, \quad Ax = b, \quad (4.1)$$

where K is closed and convex subset of H and $b \in H$, which is a special case of the SFP problem (1.9). Set

$$\varphi(x) := \frac{1}{2} \|Ax - b\|^2. \quad (4.2)$$

The least-square solution of (4.1) is the least-norm minimizer of the minimization problem (4.2). Let Ω denote the solution set of (4.2). It is known that Ω is nonempty if and only if $P_{\overline{A(K)}}(b) \in A(K)$. In this case, Ω has a unique element with minimum norm which is a least-square solution of (4.1), that is, there exists a unique point $x^* \in \Omega$ such that

$$\|x^*\| = \min\{\|x\| : x \in \Omega\}. \quad (4.3)$$

We note that $\varphi(x)$ is a quadratic function with gradient:

$$\nabla\varphi(x) = A^*(Ax - b), \quad (4.4)$$

where A^* is adjoint of A . Let $\gamma > 0$ and $x^* \in \Omega$. Thus, x^* is the minimum-norm solution of the minimization problem (4.2) if and only if x^* a solution of

$$\gamma \nabla\varphi(x) = \gamma A^*(Ax - b) = 0. \quad (4.5)$$

Now, we state applications of our theorems.

Theorem 4.1. *Assume that the solution set of convexly constrained linear inverse problem (4.1) with $K := H$, a real Hilbert space, is nonempty and that $\nabla\varphi$ is monotone. Then for $\beta \in (0, 1)$ and each $t \in (0, 1)$, there exists a sequence $\{y_t\} \subset H$ satisfying the following condition:*

$$y_t = \beta(1 - t)y_t + (1 - \beta)(y_t - \gamma A^*(Ay_t - b)), \quad (4.6)$$

where A^* is adjoint of A , and the net $\{y_t\}$ converges strongly, as $t \rightarrow 0^+$, to the minimum-norm solution of the split feasibility problem (4.1).

Proof. We note that $\varphi(x)$ is continuously differentiable function with gradient:

$$\nabla\varphi(x) = A^*(Ax - b), \quad (4.7)$$

where A^* is adjoint of A , which is Lipschitz (see Lemma 8.1 of [5]) and monotone (by hypothesis). Thus, the conclusion follows from Theorem 3.2. $\square$

Theorem 4.2. Assume that the solution set of split feasibility problem (4.1) is nonempty and that $\nabla\varphi$ with $K := H$, a real Hilbert space, is monotone. Let a sequence $\{x_n\}$ be generated from arbitrary $x_1 \in E$ by

$$x_{n+1} = x_n - \lambda_n \gamma A^*(Ax_n - b) + \lambda_n \theta_n t_n x_n, \quad (4.8)$$

for all positive integers n , where $\gamma > 0$ and A^* is adjoint of A . Then, $\{x_n\}$ converges strongly to the minimum-norm solution of the split feasibility problem (4.1).

Remark 4.3. Theorem 3.1 improves Theorem YLY1 and Theorem 3.1 of Cai et al. [16] to a more general class of pseudocontractive mappings. Moreover, Theorem 3.6 improves Theorem YLY1 and Theorem 3.2 of Cai et al. [16] in the sense that our scheme provides a minimum-norm fixed point of pseudocontractive mapping T .

References

- [1] R. P. Agarwal, D. O'Regan, and D. R. Sahu, *Fixed Point Theory for Lipschitzian-type Mappings with Applications*, Springer, Dordrecht, The Netherlands, 2000.
- [2] D. R. Sahu and A. Petrușel, "Strong convergence of iterative methods by strictly pseudocontractive mappings in Banach spaces," *Nonlinear Analysis*, vol. 74, no. 17, pp. 6012–6023, 2011.
- [3] Q. B. Zhang and C. Z. Cheng, "Strong convergence theorem for a family of Lipschitz pseudocontractive mappings in a Hilbert space," *Mathematical and Computer Modelling*, vol. 48, no. 3–4, pp. 480–485, 2008.
- [4] X. Yang, Y.-C. Liou, and Y. Yao, "Finding minimum norm fixed point of nonexpansive mappings and applications," *Mathematical Problems in Engineering*, vol. 2011, Article ID 106450, 13 pages, 2011.
- [5] C. Byrne, "A unified treatment of some iterative algorithms in signal processing and image reconstruction," *Inverse Problems*, vol. 20, no. 1, pp. 103–120, 2004.
- [6] M. I. Sezan and H. Stark, "Applications of convex projection theory to image recovery in tomography and related areas," in *Image Recovery Theory and Applications*, H. Stark, Ed., pp. 415–462, Academic Press, Orlando, Fla, USA, 1987.
- [7] D. Youla, "Mathematical theory of image restoration by the method of convex projections," in *Image Recovery Theory and Applications*, H. Stark, Ed., pp. 29–77, Academic Press, Orlando, Fla, USA, 1987.
- [8] F. E. Browder, "Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces," *Archive for Rational Mechanics and Analysis*, vol. 24, pp. 82–90, 1967.
- [9] S. Reich, "Strong convergence theorems for resolvents of accretive operators in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 75, no. 1, pp. 287–292, 1980.
- [10] W. Takahashi and Y. Ueda, "On Reich's strong convergence theorems for resolvents of accretive operators," *Journal of Mathematical Analysis and Applications*, vol. 104, no. 2, pp. 546–553, 1984.
- [11] C. H. Morales and J. S. Jung, "Convergence of paths for pseudocontractive mappings in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 128, no. 11, pp. 3411–3419, 2000.
- [12] E. U. Ofoedu and H. Zegeye, "Further investigation on iteration processes for pseudocontractive mappings with application," *Nonlinear Analysis*, vol. 75, no. 1, pp. 153–162, 2012.
- [13] E. U. Ofoedu and H. Zegeye, "Iterative algorithm for multi-valued pseudocontractive mappings in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 372, no. 1, pp. 68–76, 2010.
- [14] H. Zegeye, N. Shahzad, and T. Mekonen, "Viscosity approximation methods for pseudocontractive mappings in Banach spaces," *Applied Mathematics and Computation*, vol. 185, no. 1, pp. 538–546, 2007.
- [15] H. Zegeye, N. Shahzad, and M. A. Alghamdi, "Convergence of Ishikawa's iteration method for pseudocontractive mappings," *Nonlinear Analysis*, vol. 74, no. 18, pp. 7304–7311, 2011.
- [16] Y. Cai, Y. Tang, and L. Liu, "Iterative algorithms for minimum-norm fixed point of nonexpansive mapping in Hilbert space," *Fixed Point Theory and Applications*, vol. 2012, p. 49, 2012.
- [17] W. Takahashi, *Nonlinear Functional Analysis*, Kindikagaku, Tokyo, Japan, 1988.

- [18] C. E. Chidume, H. Zegeye, and S. J. Aneke, "Approximation of fixed points of weakly contractive nonself maps in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 270, no. 1, pp. 189–199, 2002.
- [19] H. Zegeye and N. Shahzad, "Convergence of Mann's type iteration method for generalized asymptotically nonexpansive mappings," *Computers & Mathematics with Applications*, vol. 62, no. 11, pp. 4007–4014, 2011.
- [20] K. Deimling, "Zeros of accretive operators," *Manuscripta Mathematica*, vol. 13, pp. 365–374, 1974.
- [21] C. E. Chidume and H. Zegeye, "Approximate fixed point sequences and convergence theorems for Lipschitz pseudocontractive maps," *Proceedings of the American Mathematical Society*, vol. 132, no. 3, pp. 831–840, 2004.
- [22] L. Landweber, "An iteration formula for Fredholm integral equations of the first kind," *American Journal of Mathematics*, vol. 73, pp. 615–624, 1951.