Research Article

Multiple Solutions for a Class of Differential Inclusion System Involving the $(p(x), q(x))$ -Laplacian

Bin Ge and Ji-Hong Shen

Department of Mathematics, Harbin Engineering University, Harbin 150001, China

Correspondence should be addressed to Bin Ge, gebin04523080261@163.com

Received 29 December 2011; Revised 28 February 2012; Accepted 2 April 2012

Academic Editor: Kanishka Perera

Copyright © 2012 B. Ge and J.-H. Shen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider a differential inclusion system involving the $(p(x), q(x))$ -Laplacian with Dirichlet boundary condition on a bounded domain and obtain two nontrivial solutions under appropriate hypotheses. Our approach is variational and it is based on the nonsmooth critical point theory for locally Lipschitz functions.

1. Introduction

In recent years, the study of differential equations and variational problems with $p(x)$ -growth conditions has been a new and interesting topic, which arises from nonlinear electrorheological fluids (see [1]) and elastic mechanics (see [2]). The study on variable exponent problems attracts more and more interest in recent years, and many results have been obtained on this kind of problems, for example [3–11].

Elliptic systems with standard growth conditions have been the subject of a sizeable literature. We refer to the excellent survey article of de Figueiredo [12].

In [11], the author obtained the existence and multiplicity of solutions for the following problem:

$$\begin{aligned} -\operatorname{div}\left(|\nabla u|^{p(x)-2}\nabla u\right) &= F_u(x, u, v), \quad \text{in } \Omega, \\ -\operatorname{div}\left(|\nabla v|^{q(x)-2}\nabla v\right) &= F_v(x, u, v), \quad \text{in } \Omega, \\ u &= v = 0, \quad \text{on } \partial\Omega, \end{aligned} \tag{P_1}$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with a smooth boundary $\partial\Omega$, $N \geq 2$, $(p, q) \in C(\overline{\Omega})^2$, $p(x) > 1$, $q(x) > 1$, for every $x \in \Omega$. The function F is assumed to be continuous in $x \in \Omega$ and of class C^1 in $u, v \in \mathbb{R}$. More precisely, the author was able to prove that, under suitable conditions, the system might have at least one solution or have infinite number of solutions.

Since many free boundary problems and obstacle problems may be reduced to partial differential equations with discontinuous nonlinearities, now a question arises: whether there exist solutions for system (P_1) in the case where there is no continuously differentiable hypothesis required on the potential function F with respect to $t(s)$. See, for example, $F(x, t, s)$ is locally Lipschitz with respect to $t(s)$. That is the main problem which we want to solve in the present paper.

To this end, we mainly discuss the existence and multiplicity of solutions for the following nonlinear differential inclusion system involving the $(p(x), q(x))$ -Laplacian:

$$\begin{aligned} -\operatorname{div}\left(|\nabla u|^{p(x)-2}\nabla u\right) &\in \lambda\partial_u F(x, u, v), \quad \text{in } \Omega, \\ -\operatorname{div}\left(|\nabla v|^{q(x)-2}\nabla v\right) &\in \lambda\partial_v F(x, u, v), \quad \text{in } \Omega, \\ u = v = 0, \quad &\text{on } \partial\Omega, \end{aligned} \tag{P}$$

where $\Omega \subset \mathbb{R}^N$ is a bounded domain with C^1 -boundary $\partial\Omega$, $\lambda > 0$ is the parameter, $p, q \in C(\overline{\Omega})$, $1 < p^- \leq p^+ < +\infty$, $1 < q^- \leq q^+ < +\infty$, $F : \Omega \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a function such that $F(\cdot, t, s)$ is measurable in Ω for all $(t, s) \in \mathbb{R} \times \mathbb{R}$, and $F(x, t, s)$ is locally Lipschitz with respect to $t(s)$ (in general it can be nonsmooth), $\partial_t F(x, t, s)(\partial_s F(x, t, s))$ is the subdifferential with respect to the $t(s)$ -variable in the sense of Clarke [13].

We emphasize that the operator $-\operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ is said to be $p(x)$ -Laplacian, which becomes p -Laplacian when $p(x) \equiv p$ (a constant). The $p(x)$ -Laplacian possesses more complicated nonlinearities than the p -Laplacian, for example, it is inhomogeneous and, in general, it does not have the first eigenvalue. In other words, the infimum of the eigenvalues of $p(x)$ -Laplacian equals 0 (see [14]).

Specially, if $F(x, \cdot, v), F(x, u, \cdot) \in C^1(\mathbb{R})$ for a.a. $x \in \Omega$, and $p(x) = p$, $q(x) = q$, then the problem (P) becomes the following problem:

$$\begin{aligned} -\operatorname{div}\left(|\nabla u|^{p-2}\nabla u\right) &= \lambda F_u(x, u, v), \quad \text{in } \Omega, \\ -\operatorname{div}\left(|\nabla v|^{q-2}\nabla v\right) &= \lambda F_v(x, u, v), \quad \text{in } \Omega, \\ u = v = 0, \quad &\text{on } \partial\Omega. \end{aligned} \tag{P_2}$$

There have been a large number of papers that study the existence of the solutions to (P_2) . For instance, when $p > N$, $q > N$, Li and Tang [15] ensured the existence of three solutions to this problem. In [16], Kristály studied the multiplicity of solutions of the quasilinear elliptic systems (P_1) , where Ω is a strip-like domain and $\lambda > 0$ is the parameter. Under some growth conditions on F , the author guaranteed the existence of an open interval $\Lambda \subset [0, +\infty)$, such that for each $\lambda \in \Lambda$ problem (P_2) has at least two distinct nontrivial solutions; when p and q are real numbers larger than 1, $\lambda = 1$, Boccardo and Guedes de Figueiredo [17] obtained the existence of solutions of the system (P_2) .

But up to now, to the best of our knowledge, no paper discussing the solutions of problem (P) with nonsmooth potential via nonsmooth critical point theory can be found in the existing literature. In order to fill in this gap, we study problem (P) from a more extensive viewpoint. More precisely, we would study the existence of at least two nontrivial solutions for the problem (P) as the parameter $\lambda > \lambda_0$ for some constant λ_0 .

This paper is divided into three sections: in the second section we introduce some necessary knowledge on the nonsmooth analysis, basic properties of the generalized Lebesgue-space $L^{p(x)}(\Omega)$ and the generalized Lebesgue-Sobolev space $W^{1,p(x)}(\Omega)$. In the third section, we give the assumptions on the nonsmooth potential $F(x, t, s)$ and prove the multiplicity results for problem (P).

2. Preliminary

2.1. Variable Exponent Sobolev Space

In order to discuss problem (P), we need some theories on $W_0^{k,p(x)}(\Omega)$ which we call variable exponent Sobolev space. Firstly we review some facts on variable exponent spaces $L^{p(x)}(\Omega)$ and $W^{k,p(x)}(\Omega)$. For the details see [4, 18–20].

Firstly, we need to give some notations, which we shall use through this paper:

$$\begin{aligned} C_+(\overline{\Omega}) &= \{p \in C(\overline{\Omega}) : p(x) > 1 \text{ for any } x \in \overline{\Omega}\}, \\ p^- &= \lim_{x \in \overline{\Omega}} p(x), p^+ = \max_{x \in \overline{\Omega}} p(x) \text{ for any } p \in C_+(\overline{\Omega}). \end{aligned} \quad (2.1)$$

Obviously, $1 < p^- \leq p^+ < +\infty$.

Denote by $\mathcal{U}(\Omega)$ the set of all measurable real functions defined on Ω . Two functions in $\mathcal{U}(\Omega)$ are considered to be one element of $\mathcal{U}(\Omega)$, when they are equal almost everywhere.

For $p \in C_+(\overline{\Omega})$, define

$$L^{p(x)}(\Omega) = \left\{ u \in \mathcal{U}(\Omega) : \int_{\Omega} |u(x)|^{p(x)} dx < +\infty \right\}, \quad (2.2)$$

and with the norm $|u|_{L^{p(x)}(\Omega)} = |u|_{p(x)} = \inf\{\lambda > 0 : \int_{\Omega} |u(x)/\lambda|^{p(x)} dx \leq 1\}$, $W^{1,p(x)}(\Omega) = \{u \in L^{p(x)}(\Omega) : |\nabla u| \in L^{p(x)}(\Omega)\}$, with the norm $\|u\| = \|u\|_{W^{1,p(x)}(\Omega)} = |u|_{p(x)} + |\nabla u|_{p(x)}$.

Denote $W_0^{1,p(x)}(\Omega)$ as the closure of $C_0^\infty(\Omega)$ in $W^{1,p(x)}(\Omega)$.

Hereafter, let

$$p^*(x) = \begin{cases} \frac{Np(x)}{N-p(x)}, & p(x) < N, \\ +\infty, & p(x) \geq N. \end{cases} \quad (2.3)$$

Lemma 2.1 (see [19]). (1) The spaces $L^{p(x)}(\Omega)$, $W^{1,p(x)}(\Omega)$, and $W_0^{1,p(x)}(\Omega)$ are separable and reflexive Banach spaces. Moreover, $L^{p(x)}(\Omega)$ is uniform convex.

(2) Poincare inequality in $W_0^{1,p(x)}(\Omega)$ holds; that is, there exists a positive constant C such that

$$|u|_{L^{p(x)}(\Omega)} \leq C |\nabla u|_{L^{p(x)}(\Omega)}, \quad \forall u \in W_0^{1,p(x)}(\Omega). \quad (2.4)$$

(3) If $q \in C_+(\overline{\Omega})$ and $q(x) < p^*(x)$ for any $x \in \overline{\Omega}$, then the embedding from $W^{1,p(x)}(\Omega)$ to $L^{q(x)}(\Omega)$ is compact and continuous.

By (2) of Lemma 2.1, we know that $|\nabla u|_{p(x)}$ and $\|u\|$ are equivalent norms on $W_0^{1,p(x)}(\Omega)$. We will use $|\nabla u|_{p(x)}$ to replace $\|u\|$ in the following discussions.

Lemma 2.2 (see [4]). The conjugate space of $L^{p(x)}(\Omega)$ is $L^{q(x)}(\Omega)$, where $1/p(x) + 1/q(x) = 1$. For any $u \in L^{p(x)}(\Omega)$ and $v \in L^{q(x)}(\Omega)$, one has

$$\int_{\Omega} |uv| dx \leq \left(\frac{1}{p^-} + \frac{1}{q^-} \right) |u|_{L^{p(x)}(\Omega)} |v|_{L^{q(x)}(\Omega)}. \quad (2.5)$$

Lemma 2.3 (see [4]). Set $\rho(u) = \int_{\Omega} |u(x)|^{p(x)} dx$. For $u, u_k \in L^{p(x)}(\Omega)$, one has

- (1) for $u \neq 0$, $|u|_{p(x)} = \lambda \Leftrightarrow \rho(u/\lambda) = 1$;
- (2) $|u|_{p(x)} < 1 (= 1; > 1) \Leftrightarrow \rho(u) < 1 (= 1; > 1)$;
- (3) if $|u|_{p(x)} > 1$, then $|u|_{p(x)}^{p^-} \leq \rho(u) \leq |u|_{p(x)}^{p^+}$;
- (4) if $|u|_{p(x)} < 1$, then $|u|_{p(x)}^{p^+} \leq \rho(u) \leq |u|_{p(x)}^{p^-}$;
- (5) $\lim_{k \rightarrow +\infty} |u_k|_{p(x)} = 0 \Leftrightarrow \lim_{k \rightarrow +\infty} \rho(u_k) = 0$;
- (6) $|u_k|_{p(x)} \rightarrow +\infty \Leftrightarrow \rho(u_k) \rightarrow +\infty$.

In this paper, the space $W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$ will be endowed with the following equivalent norm:

$$\|(u, v)\| = \|u\| + \|v\|, \quad (2.6)$$

where

$$\|u\| = \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{\nabla u}{\lambda} \right|^{p(x)} dx \leq 1 \right\}, \quad \|v\| = \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{\nabla v}{\lambda} \right|^{q(x)} dx \leq 1 \right\}. \quad (2.7)$$

Similar to Lemma 2.3, we have the following.

Lemma 2.4. Set $\rho(u) = \int_{\Omega} |\nabla u(x)|^{p(x)} dx$. For $u, u_k \in W^{1,p(x)}(\Omega)$, one has

- (1) for $u \neq 0$, $\|u\| = \lambda \Leftrightarrow \rho(u/\lambda) = 1$;
- (2) $\|u\| < 1 (= 1; > 1) \Leftrightarrow \rho(u) < 1 (= 1; > 1)$;

- (3) if $\|u\| > 1$, then $\|u\|^{p^-} \leq \rho(u) \leq \|u\|^{p^+}$;
- (4) if $\|u\| < 1$, then $\|u\|^{p^+} \leq \rho(u) \leq \|u\|^{p^-}$;
- (5) $\lim_{k \rightarrow +\infty} \|u_k\| = 0 \Leftrightarrow \lim_{k \rightarrow +\infty} \rho(u_k) = 0$;
- (6) $\|u_k\| \rightarrow +\infty \Leftrightarrow \rho(u_k) \rightarrow +\infty$.

Consider the following function:

$$J(u) = \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx, \quad u \in W_0^{1,p(x)}(\Omega). \quad (2.8)$$

We know that (see [21]) $J \in C^1(W_0^{1,p(x)}(\Omega), \mathbb{R})$ and $p(x)$ -Laplacian operator $-\Delta_{p(x)}u = -\operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ is the derivative operator of J in the weak sense. We denote $\mathcal{L} = J' : W_0^{1,p(x)}(\Omega) \rightarrow (W_0^{1,p(x)}(\Omega))^*$, then $\langle \mathcal{L}(u), v \rangle = \int_{\Omega} (|\nabla u(x)|^{p(x)-2} \nabla u \cdot \nabla v) dx$, for all $u, v \in W_0^{1,p(x)}(\Omega)$.

Lemma 2.5 (see [19]). Set $X = W_0^{1,p(x)}(\Omega)$, $\mathcal{L}$ is as above, then

- (1) $\mathcal{L} : X \rightarrow X^*$ is a continuous, bounded and strictly monotone operator;
- (2) $\mathcal{L}$ is a mapping of type (S_+) , if $u_n \rightharpoonup u$ (weak) in X and $\limsup_{n \rightarrow \infty} \langle \mathcal{L}(u_n), u_n - u \rangle \leq 0$, then $u_n \rightarrow u$ in X ;
- (3) $\mathcal{L} : X \rightarrow X^*$ is a homeomorphism.

2.2. Generalized Gradient

Let X be a Banach space, X^* its topological dual space and we denote $\langle \cdot, \cdot \rangle$ as the duality bracket for pair (X^*, X) . A function $\varphi : X \rightarrow \mathbb{R}$ is said to be locally Lipschitz, if for every $x \in X$ we can find a neighbourhood U of x and a constant $k > 0$ (depending on U), such that $|\varphi(y) - \varphi(z)| \leq k\|y - z\|$, for all $y, z \in U$.

The generalized directional derivative of φ at the point $u \in X$ in the direction $h \in X$ is

$$\varphi^0(u; h) = \limsup_{u' \rightarrow u; \lambda \downarrow 0} \frac{\varphi(u' + \lambda h) - \varphi(u')}{\lambda}. \quad (2.9)$$

The generalized subdifferential of φ at the point $u \in X$ is defined by the

$$\partial\varphi(u) = \left\{ u^* \in X^*; \langle u^*, h \rangle \leq \varphi^0(u; h), \quad \forall h \in X \right\}, \quad (2.10)$$

which is a nonempty, convex and w^* -compact set of X . We say that $u \in X$ is a critical point of φ , if $0 \in \partial\varphi(u)$. For further details, we refer the reader to [12].

Finally we have the following Weierstrass Theorem and Mountain Pass Theorem.

Theorem 2.6. *If X is a reflexive Banach space and $\varphi : X \rightarrow \mathbb{R}$ satisfies*

(1) *φ is weak lower semicontinuous, that is,*

$$x_n \rightharpoonup x_0 \text{ (weakly) in } X \longrightarrow \varphi(x_0) \leq \liminf_{n \rightarrow +\infty} \varphi(x_n); \quad (2.11)$$

(2) *φ is coercive, that is, $\lim_{\|x\| \rightarrow \infty} \varphi(x) = +\infty$,*

then there exists $x^ \in X$ such that $\varphi(x^*) = \min_{x \in X} \varphi(x)$.*

Theorem 2.7 (see [22]). *Let $\varphi : X \rightarrow \mathbb{R}$ be locally Lipschitz function and $x_0, x_1 \in X$. If there exists a bounded open neighbourhood U of x_0 , such that $x_1 \in X \setminus U$, $\max\{\varphi(x_0), \varphi(x_1)\} < \inf_{\partial U} \varphi$ and φ satisfies the nonsmooth C -condition at level c , where $c = \inf_{\gamma \in \mathcal{T}} \max_{t \in [0,1]} \varphi(\gamma(t))$, $\mathcal{T} = \{\gamma \in C([0,1]; X) : \gamma(0) = x_0, \gamma(1) = x_1\}$, then c is a critical value of φ and $c \geq \inf_{\partial U} \varphi$.*

3. Existence Results

For each $(u, v) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$, define

$$\begin{aligned} \Phi(u, v) &= \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{p(x)} |\nabla v|^{q(x)} dx, \\ \Psi(u, v) &= \int_{\Omega} F(x, u, v) dx. \end{aligned} \quad (3.1)$$

By a solution of (2), we mean function $(u, v) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$ to which there corresponds mapping $\Omega \ni x \rightarrow (w_1, w_2)$ with $w_1(x) \in \partial_u F(x, u, v)$, $w_2(x) \in \partial_v F(x, u, v)$ for almost every $x \in \Omega$ having the property that for every $(\xi, \eta) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$, the function $x \rightarrow (w_1(x)\xi(x), w_2(x)\eta(x)) \in L^1(\Omega) \times L^1(\Omega)$ and

$$\int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \xi dx + \int_{\Omega} |\nabla v|^{q(x)-2} \nabla v \nabla \eta dx = \lambda \int_{\Omega} w_1 \xi dx + \lambda \int_{\Omega} w_2 \eta dx. \quad (3.2)$$

Our hypotheses on nonsmooth potential $F(x, t, s)$ is as follows.

$H(F)$: $F : \Omega \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a function such that $F(x, 0, 0) = 0$ a.e. on Ω and satisfies the following facts:

- (1) for all $t \in \mathbb{R}$, $s \in \mathbb{R}$, $x \mapsto F(x, t, s)$ is measurable;
- (2) for almost all $x \in \Omega$, $t \mapsto F(x, t, s)$ and $s \mapsto F(x, t, s)$ are locally Lipschitz.

Lemma 3.1. Suppose $H(F)$ and the following conditions hold:

(f₁) there exists $\alpha \in C_+(\overline{\Omega})$ and $\alpha(x) < p^*(x)$, such that

$$|w_1| \leq c_1 \left(1 + |t|^{\alpha(x)-1} + |s|^{\beta(x)(\alpha(x)-1)/\alpha(x)} \right), \quad (3.3)$$

for almost all $x \in \Omega$, all $t, s \in \mathbb{R}$ and $w_1 \in \partial_t F(x, t, s)$;

(f₂) there exists $\beta \in C_+(\overline{\Omega})$ and $\beta(x) < q^*(x)$, such that

$$|w_2| \leq c_2 \left(1 + |t|^{(\alpha(x)(\beta(x)-1))/(\beta(x))} + |s|^{\beta(x)-1} \right), \quad (3.4)$$

for almost all $x \in \Omega$, all $t, s \in \mathbb{R}$ and $w_2 \in \partial_s F(x, t, s)$;

then $\varphi(\cdot, v)(\varphi(u, \cdot))$ is locally Lipschitz on $W_0^{1,p(x)}(\Omega)(W_0^{1,q(x)}(\Omega))$.

Proof. We need only to prove that $\varphi(\cdot, v)$ is locally Lipschitz on $W_0^{1,p(x)}(\Omega)$.

By $\Phi(\cdot, v) \in C^1(W_0^{1,p(x)}(\Omega), \mathbb{R})$, we have

$$\varphi(u_1, v) - \varphi(u_2, v) = J(u_1) - J(u_2) = J'(\bar{u}) \cdot (u_1 - u_2), \quad (3.5)$$

where $\bar{u} = tu_1 + (1-t)u_2$, $t \in (0, 1)$.

Let $B_r = \{u \in W_0^{1,p(x)}(\Omega) : \|u - u_0\|_{W_0^{1,p(x)}(\Omega)} \leq r\}$.

Note that B_r is w -compact. Then we obtain that there exists a positive constant M , such that $\|J'(\bar{u})\|_{W^{-1,p'(x)}(\Omega)} \leq M$ with $1/p(x) + 1/p'(x) = 1$, for sufficiently small r .

Therefore, for any $u_1, u_2 \in B_r$, we have

$$\begin{aligned} |\Phi(u_1, v) - \Phi(u_2, v)| &= |J'(\bar{u}) \cdot (u_1 - u_2)| \\ &\leq \|J'(\bar{u})\|_{W^{-1,p'(x)}(\Omega)} \|u_1 - u_2\|_{W_0^{1,p(x)}(\Omega)} \\ &\leq M \|u_1 - u_2\|_{W_0^{1,p(x)}(\Omega)}, \quad \forall v \in W_0^{1,q(x)}(\Omega). \end{aligned} \quad (3.6)$$

Fixing $v \in W_0^{1,q(x)}(\Omega)$, by (f₁) and Lebourg mean value theorem, we have

$$|F(x, u_1, v) - F(x, u_2, v)| \leq c_1 \left(1 + |\bar{u}|^{\alpha(x)-1} + |v|^{\beta(x)(\alpha(x)-1)/\alpha(x)} \right) |u_1 - u_2|. \quad (3.7)$$

Hence,

$$\begin{aligned} &\left| \int_{\Omega} F(x, u_1, v) dx - \int_{\Omega} F(x, u_2, v) dx \right| \\ &\leq c_1 \int_{\Omega} |u_1 - u_2| dx + c_1 \int_{\Omega} |\bar{u}|^{\alpha(x)-1} |u_1 - u_2| dx + c_1 \int_{\Omega} |v|^{\beta(x)(\alpha(x)-1)/\alpha(x)} |u_1 - u_2| dx \\ &\leq \bar{c}_1 |u_1 - u_2|_{\alpha(x)} + c_1 \left| |\bar{u}|^{\alpha(x)-1} \right|_{\alpha'(x)} |u_1 - u_2|_{\alpha(x)} + c_1 \left| |v|^{\beta(x)(\alpha(x)-1)/\alpha(x)} \right|_{\alpha'(x)} |u_1 - u_2|_{\alpha(x)}, \end{aligned} \quad (3.8)$$

where $1/\alpha'(x) + 1/\alpha(x) = 1$.

It is immediate that

$$\begin{aligned} \int_{\Omega} \left(|\bar{u}|^{\alpha(x)-1} \right)^{\alpha'(x)} dx &= \int_{\Omega} |\bar{u}|^{\alpha(x)} dx \leq \begin{cases} |\bar{u}|_{\alpha(x)}^{\alpha^+} \leq c \|\bar{u}\|^{\alpha^+}, & |\bar{u}|_{\alpha(x)} > 1, \\ |\bar{u}|_{\alpha(x)}^{\alpha^+} \leq c \|\bar{u}\|^{\alpha^-}, & |\bar{u}|_{\alpha(x)} < 1, \end{cases} \\ \int_{\Omega} \left(|v|^{\beta(x)(\alpha(x)-1)/\alpha(x)} \right)^{\alpha'(x)} dx &= \int_{\Omega} |v|^{\beta(x)} dx \leq \begin{cases} |v|_{\beta(x)}^{\beta^+} \leq c \|v\|^{\beta^+}, & |v|_{\beta(x)} > 1, \\ |v|_{\beta(x)}^{\beta^+} \leq c \|v\|^{\beta^-}, & |v|_{\beta(x)} < 1 \end{cases} \end{aligned} \quad (3.9)$$

are bounded.

So,

$$\left| \int_{\Omega} F(x, u_1, v) dx - \int_{\Omega} F(x, u_2, v) dx \right| \leq c \|u_1 - u_2\|, \quad (3.10)$$

since $W_0^{1,p(x)}(\Omega) \hookrightarrow L^{\alpha(x)}(\Omega)$ is a compact embedding.

Therefore, $\varphi(\cdot, v)$ is locally Lipschitz. Similarly, we can prove that $\varphi(u, \cdot)$ is locally Lipschitz. $\square$

Theorem 3.2. Suppose that $H(F)$, (f_1) with $\alpha^+ < p^-$, (f_2) with $\beta^+ < q^-$ and the following conditions (f_3) – (f_4) hold:

(f_3) there exists $\gamma_i \in C(\overline{\Omega})$ ($i = 1, 2$) with $p^+ < \gamma_1(x) < p^*(x)$, $q^+ < \gamma_2(x) < q^*(x)$ and $\mu_1, \mu_2 \in L^\infty(\Omega)$, such that

$$\limsup_{t \rightarrow 0, s \rightarrow 0} \frac{\langle w_1, t \rangle}{|t|^{\gamma_1(x)}} < \mu_1(x), \quad \limsup_{t \rightarrow 0, s \rightarrow 0} \frac{\langle w_2, s \rangle}{|s|^{\gamma_2(x)}} < \mu_2(x) \quad (3.11)$$

uniformly for almost all $x \in \Omega$ and all $w_1 \in \partial_t F(x, t, s)$, $w_2 \in \partial_s F(x, t, s)$;

(f_4) there exist $\xi_0 \in R$, $\eta_0 \in R$, $x_0 \in \Omega$ and $r_0 > 0$, such that

$$F(x, \xi_0, \eta_0) > \delta_0 > 0, \quad \text{a.e. } x \in B_{r_0}(x_0), \quad (3.12)$$

where $B_{r_0}(x_0) := \{x \in \Omega : |x - x_0| \leq r_0\} \subset \Omega$.

Then there exists $\lambda_* > 0$ such that, for each $\lambda > \lambda_*$, the problem (2) has at least two nontrivial solutions.

Proof. The proof is divided into four steps as follows. $\square$

Step 1. We will show that φ is coercive in the step.

Firstly, for almost all $x \in \Omega$, by $H(F)(2)$, $t \mapsto F(x, t, s)$ is differentiable almost everywhere on $\mathbb{R}$ and we have

$$\frac{d}{dt} F(x, t, s) \in \partial_t F(x, t, s). \quad (3.13)$$

Moreover, from (f_1) , (f_2) and Young inequality, we can get that

$$\begin{aligned}
 F(x, t, s) &= F(x, 0, s) + \int_0^t \frac{d}{dy} F(x, y, s) dy \\
 &= F(x, 0, 0) + \int_0^s \frac{d}{dz} F(x, 0, z) dz + \int_0^t \frac{d}{dy} F(x, y, s) dy \\
 &\leq c_1 \left[|t| + \frac{|t|^{\alpha(x)}}{\alpha(x)} + |s|^{\beta(x)(\alpha(x)-1)/\alpha(x)} |t| \right] \\
 &\quad + c_2 \left[|s| + |t|^{\alpha(x)(\beta(x)-1)/\beta(x)} |s| + \frac{1}{\beta(x)} |s|^{\beta(x)} \right] \\
 &\leq c_1 \left[|t| + |t|^{\alpha(x)} + \frac{(\alpha(x)-1)|s|^{\beta(x)}}{\alpha(x)} + \frac{|t|^{\alpha(x)}}{\beta(x)} \right] \\
 &\quad + c_2 \left[|s| + |s|^{\beta(x)} + \frac{(\beta(x)-1)|t|^{\alpha(x)}}{\beta(x)} + \frac{|s|^{\beta(x)}}{\beta(x)} \right] \\
 &\leq c_1 \left[|t| + 2|t|^{\alpha(x)} + |s|^{\beta(x)} \right] + c_2 \left[|s| + 2|s|^{\beta(x)} + |t|^{\alpha(x)} \right],
 \end{aligned} \tag{3.14}$$

for almost all $x \in \Omega$ and $t, s \in \mathbb{R}$.

Note that $1 < \alpha(x) \leq \alpha^+ < p^- < p^*(x)$ and $1 < \beta(x) \leq \beta^+ < q^- < q^*(x)$, then, by Lemma 2.1, we have $W_0^{1,p(x)}(\Omega) \hookrightarrow L^{\alpha(x)}(\Omega)$ and $W_0^{1,q(x)}(\Omega) \hookrightarrow L^{\beta(x)}(\Omega)$ (compact embedding). Furthermore, there exists $c_3, c_4 > 0$ such that $|u|_{\alpha(x)} \leq c_3 \|u\|$ and $|v|_{\beta(x)} \leq c_4 \|v\|$.

So, for any $|u|_{\alpha(x)} > 1$ and $\|u\| > 1$,

$$\int_{\Omega} |u|^{\alpha(x)} dx \leq |u|_{\alpha(x)}^{\alpha^+} \leq c_3^{\alpha^+} \|u\|^{\alpha^+} \tag{3.15}$$

and, for any $|v|_{\beta(x)} > 1$ and $\|v\| > 1$,

$$\int_{\Omega} |v|^{\beta(x)} dx \leq |v|_{\beta(x)}^{\beta^+} \leq c_4^{\beta^+} \|v\|^{\beta^+}. \tag{3.16}$$

By (3.14), (3.15), (3.31), the Hölder inequality and the Sobolev embedding theorem, we have

$$\begin{aligned}
 \varphi(u, v) &= \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v|^{q(x)} dx - \lambda \int_{\Omega} F(x, u, v) dx \\
 &\geq \frac{1}{p^+} \|u\|^{p^-} + \frac{1}{q^+} \|v\|^{q^-} - \lambda c_1 \int_{\Omega} |u| dx - 2c_1 c_3^{\alpha^+} \|u\|^{\alpha^+} - c_1 c_4^{\beta^+} \|v\|^{\beta^+} \\
 &\quad - \lambda c_2 \int_{\Omega} |v| dx - 2c_2 c_4^{\beta^+} \|v\|^{\beta^+} - c_2 c_3^{\alpha^+} \|u\|^{\alpha^+}
 \end{aligned}$$

$$\begin{aligned}
&\geq \frac{1}{p^+} \|u\|^{p^-} + \frac{1}{q^+} \|v\|^{q^-} - 2\lambda c_1 |1|_{\alpha'(x)} |u|_{\alpha(x)} - 2\lambda c_1 |1|_{\beta'(x)} |v|_{\beta(x)} \\
&\quad - 2c_1 c_3^{\alpha^+} \|u\|^{\alpha^+} - c_1 c_4^{\beta^+} \|v\|^{\beta^+} - 2c_2 c_4^{\beta^+} \|v\|^{\beta^+} - c_2 c_3^{\alpha^+} \|u\|^{\alpha^+} \\
&\geq \frac{1}{p^+} \|u\|^{p^-} + \frac{1}{q^+} \|v\|^{q^-} - 2\lambda c_1 c |1|_{\alpha'(x)} \|u\| - 2\lambda c_1 c |1|_{\beta'(x)} \|v\| \\
&\quad - 2c_1 c_3^{\alpha^+} \|u\|^{\alpha^+} - c_1 c_4^{\beta^+} \|v\|^{\beta^+} - 2c_2 c_4^{\beta^+} \|v\|^{\beta^+} - c_2 c_3^{\alpha^+} \|u\|^{\alpha^+} \\
&\longrightarrow \infty, \text{ as } \|u, v\| \longrightarrow \infty.
\end{aligned} \tag{3.17}$$

Step 2. We will show that the φ is weakly lower semicontinuous.

Letting $u_n \rightharpoonup u$ weakly in $W_0^{1,p(x)}(\Omega)$, $v_n \rightharpoonup v$ weakly in $W_0^{1,q(x)}(\Omega)$ by Lemma 2.1(3), we obtain the following results:

$$\begin{aligned}
&W_0^{1,p(x)}(\Omega) \hookrightarrow L^{p(x)}(\Omega); W_0^{1,q(x)}(\Omega) \hookrightarrow L^{q(x)}(\Omega); \\
&u_n \rightarrow u \text{ in } L^{p(x)}(\Omega); v_n \rightarrow v \text{ in } L^{q(x)}(\Omega); \\
&u_n \rightarrow u \text{ for a.a. } x \in \Omega; v_n \rightarrow v \text{ for a.a. } x \in \Omega; \\
&F(x, u_n(x), v_n(x)) \rightarrow F(x, u(x), v(x)) \text{ for a.a. } x \in \Omega.
\end{aligned}$$

By Fatou's Lemma,

$$\limsup_{n \rightarrow \infty} \int_{\Omega} F(x, u_n(x), v_n(x)) dx \leq \int_{\Omega} F(x, u(x), v(x)) dx. \tag{3.18}$$

Thus,

$$\begin{aligned}
\liminf_{n \rightarrow \infty} \varphi(u_n, v_n) &= \liminf_{n \rightarrow \infty} \left[\int_{\Omega} \frac{1}{p(x)} |\nabla u_n|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v_n|^{q(x)} dx \right] \\
&\quad - \limsup_{n \rightarrow \infty} \int_{\Omega} F(x, u_n, v_n) dx \\
&\geq \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v|^{q(x)} dx - \lambda \int_{\Omega} F(x, u, v) dx. \\
&= \varphi(u)
\end{aligned} \tag{3.19}$$

Hence, by Theorem 2.6, we deduce that there exists a global minimizer $(u_0, v_0) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$ such that

$$\varphi(u_0, v_0) = \min_{(u,v) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)} \varphi(u, v). \tag{3.20}$$

Step 3. We will show that there exists $\lambda_* > 0$ such that for each $\lambda > \lambda_*$, $\varphi(u_0, v_0) < 0$.

By the condition (f_4) , there exists $\xi_0, \eta_0 \in \mathbb{R}$ such that $F(x, \xi_0, \eta_0) > \delta_0 > 0$, a.e. $x \in B_{r_0}(x_0)$. It is clear that

$$0 < M_1 := \max_{|t| \leq \xi_0, |s| \leq \eta_0} \left\{ c_1 \left[|t| + 2|t|^{\alpha^+} + |s|^{\beta^+} \right] + c_2 \left[|s| + 2|s|^{\beta^+} + |t|^{\alpha^+} \right], \right. \\ \left. c_1 \left[|t| + 2|t|^{\alpha^-} + |s|^{\beta^-} \right] + c_2 \left[|s| + 2|s|^{\beta^-} + |t|^{\alpha^-} \right] \right\} < +\infty. \quad (3.21)$$

Now we denote

$$t_0 = \left(\frac{M_1}{\delta_0 + M_1} \right)^{1/N}, \\ K(t) := \max \left\{ \left(\frac{\xi_0}{r_0(1-t)} \right)^{p^-}, \left(\frac{\xi_0}{r_0(1-t)} \right)^{p^+}, \left(\frac{\xi_0}{r_0(1-t)} \right)^{q^-}, \left(\frac{\xi_0}{r_0(1-t)} \right)^{q^+} \right\}, \quad (3.22) \\ \lambda_* = \max_{t \in [t_1, t_2]} \frac{K(t)(1-t^N)}{[\delta_0 t^N - M_1(1-t^N)]},$$

where $t_0 < t_1 < t_2 < 1$ and δ_0 is given in the condition (f_4) . A simple calculation shows that the function $t \mapsto \delta_0 t^N - M_1(1-t^N)$ is positive whenever $t > t_0$ and $\delta_0 t_0^N - M_1(1-t_0^N) = 0$. Thus λ_* is well defined and $\lambda_* > 0$.

We will show that, for each $\lambda > \lambda_*$, the problem (P) has two nontrivial solutions. In order to do this, for $t \in [t_1, t_2]$, let us define

$$\eta_t(x) = \begin{cases} 0, & \text{if } x \in \Omega \setminus B_{r_0}(x_0), \\ \xi_0, & \text{if } x \in B_{tr_0}(x_0), \\ \frac{\xi_0}{r_0(1-t)}(r_0 - |x - x_0|), & \text{if } x \in B_{r_0}(x_0) \setminus B_{tr_0}(x_0). \end{cases} \quad (3.23)$$

By conditions (f_1) and (f_3) we have

$$\int_{\Omega} F(x, \eta_t(x), \eta_t(x)) dx = \int_{B_{tr_0}(x_0)} F(x, \eta_t(x), \eta_t(x)) dx \\ + \int_{B_{r_0}(x_0) \setminus B_{tr_0}(x_0)} F(x, \eta_t(x), \eta_t(x)) dx \quad (3.24) \\ \geq w_N r_0^N t^N \delta_0 - M_1 (1-t^N) w_N r_0^N \\ = w_N r_0^N (\delta_0 t^N - M_1 (1-t^N)).$$

Hence, for $t \in [t_1, t_2]$,

$$\begin{aligned}
\varphi(\eta_t, \eta_t) &= \int_{\Omega} \frac{1}{p(x)} |\nabla \eta_t|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla \eta_t|^{q(x)} dx - \lambda \int_{\Omega} F(x, \eta_t(x), \eta_t(x)) dx \\
&\leq \frac{1}{p^-} \int_{\Omega} |\nabla \eta_t|^{p(x)} dx + \frac{1}{q^-} \int_{\Omega} |\nabla \eta_t|^{q(x)} dx - \lambda w_N r_0^N (\delta_0 t^N - M_1 (1 - t^N)) \\
&\leq \max \left\{ \left(\frac{\xi_0}{r_0(1-t)} \right)^{p^-}, \left(\frac{\xi_0}{r_0(1-t)} \right)^{p^+}, \left(\frac{\xi_0}{r_0(1-t)} \right)^{q^-}, \left(\frac{\xi_0}{r_0(1-t)} \right)^{q^+} \right\} \\
&\quad \times w_N r_0^N (1 - t^N) - \lambda w_N r_0^N (\delta_0 t^N - M_1 (1 - t^N)) \\
&= w_N r_0^N \left[K(t) (1 - t^N) - \lambda (\delta_0 t^N - M_1 (1 - t^N)) \right],
\end{aligned} \tag{3.25}$$

so that $\varphi(\eta_t, \eta_t) < 0$ whenever $\lambda > \lambda_*$.

Step 4. We will check the C-condition in the following.

Suppose $\{(u_n, v_n)\}_{n \geq 1} \subseteq W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$ such that $\varphi(u_n, v_n) \rightarrow c$ and $(1 + \|(u_n, v_n)\|)m(u_n, v_n) \rightarrow 0$. Let $(u_n^*, v_n^*) \in \partial\varphi(u_n, v_n)$ be such that $m(u_n, v_n) = \|(u_n^*, v_n^*)\|_{(W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega))^*}$. The interpretation of $(u_n^*, v_n^*) \in \partial\varphi(u_n, v_n)$ is that $u_n^* \in \partial_u \varphi(u_n, v_n)$ and $v_n^* \in \partial_v \varphi(u_n, v_n)$. We know that

$$\begin{aligned}
u_n^* &= -\Delta_{p(x)} u_n - \lambda w_n^1, \\
v_n^* &= -\Delta_{q(x)} v_n - \lambda w_n^2
\end{aligned} \tag{3.26}$$

with $w_n^1 \in \partial_u \Psi(u_n, v_n)$ and $w_n^2 \in \partial_v \Psi(u_n, v_n)$. From Chang [23] we know that $w_n^1 \in L^{\alpha'(x)}(\Omega)$ and $w_n^2 \in L^{\beta'(x)}(\Omega)$, where $\alpha'(x) = \alpha(x)/(\alpha(x) - 1)$, $\beta'(x) = \beta(x)/(\beta(x) - 1)$.

Since φ is coercive, $\{u_n\}_{n \geq 1}$, $\{v_n\}_{n \geq 1}$ are bounded and passed to a subsequence, still denoting $\{u_n\}_{n \geq 1}$ and $\{v_n\}_{n \geq 1}$, we may assume that there exist $u \in W_0^{1,p(x)}(\Omega)$, $v \in W_0^{1,q(x)}(\Omega)$, such that $u_n \rightharpoonup u$ weakly in $W_0^{1,p(x)}(\Omega)$ and $v_n \rightharpoonup v$ weakly in $W_0^{1,q(x)}(\Omega)$. Next we will prove that

$$u_n \rightarrow u \text{ in } W_0^{1,p(x)}(\Omega), \quad v_n \rightarrow v \text{ in } W_0^{1,q(x)}(\Omega). \tag{3.27}$$

By $W_0^{1,p(x)}(\Omega) \rightarrow L^{p(x)}(\Omega)$, $W_0^{1,q(x)}(\Omega) \rightarrow L^{q(x)}(\Omega)$, we have $u_n \rightarrow u$ in $L^{p(x)}(\Omega)$ and $v_n \rightarrow v$ in $L^{q(x)}(\Omega)$. Moreover, since $\|(u_n^*, v_n^*)\|_{(W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega))^*} \rightarrow 0$, we get $\|u_n^*\|_{(W_0^{1,p(x)}(\Omega))^*} \rightarrow 0$, $\|v_n^*\|_{(W_0^{1,q(x)}(\Omega))^*} \rightarrow 0$, so $|\langle u_n^*, u_n \rangle| \leq \varepsilon_n$, $|\langle v_n^*, v_n \rangle| \leq \varepsilon_n$.

From (3.26), we have

$$\begin{aligned}
\langle -\Delta_{p(x)} u_n, u_n - u \rangle - \int_{\Omega} w_n^1 (u_n - u) dx &\leq \varepsilon_n, \quad \forall n \geq 1. \\
\langle -\Delta_{q(x)} v_n, v_n - v \rangle - \int_{\Omega} w_n^2 (v_n - v) dx &\leq \varepsilon_n, \quad \forall n \geq 1.
\end{aligned} \tag{3.28}$$

Moreover, $\int_{\Omega} w_n^1(u_n - u)dx \rightarrow 0$ and $\int_{\Omega} w_n^2(v_n - v)dx \rightarrow 0$, since $u_n \rightarrow u$ in $L^{p(x)}(\Omega)$, $v_n \rightarrow v$ in $L^{q(x)}(\Omega)$, $\{w_n^1\}_{n \geq 1}$ in $L^{p'(x)}(\Omega)$ and $\{w_n^2\}_{n \geq 1}$ in $L^{q'(x)}(\Omega)$ are bounded, where $1/p(x) + 1/(p'(x)) = 1$, $1/(q(x)) + 1/(q'(x)) = 1$. Therefore,

$$\limsup_{n \rightarrow \infty} \langle -\Delta_{p(x)} u_n, u_n - u \rangle \leq 0, \quad \limsup_{n \rightarrow \infty} \langle -\Delta_{q(x)} v_n, v_n - v \rangle \leq 0. \quad (3.29)$$

From Lemma 2.5, we have $u_n \rightarrow u$, $v_n \rightarrow v$ as $n \rightarrow \infty$. Thus φ satisfies the nonsmooth C-condition.

Step 5. We will show that there exists another nontrivial weak solution of problem (P).

From Lebourg Mean Value Theorem, we obtain

$$\begin{aligned} F(x, t, s) - F(x, 0, s) &= \langle w_1, t \rangle, \\ F(x, 0, s) - F(x, 0, 0) &= \langle w_2, s \rangle \end{aligned} \quad (3.30)$$

for some $w_1 \in \partial_t F(x, \vartheta t, s)$, $w_2 \in \partial_s F(x, 0, \tau s)$, and $0 < \vartheta, \tau < 1$. Thus by the condition (f₃), there exists $\beta \in (0, 1)$ such that

$$\begin{aligned} |F(x, t, s)| &\leq |\langle w_1, t \rangle| + |\langle w_2, s \rangle| \\ &\leq \mu_1(x) |t|^{\gamma_1(x)} + \mu_2(x) |s|^{\gamma_2(x)}, \end{aligned} \quad (3.31)$$

for all $|t|, |s| < \beta$ and a.e. $x \in \Omega$.

It follows from the conditions (f₁), (f₂) $1 < \alpha^- \leq \alpha^+ < p^- \leq p^+ < \gamma_1(x) < p^*(x)$ and $1 < \beta^- \leq \beta^+ < q^- \leq q^+ < \gamma_2(x) < p^*(x)$ that for all $|t| > \beta$, $|s| > \beta$ and a.e. $x \in \Omega$,

$$\begin{aligned} |F(x, t, s)| &\leq c_1 \left[|t| + 2|t|^{\alpha(x)} + |s|^{\beta(x)} \right] + c_2 \left[|s| + 2|s|^{\beta(x)} + |t|^{\alpha(x)} \right] \\ &\leq \left(\frac{c_1}{\beta^{\gamma_1(x)-1}} + \frac{2c_2}{\beta^{\gamma_1(x)-\alpha(x)}} + \frac{c_2}{\beta^{\gamma_1(x)-\beta(x)}} \right) |t|^{\gamma_1(x)} \\ &\quad + \left(\frac{c_2}{\beta^{\gamma_2(x)-1}} + \frac{2c_2}{\beta^{\gamma_2(x)-\beta(x)}} + \frac{c_1}{\beta^{\gamma_1(x)-\alpha(x)}} \right) |t|^{\gamma_2(x)} \\ &\leq \left(\frac{c_1}{\beta^{\gamma_1^+-1}} + \frac{2c_2}{\beta^{\gamma_1^+-\alpha^-}} + \frac{c_2}{\beta^{\gamma_1^+-\beta^-}} \right) |t|^{\gamma_1(x)} \\ &\quad + \left(\frac{c_2}{\beta^{\gamma_2^+-1}} + \frac{2c_2}{\beta^{\gamma_2^+-\beta^-}} + \frac{c_1}{\beta^{\gamma_1^+-\alpha^-}} \right) |t|^{\gamma_2(x)}, \end{aligned} \quad (3.32)$$

this together with (3.31) yields that, for all $t \in \mathbb{R}$ and a.e. $x \in \Omega$,

$$\begin{aligned} |F(x, t, s)| &\leq \left(\mu_1(x) + \frac{c_1}{\beta\gamma_1^+-1} + \frac{2c_2}{\beta\gamma_1^+-\alpha^-} + \frac{c_2}{\beta\gamma_1^+-\beta^-} \right) |t|^{\gamma_1(x)} \\ &\quad + \left(\mu_2(x) + \frac{c_2}{\beta\gamma_2^+-1} + \frac{2c_2}{\beta\gamma_2^+-\beta^-} + \frac{c_1}{\beta\gamma_1^+-\alpha^-} \right) |t|^{\gamma_2(x)} \\ &\leq c_3 |t|^{\gamma_1(x)} + c_4 |s|^{\gamma_2(x)}, \end{aligned} \quad (3.33)$$

for positive constants c_3, c_4 .

Note that $p^+ < \gamma_1(x) < p^*(x)$, $q^+ < \gamma_2(x) < q^*(x)$, then, by Lemma 2.1, we have $W_0^{1,p(x)}(\Omega) \hookrightarrow L^{\gamma_1(x)}(\Omega)$ and $W_0^{1,q(x)}(\Omega) \hookrightarrow L^{\gamma_2(x)}(\Omega)$. Furthermore, there exist c_5, c_6 such that

$$|u|_{\gamma_1(x)} \leq c_5 \|u\|, \quad \forall u \in W_0^{1,p(x)}(\Omega), \quad |v|_{\gamma_2(x)} \leq c_6 \|v\|, \quad \forall v \in W_0^{1,q(x)}(\Omega). \quad (3.34)$$

For all $\lambda > \lambda_*$, $\|(u, v)\| < 1$, $|u|_{\gamma_1(x)} < 1$ and $|v|_{\gamma_2(x)} < 1$, from (3.33) we have

$$\begin{aligned} \varphi(u, v) &= \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v|^{q(x)} dx - \lambda \int_{\Omega} F(x, u(x), v(x)) dx \\ &\geq \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v|^{q(x)} dx - \lambda c_3 \int_{\Omega} |u(x)|^{\gamma_1(x)} dx \\ &\quad - \lambda c_4 \int_{\Omega} |v(x)|^{\gamma_2(x)} dx \\ &\geq \frac{1}{p^+} \|u\|^{p^+} + \frac{1}{q^+} \|v\|^{q^+} - \lambda \left[c_3 c_5^{\gamma_1^-} \|u\|^{\gamma_1^-} + c_4 c_6^{\gamma_2^-} \|v\|^{\gamma_2^-} \right]. \end{aligned} \quad (3.35)$$

So, for $\rho > 0$ small enough, there exists a $\nu > 0$ such that

$$\varphi(u, v) > \nu, \quad \text{for } \|(u, v)\| = \rho \quad (3.36)$$

and $\|(u_0, v_0)\| > \rho$. So by the Nonsmooth Mountain Pass Theorem (cf. Theorem 2.7), we can get $(u_1, v_1) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$ which satisfies

$$\varphi(u_1, v_1) = c > 0, \quad m(u_1, v_1) = 0. \quad (3.37)$$

Therefore, (u_1, v_1) is another nontrivial solution of problem (P).

Remark 3.3. Let $p^- > \alpha^+$, $q^- > \beta^+$ and consider the following nonsmooth locally Lipschitz function:

$$F(x, t, s) = \begin{cases} t^{\gamma_1(x)} + s^{\gamma_2(x)}, & t \in (0, \delta) \text{ or } s \in (0, \delta), \\ \max\{|t - \delta|^{\theta(x)}, |t - \delta|^{\alpha(x)}\} + |\delta|^{\gamma_1(x)} \\ \quad + \max\{|s - \delta|^{\theta(x)}, |s - \delta|^{\beta(x)}\} + |\delta|^{\gamma_2(x)}, & t \in [\delta, +\infty) \text{ or } s \in [\delta, +\infty), \\ 0, & t \in (-\infty, 0] \text{ or } s \in (-\infty, 0], \end{cases} \quad (3.38)$$

where $0 < \delta < 1$, $\theta^- > 1$, $\theta^+ < \alpha^-$ and $\theta^+ < \beta^-$.

Obvious, $t \mapsto F(x, t, s)$ and $s \mapsto F(x, t, s)$ are locally Lipschitz. Then,

$$\begin{aligned} \partial_t F(x, t, s) &= \begin{cases} \gamma_1(x)t^{\gamma_1(x)-1}, & t \in (0, \delta), \\ \theta(x)(t - \delta)^{\theta(x)-1}, & t \in (\delta, 1 + \delta), \\ \alpha(x)(t - \delta)^{\alpha(x)-1}, & t \in (1 + \delta, +\infty), \\ [0, \gamma_1(x)\delta^{\gamma_1(x)-1}], & t = \delta, \\ [\theta(x), \alpha(x)], & t = 1 + \delta, \\ 0, & t \in (-\infty, 0], \end{cases} \\ \partial_s F(x, t, s) &= \begin{cases} \gamma_2(x)s^{\gamma_2(x)-1}, & s \in (0, \delta), \\ \theta(x)(s - \delta)^{\theta(x)-1}, & s \in (\delta, 1 + \delta), \\ \beta(x)(s - \delta)^{\beta(x)-1}, & s \in (1 + \delta, +\infty), \\ [0, \gamma_1(x)\delta^{\gamma_1(x)-1}], & s = \delta, \\ [\theta(x), \beta(x)], & s = 1 + \delta, \\ 0, & s \in (-\infty, 0]. \end{cases} \end{aligned} \quad (3.39)$$

Hence, for any $w_1 \in \partial_t F(x, t, s)$ and $w_2 \in \partial_s F(x, t, s)$, we have

$$\begin{aligned} |w_1| &\leq \begin{cases} \gamma_1(x)t^{\alpha(x)-1}t^{\gamma_1(x)-\alpha(x)} \leq \gamma_1^+|t|^{\alpha(x)-1}, & t \in (0, \delta), \\ \theta(x)(t - \delta)^{\theta(x)-1} < \theta^+ < \theta^+\left(\frac{1}{\delta}\right)^{\alpha^+-1}|t|^{\alpha(x)-1}, & t \in (\delta, 1 + \delta), \\ \alpha(x)(t - \delta)^{\alpha(x)-1} \leq \alpha^+|t|^{\alpha(x)-1}, & t \in (1 + \delta, +\infty), \\ \gamma_1(x)\delta^{\gamma_1(x)-1} = \gamma_1(x)\delta^{\alpha(x)-1}\delta^{\gamma_1(x)-\alpha(x)} \leq \gamma^+\delta^{\alpha(x)-1}, & t = \delta, \\ [\theta(x), \alpha(x)] \leq \alpha^+(1 + \delta)^{\alpha(x)-1}, & t = 1 + \delta, \\ 0, & t \in (-\infty, 0], \end{cases} \\ |w_2| &\leq \begin{cases} \gamma_2(x)s^{\beta(x)-1}s^{\gamma_2(x)-\beta(x)} \leq \gamma_2^+|s|^{\beta(x)-1}, & s \in (0, \delta), \\ \theta(x)(s - \delta)^{\theta(x)-1} < \theta^+ < \theta^+\left(\frac{1}{\delta}\right)^{\beta^+-1}|s|^{\beta(x)-1}, & s \in (\delta, 1 + \delta), \\ \beta(x)(s - \delta)^{\beta(x)-1} \leq \beta^+|s|^{\beta(x)-1}, & s \in (1 + \delta, +\infty), \\ \gamma_2(x)\delta^{\gamma_2(x)-1} = \gamma_2(x)\delta^{\beta(x)-1}\delta^{\gamma_2(x)-\beta(x)} \leq \gamma^+\delta^{\beta(x)-1}, & s = \delta, \\ [\theta(x), \beta(x)] \leq \beta^+(1 + \delta)^{\beta(x)-1}, & s = 1 + \delta, \\ 0, & s \in (-\infty, 0]. \end{cases} \end{aligned} \quad (3.40)$$

Therefore,

$$\begin{aligned}
 |w_1| &\leq \left(\gamma_1^+ + \alpha^+ + \theta^+ \left(\frac{1}{\delta} \right)^{(\alpha^+-1)} \right) |t|^{\alpha(x)-1}, \quad \forall w_1 \in \partial_t F(x, t, s), \\
 |w_2| &\leq \left(\gamma_2^+ + \beta^+ + \theta^+ \left(\frac{1}{\delta} \right)^{(\beta^+-1)} \right) |t|^{\beta(x)-1}, \quad \forall w_2 \in \partial_s F(x, t, s), \\
 \limsup_{t \rightarrow 0, s \rightarrow 0} \frac{\langle w_1, t \rangle}{|t|^{\gamma_1(x)}} &= \begin{cases} \lim_{t \rightarrow 0} \frac{\gamma_1(x) t^{\gamma_1(x)}}{t^{\gamma_1(x)}}, & t > 0 \\ 0, & t \leq 0 \end{cases} \leq \gamma_1(x), \\
 \limsup_{t \rightarrow 0, s \rightarrow 0} \frac{\langle w_2, s \rangle}{|s|^{\gamma_2(x)}} &= \begin{cases} \lim_{s \rightarrow 0} \frac{\gamma_2(x) s^{\gamma_2(x)}}{s^{\gamma_2(x)}}, & s > 0 \\ 0, & s \leq 0 \end{cases} \leq \gamma_2(x),
 \end{aligned} \tag{3.41}$$

uniformly for almost all $x \in \Omega$, all $w_1 \in \partial_t F(x, t, s)$ and $w_2 \in \partial_s F(x, t, s)$.

Thus far the results involved potential functions exhibiting $p(x)$ -sublinear. The next theorem concerns problems where the potential function is $p(x)$ -superlinear.

Theorem 3.4. Suppose that $H(F)$, (f_1) with $\alpha^- > p^+$, (f_2) with $\beta^- > q^+$, (f_3) , (f_4) and the following condition (f_5) hold.

(f_5) For almost all $x \in \Omega$ and all $t, s \in \mathbb{R}$, one has $F(x, t, s) \leq \kappa(x)$ with $\kappa \in L^{\gamma_3(x)}(\Omega)$, $1 \leq \gamma_3(x) < \min\{p^-, q^-\}$.

Then there exists a $\lambda_* > 0$ such that, for each $\lambda > \lambda_*$, the problem (P) has at least two nontrivial solutions.

Proof. The steps are similar to those of Theorem 3.2. In fact, we only need to modify Step 1 and Step 5 as follows: Step 6 shows, that φ is coercive under the condition (f_5) ; Step 7 shows, that there exists second nontrivial solution under the conditions (f_1) , (f_2) , and (f_3) . Then from Steps 6, 2, 3, 4, and 7 above, the problem (P) has at least two nontrivial solutions. $\square$

Step 6. By (f_5) , for all $(u, v) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$, $\|(u, v)\| > 1$, we have

$$\begin{aligned}
 \varphi(u, v) &= \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v|^{q(x)} dx - \lambda \int_{\Omega} F(x, u(x), v(x)) dx \\
 &\geq \frac{1}{p^+} \|u\|^{p^+} + \frac{1}{q^+} \|v\|^{q^+} - \lambda \int_{\Omega} \kappa(x) dx \longrightarrow \infty \longrightarrow \infty, \quad \text{as } \|(u, v)\| \rightarrow \infty.
 \end{aligned} \tag{3.42}$$

Step 7. Because of hypothesis (f_1) , (f_2) and mean value theorem for locally Lipschitz functions, we have

$$\begin{aligned}
 F(x, t, s) &\leq c_1 \left[|t| + 2|t|^{\alpha(x)} + |s|^{\beta(x)} \right] + c_2 \left[|s| + 2|s|^{\beta(x)} + |t|^{\alpha(x)} \right] \\
 &\leq c_1 \left[\left| \frac{t}{\beta} \right|^{\alpha(x)-1} |t| + 2|t|^{\alpha(x)} + |s|^{\beta(x)} \right] \\
 &\quad + c_2 \left[\left| \frac{s}{\beta} \right|^{\beta(x)-1} |s| + 2|s|^{\beta(x)} + |t|^{\alpha(x)} \right] \\
 &\leq c_1 \left[\left| \frac{1}{\beta} \right|^{\alpha^+-1} |t|^{\alpha(x)} + 2|t|^{\alpha(x)} + |s|^{\beta(x)} \right] \\
 &\quad + c_2 \left[\left| \frac{1}{\beta} \right|^{\beta^+-1} |t|^{\beta(x)} + 2|s|^{\beta(x)} + |t|^{\alpha(x)} \right] \\
 &= c_7 |t|^{\alpha(x)} + c_8 |t|^{\beta(x)}
 \end{aligned} \tag{3.43}$$

for a.e. $x \in \Omega$, all $|t| \geq \beta$, $|s| \geq \beta$ with $c_7, c_8 > 0$.

Combining (3.31) and (3.43), it follows that

$$|F(x, t, s)| \leq \left[\mu_1(x) |t|^{\gamma_1(x)} + c_7 |t|^{\alpha(x)} \right] + \left[\mu_2(x) |s|^{\gamma_2(x)} + c_8 |s|^{\alpha(x)} \right] \tag{3.44}$$

for a.e. $x \in \Omega$ and all $t, s \in \mathbb{R}$.

Thus, for all $\lambda > \lambda_*$, $\|(u, v)\| < 1$, $|u|_{\gamma_1(x)} < 1$, $|u|_{\alpha(x)} < 1$, $|v|_{\gamma_2(x)} < 1$ and $|v|_{\beta(x)} < 1$, we have

$$\begin{aligned}
 \varphi(u, v) &= \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{1}{q(x)} |\nabla v|^{q(x)} dx - \lambda \int_{\Omega} F(x, u(x), v(x)) dx \\
 &\geq \frac{1}{p^+} \|u\|^{p^+} + \frac{1}{q^+} \|v\|^{q^+} - \lambda \int_{\Omega} \mu_1(x) |u|^{\gamma_1(x)} dx - \lambda c_7 \int_{\Omega} |u|^{\alpha(x)} dx \\
 &\quad - \lambda \int_{\Omega} \mu_2(x) |v|^{\gamma_2(x)} dx - \lambda c_8 \int_{\Omega} |v|^{\beta(x)} dx \\
 &\geq \frac{1}{p^+} \|u\|^{p^+} + \frac{1}{q^+} \|v\|^{q^+} - \lambda c_9 \|u\|^{\gamma_1^-} - \lambda c_7 \|u\|^{\alpha^-} - \lambda c_{10} \|v\|^{\gamma_2^-} - \lambda c_8 \|v\|^{\beta^-}.
 \end{aligned} \tag{3.45}$$

So, for $\rho > 0$ small enough, there exists a $\nu > 0$ such that

$$\varphi(u, v) > \nu, \quad \text{for } \|u, v\| = \rho \tag{3.46}$$

and $\|(u_0, v_0)\| > \rho$. Arguing as in proof of Step 4 of Theorem 3.2, we conclude that φ satisfies the nonsmooth C-condition. So by the Nonsmooth Mountain Pass Theorem (cf. Theorem 2.7), we can get $(u_1, v_1) \in W_0^{1,p(x)}(\Omega) \times W_0^{1,q(x)}(\Omega)$ which satisfies

$$\varphi(u_1, v_1) = c > 0, \quad m(u_1, v_1) = 0. \tag{3.47}$$

Therefore, (u_1, v_1) is second nontrivial of problem (P) .

Remark 3.5. Consider the following nonsmooth locally Lipschitz function:

$$F(x, t, s) = \begin{cases} -|t|^{\gamma_1(x)} - |s|^{\gamma_2(x)}, & |t| \leq 1 \text{ or } |s| \leq 1, \\ \cos(\pi|t|) + \cos(\pi|s|), & |t| > 1; \text{ or } |s| > 1, \end{cases} \quad (3.48)$$

In the following, we will show that $F(x, t, s)$ satisfies hypotheses $H(F)$ and (f_1) – (f_5) .

It is clear that $F(x, 0, 0) = 0$ for a.e. $x \in \Omega$, then hypotheses $H(F)$ is satisfied. A direct verification shows that conditions (f_4) and (f_5) are satisfied. Note that

$$\begin{aligned} \partial_t F(x, t, s) &= \begin{cases} -\gamma_1(x)t^{\gamma_1(x)-1}, & 0 \leq t < 1, \\ \gamma_1(x)(-t)^{\gamma_1(x)-1}, & -1 < t \leq 0, \\ [-\gamma_1(x), 0], & t = 1, \\ [0, \gamma_1(x)], & t = -1, \\ \{-\pi \sin(\pi t)\}, & |t| > 1, \end{cases} \\ \partial_s F(x, t, s) &= \begin{cases} -\gamma_2(x)s^{\gamma_2(x)-1}, & 0 \leq s < 1, \\ \gamma_2(x)(-s)^{\gamma_2(x)-1}, & -1 < s \leq 0, \\ [-\gamma_2(x), 0], & s = 1, \\ [0, \gamma_2(x)], & s = -1, \\ \{-\pi \sin(\pi s)\}, & |s| > 1. \end{cases} \end{aligned} \quad (3.49)$$

So,

$$\begin{aligned} |w_1| &\leq (\gamma_1(x) + \pi)|t|^{\gamma_1(x)-1}, \quad \forall w_1 \in \partial_t F(x, t, s), \\ |w_1| &\leq (\gamma_2(x) + \pi)|s|^{\gamma_2(x)-1}, \quad \forall w_2 \in \partial_s F(x, t, s), \\ \limsup_{t \rightarrow 0, s \rightarrow 0} \frac{\langle w_1, t \rangle}{|t|^{\gamma_1(x)}} &= \begin{cases} \lim_{t \rightarrow 0} \frac{-\gamma_1(x)t^{\gamma_1(x)}}{t^{\gamma_1(x)}}, & t > 0 \\ \lim_{t \rightarrow 0} \frac{-\gamma_1(x)(-t)^{\gamma_1(x)}}{(-t)^{\gamma_1(x)}}, & t \leq 0 \end{cases} = -\gamma_1(x) \leq \gamma_1(x), \\ \limsup_{t \rightarrow 0, s \rightarrow 0} \frac{\langle w_2, s \rangle}{|s|^{\gamma_2(x)}} &= \begin{cases} \lim_{s \rightarrow 0} \frac{-\gamma_2(x)s^{\gamma_2(x)}}{s^{\gamma_2(x)}}, & s > 0 \\ \lim_{s \rightarrow 0} \frac{-\gamma_2(x)(-s)^{\gamma_2(x)}}{(-s)^{\gamma_2(x)}}, & s \leq 0 \end{cases} = -\gamma_2(x) \leq \gamma_2(x) \end{aligned} \quad (3.50)$$

which shows that assumptions (f_1) , (f_2) , and (f_3) are fulfilled.

Acknowledgments

This paper was supported by the National Natural Science Foundation of China (nos. 11126286, 11001063, 10971043), the Fundamental Research Funds for the Central Universities

(no. 2012), China Postdoctoral Science Foundation Funded Project (no. 20110491032), and the National Natural Science Foundation of Heilongjiang Province (no. A200803).

References

- [1] M. Ružicka, *Electrorheological Fluids: Modeling and Mathematical Theory*, vol. 1748 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 2000.
- [2] V. V. Zhikov, "Averaging of functionals of the calculus of variations and elasticity theory," *Izvestiya*, vol. 29, no. 1, pp. 33–66, 1987.
- [3] G. Dai and R. Hao, "Existence of solutions for a $p(x)$ -Kirchhoff-type equation," *Journal of Mathematical Analysis and Applications*, vol. 359, no. 1, pp. 275–284, 2009.
- [4] X. Fan and D. Zhao, "On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$," *Journal of Mathematical Analysis and Applications*, vol. 263, no. 2, pp. 424–446, 2001.
- [5] X. L. Fan and D. Zhao, "On the generalized Orlicz-sobolev spaces $W^{k,p(x)}(\Omega)$," *Journal of Gansu Education College*, vol. 12, no. 1, pp. 1–6, 1998.
- [6] B. Ge, X. Xue, and Q. Zhou, "The existence of radial solutions for differential inclusion problems in $\mathbb{R}^N$ involving the $p(x)$ -Laplacian," *Nonlinear Analysis*, vol. 73, no. 3, pp. 622–633, 2010.
- [7] B. Ge and X. Xue, "Multiple solutions for inequality Dirichlet problems by the $p(x)$ -Laplacian," *Nonlinear Analysis*, vol. 11, no. 4, pp. 3198–3210, 2010.
- [8] B. Ge, X.-P. Xue, and M.-S. Guo, "Three solutions to inequalities of Dirichlet problem driven by $p(x)$ -Laplacian," *Applied Mathematics and Mechanics*, vol. 31, no. 10, pp. 1283–1292, 2010.
- [9] B. Ge, X. Xue, and Q. Zhou, "Existence of at least five solutions for a differential inclusion problem involving the $p(x)$ -Laplacian," *Nonlinear Analysis*, vol. 12, no. 4, pp. 2304–2318, 2011.
- [10] B. Ge, Q.-M. Zhou, and X.-P. Xue, "Infinitely many solutions for a differential inclusion problem in $\mathbb{R}^N$ involving $p(x)$ -Laplacian and oscillatory terms," *Zeitschrift für Angewandte Mathematik und Physik*. In press.
- [11] B. Ge, Q.-M. Zhou, and X.-P. Xue, "Multiplicity of solutions for differential inclusion problems in $\mathbb{R}^N$ involving the $p(x)$ -Laplacian," *Monatshefte für Mathematik*. In press.
- [12] D. G. de Figueiredo, "Semilinear elliptic systems: a survey of superlinear problems," *Resenhas do Instituto de Matemática e Estatística da Universidade de São Paulo*, vol. 2, no. 4, pp. 373–391, 1996.
- [13] F. H. Clarke, *Optimization and Nonsmooth Analysis*, Canadian Mathematical Society Series of Monographs and Advanced Texts, John Wiley & Sons, New York, NY, USA, 1983, A Wiley-Interscience Publication.
- [14] X. Fan, Q. Zhang, and D. Zhao, "Eigenvalues of $p(x)$ -Laplacian Dirichlet problem," *Journal of Mathematical Analysis and Applications*, vol. 302, no. 2, pp. 306–317, 2005.
- [15] C. Li and C.-L. Tang, "Three solutions for a class of quasilinear elliptic systems involving the (p, q) -Laplacian," *Nonlinear Analysis*, vol. 69, no. 10, pp. 3322–3329, 2008.
- [16] A. Kristály, "Existence of two non-trivial solutions for a class of quasilinear elliptic variational systems on strip-like domains," *Proceedings of the Edinburgh Mathematical Society. Series II*, vol. 48, no. 2, pp. 465–477, 2005.
- [17] L. Boccardo and D. Guedes de Figueiredo, "Some remarks on a system of quasilinear elliptic equations," *Nonlinear Differential Equations and Applications*, vol. 9, no. 3, pp. 309–323, 2002.
- [18] X. Fan, J. Shen, and D. Zhao, "Sobolev embedding theorems for spaces $W^{k,p(x)}(\Omega)$," *Journal of Mathematical Analysis and Applications*, vol. 262, no. 2, pp. 749–760, 2001.
- [19] X.-L. Fan and Q.-H. Zhang, "Existence of solutions for $p(x)$ -Laplacian Dirichlet problem," *Nonlinear Analysis*, vol. 52, no. 8, pp. 1843–1852, 2003.
- [20] X. Fan and S.-G. Deng, "Remarks on Ricceri's variational principle and applications to the $p(x)$ -Laplacian equations," *Nonlinear Analysis*, vol. 67, no. 11, pp. 3064–3075, 2007.
- [21] K.C. Chang, *Critical Point Theory and Applications*, Shanghai Scientific and Technology Press, Shanghai, China, 1996.
- [22] N. C. Kouragenis and N. S. Papageorgiou, "Nonsmooth critical point theory and nonlinear elliptic equations at resonance," *Journal of the Australian Mathematical Society. Series A*, vol. 69, no. 2, pp. 245–271, 2000.
- [23] K. C. Chang, "Variational methods for nondifferentiable functionals and their applications to partial differential equations," *Journal of Mathematical Analysis and Applications*, vol. 80, no. 1, pp. 102–129, 1981.