Research Article

Approximately Ternary Homomorphisms and Derivations on C^* -Ternary Algebras

**M. Eshaghi Gordji,¹ A. Ebadian,² N. Ghobadipour,³
 J. M. Rassias,⁴ and M. B. Savadkouhi⁵**

¹ Department of Mathematics, Semnan University, Semnan 35195-363, Iran

² Department of Mathematics, Payame Noor University, Tabriz Branch, Tabriz, Iran

³ Department of Mathematics, Urmia University, Urmia, Iran

⁴ Section of Mathematics and Informatics, Pedagogical Department, National and Kapodistrian University of Athens, 4, Agamemnonos Street, Aghia Paraskevi, 15342 Athens, Greece

⁵ Department of Mathematics, Islamic Azad University, Mahdishahr Branch, Mahdishahr, Semnan, Iran

Correspondence should be addressed to M. Eshaghi Gordji, madjid.eshaghi@gmail.com and A. Ebadian, ebadian.ali@gmail.com

Received 4 December 2011; Accepted 17 January 2012

Academic Editor: Josip E. Pecaric

Copyright © 2012 M. Eshaghi Gordji et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the stability and superstability of ternary homomorphisms between C^* -ternary algebras and derivations on C^* -ternary algebras, associated with the following functional equation $f((x_2 - x_1)/3) + f((x_1 - 3x_3)/3) + f((3x_1 + 3x_3 - x_2)/3) = f(x_1)$.

1. Introduction

A C^* -ternary algebra is a complex Banach space A , equipped with a ternary product $(x, y, z) \mapsto [x, y, z]$ of A^3 into A , which is $\mathbb{C}$ -linear in the outer variables, conjugate $\mathbb{C}$ -linear in the middle variable, and associative in the sense that $[x, y, [z, w, v]] = [x, [w, z, y], v] = [[x, y, z], w, v]$, and satisfies $\|[x, y, z]\| \leq \|x\| \cdot \|y\| \cdot \|z\|$ and $\|[x, x, x]\| = \|x\|^3$. If a C^* -ternary algebra $(A, [\cdot, \cdot, \cdot])$ has an identity, that is, an element $e \in A$ such that $x = [x, e, e] = [e, e, x]$ for all $x \in A$, then it is routine to verify that A , endowed with $xoy := [x, e, y]$ and $x^* := [e, x, e]$, is a unital C^* -algebra. Conversely, if (A, o) is a unital C^* -algebra, then $[x, y, z] := xoy^*oz$ makes A into a C^* -ternary algebra. A $\mathbb{C}$ -linear mapping $H : A \rightarrow B$ is called a C^* -ternary algebra homomorphism if

$$H([x, y, z]) = [H(x), H(y), H(z)], \quad (1.1)$$

for all $x, y, z \in A$. A $\mathbb{C}$ -linear mapping $\delta : A \rightarrow A$ is called a C^* -ternary algebra derivation if

$$\delta([x, y, z]) = [\delta(x), y, z] + [x, \delta(y), z] + [x, y, \delta(z)], \quad (1.2)$$

for all $x, y, z \in A$.

Ternary structures and their generalization the so-called n -ary structures raise certain hopes in view of their applications in physics (see [1–8]).

We say a functional equation ζ is stable if any function g satisfying the equation ζ approximately is near to true solution of ζ . Moreover, ζ is superstable if every approximately solution of ζ is an exact solution of it.

The study of stability problems originated from a famous talk given by Ulam [9] in 1940: “Under what condition does there exist a homomorphism near an approximate homomorphism?” In the next year 1941, Hyers [10] answered affirmatively the question of Ulam for additive mappings between Banach spaces.

A generalized version of the theorem of Hyers for approximately additive maps was given by Rassias [11] in 1978 as follows.

Theorem 1.1. *Let $f : E_1 \rightarrow E_2$ be a mapping from a normed vector space E_1 into a Banach space E_2 subject to the inequality:*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p), \quad (1.3)$$

for all $x, y \in E_1$, where ϵ and p are constants with $\epsilon > 0$ and $p < 1$. Then, there exists a unique additive mapping $T : E_1 \rightarrow E_2$ such that

$$\|f(x) - T(x)\| \leq \frac{2\epsilon}{2-p} \|x\|^p, \quad (1.4)$$

for all $x \in E_1$.

The stability phenomenon that was introduced and proved by Rassias is called Hyers-Ulam-Rassias stability. And then the stability problems of several functional equations have been extensively investigated by a number of authors, and there are many interesting results concerning this problem (see [12–27]).

Throughout this paper, we assume that A is a C^* -ternary algebra with norm $\|\cdot\|_A$ and that B is a C^* -ternary algebra with norm $\|\cdot\|_B$. Moreover, we assume that $n_0 \in \mathbb{N}$ is a positive integer and suppose that $\mathbb{T}_{1/n_0}^1 := \{e^{i\theta}; 0 \leq \theta \leq 2\pi/n_0\}$.

2. Superstability

In this section, first we investigate homomorphisms between C^* -ternary algebras. We need the following Lemma in the main results of the paper.

Lemma 2.1. *Let $f : A \rightarrow B$ be a mapping such that*

$$\left\| f\left(\frac{x_2 - x_1}{3}\right) + f\left(\frac{x_1 - 3x_3}{3}\right) + f\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) \right\|_B \leq \|f(x_1)\|_B, \quad (2.1)$$

for all $x_1, x_2, x_3 \in A$. Then f is additive.

Proof. Letting $x_1 = x_2 = x_3 = 0$ in (2.1), we get

$$\|3f(0)\|_B \leq \|f(0)\|_B. \quad (2.2)$$

So $f(0) = 0$. Letting $x_1 = x_2 = 0$ in (2.1), we get

$$\|f(-x_3) + f(x_3)\|_B \leq \|f(0)\|_B = 0, \quad (2.3)$$

for all $x_3 \in A$. Hence $f(-x_3) = -f(x_3)$ for all $x_3 \in A$. Letting $x_1 = 0$ and $x_2 = 6x_3$ in (2.1), we get

$$\|f(2x_3) - 2f(x_3)\|_B \leq \|f(0)\|_B = 0, \quad (2.4)$$

for all $x_3 \in A$. Hence

$$f(2x_3) = 2f(x_3), \quad (2.5)$$

for all $x_3 \in A$. Letting $x_1 = 0$ and $x_2 = 9x_3$ in (2.1), we get

$$\|f(3x_3) - f(x_3) - 2f(x_3)\|_B \leq \|f(0)\|_B = 0, \quad (2.6)$$

for all $x_3 \in A$. Hence

$$f(3x_3) = 3f(x_3), \quad (2.7)$$

for all $x_3 \in A$. Letting $x_1 = 0$ in (2.1), we get

$$\left\| f\left(\frac{x_2}{3}\right) + f(-x_3) + f\left(x_3 - \frac{x_2}{3}\right) \right\|_B \leq \|f(0)\|_B = 0, \quad (2.8)$$

for all $x_2, x_3 \in A$. So

$$f\left(\frac{x_2}{3}\right) + f(-x_3) + f\left(x_3 - \frac{x_2}{3}\right) = 0, \quad (2.9)$$

for all $x_2, x_3 \in A$. Let $t_1 = x_3 - (x_2/3)$ and $t_2 = x_2/3$ in (2.9). Then

$$f(t_2) - f(t_1 + t_2) + f(t_1) = 0, \quad (2.10)$$

for all $t_1, t_2 \in A$, this means that f is additive. □

Now, we prove the first result in superstability as follows.

Theorem 2.2. Let $p \neq 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow B$ be a mapping such that

$$\left\| f\left(\frac{x_2 - x_1}{3}\right) + f\left(\frac{x_1 - 3\mu x_3}{3}\right) + \mu f\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) \right\|_B \leq \|f(x_1)\|_B, \quad (2.11)$$

$$\|f([x_1, x_2, x_3]) - [f(x_1), f(x_2), f(x_3)]\|_B \leq \theta(\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p}), \quad (2.12)$$

for all $\mu \in \mathbb{T}_{1/n_0}^1$ and all $x_1, x_2, x_3 \in A$. Then, the mapping $f : A \rightarrow B$ is a C^* -ternary algebra homomorphism.

Proof. Assume $p > 1$.

Let $\mu = 1$ in (2.11). By Lemma 2.1, the mapping $f : A \rightarrow B$ is additive. Letting $x_1 = x_2 = 0$ in (2.11), we get

$$\|f(-\mu x_3) + \mu f(x_3)\|_B \leq \|f(0)\|_B = 0, \quad (2.13)$$

for all $x_3 \in A$ and $\mu \in \mathbb{T}^1$. So

$$-f(\mu x_3) + \mu f(x_3) = f(-\mu x_3) + \mu f(x_3) = 0, \quad (2.14)$$

for all $x_3 \in A$ and all $\mu \in \mathbb{T}^1$. Hence $f(\mu x_3) = \mu f(x_3)$ for all $x_3 \in A$ and all $\mu \in \mathbb{T}_{1/n_0}^1$. By same reasoning as proof of Theorem 2.2 of [28], the mapping $f : A \rightarrow B$ is $\mathbb{C}$ -linear. It follows from (2.12) that

$$\begin{aligned} & \|f([x_1, x_2, x_3]) - [f(x_1), f(x_2), f(x_3)]\|_B \\ &= \lim_{n \rightarrow \infty} 8^n \left\| f\left(\frac{[x_1, x_2, x_3]}{2^n \cdot 2^n \cdot 2^n}\right) - \left[f\left(\frac{x_1}{2^n}\right), f\left(\frac{x_2}{2^n}\right), f\left(\frac{x_3}{2^n}\right)\right] \right\|_B \\ &\leq \lim_{n \rightarrow \infty} \frac{8^n \theta}{8^{np}} (\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p}) = 0, \end{aligned} \quad (2.15)$$

for all $x_1, x_2, x_3 \in A$. Thus,

$$f([x_1, x_2, x_3]) = [f(x_1), f(x_2), f(x_3)], \quad (2.16)$$

for all $x_1, x_2, x_3 \in A$. Hence, the mapping $f : A \rightarrow B$ is a C^* -ternary algebra homomorphism. Similarly, one obtains the result for the case $p < 1$. $\square$

Now, we establish the superstability of derivations on C^* -ternary algebras as follows.

Theorem 2.3. Let $p \neq 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow A$ be a mapping satisfying (2.11) such that

$$\begin{aligned} & \|f([x_1, x_2, x_3]) - [f(x_1), x_2, x_3] - [x_1, f(x_2), x_3] - [x_1, x_2, f(x_3)]\|_A \\ & \leq \theta \left(\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p} \right), \end{aligned} \quad (2.17)$$

for all $x_1, x_2, x_3 \in A$. Then the mapping $f : A \rightarrow A$ is a C^* -ternary derivation.

Proof. Assume $p > 1$.

By the Theorem 2.2, the mapping $f : A \rightarrow A$ is $\mathbb{C}$ -linear. It follows from (2.17) that

$$\begin{aligned} & \|f([x_1, x_2, x_3]) - [f(x_1), x_2, x_3] - [x_1, f(x_2), x_3] - [x_1, x_2, f(x_3)]\|_A \\ & = \lim_{n \rightarrow \infty} 8^n \left\| f\left(\frac{[x_1, x_2, x_3]}{8^n}\right) - \left[f\left(\frac{x_1}{2^n}\right), \frac{x_2}{2^n}, \frac{x_3}{2^n}\right] - \left[\frac{x_1}{2^n}, f\left(\frac{x_2}{2^n}\right), \frac{x_3}{2^n}\right] \right. \\ & \quad \left. - \left[\frac{x_1}{2^n}, \frac{x_2}{2^n}, f\left(\frac{x_3}{2^n}\right)\right] \right\|_A \\ & \leq \lim_{n \rightarrow \infty} \frac{8^n \theta}{8^{np}} \left(\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p} \right) = 0, \end{aligned} \quad (2.18)$$

for all $x_1, x_2, x_3 \in A$. So

$$f([x_1, x_2, x_3]) = [f(x_1), x_2, x_3] + [x_1, f(x_2), x_3] + [x_1, x_2, f(x_3)] \quad (2.19)$$

for all $x_1, x_2, x_3 \in A$. Thus, the mapping $f : A \rightarrow A$ is a C^* -ternary derivation. Similarly, one obtains the result for the case $p < 1$. $\square$

3. Stability

First we prove the generalized Hyers-Ulam-Rassias stability of homomorphisms in C^* -ternary algebras.

Theorem 3.1. Let $p > 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow B$ be a mapping such that

$$\begin{aligned} & \left\| f\left(\frac{x_2 - x_1}{3}\right) + f\left(\frac{x_1 - 3\mu x_3}{3}\right) + \mu f\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) - f(x_1) \right\|_B \\ & \leq \theta \left(\|x_1\|_A^p + \|x_2\|_A^p + \|x_3\|_A^p \right), \end{aligned} \quad (3.1)$$

$$\|f([x_1, x_2, x_3]) - [f(x_1), f(x_2), f(x_3)]\|_B \leq \theta \left(\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p} \right), \quad (3.2)$$

for all $\mu \in \mathbb{T}_{1/n_0}^1$, and all $x_1, x_2, x_3 \in A$. Then there exists a unique C^* -ternary homomorphism $H : A \rightarrow B$ such that

$$\|H(x_1) - f(x_1)\|_B \leq \frac{\theta(1+2^p)\|x_1\|_A^p}{1-3^{1-p}}, \quad (3.3)$$

for all $x_1 \in A$.

Proof. Let us assume $\mu = 1$, $x_2 = 2x_1$ and $x_3 = 0$ in (3.1). Then we get

$$\left\| 3f\left(\frac{x_1}{3}\right) - f(x_1) \right\|_B \leq \theta(1+2^p)\|x_1\|_A^p, \quad (3.4)$$

for all $x_1 \in A$. So by induction, we have

$$\left\| 3^n f\left(\frac{x_1}{3^n}\right) - f(x_1) \right\|_B \leq \theta(1+2^p)\|x_1\|_A^p \sum_{i=0}^{n-1} 3^{i(1-p)}, \quad (3.5)$$

for all $x_1 \in A$. Hence

$$\begin{aligned} \left\| 3^{n+m} f\left(\frac{x_1}{3^{n+m}}\right) - 3^m f\left(\frac{x_1}{3^m}\right) \right\|_B &\leq \theta(1+2^p)\|x_1\|_A^p \sum_{i=0}^{n-1} 3^{(i+m)(1-p)} \\ &\leq \theta(1+2^p)\|x_1\|_A^p \sum_{i=m}^{n+m-1} 3^{i(1-p)}, \end{aligned} \quad (3.6)$$

for all nonnegative integers m and n with $n \geq m$, and all $x_1 \in A$. It follows that the sequence $\{3^n f(x_1/3^n)\}$ is a Cauchy sequence for all $x_1 \in A$. Since B is complete, the sequence $\{3^n f(x_1/3^n)\}$ converges. Thus, one can define the mapping $H : A \rightarrow B$ by

$$H(x_1) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x_1}{3^n}\right), \quad (3.7)$$

for all $x_1 \in A$. Moreover, letting $m = 0$ and passing the limit $n \rightarrow \infty$ in (3.6), we get (3.3). It follows from (3.1) that

$$\begin{aligned} &\left\| H\left(\frac{x_2 - x_1}{3}\right) + H\left(\frac{x_1 - 3\mu x_3}{3}\right) + \mu H\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) - H(x_1) \right\|_B \\ &= \lim_{n \rightarrow \infty} 3^n \left\| f\left(\frac{x_2 - x_1}{3^{n+1}}\right) + f\left(\frac{x_1 - 3\mu x_3}{3^{n+1}}\right) + f\left(\frac{3x_1 + 3x_3 - x_2}{3^{n+1}}\right) - f\left(\frac{x_1}{3^n}\right) \right\|_B \\ &\leq \lim_{n \rightarrow \infty} \frac{3^n \theta}{3^{np}} (\|x_1\|_A^p + \|x_2\|_A^p + \|x_3\|_A^p) = 0, \end{aligned} \quad (3.8)$$

for all $\mu \in \mathbb{T}_{1/n_0}^1$, and all $x_1, x_2, x_3 \in A$. So

$$H\left(\frac{x_2 - x_1}{3}\right) + H\left(\frac{x_1 - 3\mu x_3}{3}\right) + \mu H\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) = H(x_1), \quad (3.9)$$

for all $\mu \in \mathbb{T}_{1/n_0}^1$, and all $x_1, x_2, x_3 \in A$. By the same reasoning as proof of Theorem 2.2 of [28], the mapping $H : A \rightarrow B$ is $\mathbb{C}$ -linear.

Now, let $H' : A \rightarrow B$ be another additive mapping satisfying (3.3). Then, we have

$$\begin{aligned} \|H(x_1) - H'(x_1)\|_B &= 3^n \left\| H\left(\frac{x_1}{3^n}\right) - H'\left(\frac{x_1}{3^n}\right) \right\|_B \\ &\leq 3^n \left(\left\| H\left(\frac{x_1}{3^n}\right) - f\left(\frac{x_1}{3^n}\right) \right\|_B + \left\| H'\left(\frac{x_1}{3^n}\right) - f\left(\frac{x_1}{3^n}\right) \right\|_B \right) \\ &\leq \frac{2 \cdot 3^n \theta (1 + 2^p)}{3^{np} (1 - 3^{1-p})} \|x\|_A^p, \end{aligned} \quad (3.10)$$

which tends to zero as $n \rightarrow \infty$ for all $x_1 \in A$. So we can conclude that $H(x_1) = H'(x_1)$ for all $x_1 \in A$. This proves the uniqueness of H .

It follows from (3.2) that

$$\begin{aligned} &\|H([x_1, x_2, x_3]) - [H(x_1), H(x_2), H(x_3)]\|_B \\ &= \lim_{n \rightarrow \infty} 27^n \left\| f\left(\frac{[x_1, x_2, x_3]}{3^n \cdot 3^n \cdot 3^n}\right) - \left[f\left(\frac{x_1}{3^n}\right), f\left(\frac{x_2}{3^n}\right), f\left(\frac{x_3}{3^n}\right) \right] \right\|_B \\ &\leq \lim_{n \rightarrow \infty} \frac{27^n \theta}{27^{np}} \left(\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p} \right) = 0, \end{aligned} \quad (3.11)$$

for all $x_1, x_2, x_3 \in A$.

Thus, the mapping $H : A \rightarrow B$ is a unique C^* -ternary homomorphism satisfying (3.3). $\square$

Theorem 3.2. Let $p < 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow B$ be a mapping satisfying (3.1) and (3.2). Then, there exists a unique C^* -ternary homomorphism $H : A \rightarrow B$ such that

$$\|H(x_1) - f(x_1)\|_B \leq \frac{\theta(1 + 2^p) \|x_1\|_A^p}{3^{1-p} - 1}, \quad (3.12)$$

for all $x_1 \in A$.

Proof. The proof is similar to the proof of Theorem 3.1. $\square$

Now, we prove the generalized Hyers-Ulam-Rassias stability of derivations on C^* -ternary algebras.

Theorem 3.3. Let $p > 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow A$ be a mapping such that

$$\left\| f\left(\frac{x_2 - x_1}{3}\right) + f\left(\frac{x_1 - 3\mu x_3}{3}\right) + \mu f\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) - f(x_1) \right\|_A \leq \theta (\|x_1\|_A^p + \|x_2\|_A^p + \|x_3\|_A^p), \quad (3.13)$$

$$\|f([x_1, x_2, x_3]) - [f(x_1), x_2, x_3] - [x_1, f(x_2), x_3] - [x_1, x_2, f(x_3)]\|_A \leq \theta (\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p}), \quad (3.14)$$

for all $\mu \in \mathbb{T}_{1/n_0}^1$, and all $x_1, x_2, x_3 \in A$. Then, there exists a unique C^* -ternary derivation $D : A \rightarrow A$ such that

$$\|D(x_1) - f(x_1)\|_A \leq \frac{\theta(1 + 2^p)\|x_1\|_A^p}{1 - 3^{1-p}}, \quad (3.15)$$

for all $x_1 \in A$.

Proof. By the same reasoning as in the proof of the Theorem 3.1, there exists a unique $\mathbb{C}$ -linear mapping $D : A \rightarrow A$ satisfying (3.15). The mapping $D : A \rightarrow A$ is defined by

$$D(x_1) := \lim_{n \rightarrow \infty} 3^n f\left(\frac{x_1}{3^n}\right), \quad (3.16)$$

for all $x_1 \in A$. It follows from (3.14) that

$$\begin{aligned} & \|D([x_1, x_2, x_3]) - [D(x_1), x_2, x_3] - [x_1, D(x_2), x_3] - [x_1, x_2, D(x_3)]\|_A \\ &= \lim_{n \rightarrow \infty} 27^n \left\| \frac{[x_1, x_2, x_3]}{3^n \cdot 3^n \cdot 3^n} - \left[f\left(\frac{x_1}{3^n}\right), \frac{x_2}{3^n}, \frac{x_3}{3^n} \right] - \left[\frac{x_1}{3^n}, f\left(\frac{x_2}{3^n}\right), \frac{x_3}{3^n} \right] - \left[\frac{x_1}{3^n}, \frac{x_2}{3^n}, f\left(\frac{x_3}{3^n}\right) \right] \right\|_A \\ &\leq \lim_{n \rightarrow \infty} \frac{27^n \theta}{27^{np}} (\|x_1\|_A^{3p} + \|x_2\|_A^{3p} + \|x_3\|_A^{3p}) = 0, \end{aligned} \quad (3.17)$$

for all $x_1, x_2, x_3 \in A$. So

$$D([x_1, x_2, x_3]) = [D(x_1), x_2, x_3] + [x_1, D(x_2), x_3] + [x_1, x_2, D(x_3)] \quad (3.18)$$

for all $x_1, x_2, x_3 \in A$.

Thus, the mapping $D : A \rightarrow A$ is a unique C^* -ternary derivation satisfying (3.15). $\square$

Theorem 3.4. Let $p < 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow A$ be a mapping satisfying (3.13) and (3.14). Then, there exists a unique C^* -ternary derivation $D : A \rightarrow A$ such that

$$\|D(x_1) - f(x_1)\|_A \leq \frac{\theta(1 + 2^p)\|x_1\|_A^p}{3^{1-p} - 1}, \quad (3.19)$$

for all $x_1 \in A$.

Proof. The proof is similar to the proof of Theorems 3.1 and 3.3. $\square$

4. Conclusions

In this paper, we have analyzed some detail C^* -ternary algebras and derivations on C^* -ternary algebras, associated with the following functional equation:

$$f\left(\frac{x_2 - x_1}{3}\right) + f\left(\frac{x_1 - 3x_3}{3}\right) + f\left(\frac{3x_1 + 3x_3 - x_2}{3}\right) = f(x_1). \quad (4.1)$$

A detailed study of how we can have the generalized Hyers-Ulam-Rassias stability of homomorphisms and derivations on C^* -ternary algebras is given.

Acknowledgment

This paper was partially supported by Research Program of Payame Noor University of Tabriz.

References

- [1] F. Bagarello and G. Morchio, "Dynamics of mean-field spin models from basic results in abstract differential equations," *Journal of Statistical Physics*, vol. 66, no. 3-4, pp. 849-866, 1992.
- [2] N. Bazunova, A. Borowiec, and R. Kerner, "Universal differential calculus on ternary algebras," *Letters in Mathematical Physics*, vol. 67, no. 3, pp. 195-206, 2004.
- [3] M. B. Savadkouhi, M. E. Gordji, J. M. Rassias, and N. Ghobadipour, "Approximate ternary Jordan derivations on Banach ternary algebras," *Journal of Mathematical Physics*, vol. 50, no. 4, Article ID 042303, 9 pages, 2009.
- [4] A. Ebadian, N. Ghobadipour, and M. Eshaghi Gordji, "A fixed point method for perturbation of bi-multipliers and Jordan bi-multipliers in C^* -ternary algebras," *Journal of Mathematical Physics*, vol. 51, no. 10, Article ID 103508, 2010.
- [5] M. E. Gordji, R. Khodabakhsh, and H. Khodaei, "On approximate n -ary derivations," *International Journal of Geometric Methods in Modern Physics*, vol. 8, no. 3, pp. 485-500, 2011.
- [6] L. Vainerman and R. Kerner, "On special classes of n -algebras," *Journal of Mathematical Physics*, vol. 37, no. 5, pp. 2553-2565, 1996.
- [7] S. M. Ulam, *Problems in Modern Mathematics*, John Wiley & Sons, New York, NY, USA, 1940.
- [8] Th. M. Rassias and J. Tabor, "What is left of Hyers-Ulam stability?" *Journal of Natural Geometry*, vol. 1, no. 2, pp. 65-69, 1992.
- [9] G. L. Sewell, *Quantum Mechanics and Its Emergent Macrophysics*, Princeton University Press, Princeton, NJ, USA, 2002.
- [10] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222-224, 1941.
- [11] T. M. Rassias, "On the stability of minimum points," *Mathematica*, vol. 45(68), no. 1, pp. 93-104, 2003.

- [12] R. Kerner, "The cubic chessboard," *Classical and Quantum Gravity*, vol. 14, no. 1A, pp. A203–A225, 1997.
- [13] T. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [14] A. Ebadian, S. Kaboli Gharetapeh, and M. Eshaghi Gordji, "Nearly Jordan $\ast$ -homomorphisms between unital $C^\ast$ -algebras," *Abstract and Applied Analysis*, vol. 2011, Article ID 513128, 12 pages, 2011.
- [15] A. Ebadian, A. Najati, and M. Eshaghi Gordji, "On approximate additive-quartic and quadratic-cubic functional equations in two variables on abelian groups," *Results in Mathematics*, vol. 58, no. 1-2, pp. 39–53, 2010.
- [16] M. Eshaghi Gordji, M. B. Ghaemi, S. Kaboli Gharetapeh, S. Shams, and A. Ebadian, "On the stability of $J^\ast$ -derivations," *Journal of Geometry and Physics*, vol. 60, no. 3, pp. 454–459, 2010.
- [17] M. E. Gordji, N. Ghobadipour, and C. Park, "Jordan $\ast$ -homomorphisms on $C^\ast$ -algebras," *Operators and Matrices*, vol. 5, no. 3, pp. 541–551, 2011.
- [18] M. E. Gordji and N. Ghobadipour, "Stability of (α, β, γ) -derivations on Lie $C^\ast$ -algebras," *International Journal of Geometric Methods in Modern Physics*, vol. 7, no. 7, pp. 1093–1102, 2010.
- [19] M. Eshaghi Gordji and A. Najati, "Approximately $J^\ast$ -homomorphisms: a fixed point approach," *Journal of Geometry and Physics*, vol. 60, no. 5, pp. 809–814, 2010.
- [20] D. H. Hyers, G. Isac, and T. M. Rassias, *Stability of Functional Equations in Several Variables*, Progress in Nonlinear Differential Equations and their Applications, 34, Birkhäuser, Boston, Mass, USA, 1998.
- [21] D. H. Hyers and T. M. Rassias, "Approximate homomorphisms," *Aequationes Mathematicae*, vol. 44, no. 2-3, pp. 125–153, 1992.
- [22] G. Isac and T. M. Rassias, "On the Hyers-Ulam stability of ψ -additive mappings," *Journal of Approximation Theory*, vol. 72, no. 2, pp. 131–137, 1993.
- [23] G. Isac and T. M. Rassias, "Stability of ψ -additive mappings: applications to nonlinear analysis," *International Journal of Mathematics and Mathematical Sciences*, vol. 19, no. 2, pp. 219–228, 1996.
- [24] Th. M. Rassias, Ed., *Functional Equations and Inequalities*, vol. 518 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000.
- [25] T. M. Rassias, "On the stability of functional equations and a problem of Ulam," *Acta Applicandae Mathematicae*, vol. 62, no. 1, pp. 23–130, 2000.
- [26] Th. M. Rassias, "On the stability of the quadratic functional equation and its applications," *Studia Universitatis Babeş-Bolyai. Mathematica*, vol. 43, no. 3, pp. 89–124, 1998.
- [27] Th. M. Rassias, "The problem of S. M. Ulam for approximately multiplicative mappings," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 352–378, 2000.
- [28] M. Eshaghi Gordji, "Nearly involutions on Banach algebras; A fixed point approach," to appear in *Fixed Point Theory*.