Research Article

Interpolation and Best Approximation for Spherical Radial Basis Function Networks

Shaobo Lin, Jinshan Zeng, and Zongben Xu

Institute for Information and System Sciences, School of Mathematics and statistics, Xi'an Jiaotong University, Xi'an, Shaanxi 710049, China

Correspondence should be addressed to Zongben Xu; zbxu@mail.xjtu.edu.cn

Received 1 May 2013; Revised 17 September 2013; Accepted 18 September 2013

Academic Editor: Mieczysław Mastyło

Copyright © 2013 Shaobo Lin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Within the conventional framework of a native space structure, a smooth kernel generates a small native space, and radial basis functions stemming from the smooth kernel are intended to approximate only functions from this small native space. In this paper, we embed the smooth radial basis functions in a larger native space generated by a less smooth kernel and use them to interpolate the samples. Our result shows that there exists a linear combination of spherical radial basis functions that can both exactly interpolate samples generated by functions in the larger native space and near best approximate the target function.

1. Introduction

Many scientific questions boil down to synthesizing an unknown but definite function from finitely many samples $(x_i, y_i)_{i=1}^m$. The purpose is to find a functional model that can effectively represent or approximate the underlying relation between the input x_i and the output y_i . If the unknown functions are defined on spherical domains, then data collected by satellites or ground stations are usually not restricted on any regular region and are scattered. Thus, any numerical method relying on the structure of a “grid” is doomed to fail.

The success of the radial basis function networks methodology in Euclidean space derives from its ability to generate estimators from data with essentially unstructured geometry. Therefore, it is natural to borrow this ideal to deal with spherical scattered data. This method, called the spherical radial basis function networks (SRBFNs), has been extensively used in gravitational phenomenon [1, 2], image processing [3, 4], and learning theory [5].

Mathematically, the SRBFN can be represented as

$$S_n(x) := \sum_{i=1}^n c_i \phi(\xi_i \cdot x) := \sum_{i=1}^n c_i \phi_{\xi_i}(x), \quad x \in \mathbf{S}^d, \quad (1)$$

where $c_i \in \mathbf{R}$, $\xi_i \in \mathbf{S}^d$, and $\phi : [-1, 1] \rightarrow \mathbf{R}$ are called the connection weight, center, and activation function in

the terminology of neural networks, respectively. Here and hereafter, $\mathbf{S}^d$ denotes the unit sphere embedded in the $d + 1$ -dimensional Euclidean space, $\mathbf{R}^{d+1}$. Both the connection weights c_i and the centers ξ_i are adjustable in the process of training. We denote by Φ_n the collection of functions formed as (1).

A basic and classical approach for training SRBFN is to construct exact interpolant based on the given samples $(x_i, y_i)_{i=1}^m$, that is, to find a function S_n in Φ_n such that

$$S_n(x_i) = y_i, \quad i = 1, \dots, m. \quad (2)$$

If the activation function ϕ is chosen to be positive definite [2], then the matrix $A := (\phi(x_i \cdot x_j))_{i,j=1}^m$ is nonsingular. Thus, the system of (2) can be easily solved by taking the scattered data as the centers of SRBFN. This method has been already named the spherical basis function (SBF) method. Under this circumstance, the connection weights are determined via

$$\mathbf{c} = A^{-1}\mathbf{y}, \quad (3)$$

where $\mathbf{c} = (c_1, c_2, \dots, c_m)^T$, $\mathbf{y} = (y_1, \dots, y_m)^T$, U^T denotes the transpose of the matrix (or vector) U , and A^{-1} denotes the inverse matrix of A .

For the SBF method, if the target function f belongs to the native space N_ϕ associated with the activation function ϕ ,

then the best SBF approximant of f is characterized by the exact interpolation:

$$\left\| f - c_i \sum_{i=1}^m \phi_{x_i} \right\|_{N_\phi} = \inf_{g \in \Phi_m} \|f - g\|_{N_\phi}, \quad (4)$$

where $\{c_i\}_{i=1}^m$ are determined by (3). This property makes the SBF interpolation strategy popular in spherical scattered data fitting [6–14]. However, there are also two disadvantages for the SBF method. On one hand, since the centers of SRBFN interpolants are chosen as the scattered data, the interpolation capability depends heavily on their geometric distributions. This implies that we cannot obtain a satisfactory interpolation error estimate if the data are “badly” located on the sphere. On the other hand, the well known “native space barrier” [11, 12, 15] shows that (4) only holds for a small class of smooth functions if ϕ is smooth. Therefore, for functions outside N_ϕ , the SBF interpolants are not guaranteed to be the best approximants.

Along the flavor of the previous papers [7, 11, 12], we use SRBFNs to interpolate functions in a large native space N_ψ associated with the kernel ψ which is less smooth than ϕ and study its interpolation capability. Different from the previous work, the centers are chosen in advance to be quasi-uniform located on spheres, which makes the interpolation error depend on the number rather than the geometric distribution of centers. Our purpose is not to give the detailed error estimate of the SRBFN interpolation. Instead, we focus on investigating the relation between interpolation and approximation for SRBFN. Indeed, we find that there exists an SRBFN interpolant which is also the near best approximant of functions in N_ψ , when the number of centers and geometric distribution of the scattered data satisfy a certain assumption. That is, for a suitable choice of n , there exists a function $g_n \in \Phi_n$ such that

- (1) g exactly interpolates the samples $(x_i, y_i)_{i=1}^m$;
- (2) $\|f - g_n\|_{N_\psi} \leq C \inf_{g \in \Phi_n} \|f - g\|_{N_\psi}$,

where C is a constant depending only on d, α , and β .

2. Positive Radial Basis Function on the Sphere

It is easy to deduce that the volume of $\mathbf{S}^d$, Ω_d , satisfies

$$\Omega_d := \int_{\mathbf{S}^d} d\omega = \frac{2\pi^{(d+1)/2}}{\Gamma((d+1)/2)}, \quad (5)$$

where $d\omega$ denotes the surface area element on $\mathbf{S}^d$. For integer $k \geq 0$, the restriction to $\mathbf{S}^d$ of a homogeneous harmonic polynomial of degree k on the unit sphere is called a spherical harmonic of degree k . The span of all spherical harmonics of degree k is denoted by $\mathbf{H}_k^d$, and the class of all spherical harmonics (or spherical polynomials) of degree $k \leq s$ is

denoted by Π_s^d . It is obvious that $\Pi_s^d = \oplus_{k=0}^s \mathbf{H}_k^d$. The dimension of $\mathbf{H}_k^d$ is given by

$$D_k^d := \dim \mathbf{H}_k^d = \begin{cases} \frac{2k+d-1}{k+d-1} \binom{k+d-1}{k}, & k \geq 1; \\ 1, & k = 0, \end{cases} \quad (6)$$

and that of Π_s^d is $\sum_{k=0}^s D_k^d = D_s^{d+1} \sim s^d$.

Denote by $\{Y_{k,j}\}_{j=1}^{D_k^d}$ an orthonormal basis of $\mathbf{H}_k^d$; then the following addition formula [16, 17] holds

$$\sum_{l=1}^{D_k^d} Y_{k,l}(x) Y_{k,l}(y) = \frac{D_k^d}{\Omega_d} P_k^{d+1}(x \cdot y), \quad (7)$$

where P_k^{d+1} is the Legendre polynomial with degree k and dimension $d+1$. The Legendre polynomial P_k^{d+1} can be normalized such that $P_k^{d+1}(1) = 1$ and satisfies the orthogonality relation

$$\int_{-1}^1 P_k^{d+1}(t) P_j^{d+1}(t) (1-t^2)^{(d-2)/2} dt = \frac{\Omega_d}{\Omega_{d-1} D_k^d} \delta_{k,j}, \quad (8)$$

where $\delta_{k,j}$ is the usual Kronecker symbol.

Positive definite radial basis functions on spheres were introduced and characterized by Schoenberg [18]. Namely, a radial basis function φ is positive definite if and only if its expansion

$$\varphi(x \cdot y) = \sum_{k=0}^{\infty} e_k \frac{D_k^d}{\Omega_d} P_k^d(x \cdot y) \quad (9)$$

has all Fourier-Legendre coefficients $e_k \geq 0$ and $\sum_{k=0}^{\infty} e_k (D_k^d / \Omega_d) < \infty$. We define the native space N_φ as

$$N_\varphi := \left\{ f(x) = \sum_{k=0}^{\infty} \sum_{j=1}^{D_k^d} \hat{f}_{k,j} Y_{k,j}(x) : \sum_{k=0}^{\infty} e_k^{-1} \sum_{j=1}^{D_k^d} \hat{f}_{k,j}^2 < \infty \right\}, \quad (10)$$

with its inner product

$$\langle f, g \rangle_{N_\varphi} := \sum_{k=0}^{\infty} e_k^{-1} \sum_{j=1}^{D_k^d} \hat{f}_{k,j} \hat{g}_{k,j}, \quad (11)$$

where $\hat{f}_{k,j} := \int_{\mathbf{S}^d} f(x) Y_{k,j}(x) d\omega(x)$.

3. Interpolation and Near Best Approximation

Let $\Lambda := \{\eta_j\}_{j=1}^n \subset \mathbf{S}^d$ be a set of points and $d(x, y) = \arccos x \cdot y$ be the spherical distance between x and y . We denote by $h_\Lambda := \max_{x \in \mathbf{S}^d} \min_j d(x, \eta_j)$, $q_\Lambda := (1/2) \min_{j \neq k} d(\xi_j, \eta_k)$, and $\tau_\Lambda := h_\Lambda / q_\Lambda$ the mesh norm, separation radius, and mesh ratio of Λ , respectively. It is easy to check that these three quantities describe the geometric distribution of points in Λ .

The τ -uniform set $F_\tau := F_\tau(\mathbf{S}^d)$ is defined by the family of all centers sets Ξ with $\tau_\Lambda \leq \tau$.

Let ϕ and ψ satisfy

$$\phi(x \cdot y) := \sum_{k=0}^{\infty} a_k \frac{D_k^d}{\Omega_d} P_k^d(x \cdot y), \quad (12)$$

$$\psi(x \cdot y) := \sum_{k=0}^{\infty} b_k \frac{D_k^d}{\Omega_d} P_k^d(x \cdot y) \quad (13)$$

with

$$a_k \sim (1 + \lambda_k)^{-\alpha}, \quad b_k \sim (1 + \lambda_k)^{-\beta}. \quad (14)$$

The Sobolev embedding theorem [12] implies that if $\alpha, \beta > d/2$, then N_ϕ and N_ψ are continuously embedded in $C(\mathbf{S}^d)$, and so there are reproducing kernel Hilbert spaces, with reproducing kernels being ϕ and ψ , respectively.

The aim of this section is to study the relation between the exact interpolation and best approximation for Φ_n with its centers set $\Xi_n := \{\xi_i\}_{i=1}^n$ and activation function ϕ satisfying (12) and (14). It is obvious that such a Φ_n is a linear space. The following Theorem 1 shows that there exists an SRBFN interpolant which can near best approximate $f \in N_\psi$ in the metric of N_ψ , where ψ satisfies (13) and (14).

Theorem 1. Let $X := \{x_i\}_{i=1}^m$ be the set of scattered data with separation radius q_X , and $\alpha > \beta > d/2$. If $\Xi_n \in F_\tau$ and $n \geq C_\tau q_X^{-d}$, where $C_\tau > 1$ is a constant depending only on τ and d , then, for every $f \in N_\psi$, there exists an SRBFN interpolant $S_n \in \Phi_n$ such that

(i) S_n exactly interpolates the samples $(x_i, y_i)_{i=1}^m$,

(ii) $\|f - S_n\|_{N_\psi} \leq 5 \inf_{g \in \Phi_n} \|f - g\|_{N_\psi}$.

Remark 2. Similar results have been considered for spherical polynomials both in $C(\mathbf{S}^d)$ and N_ψ . Narcowich et al. [11, 12] proved that there exists a spherical polynomial interpolant of degree at most $L \geq C q_X$ which can also best approximate the target both in $C(\mathbf{S}^d)$ and N_ψ .

To prove Theorem 1, we need the following three lemmas, which can be found in [12, Proposition 5.2], [12, Theorem 5.5], and [19, Example 2.10], respectively.

Lemma 3. Let $\mathcal{Y}$ be a (possibly complex) Banach space, $\mathcal{V}$ a subspace of $\mathcal{Y}$, and Z^* a finite-dimensional subspace of $\mathcal{Y}^*$, the dual of $\mathcal{Y}$. If for every $z^* \in Z^*$ and some $\gamma > 1$, γ independent of z^* ,

$$\|z^*\|_{\mathcal{Y}^*} \leq \gamma \|z^*|_{\mathcal{V}^*}\|_{\mathcal{V}^*}, \quad (15)$$

then for $y \in \mathcal{Y}$ there exists $v \in \mathcal{V}$ such that v interpolates y on Z^* ; that is, $z^*(y) = z^*(v)$ for all $z^* \in Z^*$. In addition, v approximates y in the sense that $\|y - v\|_{\mathcal{Y}} \leq (1 + 2\gamma) \text{dist}_{\mathcal{Y}}(y, \mathcal{V})$.

Lemma 4. Let $\beta > d/2$. If $f \in N_\phi$, then there is a $u \in \Phi_n$ such that

$$\|f - u\|_{N_\psi} \leq C h_{\Xi_n}^{\alpha-\beta} \|f\|_{N_\phi}. \quad (16)$$

Lemma 5. Let ψ be defined in (13) and (14) and $\beta > d/2$. Then for arbitrary set of real numbers $\{c_i\}_{i=1}^m$, we have

$$\left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}^2 \geq C q_X^{2\beta} \sum_{i=1}^m c_i^2, \quad (17)$$

where C is a constant depending only on d and β .

Now we provide the proof of Theorem 1.

Proof of Theorem 1. We apply Lemma 3 to the case in which the underlying space is the native space $\mathcal{Y} = N_\psi$. Let $Z^* = \text{span}\{\delta_{x_i}\}_{i=1}^m$ and $\mathcal{V} = \Phi_n$. So in order to prove Theorem 1, it suffices to prove that for arbitrary m real numbers $\{c_i\}_{i=1}^m$ such that

$$\left\| \sum_{i=1}^m c_i \delta_{x_i} \right\|_{N_\psi^*} \leq 5 \left\| \sum_{i=1}^m c_i \delta_{x_i} |_{\Phi_n} \right\|_{\Phi_n^*}. \quad (18)$$

Since N_ψ is a reproducing kernel Hilbert space with ψ its reproducing kernel, we have

$$\begin{aligned} \left\| \sum_{i=1}^m c_i \delta_{x_i} \right\|_{N_\psi^*} &= \sup_{f \in N_\psi, \|f\|_{N_\psi}=1} \left| \sum_{i=1}^m c_i f(x_i) \right| \\ &= \sup_{f \in N_\psi, \|f\|_{N_\psi}=1} \left| \left\langle \sum_{i=1}^m c_i f, \psi_{x_i} \right\rangle_{N_\psi} \right| \\ &= \sup_{f \in N_\psi, \|f\|_{N_\psi}=1} \left| \left\langle \sum_{i=1}^m c_i \psi_{x_i}, f \right\rangle_{N_\psi} \right| \\ &= \left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}. \end{aligned} \quad (19)$$

Similarly, we obtain

$$\begin{aligned} \left\| \sum_{i=1}^m c_i \delta_{x_i} |_{\Phi_n} \right\|_{\Phi_n^*} &= \sup_{g \in \Phi_n, \|g\|_{N_\psi}=1} \left| \sum_{i=1}^m c_i g(x_i) \right| \\ &= \sup_{g \in \Phi_n, \|g\|_{N_\psi}=1} \left| \left\langle \sum_{i=1}^m c_i g, \psi_{x_i} \right\rangle_{N_\psi} \right| \\ &= \sup_{g \in \Phi_n, \|g\|_{N_\psi}=1} \left| \left\langle \sum_{i=1}^m c_i \psi_{x_i}, g \right\rangle_{N_\psi} \right| \\ &= \|s\|_{N_\psi}, \end{aligned} \quad (20)$$

where s is the orthogonal projection of $\sum_{i=1}^m c_i \psi_{x_i}$ to Φ_n in the metric of N_ψ . Then the Pythagorean theorem yields that

$$\|s\|_{N_\psi}^2 = \left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}^2 - \left\| \sum_{i=1}^m c_i \psi_{x_i} - s \right\|_{N_\psi}^2. \quad (21)$$

Thus, Lemma 3 with $\gamma = 2$ yields that in order to prove (18), it suffices to prove

$$\frac{\left\| \sum_{i=1}^m c_i \psi_{x_i} - s \right\|_{N_\psi}}{\left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}} \leq \frac{\sqrt{3}}{2}. \quad (22)$$

Let $L \in \mathbb{N}$, $P \in \Pi_L$ be the best polynomial approximation of $\sum_{i=1}^m c_i \psi_{x_i}$ in the metric of N_ψ ; then for arbitrary $s' \in \Phi_n$, we have

$$\left\| \sum_{i=1}^m c_i \psi_{x_i} - s \right\|_{N_\psi} \leq \left\| \sum_{i=1}^m c_i \psi_{x_i} - P \right\|_{N_\psi} + \|P - s'\|_{N_\psi}. \quad (23)$$

It follows from [12, Page 382] that there exists a constant, κ , depending only on β such that for arbitrary $L \geq \kappa q_X^{-1}$, there holds

$$\left\| \sum_{i=1}^m c_i \psi_{x_i} - P \right\|_{N_\psi} \leq \frac{\sqrt{3}}{4} \left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}. \quad (24)$$

Let s' be the best Φ_n approximation of P in the metric of N_ψ . Then it follows from Lemma 4 and the well-known Bernstein inequality [17] that

$$\|P - s'\|_{N_\psi} \leq Ch_\Xi^{\alpha-\beta} \|P\|_{N_\psi} \leq Ch_\Xi^{\alpha-\beta} L^\alpha \|P\|_2. \quad (25)$$

Since P is the best polynomial approximation of $\sum_{i=1}^m c_i \psi_{x_i}$ in the metric of N_ψ , a simple computation yields

$$\|P\|_2^2 = \sum_{i=1}^m c_i^2. \quad (26)$$

Furthermore, it follows from Lemma 5 that

$$\left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}^2 \geq C q_X^{2\beta} \sum_{i=1}^m c_i^2. \quad (27)$$

Then, $\Xi_n \in F_\tau$ together with $L = \max\{\lceil \kappa q_X^{-1} \rceil + 1, \lceil 3^{\beta/2} / (4C)^\beta q_X \rceil\}$ yields that

$$\begin{aligned} \|P - s'\|_{N_\psi} &\leq Ch_\Xi^{\alpha-\beta} L^\alpha q_X^{-\beta} \left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi} \\ &\leq C(n^{-1/d} q_X)^{\alpha-\beta} \left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}, \end{aligned} \quad (28)$$

where C is a constant depending only on β , d , and τ . Thus, there exists a constant C_τ depending only on α , β , d , and τ such that for arbitrary $n \geq C_\tau q_X^d$, there holds

$$\|P - s'\|_{N_\psi} \leq \frac{\sqrt{3}}{4} \left\| \sum_{i=1}^m c_i \psi_{x_i} \right\|_{N_\psi}. \quad (29)$$

Inserting (24) and (29) into (23), we finish the proof of (22) and then complete the proof of Theorem 1. $\square$

Acknowledgments

An anonymous referee has carefully read the paper and has provided to us numerous constructive suggestions. As a result, the overall quality of the paper has been noticeably enhanced, to which we feel much indebted and are grateful. The research was supported by the National 973 Programming (2013CB329404), the Key Program of National Natural Science Foundation of China (Grant no. 11131006), and the National Natural Science Foundations of China (Grant no. 61075054).

References

- [1] W. Freeden, T. Gervens, and M. Schreiner, *Constructive Approximation on the Sphere*, Numerical Mathematics and Scientific Computation, The Clarendon Press, Oxford University Press, New York, NY, USA, 1998.
- [2] W. Freeden and V. Michel, "Constructive approximation and numerical methods in geodetic research today—an attempt at a categorization based on an uncertainty principle," *Journal of Geodesy*, vol. 73, no. 9, pp. 452–465, 1999.
- [3] Y. T. Tsai and Z. C. Shih, "All-frequency precomputed radiance transfer using spherical radial basis functions and clustered tensor approximation," *ACM Transactions on Graphics*, vol. 25, pp. 967–976, 2006.
- [4] Y. T. Tsai, C. C. Chang, Q. Z. Jiang, and S. C. Weng, "Importance sampling of products from illumination and BRDF using spherical radial basis functions," *The Visual Computer*, vol. 24, pp. 817–826, 2008.
- [5] H. Q. Minh, "Some properties of Gaussian reproducing kernel Hilbert spaces and their implications for function approximation and learning theory," *Constructive Approximation*, vol. 32, no. 2, pp. 307–338, 2010.
- [6] K. Jetter, J. Stöckler, and J. D. Ward, "Error estimates for scattered data interpolation on spheres," *Mathematics of Computation*, vol. 68, no. 226, pp. 733–747, 1999.
- [7] J. Levesley and X. Sun, "Approximation in rough native spaces by shifts of smooth kernels on spheres," *Journal of Approximation Theory*, vol. 133, no. 2, pp. 269–283, 2005.
- [8] H. N. Mhaskar, F. J. Narcowich, and J. D. Ward, "Approximation properties of zonal function networks using scattered data on the sphere," *Advances in Computational Mathematics*, vol. 11, no. 2–3, pp. 121–137, 1999.
- [9] H. N. Mhaskar, F. J. Narcowich, J. Prestin, and J. D. Ward, " L^p Bernstein estimates and approximation by spherical basis functions," *Mathematics of Computation*, vol. 79, no. 271, pp. 1647–1679, 2010.
- [10] T. M. Morton and M. Neamtu, "Error bounds for solving pseudodifferential equations on spheres by collocation with zonal kernels," *Journal of Approximation Theory*, vol. 114, no. 2, pp. 242–268, 2002.
- [11] F. J. Narcowich and J. D. Ward, "Scattered data interpolation on spheres: error estimates and locally supported basis functions," *SIAM Journal on Mathematical Analysis*, vol. 33, no. 6, pp. 1393–1410, 2002.
- [12] F. J. Narcowich, X. Sun, J. D. Ward, and H. Wendland, "Direct and inverse Sobolev error estimates for scattered data interpolation via spherical basis functions," *Foundations of Computational Mathematics*, vol. 7, no. 3, pp. 369–390, 2007.

- [13] X. Sun and E. W. Cheney, "Fundamental sets of continuous functions on spheres," *Constructive Approximation*, vol. 13, no. 2, pp. 245–250, 1997.
- [14] X. Sun and Z. Chen, "Spherical basis functions and uniform distribution of points on spheres," *Journal of Approximation Theory*, vol. 151, no. 2, pp. 186–207, 2008.
- [15] F. J. Narcowich, R. Schaback, and J. D. Ward, "Approximations in Sobolev spaces by kernel expansions," *Journal of Approximation Theory*, vol. 114, no. 1, pp. 70–83, 2002.
- [16] C. Müller, *Spherical Harmonics*, vol. 17 of *Lecture Notes in Mathematics*, Springer, Berlin, Germany, 1966.
- [17] K. Y. Wang and L. Q. Li, *Harmonic Analysis and Approximation on the Unit Sphere*, Science Press, Beijing, China, 2000.
- [18] I. J. Schoenberg, "Positive definite functions on spheres," *Duke Mathematical Journal*, vol. 9, pp. 96–108, 1942.
- [19] F. J. Narcowich, N. Sivakumar, and J. D. Ward, "Stability results for scattered-data interpolation on Euclidean spheres," *Advances in Computational Mathematics*, vol. 8, no. 3, pp. 137–163, 1998.