Research Article

Weighted Endpoint Estimates for Commutators of Riesz Transforms Associated with Schrödinger Operators

Yu Liu, Jielai Sheng, and Lijuan Wang

School of Mathematics and Physics, University of Science and Technology Beijing, Beijing 100083, China

Correspondence should be addressed to Yu Liu; liuyu75@pku.org.cn

Received 14 May 2013; Accepted 1 October 2013

Academic Editor: Baoxiang Wang

Copyright © 2013 Yu Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let $L = -\Delta + V$ be a Schrödinger operator, where Δ is the laplacian on $\mathbb{R}^n$ and the nonnegative potential V belongs to the reverse Hölder class B_{s_1} for some $s_1 \geq (n/2)$. Assume that $\omega \in A_1(\mathbb{R}^n)$. Denote by $H_L^1(\omega)$ the weighted Hardy space related to the Schrödinger operator $L = -\Delta + V$. Let $\mathcal{R}_b = [b, \mathcal{R}]$ be the commutator generated by a function $b \in \text{BMO}_\theta(\mathbb{R}^n)$ and the Riesz transform $\mathcal{R} = \nabla(-\Delta + V)^{-1/2}$. Firstly, we show that the operator $\mathcal{R}$ is bounded from $L^1(\omega)$ into $L_{\text{weak}}^1(\omega)$. Secondly, we obtain the endpoint estimates for the commutator $[b, \mathcal{R}]$. Namely, it is bounded from the weighted Hardy space $H_L^1(\omega)$ into $L_{\text{weak}}^1(\omega)$.

1. Introduction

Let $L = -\Delta + V$ be a Schrödinger operator, where Δ is the laplacian on $\mathbb{R}^n$ and the nonnegative potential V belongs to the reverse Hölder class B_{s_1} for some $s_1 \geq (n/2)$ and $n \geq 3$. In this paper, we consider the Riesz transform associated with the Schrödinger operator L as follows:

$$\mathcal{R} = \nabla(-\Delta + V)^{-1/2}. \quad (1)$$

Let b be a locally integrable function on $\mathbb{R}^n$ and let T be a linear operator. For a suitable function f , the commutator is defined by $T_b f \doteq [b, T]f = bT(f) - T(bf)$. It is well known that when T is a Calderón-Zygmund operator, Coifman et al. [1] proved that $[b, T]$ is a bounded operator on L^p for $1 < p < \infty$ if and only if $b \in \text{BMO}(\mathbb{R}^n)$.

Recently, some scholars have investigated the boundedness of the commutators generated by a BMO function b and Riesz transforms associated with the Schrödinger operator (cf. [2–8]). It follows from [9] that Riesz transform associated with the Schrödinger operator L is not a Calderón-Zygmund operator if the potential $V \in B_s$ ($(n/2) < s < n$). Their results imply that the boundedness of the commutators of Riesz transform associated with the Schrödinger operator L depends on the nonnegative potential V . In [5], the authors have obtained the weighted L^p , $1 < p < \infty$, and weak L log L the estimates for the commutator $\mathcal{R}_b$. In this paper we

are interested in the weighted Hardy space estimates for $\mathcal{R}_b$, which are also the weighted endpoint estimates. It is noted that our main results generalize Theorem 2.7 and Theorem 4.1 in [3] to the weighted case and the function b that we consider belongs to a larger class than the classical BMO space.

Note that a nonnegative locally L^s integrable function V on $\mathbb{R}^n$ is said to belong to B_s ($1 < s < \infty$) if there exists $C > 0$ such that the reverse Hölder inequality

$$\left(\frac{1}{|B|} \int_B V(x)^s dx \right)^{1/s} \leq \frac{C}{|B|} \int_B V(x) dx \quad (2)$$

holds for every ball B in $\mathbb{R}^n$. Obviously, $B_{s_2} \subset B_{s_1}$ if $s_2 > s_1$. But it is important that the B_q class has a property of “self-improvement”; that is, if $V \in B_s$, then $V \in B_{s+\varepsilon}$ for some $\varepsilon > 0$.

Assume $V \in B_{s_1}$ for some $s_1 > (n/2)$. Then the auxiliary function $\rho(x, V) = \rho(x)$ introduced by Shen in [9] is defined as follows:

$$\begin{aligned} \rho(x) &= \frac{1}{m(x, V)} \\ &\doteq \sup_{r>0} \left\{ r : \frac{1}{r^{n-2}} \int_{B(x,r)} V(y) dy \leq 1 \right\}, \quad x \in \mathbb{R}^n. \end{aligned} \quad (3)$$

In [4], the authors define the class $BMO_\theta(\rho)$ of locally integrable function b such that

$$\frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_B| dy \leq C \left(1 + \frac{r}{\rho(x)}\right)^\theta, \quad (4)$$

for all $x \in \mathbb{R}^n$ and $r > 0$, where $\theta > 0$ and $b_B = (1/|B|) \int_B b(y) dy$. A norm for $b \in BMO_\theta(\rho)$, denoted by $[b]_\theta$, is given by the infimum of the constants satisfying (4), after identifying functions that differ upon a constant. If we let $\theta = 0$ in (4), then $BMO_\theta(\rho)$ is exactly the John-Nirenberg space BMO . Denote $BMO_\infty(\rho) = \bigcup_{\theta > 0} BMO_\theta(\rho)$. It is easy to see that $BMO \subset BMO_\theta(\rho) \subset BMO_{\theta'}(\rho)$ for $0 < \theta \leq \theta'$. Hence, $BMO \subset BMO_\infty(\rho)$.

Throughout this paper, we set $\omega(E) = \int_E \omega(x) dx$ for any subset $E \subseteq \mathbb{R}^n$. Assume that the nonnegative function $\omega \in L^1_{\text{loc}}(\mathbb{R}^n)$. We say that $\omega \in A_p(\mathbb{R}^n)$ for $1 < p < \infty$ if there is a constant $C > 0$ such that

$$\sup_B \left(\frac{1}{|B|} \int_B \omega(x) dx \right) \left(\frac{1}{|B|} \int_B (\omega(x))^{1-p'} dx \right)^{p-1} \leq C, \quad (5)$$

for all balls B in $\mathbb{R}^n$, where $(1/p) + (1/p') = 1$.

We say that $\omega \in A_1(\mathbb{R}^n)$, if there exists a positive constant $C > 0$, such that

$$M\omega(x) \leq C\omega(x), \quad \text{a.e. } x \in \mathbb{R}^n, \quad (6)$$

where M is a Hardy-Littlewood operator.

Given a weight function $\omega \in A_p(\mathbb{R}^n)$ for $1 \leq p < \infty$, as usual we denote by $L^q(\omega)$ the space of all measurable functions satisfying

$$\|f\|_{L^q(\omega)}^q = \int_{\mathbb{R}^n} |f(x)|^q \omega(x) dx < \infty. \quad (7)$$

When $q = \infty$, $L^\infty(\omega)$ will be taken to mean L^∞ and $\|f\|_{L^\infty(\omega)} = \|f\|_{L^\infty}$. Moreover, denote by $L^1_{\text{weak}}(\omega)$ the space of all measurable functions satisfying

$$\sup_{\lambda > 0} [\lambda \omega \{x \in \mathbb{R}^n : |f(x)| \geq \lambda\}] < \infty. \quad (8)$$

In the rest of this paper, we always assume that $\omega \in A_1(\mathbb{R}^n)$.

Because $V \geq 0$ and $V \in L^{n/2}_{\text{loc}}(\mathbb{R}^n)$, the Schrödinger operator L generates a (C_0) contraction semigroup $\{T_s^L : s > 0\} = \{e^{-sL} : s > 0\}$. The maximal function associated with $\{T_s^L : s > 0\}$ is defined by $M^L f(x) = \sup_{s > 0} |T_s^L f(x)|$.

The Hardy space $H^1_L(\mathbb{R}^n)$ associated with the Schrödinger operator L is defined in terms of the maximal function mentioned above (cf. [10]). Recently, the weighted Hardy space $H^1_L(\omega)$ has been established by Liu et al. [11] and it is defined as follows.

Definition 1. A function $f \in L^1(\omega)$ is said to be in $H^1_L(\omega)$ if the maximal function $M^L f$ belongs to $L^1(\omega)$. The norm of such a function is defined by $\|f\|_{H^1_L(\omega)} = \|M^L f\|_{L^1(\omega)}$.

Definition 2. Let $1 \leq q \leq \infty$. A function $a \in L^q(\omega)$ is called an $H^{1,q}_L(\omega)$ -atom if $r < \rho(x_0)$ and the following conditions hold:

- (i) $\text{supp } a \subset B(x_0, r)$,
- (ii) $\|a\|_{L^q(\omega)} \leq \omega(B(x_0, r))^{(1/q)-1}$,
- (iii) if $r < (\rho(x_0)/4)$, then $\int_{B(x_0, r)} a(x) dx = 0$.

In [11], Liu et al. gave the following atomic decomposition for the space $H^1_L(\omega)$.

Proposition 3. Assume $V \in B_{s_1}$ for some $s_1 > (n/2)$. Let $f \in L^1(\omega)$. Then $f \in H^1_L(\omega)$ if and only if f can be written as $f = \sum_j \lambda_j a_j$, where a_j are $H^{1,\infty}_L(\omega)$ -atoms, $\sum_j |\lambda_j| < \infty$, and the sum converges in the $H^1_L(\omega)$ quasinorm. Moreover,

$$\|f\|_{H^1_L(\omega)} \sim \inf \left\{ \sum_j |\lambda_j| \right\}, \quad (9)$$

where the infimum is taken over all atomic decompositions of f into $H^{1,\infty}_L$ -atoms.

Following the above definition of atoms and the above atomic decomposition, we know that the weighted Hardy space $H^1_L(\omega)$ is not the special case of Hardy spaces established by Yang and Zhou in [12].

Now we are in a position to give the main results in this paper.

Theorem 4. Let $\omega \in A_1(\mathbb{R}^n)$. Suppose $V \in B_s$ for $s > (n/2)$. Then,

$$\omega \{x \in \mathbb{R}^n : |\nabla(-\Delta + V)^{1/2} f(x)| > \alpha\} \leq \frac{C}{\alpha} \|f\|_{L^1(\omega)}, \quad (10)$$

for every $\alpha > 0$.

Theorem 5. Let $b \in BMO_\infty(\rho)$. Then, for any $\lambda > 0$,

$$\omega \{x \in \mathbb{R}^n : |\mathcal{R}_b f(x)| > \lambda\} \leq \frac{C[b]_\theta}{\lambda} \|f\|_{H^1_L(\omega)}, \quad (11)$$

$\forall f \in H^1_L(\omega)$,

where $\theta > 0$. Namely, the commutator $\mathcal{R}_b$ is bounded from $H^1_L(\omega)$ into $L^1_{\text{weak}}(\omega)$.

This paper is organized as follows. In Section 2, we recall some basic facts to prove main results in this paper. Section 3 gives the proof of weighted estimates of Riesz transform associated with the Schrödinger operator. In Section 4, we prove Theorem 5.

Throughout this paper, the letter C stands for a constant and is not necessarily the same at each occurrence. By $B_1 \sim B_2$, we mean that there exists a constant $C > 1$ such that $(1/C) \leq (B_1/B_2) \leq C$. Moreover, for the ball $B = B(x, r)$, we denote the ball MB by $MB = B(x, Mr)$, where M is a positive constant.

2. Preliminaries

Firstly, we recall some lemmas of the auxiliary function $\rho(x)$ which have been proved by Shen in [9]. Throughout this section we always assume $V \in B_{s_1}$ for some $s_1 > (n/2)$.

Lemma 6. *The measure $V(x)dx$ satisfies the doubling condition; that is, there exists $C > 0$ such that*

$$\int_{B(x,2r)} V(y) dy \leq C \int_{B(x,r)} V(y) dy \quad (12)$$

holds for all balls $B(x, r)$ in $\mathbb{R}^n$.

Lemma 7. *For $0 < r < R < \infty$,*

$$\begin{aligned} & 1/r^{n-2} \int_{B(x,r)} V(y) dy \\ & \leq C \left(\frac{R}{r} \right)^{(n/s_1)-2} 1/R^{n-2} \int_{B(x,R)} V(y) dy. \end{aligned} \quad (13)$$

Lemma 8. *If $r = \rho(x)$, then*

$$1/r^{n-2} \int_{B(x,r)} V(y) dy = 1. \quad (14)$$

Moreover,

$$1/r^{n-2} \int_{B(x,r)} V(y) dy \sim 1 \quad \text{if and only if } r \sim \rho(x). \quad (15)$$

Lemma 9. *There exists $l_0 > 0$ such that, for any x and y in $\mathbb{R}^n$,*

$$\frac{1}{C} \left(1 + \frac{|x-y|}{\rho(x)} \right)^{-l_0} \leq \frac{\rho(y)}{\rho(x)} \leq C \left(1 + \frac{|x-y|}{\rho(x)} \right)^{l_0/(l_0+1)}. \quad (16)$$

In particular, $\rho(x) \sim \rho(y)$ if $|x-y| < C\rho(x)$.

Lemma 10. *There exists $l_1 > 0$ such that*

$$\begin{aligned} & \int_{B(x,R)} \frac{V(y)}{|x-y|^{n-2}} dy \\ & \leq \frac{C}{R^{n-2}} \int_{B(x,R)} V(y) dy \leq C \left(1 + \frac{R}{\rho(x)} \right)^{l_1}. \end{aligned} \quad (17)$$

For the proofs of Lemma 6 to Lemma 10, readers can refer to [9].

A ball $B(x, \rho(x))$ is called critical. Due to Lemma 2.3 in [10], we have Lemma 12 on $\mathbb{R}^n$.

Proposition 11. *There exists a sequence of points $\{x_k\}_{k=1}^\infty$ in $\mathbb{R}^n$, such that the family of critical balls $Q_k = B(x_k, \rho(x_k))$, $k \geq 1$, satisfies the following*

(i) $\bigcup_k Q_k = \mathbb{R}^n$.

(ii) *There exists $N = N(\rho)$ such that, for every $k \in \mathbb{N}$, card $\{j : 4Q_j \cap 4Q_k \neq \emptyset\} \leq N$.*

Secondly, we recall the estimates of the fundamental solution of the operator $-\Delta + V + i\tau$ and give the estimates of kernels of Riesz transform $\mathcal{R}$. Let $\Gamma(x, y, \tau)$ denote the fundamental solution for the operator $-\Delta + V + i\tau$, where $\tau \in \mathbb{R}$. Clearly, $\Gamma(x, y, \tau) = \Gamma(y, x, -\tau)$.

Lemma 12. *Let $N > 0$ be an integer.*

(1) *Suppose $V \in B_{s_1}$ for $s_1 > (n/2)$. Then there exists $C_N > 0$ such that, for $x \neq y$,*

$$\begin{aligned} & |\Gamma(x, y, \tau)| \\ & \leq \frac{C_N}{(1 + |x-y||\tau|^{(1/2)})^N (1 + |x-y|\rho(x)^{-1})^N} \frac{1}{|x-y|^{n-2}}. \end{aligned} \quad (18)$$

(2) *Suppose $V \in B_{s_1}$ for $s_1 > (n/2)$. Then there exists $C_N > 0$ such that, for $x \neq y$,*

$$\begin{aligned} & |\nabla_x \Gamma(x, y, \tau)| \\ & \leq \frac{C_N}{(1 + |x-y||\tau|^{(1/2)})^N (1 + |x-y|\rho(x)^{-1})^N} \\ & \quad \times \left(\frac{1}{|x-y|^{n-2}} \int_{B(y, |x-y|)} \frac{V(z) dz}{|z-y|^{n-1}} + \frac{1}{|x-y|^{n-1}} \right). \end{aligned} \quad (19)$$

By the functional calculus, we may write

$$(-\Delta + V)^{-(1/2)} = \frac{1}{2\pi} \int_{\mathbb{R}} (-i\tau)^{-(1/2)} (-\Delta + V + \tau)^{-1} d\tau. \quad (20)$$

Let $f \in C_0^\infty(\mathbb{R}^n)$. From $(-\Delta + V + \tau)^{-1} f(x) = \int_{\mathbb{R}^n} \Gamma(x, y, \tau) f(y) dy$, it follows that

$$\mathcal{R}f(x) = \int_{\mathbb{R}^n} K(x, y) f(y) dy, \quad (21)$$

where

$$K(x, y) = \frac{1}{2\pi} \int_{\mathbb{R}} (-i\tau)^{-(1/2)} \nabla_x \Gamma(x, y, \tau) d\tau. \quad (22)$$

And the adjoint operator of $\mathcal{R}$ is given by

$$\widetilde{\mathcal{R}}f(x) = \int_{\mathbb{R}^n} \widetilde{K}(x, y) f(y) dy, \quad (23)$$

where

$$\widetilde{K}(x, y) = \frac{1}{2\pi} \int_{\mathbb{R}} (-i\tau)^{-(1/2)} \nabla_y \Gamma(y, x, \tau) d\tau. \quad (24)$$

Lemma 13. Suppose $V \in B_{s_1}$ for some $s_1 > (n/2)$. Then there exists $\delta > 0$, for any integer $N > 0$, and $0 < h < (|x - y|/16)$,

$$\begin{aligned} & |K(x, y)| \\ & \leq \frac{C_N}{(1 + |x - y| \rho(x)^{-1})^N} \\ & \quad \times \left(\frac{1}{|x - y|^{n-1}} \int_{B(y, |x-y|)} \frac{V(z) dz}{|z - y|^{n-1}} + \frac{1}{|x - y|^n} \right), \\ & |K(x, y + h) - K(x, y)| \\ & \leq \frac{C_N}{(1 + |x - y| \rho(x)^{-1})^N} \frac{|h|^\delta}{|x - y|^{n-1+\delta}} \\ & \quad \times \left(\int_{B(y, |x-y|)} \frac{V(z) dz}{|z - y|^{n-1}} + \frac{1}{|x - y|^n} \right). \end{aligned} \quad (25)$$

Lemmas 12 and 13 have been proved in [9] and in [2], respectively.

Thirdly, we recall some important and useful properties of A_p weights (cf. Chapter V in [13]).

Proposition 14. (i) Let $\omega \in A_p$, $1 \leq p < \infty$. Then there exist a constant $C > 0$ and $r > 1$ depending only on p and the A_p constant of ω , such that

$$\left(\frac{1}{|B|} \int_B \omega^r dx \right)^{1/r} \leq \frac{C}{|B|} \int_B \omega dx, \quad (26)$$

for any ball B .

(ii) Let $\omega \in A_1$. If $1 \leq p < \infty$, then $\omega^{1-p} \in A_p$.

At last, we review some basic facts about the BMO space $BMO_\theta(\rho)$, which have been proved in [4].

Proposition 15. Let $\theta > 0$ and let $1 \leq p < \infty$. If $b \in BMO_\theta(\rho)$, then

$$\begin{aligned} & \left(\frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_B|^p dy \right)^{(1/p)} \\ & \leq C[b]_\theta \left(1 + \frac{r}{\rho(x)} \right)^{\theta'}, \end{aligned} \quad (27)$$

for all $B = B(x, r)$, with $x \in \mathbb{R}^n$ and $r > 0$, where $\theta' = (1 + l_0)\theta$ and l_0 is the constant appearing in Lemma 9.

Lemma 16. Let $b \in BMO_\theta(\rho)$, let $B = B(x_0, r)$, and let $s \geq 1$; then

$$\left(\frac{1}{|2^k B|} \int_{2^k B} |b(y) - b_B|^p dy \right)^{(1/p)} \leq C[b]_\theta k \left(1 + \frac{2^k r}{\rho(x_0)} \right)^{\theta'}, \quad (28)$$

for all $k \in \mathbb{N}$, with θ' as in (27).

3. Weighted Estimates of Riesz Transforms Associated with the Schrödinger Operators

In this section, we need to prove the weighted estimates of the Riesz transform associated with the Schrödinger operator, which will be used in the proof of Theorem 5.

Theorem 17. Let $\omega \in A_1(\mathbb{R}^n)$. Suppose $V \in B_s$ for some $s \geq (n/2)$. Then, for $p'_0 \leq p < \infty$,

$$\|(-\Delta + V)^{-(1/2)} \nabla f\|_{L^p(\omega)} \leq C \|f\|_{L^p(\omega)}, \quad (29)$$

where $p'_0 = p_0/(p_0 - 1)$ and $(1/p_0) = (1/s) - (1/n)$.

Proof. By the improvement of “ B_s ”, $V \in B_{s_1}$ for $s < s_1 < n$. It follows from the proof of Theorem 0.5 in [9] that

$$\begin{aligned} |(-\Delta + V)^{1/2} \nabla f(x)| & \leq C \left\{ M(|f|^{p'_1})(x) \right\}^{1/p'_1} \\ & \quad + 2 \sup_{\varepsilon > 0} \left| \int_{|y-x|>\varepsilon} K_1^0(y, x) f(y) dy \right|, \end{aligned} \quad (30)$$

where $(1/p'_1) = 1 - (1/p_1) = 1 - (1/s_1) + (1/n)$ and $K_1^0(x, y)$ is the kernel for the operator $\nabla(-\Delta)^{-1/2}$.

Since $\omega \in A_1(\mathbb{R}^n)$, then it follows from Proposition 7.2 in [14] that $\omega \in A_p(\mathbb{R}^n)$ for any $p > 1$.

By [13, Theorem 1, page 201] and [13, Corollary 2, page 205]

$$\|(-\Delta + V)^{-1/2} \nabla f\|_{L^p(\omega)} \leq C \|f\|_{L^p(\omega)}, \quad \text{for } s'_1 < p < \infty. \quad (31)$$

Since $p'_1 < p'_0$, then

$$\|(-\Delta + V)^{-1/2} \nabla f\|_{L^p(\omega)} \leq C \|f\|_{L^p(\omega)}, \quad \text{for } p'_0 < p < \infty. \quad (32)$$

□

By duality, we have the following.

Corollary 18. Let $\omega \in A_1(\mathbb{R}^n)$. Suppose $V \in B_s$ for some $s \geq (n/2)$. Then, for $1 < p \leq p_0$,

$$\|\nabla(-\Delta + V)^{-1/2} f\|_{L^p(\omega)} \leq C \|f\|_{L^p(\omega)}, \quad (33)$$

where $(1/p_0) = (1/s) - (1/n)$.

Proof. It follows from (ii) in Proposition 14 that $\omega^{1-p'} \in A_{p'}$ for $p'_0 < p' < \infty$, where $(1/p) + (1/p') = 1$. By using Theorem 17, we know that

$$\|(-\Delta + V)^{-1/2} \nabla f\|_{L^{p'}(\omega^{1-p'})} \leq C \|f\|_{L^{p'}(\omega^{1-p'})}. \quad (34)$$

Therefore, by duality and (34),

$$\begin{aligned}
& \left\| \nabla(-\Delta + V)^{-1/2} f \right\|_{L^p(\omega)} \\
&= \sup_{\|g\|_{L^{p'}(\omega)}=1} \left| \int_{\mathbb{R}^n} \nabla(-\Delta + V)^{-1/2} f(x) g(x) \omega(x) dx \right| \\
&= \sup_{\|g\|_{L^{p'}(\omega)}=1} \left| \int_{\mathbb{R}^n} f(y) ((-\Delta + V)^{-1/2} \nabla)(g\omega)(y) dy \right| \\
&\leq \sup_{\|g\|_{L^{p'}(\omega)}=1} \left(\int_{\mathbb{R}^n} |f(y)|^p \omega(y) dy \right)^{1/p} \\
&\quad \times \left(\int_{\mathbb{R}^n} |((-\Delta + V)^{-1/2} \nabla)(g\omega)(y)|^{p'} \right. \\
&\quad \left. \times (\omega(y))^{-(p'/p)} dy \right)^{1/p'} \\
&= \sup_{\|g\|_{L^{p'}(\omega)}=1} \left(\int_{\mathbb{R}^n} |f(y)|^p \omega(y) dy \right)^{1/p} \\
&\quad \times \left(\int_{\mathbb{R}^n} |((-\Delta + V)^{-1/2} \nabla)(g\omega)(y)|^{p'} \right. \\
&\quad \left. \times (\omega(y))^{1-p'} dy \right)^{1/p'} \\
&\leq C \sup_{\|g\|_{L^{p'}(\omega)}=1} \left(\int_{\mathbb{R}^n} |f(y)|^p \omega(y) dy \right)^{1/p} \\
&\quad \times \left(\int_{\mathbb{R}^n} |g(y)|^{p'} \omega(y) dy \right)^{1/p'} \\
&\leq C \|f\|_{L^p(\omega)}. \tag{35}
\end{aligned}$$

This completes the proof of the above corollary. $\square$

In order to prove Theorem 4, we need the following lemmas.

Lemma 19 (see [11, Lemma 3.12]). *Let $r = \rho(x)$. Then there exists a constant $C > 0$, such that*

$$\int_{|x-y|>r} |K(y, x)| \omega(y) dy \leq C\omega(x). \tag{36}$$

By using the proof of Lemma 3.15 in [11], we immediately have the following.

Lemma 20. *Let $r = \rho(x)$. Then there exists a constant $C > 0$, such that*

$$\int_{|x-y|\leq r} |K(y, x) - K^0(y, x)| \omega(y) dy \leq C\omega(x), \tag{37}$$

where $K^0(y, x)$ is the kernel of the operator $\nabla(-\Delta)^{-(1/2)}$.

Now we give the proof of Theorem 4.

Proof of Theorem 4. We show Theorem 4 by a method similar to the one used in the proof of Theorem 2 in [15]. By the Calderón-Zygmund decomposition in the proof of Theorem 3.5 on page 413 in [16], given $f \in L^1(\omega)$ and $\alpha > 0$, we have $f = f_1 + f_2$, with $f_2 = \sum_k b_k$, such that the following hold.

$$(1) |f_1(x)| \leq C\alpha, \text{ for a.e. } x \in \mathbb{R}^n.$$

$$(2) \text{ Each } b_k \text{ is supported in a ball } B_k,$$

$$\int_{B_k} |b_k(x)| \omega(x) dx \leq C\alpha\omega(B_k), \quad \int_{B_k} b_k(x) dx = 0. \tag{38}$$

$$(3) \sum_k \omega(B_k) \leq (C/\alpha) \|f\|_{L^1(\omega)}.$$

Because $\omega \in A_1(\mathbb{R}^n) \subseteq A_2(\mathbb{R}^n)$, by Corollary 18, we know that $\mathcal{R}$ is bounded on $L^2(\omega)$; it is clear that

$$\omega \left\{ x \in \mathbb{R}^n : |\mathcal{R}f_1(x)| > \frac{\alpha}{2} \right\} \leq \frac{C}{\alpha^2} \|f_1\|_{L^2(\omega)}^2 \leq \frac{C}{\alpha} \|f\|_{L^1(\omega)}. \tag{39}$$

Let $B_k = B(x_k, r_k)$ and $\Omega = \bigcup_k B(x_k, 2r_k)$. Then

$$\omega(\Omega) \leq C \sum_k \omega(B_k) \leq \frac{C}{\alpha} \|f\|_{L^1(\omega)}. \tag{40}$$

We only need to consider $\mathcal{R}f_2(x)$ for $x \in \Omega^c$. If $r_k \geq 1/(m(x_k, V))$, then $1/(m(y, V)) \leq Cr_k$ for any $y \in B_k$. By Lemma 19, we get

$$\begin{aligned}
& \int_{|x-x_k|\geq 2r_k} |\mathcal{R}b_k(x)| dx \\
&\leq \int_{|x-x_k|\geq 2r_k} \int_{B_k} |K(x, y)| |b_k(y)| \omega(y) dy dx \\
&\leq C \|b_k\|_{L^1(\omega)}. \tag{41}
\end{aligned}$$

If $r_k < 1/(m(x_k, V))$, then $1/(m(x_k, V)) \sim 1/(m(y, V))$ for any $y \in B_k$. Since $K_0(x, y)$ is a Calderón-Zygmund kernel, by Lemma 19, Lemma 20, and [16, Lemma 3.3, page 413], we obtain

$$\begin{aligned}
& \int_{|x-x_k|\geq 2r_k} |\mathcal{R}b_k(x)| \omega(x) dx \\
&\leq \int_{2r_k \leq |x-x_k| < 2/(m(x_k, V))} |\mathcal{R}b_k(x)| \omega(x) dx \\
&\quad + \int_{|x-x_k|\geq 2/(m(x_k, V))} |\mathcal{R}b_k(x)| \omega(x) dx
\end{aligned}$$

$$\begin{aligned}
&\leq \int_{2r_k \leq |x-x_k| < 2/(m(x_k, V))} \\
&\quad \times \int_{B_k} |K(x, y) - K_0(x, y)| |b_k(y)| dy \omega(x) dx \\
&\quad + \int_{2r_k \leq |x-x_k| < 2/(m(x_k, V))} \\
&\quad \times \int_{B_k} |K_0(x, y) - K_0(x, x_k)| |b_k(y)| dy \omega(x) dx \\
&\quad + \int_{|x-x_k| \geq 2/(m(x_k, V))} \int_{B_k} |K(x, y)| |b_k(y)| dy \omega(x) dx \\
&\leq C \|b_k\|_{L^1(\omega)}. \tag{42}
\end{aligned}$$

Then

$$\begin{aligned}
\int_{\Omega^c} |\mathcal{R}f_2(x)| dx &\leq \sum_k \int_{(B(x, 2r_k))^c} |\mathcal{R}b_k(x)| \omega(x) dx \\
&\leq C \sum_k \|b_k\|_{L^1(\omega)} \leq C \alpha \sum_k \omega(B_k) \leq C \|f\|_{L^1(\omega)}. \tag{43}
\end{aligned}$$

Therefore

$$\omega \left\{ x \in \Omega^c : |\mathcal{R}f_2(x)| > \frac{\alpha}{2} \right\} \leq \frac{C}{\alpha} \|f\|_{L^1}. \tag{44}$$

Theorem 4 is proved by combination of (39), (40), and (44). $\square$

4. Proof of Theorem 5

For $f \in H_L^1(\omega)$ we can write $f = \sum_{j=-\infty}^{\infty} \lambda_j a_j$, where each a_j is a $H_L^{1,\infty}(\omega)$ atom and $\sum_{j=-\infty}^{\infty} |\lambda_j| \leq 2\|f\|_{H_L^1(\omega)}$. Suppose that $\text{supp } a_j \subseteq B_j = B(x_j, r_j)$ with $r_j < \rho(x_j)$. Write

$$\begin{aligned}
\mathcal{R}_b f(x) &= \sum_{j=-\infty}^{\infty} \lambda_j (b(x) - b_{B_j}) \mathcal{R}a_j(x) \chi_{8B_j}(x) \\
&\quad + \sum_{j:r_j \geq (\rho(x_j)/4)} \lambda_j (b(x) - b_{B_j}) \\
&\quad \quad \times \mathcal{R}a_j(x) \chi_{(8B_j)^c}(x) \\
&\quad + \sum_{j:r_j < (\rho(x_j)/4)} \lambda_j (b(x) - b_{B_j}) \\
&\quad \quad \times \mathcal{R}a_j(x) \chi_{(8B_j)^c}(x) \\
&\quad - \mathcal{R} \left(\sum_{j=-\infty}^{\infty} \lambda_j (b - b_{B_j}) a_j \right)(x) \\
&= A_1(x) + A_2(x) + A_3(x) + A_4(x). \tag{45}
\end{aligned}$$

Using Hölder inequality, the weighted L^q boundedness of $\mathcal{R}$, Propositions 14 and 15, for $(1/q) + (1/t) + (1/\nu) = 1$ and $(1 - (1/q))t = r$ with $1 < q < p_0$, we have

$$\begin{aligned}
&\| (b(x) - b_{B_j}) \mathcal{R}a_j(x) \chi_{8B_j}(x) \|_{L^1(\omega)} \\
&\leq \left(\int_{8B_j} |b(x) - b_{B_j}|^\nu dx \right)^{1/\nu} \\
&\quad \times \| \mathcal{R}a_j \|_{L^q(\omega)} \left(\int_{8B_j} \omega(x)^r dx \right)^{1/t} \\
&\leq C |8B_j|^{1/\nu} |8B_j|^{1/t} \left(\frac{1}{|8B_j|} \int_{8B_j} |b(x) - b_{B_j}|^\nu dx \right)^{1/\nu} \\
&\quad \times \|a_j\|_{L^q(\omega)} \left(\frac{1}{|8B_j|} \int_{8B_j} \omega(x)^r dx \right)^{1/t} \\
&\leq C [b]_\theta |8B_j|^{1/\nu} |8B_j|^{1/t} \omega(8B_j)^{(1/q)-1} \\
&\quad \times \left(\frac{1}{|8B_j|} \int_{8B_j} \omega(x) dx \right)^{r/t} \\
&\leq C [b]_\theta, \tag{46}
\end{aligned}$$

since $r_j < \rho(x_j)$.

When we consider the term $A_2(x)$, we note that $\rho(x_j) > r_j \geq \rho(x_j)/4$ as following:

$$\begin{aligned}
&\| (b(x) - b_{B_j}) \mathcal{R}a_j(x) \chi_{(8B_j)^c}(x) \|_{L^1(\omega)} \\
&\leq \int_{B_j} |a_j(y)| dy \\
&\quad \times \left\{ \int_{|x-x_j| \geq 8r_j} |K(x, y)| |b(x) - b_{B_j}| \omega(x) dx \right\} \\
&\leq \int_{B_j} |a_j(y)| dy \\
&\quad \times \left\{ \int_{|x-x_j| \geq 8r_j} \frac{C_l}{(1 + |x-y|/\rho(x))^l} \frac{1}{|x-y|^{n-1}} \right. \\
&\quad \times \left(\int_{B(x, 2|x-x_j|)} \frac{V(z)}{|x-z|^{n-1}} dz + \frac{1}{|x-y|} \right) \\
&\quad \times |b(x) - b_{B_j}| \omega(x) dx \Big\} \\
&\leq \int_{B_j} |a_j(y)| \{I_1(y) + I_2(y)\} dy. \tag{47}
\end{aligned}$$

Note that $|x - x_j| \sim |x - y|$ and

$$\begin{aligned}
\left(1 + \frac{|x-y|}{\rho(x)} \right) &\geq C \left(1 + \frac{|x-x_j|}{\rho(x)} \right) \\
&\geq C \left(1 + \frac{|x-x_j|}{\rho(x_j)} \right)^{1/(k_0+1)}. \tag{48}
\end{aligned}$$

Then by Lemma 16 and Proposition 14, for any $y \in B_j$,

$$\begin{aligned}
 I_2(y) &\leq \sum_{k=1}^{\infty} \int_{2^{k+3}r_j \leq |x-x_j| < 2^{k+4}r_j} \frac{1}{\left(1 + |x-x_j| \rho(x_j)^{-1}\right)^{N/(k_0+1)}} \\
 &\quad \times \frac{1}{|x-x_j|^n} |b(x) - b_{B_j}| \omega(x) dx \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} \frac{1}{|B(x, 2^{k+3}r_j)|} \\
 &\quad \times \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}| \omega(x) dx \\
 &\leq \sum_{k=1}^{\infty} 2^{-(((k+1)N)/(k_0+1))} \left(\frac{1}{|2^{k+4}B_j|} \int_{2^{k+4}B_j} |b - b_{B_j}|^{r'} dx \right)^{1/r'} \\
 &\quad \times \left(\frac{1}{|2^{k+4}B_j|} \int_{2^{k+4}B_j} \omega(x)^r dx \right)^{1/r} \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} \\
 &\quad \times \left(\frac{1}{|2^{k+4}B_j|} \int_{2^{k+4}B_j} \omega(x) dx \right) \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} \\
 &\quad \times \left(\frac{1}{|2^{k+5}B_j|} \int_{B(y, 2^{k+5}r_j)} \omega(x) dx \right) \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} M(\omega)(y) \\
 &\leq C[b]_{\theta} M(\omega)(y),
 \end{aligned} \tag{49}$$

where we have chosen N large enough.

Similarly, for $(1/p) + (1/q) + (1/r) = 1$ and $(1/s_1) = (1/p) + (1/n)$, using Hölder inequality, Proposition 14, and the boundedness of the fractional integral $\mathcal{J}_1$, we obtain, for any $y \in B_j$,

$$\begin{aligned}
 I_1(y) &\leq \sum_{k=1}^{\infty} \int_{2^{k+3}r_j \leq |x-x_j| < 2^{k+4}r_j} \frac{1}{\left(1 + |x-x_j| \rho(x_j)^{-1}\right)^{N/(k_0+1)}} \\
 &\quad \times \frac{|b(x) - b_{B_j}|}{|x-x_j|^{n-1}} \mathcal{J}_1(V\chi_{2^{k+4}B_j})(x) \omega(x) dx
 \end{aligned}$$

$$\begin{aligned}
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} \frac{1}{(2^{k+3}r_j)^{n-1}} \\
 &\quad \times \int_{|x-x_j| < 2^{k+4}r_j} \mathcal{J}_1(V\chi_{2^{k+4}B_j})(x) |b - b_{B_j}| \omega(x) dx \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} (2^{k+3}r_j)^{1-n+(n/q)+(n/r)} \\
 &\quad \times \left(\frac{1}{(2^{k+3}r_j)^n} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\
 &\quad \times \left(\frac{1}{|2^{k+3}B_j|} \int_{2^{k+3}B_j} \omega(x)^r dx \right)^{1/r} \\
 &\quad \times \|\mathcal{J}_1(V\chi_{2^{k+4}B_j})\|_p \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} (2^{k+3}r_j)^{2-(n/s_1)} \\
 &\quad \times \left(\frac{1}{(2^{k+3}r_j)^n} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\
 &\quad \times \left(\frac{1}{|2^{k+3}B_j|} \int_{2^{k+3}B_j} \omega(x) dx \right) \|V\chi_{2^{k+4}B_j}\|_{s_1} \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} \\
 &\quad \times \left(\frac{1}{|2^{k+5}B_j|} \int_{B(y, 2^{k+5}r_j)} \omega(x) dx \right) \\
 &\quad \times (2^{k+3}r_j)^{2-n/s_1} \|V\chi_{2^{k+4}B_j}\|_{s_1} \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} \\
 &\quad \times M(\omega)(y) \frac{1}{(2^{k+3}r_j)^{n-2}} \int_{2^{k+3}B_j} V dx \\
 &\leq \sum_{k=1}^{\infty} 2^{-((k+1)N)/(k_0+1)} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} \\
 &\quad \times \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{l_1} M(\omega)(y) \\
 &\leq C[b]_{\theta} M(\omega)(y),
 \end{aligned} \tag{50}$$

where we have chosen N large enough and we have used Lemma 10 in the sixth inequality above.

Therefore,

$$\begin{aligned} & \int_{B_j} |a_j(y)| \{I_1(y) + I_2(y)\} dy \\ & \leq C[b]_\theta \int_{B_j} |a_j(y)| M(\omega)(y) dy \\ & \leq C[b]_\theta \int_{B_j} |a_j(y)| \omega(y) dy \leq C[b]_\theta, \end{aligned} \quad (51)$$

where we have used the fact that $\omega \in A_1$.

Thus, if $\rho(x_j) > r_j \geq (\rho(x_j))/4$, then

$$\left\| (b(x) - b_{B_j}) \mathcal{R}a_j(x) \chi_{(8B_j)^c}(x) \right\|_{L^1(\omega)} \leq C[b]_\theta. \quad (52)$$

For A_3 , by using the vanishing condition of a_j and Lemma 13, then

$$\begin{aligned} & \left\| (b(x) - b_{B_j}) \mathcal{R}a_j(x) \chi_{(8B_j)^c}(x) \right\|_{L^1(\omega)} \\ & \leq \int_{B_j} |a_j(y)| dy \\ & \quad \times \left\{ \int_{|x-x_j| \geq 8r_j} |K(x, y) - K(x, x_j)| \right. \\ & \quad \times |b(x) - b_{B_j}| \omega(x) dx \Big\} \\ & \leq \int_{B_j} |a_j(y)| dy \\ & \quad \times \left\{ \int_{|x-x_j| \geq 8r_j} \frac{C_l}{(1 + |x-y|/\rho(x))^N} \frac{|y-x_j|^\delta}{|x-y|^{n-1+\delta}} \right. \\ & \quad \times \left(\int_{B(x, 2|x-y|)} \frac{|V(z)|}{|x-z|^{n-1}} dz + \frac{1}{|x-y|} \right) \\ & \quad \times |b(x) - b_{B_j}| \omega(x) dx \Big\} \\ & = \int_{B_j} |a_j(y)| \{ \tilde{I}_1(y) + \tilde{I}_2(y) \} dy. \end{aligned} \quad (53)$$

First of all, we need to obtain the following estimate:

$$\begin{aligned} \left\| V \chi_{B(x_j, 2^{k+3}r_j)} \right\|_{s_1} & \leq (2^k r_j)^{-(n/s'_1)} \int_{B(x_j, 2^k r_j)} V(z) dz \\ & \leq (2^k r_j)^{n-2-(n/s'_1)} \frac{1}{(2^k r_j)^{n-2}} \\ & \quad \times \int_{B(x_j, 2^k r_j)} V(z) dz \\ & \leq (2^k r_j)^{n-2-(n/s'_1)} (1 + 2^k r_j m(x_j, |V|))^{l_1}, \end{aligned} \quad (54)$$

where we have used Lemma 10 in the last inequality above.

Similarly, for $(1/p) + (1/q) + (1/r) = 1$ and $(1/s_1) = (1/p) + (1/n)$, using Hölder inequality, Proposition 14, and the boundedness of the fractional integral $\mathcal{J}_1$, we obtain, for any $y \in B_j$,

$$\begin{aligned} \tilde{I}_1(y) & \leq \sum_{k=1}^{\infty} \int_{2^{k+3}r_j \leq |x-x_j| < 2^{k+4}r_j} \frac{\mathcal{J}_1(V \chi_{2^{k+4}B_j})(x)}{(1 + |x-x_j| \rho(x_j)^{-1})^{N/(l_0+1)}} \\ & \quad \times \frac{|x_j - y|^\delta |b(x) - b_{B_j}|}{|x-x_j|^{n-1} |x-x_j|^\delta} \omega(x) dx \\ & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \frac{(2^{k+3}r_j)^{1-n+n/q+n/r}}{(1 + 2^k r_j \rho(x_j)^{-1})^{N/(l_0+1)}} \\ & \quad \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\ & \quad \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} \omega(x)^r dx \right)^{1/r} \\ & \quad \times \left\| \mathcal{J}_1(V \chi_{2^{k+4}B_j}) \right\|_p \\ & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \frac{(2^{k+3}r_j)^{1-n/p}}{(1 + 2^k r_j \rho(x_j)^{-1})^{N/(l_0+1)}} \\ & \quad \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\ & \quad \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} \omega(x)^r dx \right)^{1/r} \\ & \quad \times \left\| V \chi_{2^{k+4}B_j} \right\|_{s_1} \\ & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \frac{(2^{k+3}r_j)^{2-n/s_1}}{(1 + 2^k r_j \rho(x_j)^{-1})^{N/(l_0+1)}} \\ & \quad \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\ & \quad \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} \omega(x) dx \right) \\ & \quad \times \left\| V \chi_{2^{k+4}B_j} \right\|_{s_1} \\ & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \frac{(2^{k+3}r_j)^2}{(1 + 2^k r_j \rho(x_j)^{-1})^{N/(l_0+1)}} \end{aligned}$$

$$\begin{aligned}
 & \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\
 & \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} \omega(x) dx \right) \\
 & \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} V(z) dz \right) \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \frac{(2^{k+3}r_j)^2}{(1 + 2^k r_j \rho(x_j)^{-1})^{N/(l_0+1)}} \\
 & \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}|^q dx \right)^{1/q} \\
 & \times \left(\frac{1}{|B(y, 2^{k+5}r_j)|} \int_{|x-y| < 2^{k+5}r_j} \omega(x) dx \right) \\
 & \times \left(\frac{1}{|B(x, 2^{k+3}r_j)|} \int_{|x-x_j| < 2^{k+4}r_j} V(z) dz \right) \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta} M(\omega)(y) \\
 & \quad \times (1 + 2^k r_j m(x_j, V))^{l_1 - (N/(l_0+1))} \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} [b]_{\theta} k M(\omega)(y) \\
 & \leq [b]_{\theta} M(\omega)(y),
 \end{aligned} \tag{55}$$

where we have chosen N large enough.

Secondly, for any $y \in B_j$,

$$\begin{aligned}
 & \tilde{I}_2(y) \\
 & \leq \int_{B_j} |a_j(y)| dy \\
 & \quad \times \sum_{k=1}^{\infty} \int_{2^{k+3}r_j \leq |x-x_j| < 2^{k+4}r_j} \frac{C_l}{(1 + |x-x_j| \rho(x_j)^{-1})^{N/(l_0+1)}} \\
 & \quad \times \frac{|x_j - y|^{\delta} |b(x) - b_{B_j}|}{|x - x_j|^n |x - x_j|^{\delta}} \omega(x) dx \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \left(1 + \frac{2^{k+3}r_j}{\rho(x_j)} \right)^{-N/(l_0+1)} \frac{1}{|B(x, 2^{k+3}r_j)|} \\
 & \quad \times \int_{|x-x_j| < 2^{k+4}r_j} |b - b_{B_j}| \omega(x) dx \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} \left(1 + \frac{2^{k+3}r_j}{\rho(x_j)} \right)^{-N/(l_0+1)}
 \end{aligned}$$

$$\begin{aligned}
 & \times \left(\frac{1}{|2^{k+4}B_j|} \int_{2^{k+4}B_j} |b - b_{B_j}|^{r'} dx \right)^{1/r'} \\
 & \times \left(\frac{1}{|2^{k+4}B_j|} \int_{2^{k+4}B_j} \omega(x)^r dx \right)^{1/r} \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} [b]_{\theta} k \left(1 + \frac{2^{k+4}r_j}{\rho(x_j)} \right)^{(k_0+1)\theta - (N/(l_0+1))} \\
 & \quad \times \left(\frac{1}{|2^{k+4}B_j|} \int_{2^{k+4}B_j} \omega(x) dx \right) \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} [b]_{\theta} k \left(\frac{1}{|2^{k+5}B_j|} \int_{B(y, 2^{k+5}r_j)} \omega(x) dx \right) \\
 & \leq \sum_{k=1}^{\infty} 2^{-(k+3)\delta} [b]_{\theta} k M(\omega)(y) \\
 & \leq [b]_{\theta} M(\omega)(y),
 \end{aligned} \tag{56}$$

where we have also chosen N large enough.

Therefore,

$$\begin{aligned}
 & \int_{B_j} |a_j(y)| \{ \tilde{I}_1(y) + \tilde{I}_2(y) \} dy \\
 & \leq C[b]_{\theta} \int_{B_j} |a_j(y)| M(\omega)(y) dy \\
 & \leq C[b]_{\theta} \int_{B_j} |a_j(y)| \omega(y) dy \leq C[b]_{\theta}.
 \end{aligned} \tag{57}$$

Therefore, if $r_j \leq \rho(x_j)/4$, then

$$\left\| (b(x) - b_{B_j}) \mathcal{R}a_j(x) \chi_{(8B_j)^c}(x) \right\|_{L^1(\omega)} \leq C[b]_{\theta}. \tag{58}$$

Thus, we have

$$\begin{aligned}
 & \omega \left\{ x \in \mathbb{R}^n : |A_i(x)| > \frac{\lambda}{4} \right\} \leq \frac{C}{\lambda} \|A_i(x)\|_{L^1(\omega)} \\
 & \leq \frac{C[b]_{\theta}}{\lambda} \sum_{j=-\infty}^{\infty} |\lambda_j|, \quad i = 1, 2, 3.
 \end{aligned} \tag{59}$$

Note that, for $(1/r') + (1/r) = 1$,

$$\begin{aligned}
 & \left\| (b - b_{B_j}) a_j \right\|_{L^1(\omega)} \\
 & \leq |8B_j| \left(\frac{1}{|8B_j|} \int_{8B_j} |b(x) - b_{B_j}|^{r'} dx \right)^{1/r'} \|a_j\|_{L^{\infty}(\omega)} \\
 & \quad \times \left(\frac{1}{|8B_j|} \int_{8B_j} \omega(x)^r dx \right)^{1/r}
 \end{aligned}$$

$$\begin{aligned}
&\leq |8B_j| \left(\frac{1}{|8B_j|} \int_{B_j} |b(x) - b_B|^{r'} dx \right)^{1/r'} \omega(B_j)^{-1} \\
&\quad \times \left(\frac{1}{|8B_j|} \int_{8B_j} \omega(x) dx \right) \\
&\leq C [b]_{\theta} \left(1 + \frac{r_j}{\rho(x_j)} \right)^{\theta'} \\
&\leq C [b]_{\theta},
\end{aligned} \tag{60}$$

where $r_j < \rho(x_j)$.

By Theorem 4, we get

$$\begin{aligned}
\omega \left\{ x \in \mathbb{R}^n : |A_4(x)| > \frac{\lambda}{4} \right\} &\leq \frac{C}{\lambda} \left\| \sum_{j=-\infty}^{\infty} \lambda_j (b - b(x_j)) a_j \right\|_{L^1(\omega)} \\
&\leq \frac{C[b]_{\theta}}{\lambda} \sum_{j=-\infty}^{\infty} |\lambda_j|.
\end{aligned} \tag{61}$$

Therefore,

$$\begin{aligned}
\omega \left\{ x \in \mathbb{R}^n : |[b, \mathcal{R}] f(x)| > \frac{\lambda}{4} \right\} \\
&\leq C \sum_{i=1}^4 \omega \left\{ x \in \mathbb{R}^n : |A_i(x)| > \frac{\lambda}{4} \right\} \\
&\leq \frac{C[b]_{\theta}}{\lambda} \sum_{j=-\infty}^{\infty} |\lambda_j| \\
&\leq \frac{C[b]_{\theta}}{\lambda} \|f\|_{H_L^1(\omega)}.
\end{aligned} \tag{62}$$

This completes the proof of Theorem 5.

References

- [1] R. R. Coifman, R. Rochberg, and G. Weiss, "Factorization theorems for Hardy spaces in several variables," *Annals of Mathematics*, vol. 103, no. 3, pp. 611–635, 1976.
- [2] Z. Guo, P. Li, and L. Peng, " L_p boundedness of commutators of Riesz transforms associated to Schrödinger operator," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 1, pp. 421–432, 2008.
- [3] P. Li and L. Peng, "Endpoint estimates for commutators of Riesz transforms associated with Schrödinger operators," *Bulletin of the Australian Mathematical Society*, vol. 82, no. 3, pp. 367–389, 2010.
- [4] B. Bongioanni, E. Harboure, and O. Salinas, "Commutators of Riesz transforms related to Schrödinger operators," *The Journal of Fourier Analysis and Applications*, vol. 17, no. 1, pp. 115–134, 2011.
- [5] B. Bongioanni, E. Harboure, and O. Salinas, "Weighted inequalities for commutators of Schrödinger-Riesz transforms," *Journal of Mathematical Analysis and Applications*, vol. 392, no. 1, pp. 6–22, 2012.
- [6] Y. Liu, "Commutators of BMO functions and degenerate Schrödinger operators with certain nonnegative potentials," *Monatshefte für Mathematik*, vol. 165, no. 1, pp. 41–56, 2012.
- [7] Y. Liu, J. Z. Huang, and J. F. Dong, "Commutators of higher order Riesz transform associated with Schrödinger operators," *Journal of Function Spaces and Applications*, vol. 2013, Article ID 842375, 15 pages, 2013.
- [8] Y. Liu, J. Z. Huang, and J. F. Dong, "Commutators of Calderón-Zygmund operators related to admissible functions on spaces of homogeneous type and applications to Schrödinger operators," *Science China Mathematics*, vol. 56, no. 9, pp. 1895–1913, 2013.
- [9] Z. W. Shen, " L_p estimates for Schrödinger operators with certain potentials," *Annales de l'Institut Fourier*, vol. 45, no. 2, pp. 513–546, 1995.
- [10] J. Dziubański and J. Zienkiewicz, "Hardy space H^1 associated to Schrödinger operator with potential satisfying reverse Hölder inequality," *Revista Matemática Iberoamericana*, vol. 15, no. 2, pp. 279–296, 1999.
- [11] H. Liu, L. Tang, and H. Zhu, "Weighted Hardy spaces and BMO spaces associated with Schrödinger operators," *Mathematische Nachrichten*, vol. 285, no. 17-18, pp. 2173–2207, 2012.
- [12] D. Yang and Y. Zhou, "Localized Hardy spaces H^1 related to admissible functions on RD-spaces and applications to Schrödinger operators," *Transactions of the American Mathematical Society*, vol. 363, no. 3, pp. 1197–1239, 2011.
- [13] E. M. Stein, *Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, vol. 43 of *Princeton Mathematical Series*, Princeton University Press, Princeton, NJ, USA, 1993.
- [14] J. Duoandikoetxea, *Fourier Analysis*, vol. 29 of *Graduate Studies in Mathematics*, American Mathematical Society, Providence, RI, USA, 2001.
- [15] C. C. Lin, H. P. Liu, and Y. Liu, "Hardy spaces associated with Schrödinger operators on the Heisenberg group," <http://arxiv.org/abs/1106.4960>.
- [16] J. García-Cuerva and J. L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, vol. 116 of *North-Holland Mathematics Studies*, North-Holland, Amsterdam, The Netherlands, 1985.