Research Article

Oscillation Criteria of Third-Order Nonlinear Impulsive Differential Equations with Delay

Xiuxiang Liu

School of Mathematical Sciences, South China Normal University, Guangzhou, Guangdong 510631, China

Correspondence should be addressed to Xiuxiang Liu; liuwx@scnu.edu.cn

Received 14 November 2012; Accepted 10 January 2013

Academic Editor: Norio Yoshida

Copyright © 2013 Xiuxiang Liu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper deals with the oscillation of third-order nonlinear impulsive equations with delay. The results in this paper improve and extend some results for the equations without impulses. Some examples are given to illustrate the main results.

1. Introduction

In this paper, we are concerned with oscillation of the third-order nonlinear impulsive equations with delay

$$\begin{aligned} x'''(t) + f(t, x(t), x(t-\sigma)) &= 0, \quad t \geq t_0, \quad t \neq \tau_k, \\ x(\tau_k^+) &= a_k x(\tau_k), \quad x'(\tau_k^+) = b_k x'(\tau_k), \\ x''(\tau_k^+) &= c_k x''(\tau_k), \quad k = 1, 2, \dots, \end{aligned} \quad (1)$$

where $\sigma > 0$ is the delay, $\{\tau_k\}$ is the sequence of impulsive moments which satisfies $0 \leq t_0 < \tau_1 < \dots < \tau_k < \dots$ and $\lim_{k \rightarrow \infty} \tau_k = \infty$, $\tau_{k+1} - \tau_k \geq \sigma$.

Throughout this paper, we will assume that the following assumptions are satisfied:

- (H1) $f(t, u, v)$ is continuous in $[t_0 - \sigma, \infty) \times \mathbb{R} \times \mathbb{R}$, $uf(t, u, v) > 0$ for $uv > 0$;
- (H2) $f(t, u, v)/\varphi(v) \geq p(t)$ for $v \neq 0$, where $p(t)$ is continuous in $[t_0 - \sigma, \infty)$, $p(t) \geq 0$ ($\neq 0$), $\varphi(x)/x \geq \mu > 0$ for all $x \neq 0$;
- (H3) a_k, b_k , and c_k are positive constants.

Our attention is restricted to those solutions of (1) which exist on half line $[t_0, \infty)$ and satisfy $\sup\{|x(t)| : t > T\} > 0$ for any $T \geq t_x$. For the general theory of impulsive differential equations with/without delay, we refer the readers to monographs or papers [1–4]. A solution of (1) is said to be oscillatory if it has arbitrarily large zeros; otherwise,

it is nonoscillatory. It is well known that there is a drastic difference in the behavior of solutions between differential equations with impulses and those without impulses. Some differential equations are nonoscillatory, but they may become oscillatory if some proper impulse controls are added to them, see [5] and Example 13 in Section 4. In the recent years, the oscillation theory and asymptotic behavior of impulsive differential equations and their applications have been and still are receiving intensive attention. For contribution, we refer to the recent survey paper by Agarwal et al. [6] and the references cited therein. But to the best of the authors' knowledge, it seems that little has been done for oscillation of third-order impulsive differential equations [7].

When $a_k = b_k = c_k = 1$, (1) reduces third-order delay equation with/without delay, which oscillatory theory has been studied by many researchers, see [8–12].

Our aim in this paper is to establish some new sufficient conditions which ensure that the solutions of (1) oscillate or converge to a finite limit as t tends to infinity. In particular, we extend some results in [9, 11] to impulsive delay differential equations. The results in this paper are more general compared by those obtained by Mao and Wan [7] and improve some of the results in [7] (see Example 13 in Section 4). The new results will be proved by making use of the techniques used in [9, 11].

The paper is organized as follows. In Section 2, we prove some lemmas which play important roles in the proof of the main results. In Section 3, some new sufficient conditions which guarantee that the solution of (1) oscillates or

converges to a finite limit are established. In Section 4, two examples are given to illustrate the main results.

2. Preliminary Results

In this section, we state and prove some lemmas which we will need in the proofs of the main results. First of all, we introduce the following notations: $\mathbb{R}^+$ and $\mathbb{N}$ are the sets of real numbers and positive integer numbers, respectively, PC^1 is defined by

$$\begin{aligned} PC^1(\mathbb{R}^+, \mathbb{R}) \\ = \{x : \mathbb{R}^+ \rightarrow \mathbb{R} : x(t) \text{ is differentiable for} \\ t \geq 0 \text{ and } t \neq \tau_k, x(\tau_k^+) \text{ and } x'(\tau_k^+) \text{ exist,} \\ \text{and } x(\tau_k^-) = x(\tau_k), x'(\tau_k^-) = x'(\tau_k)\}. \end{aligned} \quad (2)$$

The following lemma is from Lakshmikantham et al. [3, Page 32, Theorem 1.4.1].

Lemma 1. Assume that

- (i) $\{\tau_k\}_{k \in \mathbb{N}}$ is the impulse moments sequence with $0 \leq t_0 < \tau_1 < \dots < \tau_k < \dots$, $\lim_{k \rightarrow \infty} \tau_k = \infty$;
- (ii) $m \in PC^1(\mathbb{R}^+, \mathbb{R})$, and for $t \geq t_0$, $k \in \mathbb{N}$, it holds that

$$\begin{aligned} m'(t) &\leq u(t)m(t) + v(t), \quad t \neq \tau_k, \\ m(\tau_k^+) &\leq d_k m(\tau_k) + e_k, \end{aligned} \quad (3)$$

where $u, v \in C(\mathbb{R}^+, \mathbb{R})$, $d_k \geq 0$, and e_k are real constants. Then,

$$\begin{aligned} m(t) &\leq m(t_0) \prod_{t_0 < \tau_k < t} d_k \exp\left(\int_{t_0}^t u(s)ds\right) \\ &+ \int_{t_0}^t \prod_{s < \tau_k < t} d_k \exp\left(\int_s^t u(\sigma)d\sigma\right) v(s)ds \\ &+ \sum_{t_0 < \tau_k < t} \prod_{\tau_k < \tau_j < t} d_j \exp\left(\int_{\tau_k}^t u(s)ds\right) e_k. \end{aligned} \quad (4)$$

Motivated by the ideas of Chen and Feng [5], we present the following key lemma which determines the sign of $x'(t)$ and $x''(t)$ of the nonoscillation solution $x(t)$ of (1).

Lemma 2. Suppose that $x(t)$ is an eventually positive solution of (1), and

$$\lim_{t \rightarrow \infty} \int_{t_0}^t \prod_{t_0 < \tau_k < s} \frac{b_k}{a_k} ds = \infty, \quad (5)$$

$$\lim_{t \rightarrow \infty} \int_{t_0}^t \prod_{t_0 < \tau_k < s} \frac{c_k}{b_k} ds = \infty. \quad (6)$$

Then, it holds that one of the following two cases for sufficiently large T :

- (i) $x''(\tau_k^+) > 0$, $x''(t) > 0$ and $x'(\tau_k^+) > 0$, $x'(t) > 0$,
- (ii) $x''(\tau_k^+) > 0$, $x''(t) > 0$ and $x'(\tau_k^+) < 0$, $x'(t) < 0$,

with $t \in (\tau_k, \tau_{k+1}]$ and $\tau_k \geq T$.

Proof. Assume that $x(t)$ is an eventually positive solution of (1). We may assume that there exists $t_1 > t_0$ such that $x(t) > 0$ and $x(t - \sigma) > 0$ for $t \geq t_1$. First, we assert that $x''(\tau_k) > 0$ for any $k \in \mathbb{N}$. Suppose not, there exists some $\tau_j \geq t_1$ such that $x''(\tau_j) \leq 0$. By

$$x'''(t) = -f(t, x(t), x(t - \sigma)) \leq 0, \neq 0, \quad (7)$$

we have $x''(t)$ monotonically decreasing in $(\tau_i, \tau_{i+1}]$, $i = j, j+1, \dots$. Thus, $x''(\tau_i^+) = c_i x''(\tau_i) < 0$, $i = j+1, j+2, \dots$. Consider the impulsive differential inequalities

$$\begin{aligned} x'''(t) &< 0, \quad t \geq \tau_{j+1}, \quad t \neq \tau_k, \\ x''(\tau_k^+) &\leq c_k x''(\tau_k), \quad k = j+1, j+2, \dots \end{aligned} \quad (8)$$

By Lemma 1, we have

$$x''(t) \leq x''(\tau_{j+1}^+) \prod_{\tau_{j+1} < \tau_k < t} c_k := -\alpha \prod_{\tau_{j+1} < \tau_k < t} c_k < 0. \quad (9)$$

There are two cases of the sign of $x'(\tau_k)$.

Case 1. If there exists some $\tau_n \geq \tau_{j+1}$ such that $x'(\tau_n) \leq 0$, since $x''(t) < 0$, then $x'(\tau_{n+1}) < x'(\tau_n^+) \leq 0$ and $x'(\tau_{n+1}^+) := \beta = b_{n+1} x'(\tau_{n+1}) < 0$. By induction it easily show $x'(\tau_k) < 0$, and hence $x'(\tau_k^+) \leq b_k x'(\tau_k) < 0$ for $k = n+1, n+2, \dots$. So, we obtain the following impulsive differential inequalities:

$$\begin{aligned} x''(t) &\leq 0, \quad t \geq \tau_{n+1}, \quad t \neq \tau_k, \\ x'(\tau_k^+) &\leq b_k^* x'(\tau_k), \quad k = n+1, n+2, \dots, \end{aligned} \quad (10)$$

which follows from Lemma 1 that

$$x'(t) \leq x'(\tau_{n+1}^+) \prod_{\tau_{n+1} < \tau_k < t} b_k = -\beta \prod_{\tau_{n+1} < \tau_k < t} b_k. \quad (11)$$

From (11) and applying Lemma 1, noting that $x(t) > 0$ for $t > t_1$ and $x(\tau_k^+) \leq a_k x(\tau_k)$ for any $k \in \mathbb{N}$, we have

$$\begin{aligned} x(t) &\leq x(\tau_{n+1}^+) \prod_{\tau_{n+1} < \tau_k < t} a_k - \beta \int_{\tau_{n+1}}^t \prod_{s < \tau_k < t} a_k \prod_{\tau_{n+1} < \tau_k < s} b_k ds \\ &= \prod_{\tau_{n+1} < \tau_k < t} a_k \left(x(\tau_{n+1}^+) - \beta \int_{\tau_{n+1}}^t \prod_{\tau_{n+1} < \tau_k < s} \frac{b_k}{a_k} ds \right). \end{aligned} \quad (12)$$

Thus, by (5) we have $x(t) < 0$ for t sufficiently large which is a contradiction.

Case 2. If $x'(\tau_k) > 0$ for any $k \geq j$, noting that $x''(t) < 0$ for $t \in (\tau_{j+1}, \tau_{j+2}]$, we have $x'(t) > x'(\tau_{j+1}) > 0$. By induction,

we get that $x'(t) > 0$ for any $t \in (\tau_k, \tau_{k+1}]$, $k = j+1, j+2, \dots$. So the following impulsive differential inequalities hold:

$$x''(t) \leq -\alpha \prod_{\tau_{j+1} < \tau_k < t} c_k, \quad t > \tau_{j+1}, \quad t \neq \tau_k, \quad (13)$$

$$x'(\tau_k^+) \leq b_k x(\tau_k), \quad k = j+1, j+2, \dots$$

According to Lemma 1, we get

$$\begin{aligned} x'(t) &\leq x'(\tau_{j+1}^+) \prod_{\tau_{j+1} < \tau_k < t} b_k - \alpha \int_{\tau_{j+1}}^t \prod_{s < \tau_k < t} b_k \prod_{\tau_{j+1} < \tau_k < s} c_k ds \\ &= \prod_{\tau_{j+1} < \tau_k < t} b_k \left(x'(\tau_{j+1}^+) - \alpha \int_{\tau_{j+1}}^t \prod_{\tau_{j+1} < \tau_k < s} \frac{c_k}{b_k} ds \right). \end{aligned} \quad (14)$$

Hence, the condition (6) implies that $x'(t) < 0$ when t is sufficiently large, which contradicts to $x'(t) > 0$ for $t > \tau_{j+1}$ again. In terms of the above discussion, we see that $x''(\tau_k) > 0$ for any $\tau_k > T$ with sufficiently large T . Consequently, noting that $x'''(t) < 0$ for any $t \in (\tau_k, \tau_{k+1}]$, we have $x''(t) > x''(\tau_{k+1}) > 0$.

Next, if there exists a $\tau_j > T$ such that $x'(\tau_j) \geq 0$, then $x'(\tau_j^+) = b_j(x(\tau_j^+) \geq 0$, $x'(\tau_{j+1}) > x'(\tau_j^+) \geq 0$. Therefore, by induction, we have $x'(t) > x'(\tau_i^+) > 0$ for $t \in (\tau_i, \tau_{i+1}]$, $i = j+1, j+2, \dots$. So case (i) is satisfied. Otherwise, if $x'(\tau_k) < 0$ for all $\tau_k \geq T$, then $x'(\tau_k^+) = b_k x(\tau_k^+) < 0$. Thus, for $t \in (\tau_k, \tau_{k+1}]$, using $x''(t) > 0$, we have $x'(t) < x'(\tau_{k+1}) < 0$; hence, case (ii) is satisfied. This completes the proof. $\square$

Remark 3. Suppose that $x(t)$ is an eventually negative solution of (1). If (5) and (6) hold, one can prove it holds that one of the following two cases in a similar way as Lemma 2:

- (i) $x''(\tau_k^+) < 0$, $x''(t) < 0$ and $x'(\tau_k^+) < 0$, $x'(t) < 0$,
- (ii) $x''(\tau_k^+) < 0$, $x''(t) < 0$ and $x'(\tau_k^+) > 0$, $x'(t) > 0$, with $t \in (\tau_k, \tau_{k+1}]$ and $\tau_k \geq T$.

Lemma 4. Let $x(t)$ be a piecewise continuous function on $\cup_{k \in \mathbb{N}} (\tau_k, \tau_{k+1}]$, which is continuous at $t \neq \tau_k$ and is left continuous at $t = \tau_k$. If

- (1) $x(t) \geq 0$ (≤ 0) for $t \geq t_0$;
- (2) $x(t)$ is monotone nonincreasing (monotone nondecreasing) on $(\tau_k, \tau_{k+1}]$ ($\tau_k \geq T$) for T large enough;
- (3) $\sum_{k=1}^{\infty} [x(\tau_k^+) - x(\tau_k)]$ converges,

then $\lim_{t \rightarrow \infty} x(t) = a \geq 0$ (≤ 0).

The proof of Lemma 4 is similar to that of [13, Theorem 5], and hence is omitted.

Lemma 5. Assume that $x(t)$ is a solution of (1) which satisfies case (ii) in Lemma 2. In addition, if

$$\sum_{k=1}^{\infty} |a_k - 1| < \infty, \quad \prod_{k=1}^{\infty} a_k \text{ is bounded}, \quad (15)$$

then $\lim_{t \rightarrow \infty} x(t)$ exists (finite).

Proof. First, we claim that $\sum_{k=1}^{\infty} [x(\tau_k^+) - x(\tau_k)]$ is convergence. In fact, since $x(t)$ is decreasing on $(\tau_k, \tau_{k+1}]$ ($\tau_k \geq T$, and T is defined in Lemma 2), then

$$x(\tau_{k+1}) \leq x(\tau_k^+), \quad x(\tau_{k+1}^+) \leq a_{k+1} x(\tau_{k+1}) \leq a_{k+1} x(\tau_k^+). \quad (16)$$

Obviously, by induction, we can get

$$\begin{aligned} x(\tau_{k+n}) &\leq a_{k+n-1} \cdots a_{k+1} x(\tau_k^+), \\ x(\tau_{k+n}^+) &\leq a_{k+n} \cdots a_{k+1} x(\tau_k^+). \end{aligned} \quad (17)$$

Since $\prod_{k=1}^{\infty} a_k$ is bounded, we conclude that $\{x(\tau_k)\}$ is bounded, which follows that there exists $M_1 > 0$ such that

$$|x(\tau_k^+) - x(\tau_k)| = |a_k - 1| x(\tau_k) \leq M_1 |a_k - 1|. \quad (18)$$

Hence, $\sum_{k=1}^{\infty} [x(\tau_k^+) - x(\tau_k)]$ is convergence since $\sum_{k=1}^{\infty} |a_k - 1|$ is convergence, which follows from Lemma 4 that $\lim_{t \rightarrow \infty} x(t)$ exists. The proof is complete. $\square$

3. Main Results

In this section, we establish some sufficient conditions which guarantee that every solution $x(t)$ of (1) either oscillates or has a finite limit. Occasionally, we will make the additional assumption

$$\limsup_{t \rightarrow \infty} \int_{\eta}^t \int_u^{\infty} \int_s^{\infty} p(\theta) d\theta ds du = \infty, \quad (19)$$

where here it is understood that

$$\int_{t_0}^{\infty} p(t) dt < \infty, \quad \int_{t_0}^{\infty} \int_u^{\infty} p(\theta) d\theta du < \infty. \quad (20)$$

Now we are ready to state and prove the main results in this paper. The results will be proved by making use of the technique in [11].

Theorem 6. Assume that (5), (6), and (19) hold, and $x(t)$ is a solution of (1). Furthermore, assume that $a_k \leq 1$, $b_k \geq 1$, and $c_k \leq 1$ for $k \in \mathbb{N}$. If there exists a positive differentiable function r such that

$$\lim_{t \rightarrow \infty} \int_{t_0}^t [p(s) r(s) - A(s, \tau_{k(s)})] \prod_{t_0 < \tau_k < s} \frac{1}{c_k} \prod_{t_0 < \tau_k < \sigma < s} a_k ds = \infty, \quad (21)$$

where $k(s) = \max\{k : \tau_k < s\}$, and

$$\begin{aligned} A(s, \tau_{k(s)}) &= \begin{cases} \frac{c_{k(s)} [r'(s)]^2}{4\mu r(s) (s - \tau_{k(s)})}, & s \in (\tau_k, \tau_k + \sigma], \\ \frac{[r'(s)]^2}{4\mu r(s) (s - \tau_{k(s)} - ((c_{k(s)} - 1)/c_{k(s)})\sigma)}, & s \in (\tau_k + \sigma, \tau_{k+1}]. \end{cases} \end{aligned} \quad (22)$$

Then $x(t)$ is oscillatory.

Proof. Let $x(t)$ be a nonoscillatory solution of (1), without loss of generality, we may assume that $x(t) > 0$ eventually ($x(t) < 0$ eventually can be achieved in the similar way). By Lemma 2, either case (i) or case (ii) in Lemma 2 holds. Assume that $x(t)$ satisfies case (i), then $x''(t) > 0$, $x'(t) > 0$ for $t \in (\tau_j, \tau_{j+1}]$, $\tau_j \geq T$ (T is defined in Lemma 2). Define the Riccati transformation u by

$$u(t) = \frac{r(t)x''(t)}{\varphi(x(t-\sigma))}, \quad t \geq \tau_j, \quad t \neq \tau_k. \quad (23)$$

Thus, $u(t) > 0$ for $t \in (\tau_k, \tau_{k+1}]$, $k = j, j+1, \dots$, and

$$\begin{aligned} u'(t) &= \frac{r(t)}{\varphi(x(t-\sigma))} x'''(t) \\ &\quad + \frac{r'(t)\varphi(x(t-\sigma)) - r(t)\varphi'(x(t-\sigma))x'(t-\sigma)}{\varphi^2(x(t-\sigma))} \\ &\quad \times x''(t) \\ &\leq -p(t)r(t) + \frac{r'(t)}{r(t)}u(t) \\ &\quad - \frac{r(t)\mu x'(t-\sigma)}{\varphi^2(x(t-\sigma))}x''(t). \end{aligned} \quad (24)$$

If $t \in (\tau_k, \tau_k + \sigma] \subset (\tau_k, \tau_{k+1}]$, namely, $t - \sigma \leq \tau_k < t$, $x''(t)$ is decreasing in $(t - \sigma, \tau_k]$ and $(\tau_k, t]$, respectively. In view of the following

$$x''(t - \sigma) > x''(\tau_k) \geq \frac{x''(\tau_k^+)}{c_k} > \frac{x''(t)}{c_k}, \quad (25)$$

we have

$$\begin{aligned} x'(t - \sigma) &= x'(\tau_k - \sigma) + \int_{\tau_k - \sigma}^{t - \sigma} x''(s) ds \\ &> x''(t - \sigma)(t - \tau_k) > \frac{x''(t)}{c_k}(t - \tau_k). \end{aligned} \quad (26)$$

Thus,

$$\begin{aligned} u'(t) &\leq -p(t)r(t) + \frac{r'(t)}{r(t)}u(t) - \frac{\mu r(t)(t - \tau_k)}{c_k \varphi^2(x(t - \sigma))} [x''(t)]^2 \\ &= -p(t)r(t) + \frac{r'(t)}{r(t)}u(t) - \frac{\mu(t - \tau_k)}{c_k r(t)} [u(t)]^2 \\ &= -\left(\sqrt{\frac{\mu(t - \tau_k)}{c_k r(t)}} u(t) - \sqrt{\frac{c_k}{4\mu r(t)(t - \tau_k)}} r'(t) \right)^2 \\ &\quad - \left[p(t)r(t) - \frac{c_k [r'(t)]^2}{4\mu r(t)(t - \tau_k)} \right] \\ &\leq -\left[p(t)r(t) - \frac{c_k [r'(t)]^2}{4\mu r(t)(t - \tau_k)} \right]. \end{aligned} \quad (27)$$

If $t \in (\tau_k + \sigma, \tau_{k+1}] \subset (\tau_k, \tau_{k+1}]$, that is, $\tau_k < t - \sigma < t \leq \tau_{k+1}$, then

$$\begin{aligned} x'(t - \sigma) &= x'(\tau_k - \sigma) + \int_{\tau_k - \sigma}^{\tau_k} x''(s) ds + \int_{\tau_k}^{t - \sigma} x''(s) ds \\ &\geq x''(\tau_k)\sigma + x''(t - \sigma)(t - \tau_k - \sigma) \\ &\geq \frac{x''(t)}{c_k}\sigma + x''(t)(t - \tau_k - \sigma) \\ &\geq x''(t)\left(t - \tau_k - \frac{c_k - 1}{c_k}\sigma\right). \end{aligned} \quad (28)$$

Similarly, we have

$$u'(t) \leq -\left[p(t)r(t) - \frac{[r'(t)]^2}{4\mu r(t)(t - \tau_k - ((c_k - 1)/c_k)\sigma)} \right]. \quad (29)$$

Thus, we obtain

$$u'(t) \leq -[p(t)r(t) - A(t, \tau_{k(t)})] \quad \text{for } t \in (\tau_k, \tau_{k+1}]. \quad (30)$$

On the other hand,

$$u(\tau_k^+) = \frac{r(\tau_k)x''(\tau_k^+)}{\varphi(x(\tau_k - \sigma))} \leq c_k u(\tau_k). \quad (31)$$

Observing that $\varphi(u) \geq \mu u$, we have

$$\begin{aligned} u(\tau_k^+ + \sigma) &= \frac{r(\tau_k + \sigma)x''(\tau_k + \sigma)}{\varphi(x(\tau_k^+))} \\ &\leq \frac{r(\tau_k + \sigma)x''(\tau_k + \sigma)}{\varphi(a_k x(\tau_k))} \leq \frac{u(\tau_k + \sigma)}{\mu a_k}. \end{aligned} \quad (32)$$

Applying Lemma 1, it follows from (30), (31), and (32) that

$$\begin{aligned} u(t) &\leq u(\tau_j^+) \prod_{\tau_j < \tau_k < t} c_k \prod_{\tau_j < \tau_k + \sigma < t} \frac{1}{\mu a_k} \\ &\quad - \int_{\tau_j}^t [p(s)r(s) - A(s, \tau_{k(s)})] \\ &\quad \cdot \prod_{s < \tau_k < t} c_k \prod_{s < \tau_k + \sigma < t} \frac{1}{\mu a_k} ds \\ &\leq \prod_{\tau_j < \tau_k < t} c_k \prod_{\tau_j < \tau_k + \sigma < t} \frac{1}{\mu a_k} \\ &\quad \times \left(u(\tau_j^+) - \mu \int_{\tau_j}^t [p(s)r(s) - A(s, \tau_{k(s)})] \right. \\ &\quad \cdot \left. \prod_{\tau_j < \tau_k < s} \frac{1}{c_k} \prod_{\tau_j < \tau_k + \sigma < s} a_k ds \right), \end{aligned} \quad (33)$$

which yields $u(t) < 0$ for all large t . This is contrary to $u(t) > 0$, and so, case (i) in Lemma 2 is not possible.

If $x(t)$ satisfies the case (ii) in Lemma 2, that is, $x''(\tau_k^+) > 0$, $x''(t) > 0$ and $x'(\tau_k^+) < 0$, $x'(t) < 0$, which proves that the solution $x(t)$ is positive and decreasing. Integrating (1) from s to t ($t \geq s \geq T$), we obtain

$$\begin{aligned} x''(t) - \sum_{s < \tau_k < t} (c_k - 1) x''(\tau_k) - x''(s) \\ + \int_s^t p(\theta) \varphi(x(\theta - \sigma)) d\theta \leq 0. \end{aligned} \quad (34)$$

Noting $c_k \leq 1$ and $x''(t) > 0$, then it holds that

$$x''(t) - x''(s) + \int_s^t p(\theta) \varphi(x(\theta - \sigma)) d\theta \leq 0, \quad (35)$$

which leads to

$$-x''(s) + \int_s^t p(\theta) \varphi(x(\theta - \sigma)) d\theta \leq 0, \quad (36)$$

and hence

$$-x''(s) + \int_s^\infty p(\theta) \varphi(x(\theta - \sigma)) d\theta \leq 0. \quad (37)$$

Integrating the above inequality again from u to t ($t \geq u \geq T$), one has

$$\begin{aligned} -x'(t) + \sum_{s < \tau_k < t} (b_k - 1) x'(\tau_k) + x'(u) \\ + \int_u^t \int_s^\infty p(\theta) \varphi(x(\theta - \sigma)) d\theta ds \leq 0. \end{aligned} \quad (38)$$

Using $x'(t) < 0$ and $b_k \geq 1$, we have

$$x'(u) + \int_u^\infty \int_s^\infty p(\theta) \varphi(x(\theta - \sigma)) d\theta ds \leq 0. \quad (39)$$

Now, we integrate the last inequality from η to t ($t \geq \eta \geq T$) to obtain

$$\begin{aligned} x(t) - \sum_{\eta < \tau_k < t} (a_k - 1) x(\tau_k) - x(\eta) \\ + \int_\eta^t \int_u^\infty \int_s^\infty p(\theta) \varphi(x(\theta - \sigma)) d\theta ds du \leq 0. \end{aligned} \quad (40)$$

Since $a_k \leq 1$ ($k \in \mathbb{N}$) and $x(t)$ is decreasing, then for $t \in (\tau_k, \tau_{k+1}]$, $x(\tau_{k+1}) \leq x(t) \leq x(\tau_k^+) = a_k x(\tau_k) \leq x(\tau_k)$, $k \in \mathbb{N}$. Thus, we get

$$\mu x(\eta) \int_\eta^t \int_u^\infty \int_s^\infty p(\theta) d\theta ds du \leq -x(t) + x(\eta) \leq x(\eta), \quad (41)$$

and then,

$$\int_\eta^t \int_u^\infty \int_s^\infty p(\theta) d\theta ds du \leq \frac{1}{\mu}, \quad (42)$$

which contradicts the condition (19). The proof is complete. $\square$

Replace the condition (19) with (15), we may obtain the following asymptotic results.

Theorem 7. Assume that (5), (6), and (15) hold, and $x(t)$ is a solution of (1). If there exists a positive differentiable function r such that (21) hold, then $x(t)$ is either oscillatory or has a finite limit.

Proof. By the proof of Theorem 6, we know the case (i) in Lemma 2 is not possible, too, since the condition (19) is not required to prove it. So it suffices to show if there is a solution satisfying case (ii) in Lemma 2, that is, if

$$x''(\tau_k^+) > 0, \quad x''(t) > 0, \quad x'(\tau_k^+) < 0, \quad x'(t) < 0 \quad (43)$$

with $t \in (\tau_k, \tau_{k+1}]$ and $\tau_k \geq T$. then $\lim_{t \rightarrow \infty} x(t)$ exists. This is obtained by applying Lemma 5 which leads to $\lim_{t \rightarrow \infty} x(t)$ exists. The proof is complete. $\square$

Corollary 8. In addition to the assumption of Theorem 7, assume that

$$\lim_{t \rightarrow \infty} \int_{t_0}^t \prod_{t_0 < \tau_k < s} \frac{1}{c_k} p(s) ds = \infty. \quad (44)$$

Then, solution $x(t)$ of (1) either oscillates or satisfies $\lim_{t \rightarrow \infty} x(t) = 0$.

Proof. By the proof of Theorem 7, $\lim_{t \rightarrow \infty} x(t)$ exists, and we define it by $\lim_{t \rightarrow \infty} x(t) = \gamma \geq 0$. We now show $\gamma = 0$. If not, then $\gamma > 0$. So, $\lim_{t \rightarrow \infty} \varphi(x(t - \sigma)) = \varphi(\gamma) =: \kappa > 0$. Hence, there exists $\tau_j > T$ such that $\varphi(x(t - \sigma)) > \kappa/2$ for $t > \tau_j$. Then

$$\begin{aligned} x'''(t) &= -f(t, x(t), x(t - \sigma)) \\ &\leq -p(t) \varphi(x(t - \sigma)) \leq -\frac{\kappa}{2} p(t), \quad t \geq \tau_j, \end{aligned} \quad (45)$$

and note that $x''(\tau_k^+) \leq c_k x''(\tau_k)$ since $x''(t) > 0$, which imply that

$$\begin{aligned} x''(t) &\leq x''(\tau_j^+) \prod_{\tau_j < \tau_k < t} c_k - \frac{\kappa}{2} \int_{\tau_j}^t \prod_{s < \tau_k < t} c_k p(s) ds \\ &\leq \prod_{\tau_j < \tau_k < t} c_k \left[x''(\tau_j^+) - \frac{\kappa}{2} \int_{\tau_j}^t \prod_{\tau_j < \tau_k < s} \frac{1}{c_k} p(s) ds \right]. \end{aligned} \quad (46)$$

Thus, in virtue of (44) it holds that $x''(t) < 0$ and contradicts $x''(t) > 0$ for t large enough, the proof is complete. $\square$

Remark 9. Theorem 6 and Corollary 8 extend the results in [11, Theorem 3.1] and [9, Corollary 1], respectively. In fact, when $a_k = b_k = c_k = 1$ for $k \in \mathbb{N}$ which implies that the impulses in (1) disappear. In such a case, (5) and (6) hold naturally, and (21) and (44) are reduced to

$$\lim_{t \rightarrow \infty} \int_{t_0}^t \left[p(s) r(s) - \frac{[r'(s)]^2}{4\mu r(s)(s - T)} \right] ds = \infty, \quad (47)$$

$$\lim_{t \rightarrow \infty} \int_{t_0}^t p(s) ds = \infty,$$

which are similar to those in [11, Theorem 3.1] and [9, Corollary 1], respectively.

Next, we present some new oscillation results for (1), by using an integral averaging condition of Kamenev's type.

Theorem 10. Assume (5), (6), and (19) hold. Furthermore, $a_k \leq 1/\mu \leq 1$, $b_k \geq 1$ and $c_k \leq 1$, $k \in \mathbb{N}$. If there exists a positive differentiable function r such that

$$\limsup_{t \rightarrow \infty} \frac{1}{t^m} \int_T^t (t-s)^m [p(s)r(s) - A(s, \tau_{k(s)})] ds = \infty, \quad (48)$$

where $A(s, \tau_{k(s)})$ is defined by (21) and $m \geq 1$. Then every solution of (1) is oscillatory.

Proof. We choose T large enough such that Lemma 2 holds. By Lemma 2 there are two possible cases. First, if the case (i) holds, proceeding as in the proof of Theorem 6, we will end up with (32). By (30), we have

$$p(t)r(t) - A(t, \tau_{k(t)}) \leq -u'(t), \quad t \geq T, \quad t \neq \tau_k. \quad (49)$$

If $t \in (\tau_k + \sigma, \tau_{k+1}] \subset (\tau_k, \tau_{k+1}]$, for $\tau_j \geq T$, we obtain

$$\begin{aligned} & \int_{\tau_j}^t (t-s)^m [p(s)r(s) - A(s, \tau_{k(s)})] ds \\ & \leq - \int_{\tau_j}^t (t-s)^m u'(s) ds. \end{aligned} \quad (50)$$

An integration by parts of the right-hand side leads to

$$\begin{aligned} & \int_{\tau_j}^t (t-s)^m u'(s) ds \\ & = \left(\int_{\tau_j}^{\tau_j+\sigma} + \int_{\tau_j+\sigma}^{\tau_{j+1}} + \cdots + \int_{\tau_k}^{\tau_k+\sigma} + \int_{\tau_k+\sigma}^t \right) (t-s)^m u'(s) ds \\ & = \int_{\tau_j}^t \frac{u(s)}{m} (t-s)^{m-1} ds \\ & \quad + \sum_{i=j}^k [t - (\tau_i + \sigma)]^m [u(\tau_i + \sigma) - u(\tau_i^+ + \sigma)] \\ & \quad + \sum_{i=j+1}^k (t - \tau_i)^m [u(\tau_i) - u(\tau_i^+)] - (t - \tau_j)^m u(\tau_j^+). \end{aligned} \quad (51)$$

Take into account (31), (32), $a_k \leq 1/\mu$, and $c_k \leq 1$, we have

$$\begin{aligned} & \int_{\tau_j}^t (t-s)^m u'(s) ds \\ & \geq \sum_{i=j+1}^k (t - \tau_i)^m (1 - c_i) u(\tau_i) - (t - \tau_j)^m u(\tau_j^+) \quad (52) \\ & \geq -(t - \tau_j)^m u(\tau_j^+). \end{aligned}$$

If $t \in (\tau_k, \tau_k + \sigma]$, similarly we also get

$$\int_{\tau_j}^t (t-s)^m u'(s) ds \geq -(t - \tau_j)^m u(\tau_j^+). \quad (53)$$

So, it yields

$$\int_{\tau_j}^t (t-s)^m [p(s)r(s) - A(s, \tau_{k(s)})] ds \leq (t - \tau_j)^m u(\tau_j^+), \quad (54)$$

which follows that

$$\begin{aligned} & \frac{1}{t^m} \int_{\tau_j}^t (t-s)^m [p(s)r(s) - A(s, \tau_{k(s)})] ds \\ & \leq \left(\frac{t - \tau_j}{t} \right)^m u(\tau_j^+). \end{aligned} \quad (55)$$

Hence,

$$\limsup_{t \rightarrow \infty} \frac{1}{t^m} \int_{\tau_j}^t (t-s)^m [p(s)r(s) - A(s, \tau_{k(s)})] ds \leq u(\tau_j^+), \quad (56)$$

which is a contradiction of (48).

If case (ii) holds, then as a manner with case (i) in Theorem 6, it is not possible, too. The proof is complete. $\square$

Corollary 11. Assume (19) holds and $a_k = b_k = c_k = 1$, for $k \in \mathbb{N}$. If there exists a positive differential function r such that

$$\limsup_{t \rightarrow \infty} \frac{1}{t^m} \int_T^t (t-s)^m \left[p(s)r(s) - \frac{[r'(s)]^2}{4\mu r(s)(s-T)} \right] ds = \infty, \quad (57)$$

where T is large enough such that Lemma 2 holds. Then every solution of (1) is oscillatory.

Remark 12. Corollary 11 is an extension of [11, Theorem 3.2] into impulsive case. Especially, let $r(t) \equiv 1$ in (48), it reduces to

$$\limsup_{t \rightarrow \infty} \frac{1}{t^m} \int_T^t (t-s)^m p(s) ds = \infty, \quad (58)$$

naturally, which can be considered as the extension of Kamenev-type oscillation criteria for third-order impulsive differential equations with delay (see [8, 14, 15]).

4. Examples

Example 13. Consider the third-order impulsive differential equation with delay

$$\begin{aligned} & x'''(t) + (1 + \alpha x^2(t))x(t - \sigma) = 0, \quad t > t_0, \quad t \neq \tau_k, \\ & x(\tau_k^+) = a_k x(\tau_k), \quad x'(\tau_k^+) = b_k x'(\tau_k), \\ & x''(\tau_k^+) = c_k x''(\tau_k), \quad k \in \mathbb{N}, \end{aligned} \quad (59)$$

where $\sigma > 0$, $\alpha \geq 0$ are constants, $\tau_k - \tau_{k-1} > \sigma$ for any $k \in \mathbb{N}$.

When $a_k = b_k = c_k = 1$ for any $k \in \mathbb{N}$, the impulses in (59) disappear, by [16, Theorem 4], (59) is nonoscillatory if $\sigma < e/3$ and $\alpha = 0$. However, we may change its oscillation by proper impulsive control. In fact, let $\sigma < e/3$ and $\alpha = 0$ and $\tau_k = t_0 + k\sigma$ ($k \in \mathbb{N}$) in (59); choose $\varphi(x) = x$, $p(t) \equiv 1$, and $r(t) = 1$; a simple calculation leads to

$$\begin{aligned} & \int_{t_0}^{\tau_n} \prod_{t_0 < \tau_k < s} \frac{1}{c_k} \prod_{t_0 < \tau_k + \sigma < s} a_k p(s) ds \\ &= \left[\int_{t_0}^{\tau_1} + \left(\int_{\tau_1}^{\tau_1 + \sigma} + \int_{\tau_1 + \sigma}^{\tau_2} \right) + \dots + \left(\int_{\tau_{n-1}}^{\tau_{n-1} + \sigma} + \int_{\tau_{n-1} + \sigma}^{\tau_n} \right) \right] \\ & \cdot \prod_{t_0 < \tau_k < s} \frac{1}{c_k} \prod_{t_0 < \tau_k + \sigma < s} a_k ds \\ &= (\tau_1 - t_0) + \left[\frac{\sigma}{c_1} + \frac{a_1}{c_1} (\tau_2 - \tau_1 - \sigma) \right] + \dots + \frac{a_1 a_2 \dots a_{n-2} \sigma}{c_1 c_2 \dots c_{n-1}} \\ & + \frac{a_1 a_2 \dots a_{n-1}}{c_1 c_2 \dots c_{n-1}} (\tau_n - \tau_{n-1} - \sigma) \\ &= \sigma \left(1 + \frac{1}{c_1} + \frac{1}{c_2} + \dots + \frac{1}{c_{n-1}} \right). \end{aligned} \quad (60)$$

Then, let

$$a_k = c_k = \frac{k}{k+1}, \quad b_k = 1, \quad k \in \mathbb{N}. \quad (61)$$

Obviously, (5), (6), and (19) hold, and

$$\begin{aligned} & \int_{t_0}^{\tau_n} \prod_{t_0 < \tau_k < s} \frac{1}{c_k} \prod_{t_0 < \tau_k + \sigma < s} a_k p(s) ds \\ &= \sigma \left(1 + \frac{2}{1} + \frac{3}{2} + \dots + \frac{n}{n-1} \right) \rightarrow \infty \quad (n \rightarrow \infty). \end{aligned} \quad (62)$$

Thus, (21) is also satisfied. By Theorem 6, every solution of (59) is oscillatory.

If we let

$$a_k = 1 + \frac{1}{k^2}, \quad b_k = c_k = \frac{k+1}{k}, \quad k \in \mathbb{N}. \quad (63)$$

In this case, it is easily to verify (5), (6), (15), (44), and (21) hold. By Corollary 8, every solution of (59) is either oscillatory or tends to zero.

Remark 14. It is easy to verify that in [7, Theorems 1, 2, and 3], cannot be applied to (59). On the other hand, Theorem 7 is not applicable for the condition (61) since $\sum_{n=1}^{\infty} |a_k - 1|$ does not convergence.

Example 15. Consider the third-order impulsive differential equation with delay

$$\begin{aligned} x'''(t) + e^t \cosh(|x(t)|^{\alpha-1} x(t)) x(t-1) &= 0, \\ t &\geq 0, \quad t \neq 2k, \\ x(\tau_k^+) &= a_k x(\tau_k), \quad x'(\tau_k^+) = b_k x'(\tau_k), \\ x''(\tau_k^+) &= c_k x''(\tau_k), \quad \tau_k = 2k, \quad k \in \mathbb{N}, \end{aligned} \quad (64)$$

where $\alpha > 0$, $a_k = 1$, $b_k = c_k = k/(k+1)$, $k \in \mathbb{N}$.

Let $\varphi(x) = x$, $p(t) = e^t$, it is easy to verify that (5), (6), and (19) hold. Choose $r(t) \equiv 1$ and $m = 1$, we have

$$\frac{1}{t} \int_T^t (t-s) e^s ds = \frac{e^t}{t} + \left(\frac{T-1}{t} - 1 \right) e^T \rightarrow \infty \quad (t \rightarrow \infty). \quad (65)$$

Acknowledgments

The author is very grateful to Professor H. Saker who presents the references [8–12, 17] and gives many helpful suggestions, which leads to an improvement of this paper. The work is supported in part by the NSF of Guangdong province (S2012010010034).

References

- [1] D. D. Bainov and P. S. Simeonov, *Impulsive Differential Equations: Periodic Solutions and Applications*, Longman Scientific and Technical, 1993.
- [2] G. Ballinger and X. Liu, “Existence, uniqueness and boundedness results for impulsive delay differential equations,” *Applicable Analysis*, vol. 74, no. 1-2, pp. 71–93, 2000.
- [3] V. Lakshmikantham, D. D. Bainov, and P. S. Simeonov, *Theory of Impulsive Differential Equations*, vol. 6, World Scientific Publishing, New Jersey, NJ, USA, 1989.
- [4] A. M. Samoilenko and N. A. Perestyuk, *Impulsive Differential Equations*, vol. 14, World Scientific Publishing, New Jersey, NJ, USA, 1995.
- [5] Y. S. Chen and W. Z. Feng, “Oscillations of second order nonlinear ODE with impulses,” *Journal of Mathematical Analysis and Applications*, vol. 210, no. 1, pp. 150–169, 1997.
- [6] R. P. Agarwal, F. Karakoc, and A. Zafer, “A survey on oscillation of impulsive ordinary differential equations,” *Advances in Differential Equations*, vol. 2010, pp. 1–52, 2010.
- [7] W.-H. Mao and A.-H. Wan, “Oscillatory and asymptotic behavior of solutions for nonlinear impulsive delay differential equations,” *Acta Mathematicae Applicatae Sinica*, vol. 22, English Series, no. 3, pp. 387–396, 2006.
- [8] L. Erbe, A. Peterson, and S. H. Saker, “Asymptotic behavior of solutions of a third-order nonlinear dynamic equation on time scales,” *Journal of Computational and Applied Mathematics*, vol. 181, no. 1, pp. 92–102, 2005.
- [9] L. Erbe, A. Peterson, and S. H. Saker, “Oscillation and asymptotic behavior of a third-order nonlinear dynamic equation,” *The Canadian Applied Mathematics Quarterly*, vol. 14, no. 2, pp. 124–147, 2006.

- [10] L. Erbe, A. Peterson, and S. H. Saker, "Hille and Nehari type criteria for third-order dynamic equations," *Journal of Mathematical Analysis and Applications*, vol. 329, no. 1, pp. 112–131, 2007.
- [11] S. H. Saker, "Oscillation criteria of third-order nonlinear delay differential equations," *Mathematica Slovaca*, vol. 56, no. 4, pp. 433–450, 2006.
- [12] S. H. Saker and J. Džurina, "On the oscillation of certain class of third-order nonlinear delay differential equations," *Mathematica Bohemica*, vol. 135, no. 3, pp. 225–237, 2010.
- [13] J. Yu and J. Yan, "Positive solutions and asymptotic behavior of delay differential equations with nonlinear impulses," *Journal of Mathematical Analysis and Applications*, vol. 207, no. 2, pp. 388–396, 1997.
- [14] T. Candan and R. S. Dahiya, "Oscillation of third order functional differential equations with delay," in *Proceedings of the 5th Mississippi State Conference on Differential Equations and Computational Simulations*, vol. 10, p. 7988, 2003.
- [15] I. V. Kamenev, "An integral criterion for oscillation of linear differential equations of second order," *Matematicheskie Zametki*, vol. 23, pp. 136–138, 1978.
- [16] G. Ladas, Y. G. Sficas, and I. P. Stavroulakis, "Necessary and sufficient conditions for oscillations of higher order delay differential equations," *Transactions of the American Mathematical Society*, vol. 285, no. 1, pp. 81–90, 1984.
- [17] B. Baculiková, E. M. Elabbasy, S. H. Saker, and J. Džurina, "Oscillation criteria for third-order nonlinear differential equations," *Mathematica Slovaca*, vol. 58, no. 2, pp. 201–220, 2008.