Research Article

On Modified Mellin Transform of Generalized Functions

S. K. Q. Al-Omari¹ and Adem Kılıçman²

¹ Department of Applied Sciences, Faculty of Engineering Technology, Al-Balqa Applied University, Amman 11134, Jordan

² Department of Mathematics and Institute of Mathematical Research, Universiti Putra Malaysia (UPM), 43400 Serdang, Selangor, Malaysia

Correspondence should be addressed to Adem Kılıçman; kilicman@yahoo.com

Received 10 September 2013; Accepted 14 November 2013

Academic Editor: S. A. Mohiuddine

Copyright © 2013 S. K. Q. Al-Omari and A. Kılıçman. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We investigate the modified Mellin transform on certain function space of generalized functions. We first obtain the convolution theorem for the classical and distributional modified Mellin transform. Then we describe the domain and range spaces where the extended modified transform is well defined. Consistency, convolution, analyticity, continuity, and sufficient theorems for the proposed transform have been established. An inversion formula is also obtained and many properties are given.

1. Introduction

The Mellin transform μ of a suitably restricted function over $\mathbb{R}_+((0, \infty))$ was defined on some strip in the complex plane [1], where many of the transform properties are obtained by applying change of variables to various properties of the Laplace transformation. The Mellin transform is extended to distributions in [1] and to Boehmians in [2].

By combining Fourier and Mellin transforms, the obtained transform is called Fourier-Mellin transform which has many applications in digital signals, image processing, and ship target recognition by sonar system and radar signals as well.

The modified Mellin transform of a suitably restricted function f over $\mathbb{R}_+$ was introduced by [3]

$$\mu_f^m(y) =: \mu^m f(y) =: \int_{\mathbb{R}_+} f(x) yx^{y-1} dx \quad (1)$$

as a scale-invariant transform. Then, as earlier, combining the modified Mellin transform with the Fourier transform gives the Fourier-modified Mellin transform [3, Equation 16].

The Mellin-type convolution product of two functions f and g is given by

$$(f \vee g)(x) = \int_{\mathbb{R}_+} f(\tau) g\left(\frac{x}{\tau}\right) \tau^{-1} d\tau. \quad (2)$$

From [1] it has been noted that

$$\mu(f \vee g)(y) = \mu_f(y) \mu_g(y). \quad (3)$$

Utilizing the Mellin-type convolution product the following theorem is essential for our next investigations.

Theorem 1 (convolution theorem). *Let $\mathcal{L}^1$ be the Lebesgue space of integrable functions and $f, g \in \mathcal{L}^1$; then*

$$\mu^m(f \vee g)(y) = \mu_f^m(y) \mu_g(y), \quad (4)$$

where μ_f^m and μ_g are the Mellin-type and Mellin transforms of f and g , respectively.

Proof. By the definition of the Mellin and modified Mellin transforms we have

$$\begin{aligned} \mu^m(f \vee g)(y) &= \int_{\mathbb{R}_+} (f \vee g)(x) yx^{y-1} dx \\ &= \int_{\mathbb{R}_+} \left(\int_{\mathbb{R}_+} f(\tau) g\left(\frac{x}{\tau}\right) \tau^{-1} d\tau \right) yx^{y-1} dx \\ &= \int_{\mathbb{R}_+} yf(\tau) \tau^{-1} \left(\int_{\mathbb{R}_+} g\left(\frac{x}{\tau}\right) x^{y-1} dx \right) d\tau. \end{aligned} \quad (5)$$

Employing Fubini's theorem, then the substitution $z = x/\tau$ together with simple computations establishes that

$$\mu^m(f \vee g)(y) = \mu_f^m(y) \mu_g(y). \quad (6)$$

Hence the theorem is proved. $\square$

2. Modified Mellin Transform of Distribution

Let $\mu_{a,b}$ be the space of smooth functions φ over $\mathbb{R}_+$ such that [1]

$$\gamma_k(\varphi) = \sup_{x \in \mathbb{R}_+} \left| \xi_{a,b}(x) x^{k-1} \frac{d^k}{dx^k} \varphi(x) \right| \quad (7)$$

is finite, $k = 0, 1, 2, \dots$, where

$$\xi_{a,b}(x) \triangleq \begin{cases} x^{-a}, & \text{for } 0 < x \leq 1, \\ x^{-b}, & \text{for } 1 < x < \infty, \end{cases} \quad (8)$$

and $a, b \in \mathbb{R}_+$.

Then $\mu_{a,b}$ is linear space under addition and multiplication by complex numbers. $\mu_{a,b}$ can also be generated by the multinorms $(\gamma_k)_{k=0}^\infty$ which turns to be a countably multinormed space.

Denote by $\dot{\mu}_{a,b}$ the complete strong dual space of $\mu_{a,b}$; then it is assigned the weak topology. Let $\xi(\mathbb{R}_+)$ be the space of smooth functions over $\mathbb{R}_+$; then for any $a, b \in \mathbb{R}$, $\mu_{a,b}$ is dense in $\xi(\mathbb{R}_+)$ and the topology assigned to $\mu_{a,b}$ is stronger than that induced on $\mu_{a,b}$ by $\xi(\mathbb{R}_+)$ and is identified with a subspace of $\dot{\mu}_{a,b}$.

The straightforward conclusion is that the kernel function (yx^{y-1}) of μ_f^m is a member of $\mu_{a,b}$ for $a \leq \operatorname{Re} y \leq b$.

This usually leads to the following definition: let $f \in \dot{\mu}_{a,b}$; then the distributional transform $\widehat{\mu_f^m}$ of f is defined as

$$\widehat{\mu_f^m}(y) \triangleq \langle f(x), yx^{y-1} \rangle, \quad y \in \Omega_f, \quad (9)$$

where $f \in \dot{\mu}_{a,b}$ and $\Omega_f = \{y \in \mathbb{C} : a < \operatorname{Re} y < b\}$.

Theorem 2 (analyticity theorem). *Let $f \in \dot{\mu}_{a,b}$; then $\widehat{\mu_f^m}$ is analytic and*

$$\frac{d^k}{dy^k} \widehat{\mu_f^m}(y) = \langle f(x), yx^{y-1} \ln^k x + kx^{y-1} \ln x^{k-1} \rangle, \quad (10)$$

where k is nonnegative integer and $y \in \Omega_f$.

It is easy for reader to see that $\widehat{\mu_f^m}$ is injective and linear from $\dot{\mu}_{a,b}$ into $\dot{\mu}_{a,b}$.

The Mellin-type convolution product of $f, g \in \dot{\mu}_{a,b}$ is given as

$$\langle f \vee g, \varphi \rangle \triangleq \langle f(x), \langle g(t), \varphi(tx) \rangle \rangle, \quad (11)$$

where $\varphi \in \mu_{a,b}$.

From (11) it is clear that $f \vee g$ is a member of $\dot{\mu}_{a,b}$, for $a \leq b$.

Therefore, denote by $\mathcal{D}(\mathbb{R}_+)$ the space of test functions of bounded support over $\mathbb{R}_+$; then the convolution product of $f \in \dot{\mu}_{a,b}$ and $g \in \mathcal{D}(\mathbb{R}_+)$ can be given as

$$(f \vee g)(x) = \left\langle f(t), \frac{1}{t} g\left(\frac{x}{t}\right) \right\rangle, \quad (12)$$

where $x \in \mathbb{R}_+$.

3. Boehmians

Let $\mathcal{G}$ a group and $\mathcal{S}$ a subgroup of $\mathcal{G}$. We assume that to each pair of elements $f \in \mathcal{G}$ and $\omega \in \mathcal{S}$ is assigned the product $f * g$ such that

- (1) if $\omega, \psi \in \mathcal{S}$, then $\omega * \psi \in \mathcal{S}$ and $\omega * \psi = \psi * \omega$;
- (2) if $f \in \mathcal{G}$ and $\omega, \psi \in \mathcal{S}$, then $(f * \omega) * \psi = f * (\omega * \psi)$;
- (3) if $f, g \in \mathcal{G}$, $\omega \in \mathcal{S}$, and $\lambda \in \mathbb{R}$, then

$$(f + g) * \omega = f * \omega + g * \omega, \quad \lambda(f * \omega) = (\lambda f) * \omega. \quad (13)$$

Let Δ be a family of sequences from $\mathcal{S}$ such that

- (1) if $f, g \in \mathcal{G}$, $(\delta_n) \in \Delta$, and $f * \delta_n = g * \delta_n$ ($n = 1, 2, \dots$), then $f = g$, for all n ;
- (2) if $(\omega_n), (\delta_n) \in \Delta$, then $(\omega_n * \psi_n) \in \Delta$.

Elements of Δ will be called delta sequences.

Consider the class $\mathcal{A}$ of pair of sequences defined by

$$\mathcal{A} = \{((f_n), (\omega_n)) : (f_n) \in \mathcal{G}^\mathbb{N}, (\omega_n) \in \Delta\}, \quad (14)$$

for each $n \in \mathbb{N}$.

An element $((f_n), (\omega_n)) \in \mathcal{A}$ is called a quotient of sequences, denoted by $[f_n/\omega_n]$, if $f_n * \omega_m = f_m * \omega_n$, for all $n, m \in \mathbb{N}$.

Two quotients of sequences f_n/ω_n and g_n/ψ_n are said to be equivalent, $f_n/\omega_n \sim g_n/\psi_n$, if $f_n * \psi_m = g_m * \omega_n$, for all $n, m \in \mathbb{N}$.

The relation $\sim$ is an equivalent relation on $\mathcal{A}$. The equivalence class containing f_n/ω_n is denoted by $[f_n/\omega_n]$. These equivalence classes are called Boehmians. The space of all Boehmians is denoted by β_1 .

The sum of two Boehmians and multiplication by a scalar can be defined in a natural way

$$\begin{aligned} \left[\frac{f_n}{\omega_n} \right] + \left[\frac{g_n}{\psi_n} \right] &= \left[\frac{f_n * \psi_n + g_n * \omega_n}{\omega_n * \psi_n} \right], \\ \alpha \left[\frac{f_n}{\omega_n} \right] &= \left[\frac{\alpha f_n}{\omega_n} \right] = \left[\frac{\alpha f_n}{\omega_n} \right], \end{aligned} \quad (15)$$

$\alpha \in \mathbb{C}$, space of complex numbers.

The operation $*$ and the differentiation are defined by $[f_n/\omega_n] * [g_n/\psi_n] = [(f_n * g_n)/(\omega_n * \psi_n)]$ and $\mathcal{D}^\alpha[f_n/\omega_n] = [\mathcal{D}^\alpha f_n/\omega_n]$.

The operation $*$ can be extended to $\beta \times \mathcal{S}$ as follows. If $[f_n/\omega_n] \in \beta_1$ and $\omega \in \mathcal{S}$, then

$$\left[\frac{f_n}{\omega_n} \right] * \omega = \left[\frac{f_n * \omega}{\omega_n} \right]. \quad (16)$$

δ -Convergence. A sequence of Boehmians (β_n) in β_1 is said to be δ convergent to a Boehmian β in β_1 , denoted by $\beta_n \xrightarrow{\delta} \beta$, if there exists a delta sequence (ω_n) such that

$$\begin{aligned} (\beta_n * \omega_n), (\beta * \omega_n) &\in \mathcal{E}, \quad \forall k, n \in \mathbb{N}, \\ (\beta_n * \omega_k) &\longrightarrow (\beta * \omega_k) \quad \text{as } n \longrightarrow \infty, \text{ in } \mathcal{E}, \quad (17) \\ &\text{for every } k \in \mathbb{N}. \end{aligned}$$

The following is equivalent for the statement of δ -convergence: $\beta_n \xrightarrow{\delta} \beta$ ($n \rightarrow \infty$) in β_1 if and only if there is $f_{n,k}, f_k \in \mathcal{E}$ and $(\omega_k) \in \Delta$ such that $\beta_n = [f_{n,k}/\omega_k]$, $\beta = [f_k/\omega_k]$ and for each $k \in \mathbb{N}$, $f_{n,k} \rightarrow f_k$ as $n \rightarrow \infty$ in $\mathcal{E}$.

A sequence of Boehmians (β_n) in β_1 is said to be Δ convergent to a Boehmian β in β_1 , denoted by $\beta_n \xrightarrow{\Delta} \beta$, if there exists a $(\omega_n) \in \Delta$ such that $(\beta_n - \beta) * \omega_n \in \beta_1$, for all $n \in \mathbb{N}$, and $(\beta_n - \beta) * \omega_n \rightarrow 0$ as $n \rightarrow \infty$ in β_1 . See [2, 5–15].

4. Modified Mellin Transform of Boehmian

In this section we discuss the modified Mellin transform on spaces of Boehmians. Consider the group $\dot{\mu}_{a,b}(\mathbb{R}_+)$ and $\mathcal{V}(\mathbb{R}_+)$ as a subgroup of $\dot{\mu}_{a,b}(\mathbb{R}_+)$. Let $\vee$ be the operation between $\dot{\mu}_{a,b}(\mathbb{R}_+)$ and $\mathcal{V}(\mathbb{R}_+)$ and Δ the set of delta sequences given by [2]

- (1) $\int_{\mathbb{R}_+} \varphi_n(t) dt = 1$, for all $n \in \mathbb{N}$;
- (2) $\int_{\mathbb{R}_+} |\varphi_n(t)| dt \leq m$, for all $n \in \mathbb{N}$, for some $m > 0$;
- (3) $\text{supp } \varphi_n \subset (a_n, b_n)$, for all $n \in \mathbb{N}$ for some $0 < a_n < b_n < \infty$ with $a_n \rightarrow 1, b_n \rightarrow 1$ as $n \rightarrow \infty$.

Let β_1 be the Boehmian space obtained from $\dot{\mu}_{a,b}, \mathcal{V}(\mathbb{R}_+)$ and Δ ; then β_1 will serve as the domain space of $\hat{\mu}^m$.

Our next objective is to construct a range space, say β_2 , for $\hat{\mu}^m$.
Let

$$\begin{aligned} \Delta_{\mathbb{R}_+}^e &= \{\mu_{\varphi_n} : (\varphi_n) \in \Delta\}, \quad \mathcal{V}^e = \{\mu_{\varphi} : \varphi \in \mathcal{V}\}, \\ \ell^m &= \{\widehat{\mu}_f^m : f \in \dot{\mu}_{a,b}\}. \end{aligned} \quad (18)$$

For $g \in \ell^m, \psi \in \mathcal{V}^e$, define

$$(g \wedge \psi)(y) = g(y) \psi(y). \quad (19)$$

We have the following theorem.

Theorem 3. Let $g \in \ell^m$ and $\psi \in \mathcal{V}^e$, then $g \wedge \psi \in \ell^m$.

Proof. Let g and ψ belong to ℓ^m and $\mathcal{V}^e$, respectively. Then there are $f \in \dot{\mu}_{a,b}$ and $\varphi \in \mathcal{V}$ such that $g = \hat{\mu}_f^m$ and $\psi = \mu_{\varphi}$,

respectively. Therefore, by the convolution theorem and (19) we get that

$$\begin{aligned} (g \wedge \psi)(y) &= g(y) \psi(y) \\ &= (\widehat{\mu}_f^m \mu_{\varphi})(y) \\ &= \mu^m(f \vee \varphi)(y). \end{aligned} \quad (20)$$

Since $f \vee \varphi \in \dot{\mu}_{a,b}$, it follows $g \wedge \psi \in \ell^m$.

Hence we have proved the theorem. $\square$

Theorem 4. Let $\psi_1, \psi_2 \in \mathcal{V}^e$; then $\psi_1 \wedge \psi_2 \in \mathcal{V}^e$.

Proof. By definition of $\mathcal{V}^e$ we can find $\varphi_1, \varphi_2 \in \mathcal{V}$ such that $\psi_1 = \mu_{\varphi_1}$ and $\psi_2 = \mu_{\varphi_2}$.

Therefore, by [1],

$$\begin{aligned} \psi_1 \wedge \psi_2 &= \psi_1(y) \psi_2(y) \\ &= \mu_{\varphi_1}(y) \mu_{\varphi_2}(y) \\ &= \mu(\varphi_1 \vee \varphi_2)(y). \end{aligned} \quad (21)$$

But since $\varphi_1 \vee \varphi_2 \in \mathcal{V}$, we get $\psi_1 \wedge \psi_2 \in \mathcal{V}^e$. Thus we have the theorem. $\square$

Theorem 5. Let $g_1, g_2 \in \ell^m$ and $\psi \in \mathcal{V}^e$; then $(g_1 + g_2) \wedge \psi = g_1 \wedge \psi + g_2 \wedge \psi$ and $(\alpha g) \wedge \psi = g \wedge (\alpha \psi) = \alpha(g \wedge \psi)$.

Proof. Is straightforward. $\square$

Theorem 6. Let $g_n \rightarrow g$ in $\ell^m, \psi \in \mathcal{V}^e$; then $g_n \wedge \psi \rightarrow g \wedge \psi$ as $n \rightarrow \infty$ in ℓ^m .

Proof. Can easily be checked. $\square$

Theorem 7. Let $g \in \ell^m$ and $(\psi_n) \in \Delta_{\mathbb{R}_+}^e$; then $g \wedge \psi_n \rightarrow g$ as $n \rightarrow \infty$.

Proof. By (19) we have

$$(g \wedge \psi_n)(y) = (\widehat{\mu}_f^m \mu_{\varphi_n})(y), \quad (22)$$

where $f \in \dot{\mu}_{a,b}$ and $(\varphi_n) \in \Delta$ are such that $\widehat{\mu}_f^m = g$ and $\mu_{\varphi_n} = \psi_n$, for all $n \in \mathbb{N}$.

Since $\mu_{\varphi_n}(y) \rightarrow 1$ as $n \rightarrow \infty$ on compact subsets of $\mathbb{R}_+$, (22) implies that $(g \wedge \psi_n)(y) \rightarrow \widehat{\mu}_f^m(y) = g(y)$, for all y , as $n \rightarrow \infty$. Hence we obtain the theorem. $\square$

Theorem 8. Let $(\psi_n), (\theta_n) \in \Delta_{\mathbb{R}_+}^e$; then $\psi_n \wedge \theta_n \in \Delta_{\mathbb{R}_+}^e$.

Let $\psi_n = \mu_{\alpha_n}, \theta_n = \mu_{\sigma_n}$; then taking into account the fact that $(\psi_n \wedge \theta_n)(y) = (\mu_{\alpha_n} \wedge \mu_{\sigma_n})(y) = \mu(\alpha_n \vee \sigma_n)(y)$, since $\alpha_n \vee \sigma_n \in \Delta$, this theorem follows.

The Boehmian space β_2 is therefore constructed.

In addition, scalar multiplication, differentiation, and the operation $\wedge$ in β_2 are defined similar to that of usual Boehmian spaces.

Each $g \in \ell^m$ can be identified by a member of β_2 given as

$$g \longrightarrow \left[\frac{g \wedge \psi_n}{\psi_n} \right] \quad \text{as } n \longrightarrow \infty, \quad (23)$$

where $\psi_n \in \Delta_{\mathbb{R}_+}^e$.

Definition 9. The extended modified Mellin transform $\tilde{\mu}^m : \beta_1 \rightarrow \beta_2$ is defined by

$$\tilde{\mu}^m(\beta) = \left[\frac{\widehat{\mu}_{f_n}^m}{\mu_{\varphi_n}} \right], \quad \forall \beta = \left[\frac{f_n}{\varphi_n} \right] \in \beta_1. \quad (24)$$

Theorem 10. The extended modified Mellin transform is well defined.

Proof. The proof of this theorem is straightforward. See [11–13]. $\square$

Theorem 11 (consistency theorem). The extended modified Mellin transform $\tilde{\mu}^m$ is consistent with the distributional $\widehat{\mu}_f^m (\widehat{\mu}_f^m : \dot{\mu}_{a,b} \rightarrow \dot{\mu}_{a,b})$.

Proof. For every $f \in \dot{\mu}_{a,b}$, let β be its representative in β_1 ; then $\beta = [(f \vee \varphi_n)/\varphi_n]$, where $\varphi_n \in \Delta$, for all n . Then it is clear that φ_n is independent of the representative, for all n .

Therefore

$$\begin{aligned} \tilde{\mu}^m(\beta) &= \tilde{\mu}^m \left(\left[\frac{f \vee \varphi_n}{\varphi_n} \right] \right) \\ \text{i.e.} &= \left[\frac{\widehat{\mu}_{f \vee \varphi_n}^m}{\mu_{\varphi_n}} \right] \\ \text{i.e.} &= \left[\frac{(\widehat{\mu}_f^m) \mu_{\varphi_n}}{\mu_{\varphi_n}} \right] \end{aligned} \quad (25)$$

which is the representative of $\widehat{\mu}_f^m$ in $\dot{\mu}_{a,b}$.

Hence we have the proof. $\square$

Theorem 12 (necessity theorem). Let $[g_n/\psi_n] \in \beta_2$; then the necessary and sufficient condition that $[g_n/\psi_n]$ is to be in the range of $\tilde{\mu}^m$ is that g_n belongs to range of $\widehat{\mu}_f^m$ for every $n \in \mathbb{N}$.

Proof. Let $[g_n/\psi_n]$ be in the range of $\tilde{\mu}^m$; then of course g_n belongs to the range of $(\widehat{\mu}_f^m)$, for all $n \in \mathbb{N}$.

To establish the converse, let g_n be in the range of $(\widehat{\mu}_f^m)$, for all $n \in \mathbb{N}$. Then there is $f_n \in \dot{\mu}_{a,b}$ such that $\widehat{\mu}_{f_n}^m = g_n$, $n \in \mathbb{N}$.

Since $[g_n/\psi_n] \in \beta_2$,

$$g_n \vee \psi_m = g_m \vee \psi_n, \quad (26)$$

for all $m, n \in \mathbb{N}$. Therefore,

$$\mu^m(f_n \wedge \varphi_n) = \mu^m(f_m \wedge \varphi_n), \quad \forall m, n \in \mathbb{N}, \quad (27)$$

where $f_n \in \dot{\mu}_{a,b}$ and $\varphi_n \in \Delta$.

The fact that $\widehat{\mu}_f^m$ is injective, $\widehat{\mu}_f^m : \dot{\mu}_{a,b} \rightarrow \dot{\mu}_{a,b}$, implies that $f_n \wedge \varphi_m = f_m \wedge \varphi_n$, $m, n \in \mathbb{N}$.

Thus f_n/φ_n is quotient of sequences in β_1 . Hence, $[f_n/\varphi_n] \in \beta_1$ and

$$\tilde{\mu}^m \left(\left[\frac{f_n}{\varphi_n} \right] \right) = \left[\frac{g_n}{\psi_n} \right]. \quad (28)$$

Hence the theorem is proved. $\square$

Theorem 13 (generalized convolution theorem). Let $\beta = [f_n/\varphi_n] \in \beta_1$ and $\gamma = [\kappa_n/\phi_n] \in \beta_1$; then

$$\tilde{\mu}^m(\beta \vee \gamma) = \tilde{\mu}^m \left(\left[\frac{f_n}{\varphi_n} \right] \right) \wedge \tilde{\mu}^m \left(\left[\frac{\kappa_n}{\phi_n} \right] \right). \quad (29)$$

Proof. Assume that the requirements of the theorem satisfy for some β and $\gamma \in \beta_1$; then using Definition 9 and the operation $\wedge$ yields

$$\begin{aligned} \tilde{\mu}^m(\beta \vee \gamma) &= \tilde{\mu}^m \left(\left[\frac{f_n \vee \kappa_n}{\varphi_n \vee \phi_n} \right] \right) = \left[\frac{\mu^m(f_n \vee \kappa_n)}{\mu(\varphi_n \vee \phi_n)} \right] \\ &= \left[\frac{\widehat{\mu}_{f_n}^m \wedge \mu_{\kappa_n}^m}{\mu_{\varphi_n} \wedge \mu_{\phi_n}} \right] = \left[\frac{\widehat{\mu}_{f_n}^m}{\mu_{\varphi_n}} \right] \wedge \left[\frac{\mu_{\kappa_n}^m}{\mu_{\phi_n}} \right]. \end{aligned} \quad (30)$$

Therefore

$$\tilde{\mu}^m(\beta \vee \gamma) = \tilde{\mu}^m \left(\left[\frac{f_n}{\varphi_n} \right] \right) \wedge \tilde{\mu}^m \left(\left[\frac{\kappa_n}{\phi_n} \right] \right). \quad (31)$$

This completes the proof. $\square$

Theorem 14. The extended modified Mellin transform $\tilde{\mu}^m : \beta_1 \rightarrow \beta_2$ is bijective.

Proof. Assume $\tilde{\mu}^m [f_n/\varphi_n] = \tilde{\mu}^m [\kappa_n/\phi_n]$; then it follows from the concept of quotients of sequences that $\widehat{\mu}_{f_n}^m \wedge \mu_{\phi_n}^m = \widehat{\mu}_{\kappa_n}^m \wedge \mu_{\varphi_n}^m$. Therefore, $\mu^m(f_n \vee \phi_m) = \mu^m(\kappa_m \vee \varphi_n)$. The property that μ^m is one to one implies $f_n \vee \phi_m = \kappa_m \vee \varphi_n$. Therefore,

$$\left[\frac{f_n}{\varphi_n} \right] = \left[\frac{\kappa_n}{\phi_n} \right]. \quad (32)$$

Next to establish that $\tilde{\mu}^m$ is onto, let $[\widehat{\mu}_{f_n}^m/\mu_{\varphi_n}] (\in \beta_2)$ be arbitrary; then $\widehat{\mu}_{f_n}^m \wedge \mu_{\varphi_m}^m = \widehat{\mu}_{f_m}^m \wedge \mu_{\varphi_n}^m$ for every $m, n \in \mathbb{N}$. Hence $f_n, f_m \in \dot{\mu}_{a,b}$ are such that $\mu^m(f_n \vee \varphi_m) = \mu^m(f_m \vee \varphi_n)$, for all $m, n \in \mathbb{N}$.

Hence, the Boehmian $[f_n/\varphi_n]$ belongs to β_1 and satisfies

$$\tilde{\mu}^m \left[\frac{f_n}{\varphi_n} \right] = \left[\frac{\widehat{\mu}_{f_n}^m}{\mu_{\varphi_n}} \right]. \quad (33)$$

This completes the proof of the theorem. $\square$

Now we introduce $(\widehat{\mu}^m)^{-1}$ as the inverse transform of $\widehat{\mu}^m$, where

$$\left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right]\right)=\left[\frac{\left(\widehat{\mu}_{f_n}^m\right)^{-1}\left(\widehat{\mu}_{f_n}^m\right)}{\left(\widehat{\mu}_{\varphi_n}\right)^{-1}\left(\widehat{\mu}_{\varphi_n}\right)}\right], \quad (34)$$

for every $[f_n/\varphi_n] \in \beta_1$.

Theorem 15. Let $[\widehat{\mu}_{f_n}^m/\mu_{\varphi_n}] \in \beta_2$ and $\phi \in \mathcal{G}^e$, then

$$\left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right] \wedge \phi\right)=\left[\frac{f_n}{\varphi_n}\right] \vee \phi, \quad (35)$$

$$\widehat{\mu}^m\left(\left[\frac{f_n}{\varphi_n}\right] \vee \phi\right)=\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right] \wedge \phi. \quad (36)$$

Proof. We prove (35) and omit the proof of (36) due to its similarity. Given $[\widehat{\mu}_{f_n}^m/\mu_{\varphi_n}] \in \beta_2$ and $\psi \in \mathcal{G}$ such that $\phi = \mu_{\psi}^m$ then employing (34) yields

$$\begin{aligned} \left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right] \wedge \phi\right) &= \left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m \wedge \phi}{\widehat{\mu}_{\varphi_n}}\right]\right) \\ &= \left[\frac{\left(\mu_{f_n}^m\right)^{-1}\left(\widehat{\mu}_{f_n}^m \wedge \mu_{\psi}^m\right)}{\left(\mu_{\varphi_n}\right)^{-1}\left(\mu_{\varphi_n}\right)}\right]. \end{aligned} \quad (37)$$

Using (19) gives

$$\left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right] \wedge \phi\right)=\left[\frac{\left(\mu_{f_n}^m\right)^{-1}\left(\widehat{\mu}_{f_n}^m \mu_{\psi}^m\right)}{\left(\mu_{\varphi_n}\right)^{-1}\left(\mu_{\varphi_n}\right)}\right]. \quad (38)$$

Hence the convolution theorem gives

$$\left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right] \wedge \phi\right)=\left[\frac{\left(\mu_{f_n}^m\right)^{-1}\left(\mu_{f_n}^m\left(f_n \vee \phi\right)\right)}{\left(\mu_{\varphi_n}\right)^{-1}\left(\mu_{\varphi_n}\right)}\right]. \quad (39)$$

Thus

$$\left(\widehat{\mu}^m\right)^{-1}\left(\left[\frac{\widehat{\mu}_{f_n}^m}{\widehat{\mu}_{\varphi_n}}\right] \wedge \phi\right)=\left[\frac{f_n}{\varphi_n}\right] \vee \phi. \quad (40)$$

Proof of the second part is similar.

This completes the proof of the theorem. $\square$

Theorem 16. $\widehat{\mu}^m : \beta_1 \rightarrow \beta_2$ and $(\widehat{\mu}^m)^{-1} : \beta_2 \rightarrow \beta_1$ are continuous with respect to δ -convergence.

Proof. Let $\beta_n \xrightarrow{\delta} \beta$ in β_1 as $n \rightarrow \infty$; then we establish that $\widehat{\mu}^m \beta_n \rightarrow \widehat{\mu}^m \beta$ as $n \rightarrow \infty$. Let $f_{n,k}$ and f_k be in $\mu_{a,b}$ such that

$$\beta_n = \left[\frac{f_{n,k}}{\varphi_k}\right], \quad \beta = \left[\frac{f_k}{\varphi_k}\right] \quad (41)$$

and $f_{n,k} \rightarrow f_k$ as $n \rightarrow \infty$ for every $k \in \mathbb{N}$.

The continuity of $\widehat{\mu}_{f_{n,k}}^m$ implies $\widehat{\mu}_{f_{n,k}}^m \rightarrow \widehat{\mu}_{f_k}^m$ as $n \rightarrow \infty$. Thus,

$$\left[\frac{\widehat{\mu}_{f_{n,k}}^m}{\widehat{\mu}_{\varphi_k}}\right] \rightarrow \left[\frac{\widehat{\mu}_{f_k}^m}{\widehat{\mu}_{\varphi_k}}\right] \quad (42)$$

as $n \rightarrow \infty$ in β_2 . This proves continuity of $\widehat{\mu}^m$.

Next, let $g_n \xrightarrow{\delta} g \in \beta_2$ as $n \rightarrow \infty$; then we have $g_n = [\widehat{\mu}_{f_{n,k}}^m/\mu_{\varphi_k}]$ and $g = [\widehat{\mu}_{f_k}^m/\mu_{\varphi_k}]$ for some $\widehat{\mu}_{f_{n,k}}^m, \widehat{\mu}_{f_k}^m \in \ell^m$, where $\widehat{\mu}_{f_{n,k}}^m \rightarrow \widehat{\mu}_{f_k}^m$ as $n \rightarrow \infty$. Hence

$$\left(\widehat{\mu}_{f_{n,k}}^m\right)^{-1}\left(\widehat{\mu}_{f_{n,k}}^m\right) \rightarrow \left(\widehat{\mu}_{f_k}^m\right)^{-1}\left(\widehat{\mu}_{f_k}^m\right) \quad (43)$$

as $n \rightarrow \infty$ in β_1 . That is,

$$\begin{aligned} \left[\frac{\left(\widehat{\mu}_{f_{n,k}}^m\right)^{-1}\left(\widehat{\mu}_{f_{n,k}}^m\right)}{\varphi_k}\right] &= \left[\frac{\left(\widehat{\mu}_{f_{n,k}}^m\right)^{-1}\left(\widehat{\mu}_{f_{n,k}}^m\right)}{\left(\mu_{\varphi_k}\right)^{-1}\left(\mu_{\varphi_k}\right)}\right] \\ &\rightarrow \left[\frac{\left(\widehat{\mu}_{f_k}^m\right)^{-1}\left(\widehat{\mu}_{f_k}^m\right)}{\varphi_k}\right] \end{aligned} \quad (44)$$

as $n \rightarrow \infty$.

Hence

$$\left[\frac{\left(\widehat{\mu}_{f_{n,k}}^m\right)^{-1}\left(\widehat{\mu}_{f_{n,k}}^m\right)}{\varphi_k}\right] \rightarrow \left[\frac{\left(\widehat{\mu}_{f_k}^m\right)^{-1}\left(\widehat{\mu}_{f_k}^m\right)}{\left(\mu_{\varphi_k}\right)^{-1}\left(\mu_{\varphi_k}\right)}\right] \quad (45)$$

as $n \rightarrow \infty$.

That is,

$$\left(\widehat{\mu}^m\right)^{-1} g_n \rightarrow \left(\widehat{\mu}^m\right)^{-1} g \quad (46)$$

as $n \rightarrow \infty$. This completes the proof. $\square$

Theorem 17. $\widehat{\mu}^m : \beta_1 \rightarrow \beta_2$ and $(\widehat{\mu}^m)^{-1} : \beta_2 \rightarrow \beta_1$ are continuous with respect to Δ -convergence.

Proof. Let $\beta_n \xrightarrow{\Delta} \beta$ in β_1 as $n \rightarrow \infty$. Then, we find $f_n \in \mu_{a,b}$ and $(\varphi_k) \in \Delta$ such that $(\beta_n - \beta) \wedge \varphi_k = [(f_n \wedge \varphi_k)/\varphi_k]$ and $f_n \rightarrow 0$ as $n \rightarrow \infty$. Therefore

$$\widehat{\mu}^m((\beta_n - \beta) \vee \varphi_k) = \left[\frac{\mu^m(f_n \vee \varphi_k)}{\mu_{\varphi_k}}\right]. \quad (47)$$

Hence, $\widehat{\mu}^m((\beta_n - \beta) \vee \varphi_k) = [(\widehat{\mu}_{f_n}^m \wedge \mu_{\varphi_k})/\mu_{\varphi_k}] = \widehat{\mu}_{f_n}^m \rightarrow 0$ as $n \rightarrow \infty$ in ℓ^m .

Therefore

$$\begin{aligned} \widehat{\mu}^m((\beta_n - \beta) \vee \varphi_n) &= \left(\widehat{\mu}^m \beta_n - \widehat{\mu}^m \beta\right) \wedge \mu_{\varphi_k} \rightarrow \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (48)$$

Hence, $\widehat{\mu}^m \beta_n \xrightarrow{\Delta} \widehat{\mu}^m \beta$ as $n \rightarrow \infty$.

Proof of the second part is analogous. Detailed proof is as follows.

Finally, let $g_n \xrightarrow{\Delta} g$ in β_2 as $n \rightarrow \infty$; then we can find $\widehat{\mu_{f_k}^m} \in \ell^m$ such that $(g_n - g) \wedge \mu_{\varphi_k} = [(\widehat{\mu_{f_k}^m} \wedge \mu_{\varphi_k}) / \mu_{\varphi_k}]$ and $\widehat{\mu_{f_k}^m} \rightarrow 0$ as $n \rightarrow \infty$ for some $(\mu_{\varphi_k}) \in \Delta_{\mathbb{R}_+}^e$.

Next, we have

$$\left(\widehat{\mu^m}\right)^{-1} ((g_n - g) \wedge \mu_{\varphi_k}) = \left[\frac{(\mu_{f_k}^m)^{-1} (\widehat{\mu_{f_k}^m} \wedge \mu_{\varphi_k})}{(\mu_{\varphi_k})^{-1} (\mu_{\varphi_k})} \right]. \quad (49)$$

Thus, by (34) we get

$$\left(\widehat{\mu^m}\right)^{-1} ((g_n - g) \wedge \mu_{\varphi_k}) = \left[\frac{f_n \vee \varphi_k}{\varphi_k} \right] = f_n \rightarrow 0 \quad (50)$$

as $n \rightarrow \infty$ in $\dot{\mu}_{a,b}$.

Therefore

$$\begin{aligned} & \left(\widehat{\mu^m}\right)^{-1} ((g_n - g) \wedge \mu_{\varphi_k}) \\ &= \left(\left(\widehat{\mu^m}\right)^{-1} g_n - \left(\widehat{\mu^m}\right)^{-1} g \right) \vee \varphi_k \rightarrow 0 \end{aligned} \quad (51)$$

as $n \rightarrow \infty$.

Thus, we have

$$\left(\widehat{\mu^m}\right)^{-1} g_n \xrightarrow{\Delta} \left(\widehat{\mu^m}\right)^{-1} g \quad (52)$$

as $n \rightarrow \infty$ in β_1 .

This completes the proof of the theorem. $\square$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

The authors acknowledge that this research was partially supported by Universiti Putra Malaysia under the ERGS 1-2013/5527179.

References

- [1] A. H. Zemanian, *Generalized Integral Transformations*, Dover, New York, NY, USA, 1987.
- [2] S. K. Q. Al-Omari, "On the distributional Mellin transformation and its extension to Boehmian spaces," *International Journal of Contemporary Mathematical Sciences*, vol. 6, no. 17–20, pp. 801–810, 2011.
- [3] J. Yang, T. K. Sarkar, and P. Antonik, "Applying the Fourier-modified Mellin transform to Doppler-distorted waveforms," *Digital Signal Processing*, vol. 17, no. 6, pp. 1030–1039, 2007.
- [4] S. K. Q. Al-Omari, "A Mellin transform for a space of Lebesgue integrable Boehmians," *International Journal of Contemporary Mathematical Sciences*, vol. 6, no. 29–32, pp. 1597–1606, 2011.
- [5] S. K. Q. Al-Omari, "Distributional and tempered distributional diffraction Fresnel transforms and their extension to Boehmian spaces," *Italian Journal of Pure and Applied Mathematics*, no. 30, pp. 179–194, 2013.
- [6] P. K. Banerji, S. K. Al-Omari, and L. Debnath, "Tempered distributional Fourier sine (cosine) transform," *Integral Transforms and Special Functions*, vol. 17, no. 11, pp. 759–768, 2006.
- [7] P. Mikusiński, "Fourier transform for integrable Boehmians," *The Rocky Mountain Journal of Mathematics*, vol. 17, no. 3, pp. 577–582, 1987.
- [8] P. Mikusiński, "Tempered Boehmians and ultradistributions," *Proceedings of the American Mathematical Society*, vol. 123, no. 3, pp. 813–817, 1995.
- [9] P. Mikusiński, "Convergence of Boehmians," *Japanese Journal of Mathematics*, vol. 9, no. 1, pp. 159–179, 1983.
- [10] S. K. Q. Al-Omari, D. Loonker, P. K. Banerji, and S. L. Kalla, "Fourier sine (cosine) transform for ultradistributions and their extensions to tempered and ultraBoehmian spaces," *Integral Transforms and Special Functions*, vol. 19, no. 5–6, pp. 453–462, 2008.
- [11] S. K. Q. Al-Omari, "Hartley transforms on a certain space of generalized functions," *Georgian Mathematical Journal*, vol. 20, no. 3, pp. 415–426, 2013.
- [12] S. K. Q. Al-Omari and A. Kılıçman, "Note on Boehmians for class of optical Fresnel wavelet transforms," *Journal of Function Spaces and Applications*, vol. 2012, Article ID 405368, 14 pages, 2012.
- [13] S. K. Q. Al-Omari and A. Kılıçman, "On generalized hartley-Hilbert and Fourier-Hilbert transforms," *Advances in Difference Equations*, vol. 2012, article 232, 12 pages, 2012.
- [14] T. K. Boehme, "The support of Mikusiński operators," *Transactions of the American Mathematical Society*, vol. 176, pp. 319–334, 1973.
- [15] S. K. Q. Al-Omari and A. Kılıçman, "On diffraction Fresnel transforms for Boehmians," *Abstract and Applied Analysis*, vol. 2011, Article ID 712746, 11 pages, 2011.