Research Article

Generalizations of Fixed-Point Theorems of Altman and Rothe Types

Li Sun,^{1,2} Jingxian Sun,² and Guangwa Wang²

¹ School of Mechanics and Civil Engineering, China University of Mining & Technology, Xuzhou, Jiangsu 221008, China

² School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou, Jiangsu 221116, China

Correspondence should be addressed to Guangwa Wang; wanggw7653@163.com

Received 19 June 2013; Revised 9 September 2013; Accepted 27 September 2013

Academic Editor: Lishan Liu

Copyright © 2013 Li Sun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

It is intended to present some extensions of the famous Altman and Rothe types fixed-point theorems. The inequality conditions are relaxed to the α -positive-homogeneous operator. Some new fixed-point theorems are obtained with the help of the theory of topological degree.

1. Introduction

The topological degree theory and fixed-point theorems play an important role in the study of fixed points for various classes of nonlinear operators in Banach spaces. Many important results in mathematics are obtained by the use of these theories (see [1–12]). For example, Altman theorem is an important result from the theoretical as well as the applied point of view and is one of the core issues of mathematics. This fixed-point theorem has plenty of extensions and generalizations in the framework of nonlinear functional analysis and finds a wide application. In recent years, some researchers have focused on the inequality condition in Altman theorem and gained a lot of different generalized forms (see [2–6]). In this paper, we continue to study Altman theorem and relax the condition of square function to α -positive-homogeneous operator. It is well known that α -positive-homogeneous operator is a class of important nonlinear operators. Linear operator is an α -positive-homogeneous operator when $\alpha = 1$. Our main aim is to present and prove a collection of new fixed-point theorems for α -positive-homogeneous operator. Many existing results in the references are extended. On the other hand, it is well known that semiclosed 1-set-contraction mapping is a significant operator and research object in nonlinear functional analysis (see [3–5, 7, 8]). Clearly, this class of 1-set-contractive operators includes strict set-contractive operators, condensing

operators, semicontractive operators, completely continuous operators, and some others. The fixed-point theory of 1-set-contractive operators plays an important role in the study of the existence of the solution of some operator equations. In spite of this, completely continuous operators still have a pivotal position in modern nonlinear functional analysis. They have a widely and profoundly applicative prospect. Therefore, in this paper, we mainly investigate the fixed-point theorems about completely continuous operators and get some new results. Our conclusions are different from the ones in many recent works and extend some commonly known theorems. As far as we know, the cases in our paper have scarcely been seen in the available reference materials.

This paper contains three sections besides the introductory one. In Section 2, we give some basic concepts and some preparatory theorems. The main results will be presented and proved in Section 3. Then, we will list some existing results in the reference as our corollaries and this is the content of Section 4.

2. Preliminaries

In this section, we review some basic concepts and results which will be used later. For convenience, throughout all the paper, we let E, F be two real Banach spaces and let Ω be a bounded open subset of E and $\theta \in \Omega$.

Definition 1 (α -positive-homogeneous operator). Given $\alpha \in \mathbb{R}$, the mapping $B : E \rightarrow F$ is called an α -positive-homogeneous operator if for any $x \in E, t \geq 0$, the equation

$$B(tx) = t^\alpha Bx \quad (1)$$

always holds.

Definition 2 (cone). If P is a subset of Banach space F and satisfies the following three conditions:

- (i) P is a closed and convex set;
- (ii) $x \in P, \alpha \geq \theta \Rightarrow \alpha x \in P$;
- (iii) $x \in P, x \neq \theta \Rightarrow -x \notin P$,

then P is called a cone in F . Furthermore, if $x, y \in F$, then $y - x \in P$ defines a semiorder, denoted by $x \leq y$.

Lemma 3 (see [9], Leray-Schauder). Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. If

$$Ax \neq \mu x, \quad \forall x \in \partial\Omega, \mu \geq 1, \quad (2)$$

then $\deg(I - A, \Omega, \theta) = 1$, and hence A has at least one fixed point in $\overline{\Omega}$.

Lemma 4 (see [3, 5], Leray-Schauder). Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator; that is, A is 1-set contraction and $I - A$ is closed. If

$$Ax \neq \mu x, \quad \forall x \in \partial\Omega, \mu \geq 1, \quad (3)$$

then $\deg(I - A, \Omega, \theta) = 1$, and hence A has at least one fixed point in $\overline{\Omega}$.

In Lemmas 3 and 4, the same condition that

$$Ax \neq \mu x, \quad \forall x \in \partial\Omega, \mu \geq 1 \quad (4)$$

is called (L-S) condition.

Theorem 5 (see [9]). Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. If one of the following condition is satisfied:

- (i) (Altman) $\|Ax - x\|^2 \geq \|Ax\|^2 - \|x\|^2$, for all $x \in \partial\Omega$,
- (ii) (Rothe) $\|Ax\| \leq \|x\|$, for all $x \in \partial\Omega$,
- (iii) (Petryshyn) $\|Ax\| \leq \|Ax - x\|$, for all $x \in \partial\Omega$,

then $\deg(I - A, \Omega, \theta) = 1$, and hence A has at least one fixed point in $\overline{\Omega}$.

Theorem 6 (see [5]). Suppose that $A : \overline{\Omega} \rightarrow E$ is a semi-closed 1-set-contractive operator. If one of the following conditions is satisfied:

- (i) (Altman) $\|Ax - x\|^2 \geq \|Ax\|^2 - \|x\|^2$, for all $x \in \partial\Omega$,
- (ii) (Rothe) $\|Ax\| \leq \|x\|$, for all $x \in \partial\Omega$,
- (iii) (Petryshyn) $\|Ax\| \leq \|Ax - x\|$, for all $x \in \partial\Omega$,

then $\deg(I - A, \Omega, \theta) = 1$, and hence A has at least one fixed point in $\overline{\Omega}$.

3. Main Results

Now, we present and prove our main results.

Theorem 7. Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. Let P be a cone of F and let the operator $B : E \rightarrow P$ be α -positive-homogeneous such that $Bx > \theta$ for $x \neq \theta$. If $\alpha > 1$ and B satisfies the following condition:

$$B(Ax - x) \geq BAx - Bx, \quad \forall x \in \partial\Omega, \quad (5)$$

then A has at least one fixed point in $\overline{\Omega}$.

Proof. If the operator A has a fixed point on $\partial\Omega$, then A has at least one fixed point in $\overline{\Omega}$. Now, suppose that A has no fixed points on $\partial\Omega$. It suffices to prove that (5) implies the condition (L-S). Suppose the contrary, then there exist $x_1 \in \partial\Omega$ and $\mu_1 \geq 1$ such that $Ax_1 = \mu_1 x_1$; it is easy to see that $\mu_1 > 1$.

Now, consider the function defined by

$$f(t) = (t - 1)^\alpha - t^\alpha + 1, \quad \forall t \geq 1, \alpha > 1. \quad (6)$$

Since $f'(t) = \alpha(t - 1)^{\alpha-1} - \alpha t^{\alpha-1} < 0$ by formal differentiation, $f(t)$ is strictly decreasing in $[1, +\infty)$, and so $f(t) < f(1) = 0$, for all $t > 1$; that is, $(t - 1)^\alpha < t^\alpha - 1$, for all $t > 1, \alpha > 1$. Then, we can get from (5)

$$\begin{aligned} B(Ax_1 - x_1) &= B(\mu_1 x_1 - x_1) \\ &= (\mu_1 - 1)^\alpha Bx_1 \\ &\geq BAx_1 - Bx_1 \\ &= B(\mu_1 x_1) - Bx_1 \\ &= \mu_1^\alpha Bx_1 - Bx_1 \\ &= (\mu_1^\alpha - 1) Bx_1; \end{aligned} \quad (7)$$

that is, $(\mu_1 - 1)^\alpha Bx_1 \geq (\mu_1^\alpha - 1) Bx_1$. Thus $[(\mu_1 - 1)^\alpha - (\mu_1^\alpha - 1)] Bx_1 \leq \alpha_1 Bx_1 \in P$. Noticing $\alpha_1 < 0$ and $Bx_1 \neq \theta$, one can have that this contradicts the condition (iii) in Definition 2, and so the condition (L-S) is satisfied. Therefore, it follows from Lemma 3 that the conclusion of Theorem 7 holds. $\square$

Remark 8. Obviously, we only need to take $B = \|\cdot\|^2$ in Theorem 7 and we can get the famous fixed-point theorem of Altman in Theorem 5 (i). Consequently, Theorem 7 generalizes the classical Altman theorem.

Theorem 9. Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. Let P be a cone of F and let the operator $B : E \rightarrow P$ be α -positive-homogeneous such that $Bx > \theta$ for $x \neq \theta$. If $\alpha > 1$ and B satisfies the following condition:

$$BAx \leq Bx, \quad \forall x \in \partial\Omega, \quad (8)$$

then A has at least one fixed point in $\overline{\Omega}$.

Proof. We will also prove that (L-S) is satisfied under the condition (8). Suppose this is not true. Then there exist

$x_2 \in \partial\Omega$ and $\mu_2 \geq 1$ such that $Ax_2 = \mu_2 x_2$. It is easy to see that $\mu_2 > 1$. Then we can get from (8)

$$B(Ax_2) = B(\mu_2 x_2) = \mu_2^\alpha Bx_2 \leq Bx_2; \quad (9)$$

that is, $Bx_2 \geq \mu_2^\alpha Bx_2$, thus $(1 - \mu_2^\alpha)Bx_2 \doteq \alpha_2 Bx_2 \in P$. This contradicts the condition (iii) in Definition 2 on account of $\alpha_2 < 0$, $Bx_2 \neq \theta$, and so the condition (L-S) is satisfied. Therefore, it follows from Lemma 3 that the conclusion of Theorem 9 holds. $\square$

Remark 10. Taking $B = \|\cdot\|^2$ in Theorem 9, we can obtain the Rothe theorem in Theorem 5 (ii) immediately.

Theorem 11. Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. Let P be a cone of F and let $B : E \rightarrow P$ be an α -positive-homogeneous operator such that $Bx > \theta$ for $x \neq \theta$. If $\alpha > 1$ and B satisfies the following condition:

$$BAx \leq B(Ax - x), \quad \forall x \in \partial\Omega, \quad (10)$$

then A has at least one fixed point in $\overline{\Omega}$.

Proof. Similarly, we will also prove that (L-S) is satisfied. Suppose the contrary, then there exist $x_3 \in \partial\Omega$ and $\mu_3 \geq 1$ such that $Ax_3 = \mu_3 x_3$. Obviously, $\mu_3 > 1$. Then we can get from (10)

$$\begin{aligned} B(Ax_3) &= B(\mu_3 x_3) = \mu_3^\alpha Bx_3 \\ &\leq B(Ax_3 - x_3) = B(\mu_3 x_3 - x_3) \\ &= (\mu_3 - 1)^\alpha Bx_3, \end{aligned} \quad (11)$$

thus $(\mu_3 - 1)^\alpha Bx_3 \geq \mu_3^\alpha Bx_3$ and $[(\mu_3 - 1)^\alpha - \mu_3^\alpha]Bx_3 \doteq \alpha_3 Bx_3 \in P$. It also contradicts the condition (iii) in Definition 2 for $\alpha_3 < 0$, $Bx_3 \neq \theta$, and so the condition (L-S) is satisfied. Therefore, it follows from Lemma 3 that the conclusion of Theorem 11 holds. $\square$

Remark 12. (i) Theorem 11 is a generalization of the Petryshyn theorem.

(ii) In fact, Theorems 9 and 11 are actually the direct consequences because (8) and (10) immediately imply (5) due to $B : E \rightarrow P$.

Theorem 13. Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. Let P be a cone of F and let the operator $B : E \rightarrow P$ be α -positive-homogeneous operator such that $Bx > \theta$ for $x \neq \theta$. If $\alpha > 1$ and B satisfies the following condition:

$$B(Ax + x) \leq BAx + Bx, \quad \forall x \in \partial\Omega, \quad (12)$$

then A has at least one fixed point in $\overline{\Omega}$.

Proof. Suppose that (L-S) is not satisfied under the condition of (12). Then there exist $x_4 \in \partial\Omega$ and $\mu_4 \geq 1$ such that $Ax_4 = \mu_4 x_4$. It is easy to see that $\mu_4 > 1$. Now, we consider the function defined by

$$f(t) = t^\alpha + 1 - (t + 1)^\alpha, \quad \forall t \geq 1, \alpha > 1. \quad (13)$$

For $f'(t) = \alpha t^{\alpha-1} - \alpha(t + 1)^{\alpha-1} < 0$, for all $t > 1, \alpha > 1$. So $f(t)$ is a strictly decreasing function in $[1, +\infty)$, and then $f(t) < f(1) = 0$, for all $t > 1$, or $t^\alpha + 1 < (t + 1)^\alpha$, for all $t > 1, \alpha > 1$. Consequently, we can get from (12)

$$\begin{aligned} B(Ax_4 + x_4) &= B(\mu_4 x_4 + x_4) \\ &= (\mu_4 + 1)^\alpha Bx_4 \\ &\leq BAx_4 + Bx_4 \\ &= B(\mu_4 x_4) + Bx_4 \\ &= \mu_4^\alpha Bx_4 + Bx_4 \\ &= (\mu_4^\alpha + 1) Bx_4, \end{aligned} \quad (14)$$

then $[(\mu_4^\alpha + 1) - (\mu_4 + 1)^\alpha]Bx_4 \doteq \alpha_4 Bx_4 \in P$, which contradicts the condition (iii) in Definition 2. Hence, the condition (L-S) is satisfied and the conclusion of Theorem 13 holds. $\square$

In the same way, if $0 < \alpha < 1$, then some new fixed-point theorems can be obtained by taking the inverse direction in the inequalities in conditions (5), (8), (10), and (12).

Theorem 14. Suppose that $A : \overline{\Omega} \rightarrow E$ is a completely continuous operator. Let P be a cone of F , and let the operator $B : E \rightarrow P$ be α -positive-homogeneous operator such that $Bx > \theta$ for $x \neq \theta$. If $0 < \alpha < 1$ and B satisfies one of the following conditions:

- (i) $B(Ax - x) \leq BAx - Bx$, for all $x \in \partial\Omega$,
- (ii) $BAx \geq Bx$, for all $x \in \partial\Omega$,
- (iii) $BAx \geq B(Ax - x)$, for all $x \in \partial\Omega$,
- (iv) $B(Ax + x) \geq BAx + Bx$, for all $x \in \partial\Omega$,

then A has at least one fixed point in $\overline{\Omega}$.

Similarly, we can apply the above definitions and Lemma 4 to derive the following fixed point theorems.

Theorem 15. Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator. Let P be a cone of F , and let the operator $B : E \rightarrow P$ be α -positive-homogeneous such that $Bx > \theta$ for $x \neq \theta$. If $\alpha > 1$ and B satisfies one of the following conditions:

- (i) $B(Ax - x) \geq BAx - Bx$, for all $x \in \partial\Omega$,
- (ii) $BAx \leq Bx$, for all $x \in \partial\Omega$,
- (iii) $BAx \leq B(Ax - x)$, for all $x \in \partial\Omega$,
- (iv) $B(Ax + x) \leq BAx + Bx$, for all $x \in \partial\Omega$,

then A has at least one fixed point in $\overline{\Omega}$.

Theorem 16. Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator. Let P be a cone of F , and let the operator $B : E \rightarrow P$ be α -positive-homogeneous operator such that $Bx > \theta$ for $x \neq \theta$. If $0 < \alpha < 1$ and B satisfies one of the following conditions:

- (i) $B(Ax - x) \leq BAx - Bx$, for all $x \in \partial\Omega$,

- (ii) $B Ax \geq Bx$, for all $x \in \partial\Omega$,
- (iii) $B Ax \geq B(Ax - x)$, for all $x \in \partial\Omega$,
- (iv) $B(Ax + x) \geq B Ax + Bx$, for all $x \in \partial\Omega$,

then A has at least one fixed point in $\overline{\Omega}$.

Remark 17. Because completely continuous operators belong to the class of semiclosed 1-set-contractive operators, Theorems 15 and 16 are more general than Theorems 7–14.

4. Some Corollaries

Now we will list some existing results in the references.

Corollary 18 (see [4]). Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator. If there exists $\alpha > 1$ such that

$$\|Ax - x\|^\alpha \geq \|Ax\|^\alpha - \|x\|^\alpha, \quad \forall x \in \partial\Omega, \quad (15)$$

then $\deg(I - A, \Omega, \theta) = 1$, and so A has at least one fixed point in $\overline{\Omega}$.

In fact, if we take the operator $B = \|\cdot\|^\alpha$ in Theorem 15 (i), the Corollary 18 can be obtained immediately. So it is the special case of our main results and also the following corollaries are similar.

Corollary 19 (see [5]). Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator. If $0 < \alpha < 1$ such that

$$\|Ax - x\|^\alpha \leq \|Ax\|^\alpha - \|x\|^\alpha, \quad \forall x \in \partial\Omega, \quad (16)$$

then $\deg(I - A, \Omega, \theta) = 1$, and so A has at least one fixed point in $\overline{\Omega}$.

Corollary 20 (see [4]). Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator. If $\alpha > 1$ such that

$$\|Ax + x\|^\alpha \leq \|Ax\|^\alpha + \|x\|^\alpha, \quad \forall x \in \partial\Omega, \quad (17)$$

then $\deg(I - A, \Omega, \theta) = 1$, and so A has at least one fixed point in $\overline{\Omega}$.

Corollary 21 (see [4]). Suppose that $A : \overline{\Omega} \rightarrow E$ is a semiclosed 1-set-contractive operator. If there exists $0 < \alpha < 1$ such that

$$\|Ax + x\|^\alpha \geq \|Ax\|^\alpha + \|x\|^\alpha, \quad \forall x \in \partial\Omega, \quad (18)$$

then $\deg(I - A, \Omega, \theta) = 1$, and so A has at least one fixed point in $\overline{\Omega}$.

Acknowledgments

The authors would like to thank the referees for the thorough reading and excellent comments which led to the improvement of the original paper. This work was supported by the Special Funds of the National Natural Science Foundation of China (Grant no. 51227003), the National Nature Science Foundation of China (Grant nos. 10971179, 11026203), the Nature Science Foundation of Jiangsu Province (Grant no. BK2011202), and the Priority Academic Program Development (PAPD) of Jiangsu Higher Education Institutions.

References

- [1] M. Altman, "A fixed point theorem in Banach space," *Polish Academy of Sciences*, vol. 5, pp. 89–92, 1957.
- [2] G. Zhang, T. Zhang, and T. Zhang, "Fixed point theorems of Rothe and Altman types about convex-power condensing operator and application," *Applied Mathematics and Computation*, vol. 214, no. 2, pp. 618–623, 2009.
- [3] G. Li, "The fixed-point index and the fixed-point theorems of k -set-contraction mappings," *Proceedings of the American Mathematical Society*, vol. 104, no. 4, pp. 1163–1170, 1988.
- [4] G.-Z. Li, S.-Y. Xu, and H.-G. Duan, "Fixed point theorems of 1-set-contractive operators in Banach spaces," *Applied Mathematics Letters*, vol. 19, no. 5, pp. 403–412, 2006.
- [5] S. Xu, "New fixed point theorems for 1-set-contractive operators in Banach spaces," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 67, no. 3, pp. 938–944, 2007.
- [6] J. X. Sun and X. Y. Zhang, "A fixed point theorem for convex-power condensing operators and its applications to abstract semilinear evolution equations," *Acta Mathematica Sinica. Chinese Series*, vol. 48, no. 3, pp. 439–446, 2005.
- [7] R. D. Nussbaum, "Degree theory for local condensing maps," *Journal of Mathematical Analysis and Applications*, vol. 37, pp. 741–766, 1972.
- [8] R. D. Nussbaum, "The fixed point index and asymptotic fixed point theorems for k -set-contractions," *Bulletin of the American Mathematical Society*, vol. 75, pp. 490–495, 1969.
- [9] J. X. Sun, *Nonlinear Functional Analysis and Its Application*, Science Press, Beijing, China, 2007, Chinese.
- [10] H. Amann, "Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces," *SIAM Review*, vol. 18, no. 4, pp. 620–709, 1976.
- [11] D. J. Guo and J. X. Sun, "The calculation of topological degree and its applications," *Journal of Mathematical Research and Exposition*, vol. 8, no. 3, pp. 469–480, 1988.
- [12] W. V. Petryshyn, "Remarks on condensing and k -set-contractive mappings," *Journal of Mathematical Analysis and Applications*, vol. 39, pp. 717–741, 1972.