Research Article

Explicit Multistep Mixed Finite Element Method for RLW Equation

Yang Liu,¹ Hong Li,¹ Yanwei Du,¹ and Jinfeng Wang²

¹ School of Mathematical Sciences, Inner Mongolia University, Hohhot 010021, China

² School of Statistics and Mathematics, Inner Mongolia University of Finance and Economics, Hohhot 010070, China

Correspondence should be addressed to Yang Liu; mathliuyang@aliyun.com and Hong Li; smslh@imu.edu.cn

Received 15 February 2013; Revised 22 April 2013; Accepted 30 April 2013

Academic Editor: Marco Donatelli

Copyright © 2013 Yang Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An explicit multistep mixed finite element method is proposed and discussed for regularized long wave (RLW) equation. The spatial direction is approximated by the mixed Galerkin method using mixed linear space finite elements, and the time direction is discretized by the explicit multistep method. The optimal error estimates in L^2 and H^1 norms for the scalar unknown u and its flux $q = u_x$ based on time explicit multistep method are derived. Some numerical results are given to verify our theoretical analysis and illustrate the efficiency of our method.

1. Introduction

In this paper, we consider the following initial boundary problem of RLW equation:

$$\begin{aligned} u_t + \gamma u_x + \delta u u_x - \beta u_{xxt} &= 0, \quad (x, t) \in I \times J, \\ u(a, t) = u(b, t) &= 0, \quad t \in \bar{J}, \\ u(x, 0) &= u_0(x), \quad x \in \bar{I}, \end{aligned} \quad (1)$$

where $I = (a, b)$ is a bounded open interval, $J = (0, T]$ with $0 < T < \infty$. The initial value $u_0(x)$ is given function, and the coefficients β, γ, δ are all positive constants.

In recent years, large nonlinear phenomena are found in many research fields, for example, physics, biology, fluid dynamics, and so forth. These phenomena can be described by the mathematical model of some nonlinear evolution equations. In particular, some attention has also been paid to nonlinear RLW equations [1, 2] which play a very important role in the study of nonlinear dispersive waves. Solitary waves are wave packet or pulses, which propagate in nonlinear dispersive media. Due to dynamical balance between the nonlinear and dispersive effects, these waves retain an unchanged waveform. A soliton is a very special type of solitary wave, which also keeps its waveform after collision with other solitons. The regularized long wave

(RLW) equation is an alternative description of nonlinear dispersive waves to the more usual Korteweg-de Vries (KdV) equation. Mathematical theories and numerical methods for (1) were considered in [1–24]. The existence and uniqueness of the solution of RLW equation are discussed in [5]. Their analytical solutions were found under restricted initial and boundary conditions, and therefore they got interest from a numerical point of view. Several numerical methods for the solution of the RLW equation have been introduced in the literature. These include a variety of difference methods [5–8, 17], finite element methods based on Galerkin and collocation principles [9–13], mixed finite element methods [14–16], meshfree method [18], adomian decomposition method [19], and so on.

In [25], Chatzipantelidis studied the explicit multistep methods for some nonlinear partial differential equations and discussed some mathematical theories. Akrivis et al. [26] studied the multistep method for some nonlinear evolution equations. Mei and Chen [20] presented the explicit multistep method based on Galerkin method for regularized long wave (RLW) equation. In this paper, our purpose is to propose and study an explicit multistep mixed method, which combines a mixed Galerkin method in the spatial direction and the explicit multistep method in the time direction, for RLW equation. We derive optimal error estimates in L^2 and H^1 norms for the scalar unknown u and its flux $q = u_x$

for the fully discrete explicit multistep mixed scheme and compare our method's accuracy with some other numerical schemes. Compared to the numerical methods in [20, 25, 26], we not only obtain the approximation solution for u , but also get the approximation solution for $q = u_x$.

The layout of the paper is as follows. In Section 2, an explicit multistep mixed scheme and numerical process are given. The optimal error estimates in L^2 and H^1 norms for the scalar unknown u and its flux $q = u_x$ for the fully discrete explicit multistep mixed scheme are proved in Section 3. In Section 4, some numerical results are shown to confirm our theoretical analysis. Finally, some concluding remarks are given in Section 5. Throughout this paper, C will denote a generic positive constant which does not depend on the spatial mesh parameter h or time discretization parameter Δt .

2. The Mixed Numerical Scheme

With the auxiliary variable $q = u_x$, we reformulate (1) as the following first-order coupled system:

$$\begin{aligned} u_x &= q, \\ u_t + \gamma q + \delta u q - \beta q_{xt} &= 0. \end{aligned} \quad (2)$$

We consider the following mixed weak formulation of (2). Find $\{u, q\} : [0, T] \mapsto H_0^1 \times H^1$ satisfying:

$$(u_x, v_x) = (q, v_x), \quad \forall v \in H_0^1, \quad (3)$$

$$(q_t, w) + \beta (q_{xt}, w_x) = \gamma (q, w_x) + \delta (uq, w_x), \quad \forall w \in H^1. \quad (4)$$

Noting the Dirichlet boundary conditions $u_t(a, t) = u_t(b, t) = 0$ and $q_t = u_{xt}$, we can get $(u_t, -w_x) = (u_{xt}, w) = (q_t, w)$ easily and then get the scheme (4).

Let V_h and W_h be finite dimensional subspaces of H_0^1 and H^1 , respectively, defined by

$$\begin{aligned} V_h &= \left\{ v_h \mid v_h \in C^0(\bar{I}), v_h|_{I_j} \in P_k(I_j), \right. \\ &\quad \left. \forall I_j \in T_h, v_h(a) = v_h(b) = 0 \right\} \\ &\subset H_0^1, \\ W_h &= \left\{ w_h \mid w_h \in C^0(\bar{I}), w_h|_{I_j} \in P_r(I_j), \forall I_j \in T_h \right\} \subset H^1, \end{aligned} \quad (5)$$

where T_h is a partition of $\bar{I} = [a, b]$ into N subintervals $I_j = [x_j, x_{j+1}]$, $j = 0, 1, 2, \dots, (N-1)$, $h_j = x_{j+1} - x_j$, $h = \max_{0 \leq j \leq N-1} h_j$, and $P_m(I_j)$ denotes the polynomials of degree less than or equal to m in I_j .

The semidiscrete mixed finite element method for (3) and (4) consists in determining $\{u^h, q^h\} : [0, T] \mapsto V_h \times W_h$ such that

$$(u_x^h, v_x^h) = (q^h, v_x^h), \quad \forall v^h \in V_h, \quad (6)$$

$$\begin{aligned} (q_t^h, w^h) + \beta (q_{xt}^h, w_x^h) &= \gamma (q^h, w_x^h) + \delta (u^h q^h, w_x^h), \\ \forall w^h &\in W_h. \end{aligned} \quad (7)$$

In the following discussion, we will give an explicit multistep mixed scheme. We take linear finite element spaces $V_h = \text{span}\{\varphi_0, \varphi_1, \dots, \varphi_N\}$ and $W_h = \text{span}\{\varphi_0, \varphi_1, \dots, \varphi_N\}$, and then u_h and q_h can be expressed as the following formulation:

$$\begin{aligned} u_h(x, t) &= \sum_{i=0}^N u_i(t) \varphi_i(x), \quad (x, t) \in \Omega \times J, \\ q_h(x, t) &= \sum_{i=0}^N q_i(t) \varphi_i(x), \quad (x, t) \in \Omega \times J. \end{aligned} \quad (8)$$

Substitute (8) into (6) and (7), and take $v^h = \varphi_j$ and $w^h = \varphi_j$ in (6) and (7), respectively, to obtain

$$\begin{aligned} \sum_{i=0}^N \left[\left(\int_a^b \varphi_i \cdot \varphi_j dx + \beta \int_a^b \varphi_{ix} \cdot \varphi_{jx} dx \right) \frac{\partial q_i}{\partial t} \right. \\ \left. - \left(\gamma \int_a^b \varphi_i \cdot \varphi_{jx} dx + \delta \int_a^b \left(\sum_{i=0}^N u_i \varphi_i \right) \varphi_i \cdot \varphi_{jx} dx \right) q_i \right] \\ = 0, \\ \sum_{i=0}^N \left[\left(\int_a^b \varphi_{ix} \cdot \varphi_{jx} dx \right) u_i - \left(\int_a^b \varphi_i \cdot \varphi_{jx} dx \right) q_i \right] = 0, \end{aligned} \quad (9)$$

where $j = 0, 1, 2, \dots, N$.

We subdivide the space variable domain $[a, b]$ into uniform subintervals with $N+1$ grid points x_k , $k = 0, \dots, N$, such that $a = x_0 < x_1 < \dots < x_N = b$, $h = x_{k+1} - x_k = (b-a)/N$. Using the local coordinate transformation $x = x_k + \mu h$, $0 \leq \mu \leq 1$, we transform a subinterval $[x_k, x_{k+1}]$ into a standard interval $[0, 1]$. Furthermore, we have

$$\begin{aligned} \sum_{i=k}^{k+1} \left[\left(\int_0^1 \varphi_i \cdot \varphi_j dx + \frac{\beta}{h^2} \int_0^1 \varphi_{ix} \cdot \varphi_{jx} dx \right) \frac{\partial q_i}{\partial t} \right. \\ \left. - \frac{1}{h} \left(\gamma \int_0^1 \varphi_i \cdot \varphi_{jx} dx + \delta \int_0^1 \left(\sum_{i=0}^N u_i \varphi_i \right) \varphi_i \cdot \varphi_{jx} dx \right) q_i \right] \\ = 0, \\ \sum_{i=k}^{k+1} \left[\frac{1}{h^2} \left(\int_0^1 \varphi_{ix} \cdot \varphi_{jx} dx \right) u_i - \frac{1}{h} \left(\int_a^b \varphi_i \cdot \varphi_{jx} dx \right) q_i \right] = 0. \end{aligned} \quad (10)$$

TABLE 1: Solitary wave Amp. 0.3 and the errors in L^2 and L^∞ norms for u , Q_1 , Q_2 , and Q_3 at $t = 20$, $h = 0.125$, $\Delta t = 0.1$, and $-40 \leq x \leq 60$.

Method	Time	Q_1	Q_2	Q_3	L^2 for u	L^∞ for u
Our method	0	3.9797	0.8104	2.5787	0	0
	4	3.9797	0.8104	2.5786	$3.6304e-004$	$5.2892e-005$
	8	3.9797	0.8104	2.5786	$7.2873e-004$	$5.8664e-005$
	12	3.9797	0.8104	2.5787	$1.0817e-003$	$6.3283e-005$
	16	3.9795	0.8104	2.5787	$1.4186e-003$	$6.8001e-005$
	20	3.9790	0.8103	2.5785	$1.7396e-003$	$7.7154e-005$
[20]	20	3.9800	0.8104	2.5792	$1.7569e-003$	$6.8432e-004$
[21]	20	3.9820	0.8087	2.5730	$4.688e-003$	$1.755e-003$
[22]	20	3.9905	0.8235	2.6740	$2.157e-003$	—
[23]	20	3.9616	0.8042	2.5583	$0.018e-003$	$1.566e-003$
[24]	20	3.9821	0.8112	2.5813	$0.511e-003$	$0.198e-003$

TABLE 2: Convergence order and error in L^2 norm for u of time with $h = 0.125$ and $c = 0.1$.

Time	$\Delta t = 0.4$	$\Delta t = 0.2$	$\Delta t = 0.1$	Order (0.2/0.4)	Order (0.1/0.2)
4	$5.3805e-003$	$1.4267e-003$	$3.6304e-004$	1.9151	1.9745
8	$1.1688e-002$	$2.9411e-003$	$7.2873e-004$	1.9906	2.0129
12	$1.7830e-002$	$4.3997e-003$	$1.0817e-003$	2.0188	2.0241
16	$2.3751e-002$	$5.7916e-003$	$1.4186e-003$	2.0360	2.0295
20	$2.9434e-002$	$7.1148e-003$	$1.7396e-003$	2.0486	2.0321

We take linear basis functions defined as follows:

$$L_1 = 1 - \mu, \quad L_2 = \mu, \quad (11)$$

and then the variables u and q over the element $[x_k, x_{k+1}]$ are written as

$$u^e = \sum_{j=1}^2 L_j u_j, \quad q^e = \sum_{j=1}^2 L_j q_j. \quad (12)$$

Then, we get the following equations:

$$\begin{aligned} & \sum_{i=1}^2 \left[\left(\int_0^1 L_i \cdot L_j d\mu + \frac{\beta}{h^2} \int_0^1 L_{i\mu} \cdot L_{j\mu} d\mu \right) \frac{\partial q_i}{\partial t} \right. \\ & \quad \left. - \frac{1}{h} \left(\gamma \int_0^1 L_i \cdot L_{j\mu} d\mu + \delta \int_0^1 \left(\sum_{i=1}^2 u_i L_i \right) L_i \cdot L_{j\mu} d\mu \right) q_i \right] \\ & = 0, \\ & \sum_{i=1}^2 \left[\frac{1}{h^2} \left(\int_0^1 L_{i\mu} \cdot L_{j\mu} d\mu \right) u_i - \frac{1}{h} \left(\int_a^b L_i \cdot L_{j\mu} d\mu \right) q_i \right] = 0. \end{aligned} \quad (13)$$

Then, the system (13) has the following matrix form:

$$\begin{aligned} (A_{ij}^e + \beta B_{ij}^e) \frac{\partial \mathbf{q}^e}{\partial t} - (\gamma C_{ij}^e + \delta D_{ij}^e(u^e)) \mathbf{q}^e &= 0, \\ B_{ij}^e \mathbf{u}^e - C_{ij}^e \mathbf{q}^e &= 0, \end{aligned} \quad (14)$$

with the following element matrices:

$$\begin{aligned} \mathbf{u}^e &= (u_1, u_2)^T, \quad \mathbf{q}^e = (q_1, q_2)^T, \\ A_{ij}^e &= \int_0^1 L_i \cdot L_j d\mu, \quad B_{ij}^e = \frac{1}{h^2} \int_0^1 L_{i\mu} \cdot L_{j\mu} d\mu, \\ C_{ij}^e &= \frac{1}{h} \int_0^1 L_i \cdot L_{j\mu} d\mu, \\ D_{jk}^e &= \frac{\delta}{h} \int_0^1 \left(\sum_{i=1}^2 u_i L_i \right) L_{i\mu} \cdot L_{j\mu} d\mu. \end{aligned} \quad (15)$$

Assembling contributions from all elements, we obtain the following coupled system of nonlinear matrix equations:

$$\begin{aligned} (A + \beta B) \frac{\partial \mathbf{q}_h}{\partial t} - (\gamma C + \delta D(\mathbf{u}_h)) \mathbf{q}_h &= 0, \\ B \mathbf{u}_h - C \mathbf{q}_h &= 0. \end{aligned} \quad (16)$$

To formulate a fully discrete scheme, we consider a uniform partition of $\bar{J} = [0, T]$ with time step length $\Delta t = T/N$, $N \in \mathbf{Z}^+$, and time levels $t^n = n\Delta t$, $n = 0, \dots, N$. We now discuss a fully discrete scheme based on a linear explicit multistep method. We now define $U^n \in V_h$ and $Z^n \in W_h$ as approximations to $u(t^n)$ and $q(t^n)$, respectively, and formulate the following fully discrete linear explicit multistep mixed scheme:

$$\begin{aligned} (A + \beta B) \sum_{i=0}^p \alpha_i Z^{n+i} &= \Delta t \sum_{i=0}^{p-1} \sigma_i \left[(\gamma C + \delta D(U^{n+i})) Z^{n+i} \right], \\ B U^{n+p} &= C Z^{n+p}, \end{aligned} \quad (17)$$

TABLE 3: Convergence order and error in L^∞ norm for u of time with $h = 0.125$ and $c = 0.1$.

Time	$\Delta t = 0.4$	$\Delta t = 0.2$	$\Delta t = 0.1$	Order (0.2/0.4)	Order (0.1/0.2)
4	$8.0743e-004$	$2.0299e-004$	$5.2892e-005$	1.9919	1.9403
8	$9.1778e-004$	$2.2908e-004$	$5.8664e-005$	2.0023	1.9653
12	$9.9866e-004$	$2.4966e-004$	$6.3283e-005$	2.0000	1.9801
16	$1.0718e-003$	$2.6685e-004$	$6.8001e-005$	2.0059	1.9724
20	$1.1277e-003$	$2.8609e-004$	$7.7154e-005$	1.9788	1.8907

TABLE 4: Convergence order and error in L^2 norm for q of time with $h = 0.125$ and $c = 0.1$.

Time	$\Delta t = 0.4$	$\Delta t = 0.2$	$\Delta t = 0.1$	Order (0.2/0.4)	Order (0.1/0.2)
4	$2.4430e-003$	$6.4427e-004$	$1.6260e-004$	1.9229	1.9863
8	$5.1823e-003$	$1.2934e-003$	$3.1976e-004$	2.0024	2.0161
12	$7.6616e-003$	$1.8713e-003$	$4.5963e-004$	2.0336	2.0255
16	$9.8719e-003$	$2.3787e-003$	$5.8232e-004$	2.0532	2.0303
20	$1.1843e-002$	$2.8251e-003$	$6.8991e-004$	2.0677	2.0338

with given the initial approximations $U^0, \dots, U^{p-1}$ and $Z^0, \dots, Z^{p-1}$. In the explicit multistep mixed system (17), the parameter variable α_i and σ_i is described by the coefficients of the term χ^i , for the following polynomials $\alpha(\chi)$ and $\sigma(\chi)$, respectively:

$$\alpha(\chi) := \sum_{j=1}^p \frac{1}{j} \chi^{p-j} (\chi - 1)^j, \quad (18)$$

$$\sigma(\chi) := \chi^p - (\chi - 1)^p.$$

In this paper, we consider the explicit 2-step mixed method for the RLW equation. For $p = 2$, we obtained easily

$$\begin{aligned} \alpha_0 &= \frac{3}{2}, & \alpha_1 &= -2, & \alpha_2 &= \frac{1}{2}, \\ \sigma_0 &= -1, & \sigma_1 &= 2. \end{aligned} \quad (19)$$

Substituting (19) into (17), we obtain the following 2-step mixed scheme:

$$\begin{aligned} (A + \beta B) \left(\frac{3}{2} Z^{n+2} - 2Z^{n+1} + \frac{1}{2} Z^n \right) \\ = \Delta t \left[\gamma C (2Z^{n+1} - Z^n) + \delta (2D(U^{n+1}) Z^{n+1} \right. \\ \left. - D(U^n) Z^n) \right], \\ BU^{n+2} = CZ^{n+2}. \end{aligned} \quad (20)$$

Remark 1. There have been many numerical schemes for the RLW equation, but we have not seen the related research on explicit multistep mixed element method for RLW equation in the literature. From the viewpoint of numerical theory, we propose a mixed element scheme (6) and (7), which is different from some other mixed finite element methods in [14–16], for the RLW equation and derive some a priori error estimates based on the explicit multistep mixed element method. From the perspective of numerical calculation, our method is efficient for RLW equation.

3. Two-Step Mixed Scheme and Optimal Error Estimates

3.1. Two-Step Mixed Scheme and Some Lemmas. In this section, we will discuss some a priori error estimates based on explicit 2-step mixed finite element method for the RLW equation. For the fully discrete procedure, let $0 = t_0 < t_1 < \dots < t_N = T$ be a given partition of the time interval $[0, T]$ with step length $\Delta t = T/N$, for some positive integer N . For a smooth function ϕ on $[0, T]$, define $\phi^n = \phi(t_n)$.

The system (3) and (4) has the following formulation at $t = t_{n+1}$:

$$\begin{aligned} (u_x^{n+1}, v_x) &= (q^{n+1}, v_x), \quad \forall v \in H_0^1, \\ (q_t^{n+1}, w) + \beta (q_{xt}^{n+1}, w_x) &= \gamma (q^{n+1}, w_x) + \delta (u^{n+1} q^{n+1}, w_x), \\ \forall w &\in H^1. \end{aligned} \quad (21)$$

Based on system (17), we get an equivalent formulation for system (21) as

$$\begin{aligned} (u_x^{n+1}, v_x) &= (q^{n+1}, v_x), \quad \forall v \in H_0^1, \\ \left(\frac{3q^{n+1} - 4q^n + q^{n-1}}{2\Delta t}, w \right) + \beta \left(\frac{3q_x^{n+1} - 4q_x^n + q_x^{n-1}}{2\Delta t}, w_x \right) \\ &= -(\tau^{n+1}, w) - \beta (\kappa_1^{n+1}, w_x) + \gamma (2q^n - q^{n-1}, w_x) \\ &\quad + \delta (2u^n q^n - u^{n-1} q^{n-1}, w_x) + \gamma (R_1^{n+1}, w_x) \\ &\quad + \delta (R_2^{n+1}, w_x), \quad \forall w \in H^1, \end{aligned} \quad (22)$$

(23)

TABLE 5: Convergence order and error in L^∞ norm for q of time with $h = 0.125$ and $c = 0.1$.

Time	$\Delta t = 0.4$	$\Delta t = 0.2$	$\Delta t = 0.1$	Order (0.2/0.4)	Order (0.1/0.2)
4	$1.1595e-004$	$2.7265e-005$	$6.3127e-006$	2.0884	2.1107
8	$1.9712e-004$	$4.6625e-005$	$1.0725e-005$	2.0799	2.1201
12	$2.5295e-004$	$5.9837e-005$	$1.3579e-005$	2.0797	2.1397
16	$2.9012e-004$	$6.8641e-005$	$1.5594e-005$	2.0795	2.1381
20	$3.1768e-004$	$7.5864e-005$	$1.7195e-005$	2.0661	2.1414

TABLE 6: Convergence order and error in L^2 norm for u of space with $\Delta t = 0.01$ and $c = 0.1$.

Time	$h = 0.8$	$h = 0.4$	$h = 0.2$	Order (0.4/0.8)	Order (0.2/0.4)
4	$1.7109e-003$	$4.2677e-004$	$1.1449e-004$	1.9940	1.8982
8	$1.8945e-003$	$4.6862e-004$	$1.1924e-004$	2.0195	1.9746
12	$2.1232e-003$	$5.2130e-004$	$1.3025e-004$	2.0102	2.0008
16	$2.3559e-003$	$5.7598e-004$	$1.4421e-004$	2.0589	1.9978
20	$2.5778e-003$	$6.3407e-004$	$1.7877e-004$	2.0358	1.8265

where

$$\begin{aligned}\tau^{n+1} &= q_t^{n+1} - \frac{3q^{n+1} - 4q^n + q^{n-1}}{2\Delta t}, \\ \kappa_1^{n+1} &= q_{xt}(t_{n+1}) - \frac{3q_x^{n+1} - 4q_x^n + q_x^{n-1}}{2\Delta t},\end{aligned}\quad (24)$$

$$\begin{aligned}R_1^{n+1} &= q^{n+1} - (2q^n - q^{n-1}), \\ R_2^{n+1} &= u^{n+1}q^{n+1} - (2u^nq^n - u^{n-1}q^{n-1}).\end{aligned}$$

We now find a pair $\{U^{n+1}, Z^{n+1}\}$ in $V_h \times W_h$ satisfying

$$\begin{aligned}(U_x^{n+1}, v_x^h) &= (Z^{n+1}, v_x^h), \quad \forall v^h \in V_h, \\ \left(\frac{3Z^{n+1} - 4Z^n + Z^{n-1}}{2\Delta t}, w^h\right) &+ \beta \left(\frac{3Z_x^{n+1} - 4Z_x^n + Z_x^{n-1}}{2\Delta t}, w_x^h\right) \\ &= \gamma(2Z^n - Z^{n-1}, w_x^h) + \delta(2U^n Z^n - U^{n-1} Z^{n-1}, w_x^h), \\ &\forall w^h \in W_h.\end{aligned}\quad (25)$$

For the theoretical analysis of a priori error estimates, we define the following projections.

Lemma 2 (see [15, 27, 28]). *One defines the elliptic projection $\tilde{u}^h \in V_h$ by*

$$(u_x - \tilde{u}_x^h, v_x^h) = 0, \quad v^h \in V_h. \quad (27)$$

With $\eta = u - \tilde{u}^h$, the following estimates are well known for $j = 0, 1$:

$$\|\eta\|_j \leq Ch^{k+1-j} \|u\|_{k+1}. \quad (28)$$

Lemma 3 (see [15, 27, 28]). *Furthermore, one also defines a Ritz projection $\tilde{q}^h \in W_h$ of q as the solution of*

$$A(q - \tilde{q}^h, w^h) = 0, \quad w^h \in W_h, \quad (29)$$

where $A(q, w) = (q_x, w_x) + \lambda(q, w)$, and λ is taken appropriately so that

$$A(w, w) \geq \mu_0 \|w\|_1^2, \quad w \in H^1, \quad (30)$$

where μ_0 is a positive constant. Moreover, it is easy to verify that $A(\cdot, \cdot)$ is bounded.

With $\rho = q - \tilde{q}^h$, the following estimates hold:

$$\left\| \frac{\partial^j \rho}{\partial t^i} \right\|_j \leq Ch^{r+1-j} \left\| \frac{\partial^j q}{\partial t^i} \right\|_{r+1}, \quad i = 0, 1, 2, 3; \quad j = 0, 1. \quad (31)$$

For fully discrete error estimates, we now write the errors as

$$\begin{aligned}u(t_n) - U^n &= (u(t_n) - \tilde{u}^h(t_n)) + (\tilde{u}^h(t_n) - U^n) = \eta^n + \zeta^n, \\ q(t_n) - Z^n &= (q(t_n) - \tilde{q}^h(t_n)) \\ &\quad + (\tilde{q}^h(t_n) - Z^n) = \rho^n + \xi^n.\end{aligned}\quad (32)$$

Combine (27), (29), (22), (23), (25), and (26) at $t = t_{n+1}$ to get the following error equations:

$$\begin{aligned}(\zeta_x^{n+1}, v_x^h) &= (\rho^{n+1} + \xi^{n+1}, v_x^h), \quad \forall v^h \in V_h, \\ \left(\frac{3\zeta^{n+1} - 4\zeta^n + \zeta^{n-1}}{2\Delta t}, w^h\right) &+ \beta \left(\frac{3\zeta_x^{n+1} - 4\zeta_x^n + \zeta_x^{n-1}}{2\Delta t}, w_x^h\right) \\ &= - \left((1 - \beta\lambda) \frac{3\rho^{n+1} - 4\rho^n + \rho^{n-1}}{2\Delta t} + \tau^{n+1}, w^h \right) \\ &\quad - \beta(\kappa_2^{n+1}, w_x^h) + \gamma(2\xi^n - \xi^{n-1}, w_x^h) \\ &\quad + \delta(2(u(t_n)q(t_n) - U^n Z^n) \\ &\quad \quad - (u(t_{n-1})q(t_{n-1}) - U^{n-1} Z^{n-1}), w_x^h) \\ &\quad + \gamma(R_1^{n+1}, w_x^h) + \delta(R_2^{n+1}, w_x^h), \quad \forall w^h \in W_h,\end{aligned}\quad (33)$$

(34)

TABLE 7: Convergence order and error in L^∞ norm for u of space with $\Delta t = 0.01$ and $c = 0.1$.

Time	$h = 0.8$	$h = 0.4$	$h = 0.2$	Order (0.4/0.8)	Order (0.2/0.4)
4	$2.1763e-003$	$5.8447e-004$	$1.5502e-004$	1.8967	1.9147
8	$4.7892e-003$	$1.2098e-003$	$3.0533e-004$	1.9850	1.9863
12	$7.2371e-003$	$1.7871e-003$	$4.4434e-004$	2.0178	2.0079
16	$9.4746e-003$	$2.3084e-003$	$5.7081e-004$	2.0372	2.0158
20	$1.1534e-002$	$2.7859e-003$	$6.8980e-004$	2.0497	2.0139

TABLE 8: Convergence order and error in L^2 norm for q of space with $\Delta t = 0.01$ and $c = 0.1$.

Time	$h = 0.8$	$h = 0.4$	$h = 0.2$	Order (0.4/0.8)	Order (0.2/0.4)
4	$2.3426e-004$	$5.4759e-005$	$1.2581e-005$	2.0969	2.1218
8	$4.2831e-004$	$1.0038e-004$	$2.2853e-005$	2.0932	2.1350
12	$5.7270e-004$	$1.3452e-004$	$3.0443e-005$	2.0900	2.1436
16	$6.7812e-004$	$1.5951e-004$	$3.5927e-005$	2.0879	2.1505
20	$7.5693e-004$	$1.7832e-004$	$4.0176e-005$	2.0857	2.1501

where

$$\kappa_2^{n+1} = \tilde{q}_{xt}^h(t_{n+1}) - \frac{3\tilde{q}_{xx}^{h,n+1} - 4\tilde{q}_{xx}^{h,n} + \tilde{q}_{xx}^{h,n-1}}{2\Delta t}. \quad (35)$$

Lemma 4. For τ^{n+1} , κ_2^{n+1} , R_1^{n+1} , and R_2^{n+1} , the following estimates hold:

$$\begin{aligned} \|\tau^{n+1}\| &\leq C\Delta t^2 \|q_{ttt}\|_{L^\infty(L^2)}, \\ \|\kappa_2^{n+1}\| &\leq C\Delta t^2 (h^r \|q_{xttt}\|_{L^\infty(H^{r+1})} + \|q_{xttt}\|_{L^\infty(L^2)}), \\ \|R_1^{n+1}\| &\leq C\Delta t^2 \|q_{tt}\|_{L^\infty(L^2)}, \\ \|R_2^{n+1}\| &\leq C\Delta t^2 (\|uq_{tt}\|_{L^\infty(L^2)} + \|u_t q_t\|_{L^\infty(L^2)} + \|u_{tt} q\|_{L^\infty(L^2)}). \end{aligned} \quad (36)$$

Proof. Using the Taylor expansion, we have

$$\begin{aligned} q(t_n) &= q(t_{n+1}) + q_t(t_{n+1})(t_n - t_{n+1}) \\ &\quad + \frac{q_{tt}(t_{\Delta 1})}{2}(t_n - t_{n+1})^2, \quad t_n < t_{\Delta 1} < t_{n+1}, \end{aligned} \quad (37)$$

$$\begin{aligned} q(t_{n-1}) &= q(t_{n+1}) + q_t(t_{n+1})(t_{n-1} - t_{n+1}) \\ &\quad + \frac{q_{tt}(t_{\Delta 2})}{2}(t_{n-1} - t_{n+1})^2, \quad t_{n-1} < t_{\Delta 2} < t_{n+1}. \end{aligned} \quad (38)$$

Combining (37) and (38) and noting that $-2\Delta t = 2(t_n - t_{n+1}) = t_{n-1} - t_{n+1}$, we obtain

$$q(t_{n+1}) = 2q(t_n) - q(t_{n-1}) + (q_{tt}(t_{\Delta 1}) - 2q_{tt}(t_{\Delta 2}))\Delta t^2. \quad (39)$$

From (39), we have

$$\|q(t_{n+1}) - (2q(t_n) - q(t_{n-1}))\| \leq C\Delta t^2 \|q_{tt}\|_{L^\infty(L^2)}. \quad (40)$$

Using a similar estimate as the one for $\|R_1^{n+1}\|$, we have

$$\begin{aligned} u(t_{n+1})q(t_{n+1}) &= 2u(t_n)q(t_n) - u(t_{n-1})q(t_{n-1}) \\ &\quad + ((uq)_{tt}(t_{\Delta 3}) - 2(uq)_{tt}(t_{\Delta 4}))\Delta t^2, \end{aligned} \quad (41)$$

where $t_n < t_{\Delta 3} < t_{n+1}$, $t_{n-1} < t_{\Delta 4} < t_{n+1}$.

From (41), we have

$$\begin{aligned} \|u(t_{n+1})q(t_{n+1}) - (2u(t_n)q(t_n) - u(t_{n-1})q(t_{n-1}))\| \\ \leq C\Delta t^2 (\|uq_{tt}\|_{L^\infty(L^2)} + \|u_t q_t\|_{L^\infty(L^2)} + \|u_{tt} q\|_{L^\infty(L^2)}). \end{aligned} \quad (42)$$

Using the Taylor expansion and noting that $-2\Delta t = 2(t_n - t_{n+1}) = t_{n-1} - t_{n+1}$, we have

$$\begin{aligned} 4q(t_n) &= 4q(t_{n+1}) - 4q_t(t_{n+1})\Delta t + 2q_{tt}(t_{n+1})\Delta t^2 \\ &\quad - \frac{2q_{ttt}(t_{\Delta 5})}{3}\Delta t^3, \quad t_n < t_{\Delta 5} < t_{n+1}, \\ q(t_{n-1}) &= q(t_{n+1}) - 2q_t(t_{n+1})\Delta t + 2\frac{q_{tt}(t_{n+1})}{2}\Delta t^2 \\ &\quad - \frac{4q_{ttt}(t_{\Delta 6})}{3}\Delta t^3, \quad t_{n-1} < t_{\Delta 6} < t_{n+1}. \end{aligned} \quad (43)$$

Using (43), we obtain

$$q_t^{n+1} = \frac{3q^{n+1} - 4q^n + q^{n-1}}{2\Delta t} - \left(\frac{2q_{ttt}(t_{\Delta 5})}{3} - \frac{4q_{ttt}(t_{\Delta 6})}{3} \right) \Delta t^2. \quad (44)$$

By (44), we have

$$\|\tau^{n+1}\| \leq C\Delta t^2 \|q_{ttt}(t)\|_{L^\infty(L^2)}. \quad (45)$$

TABLE 9: Convergence order and error in L^∞ norm for q of space with $\Delta t = 0.01$ and $c = 0.1$.

Time	$h = 0.8$	$h = 0.4$	$h = 0.2$	Order (0.4/0.8)	Order (0.2/0.4)
4	$1.1957e - 003$	$3.1367e - 004$	$7.8812e - 005$	1.9305	1.9928
8	$2.4344e - 003$	$6.0206e - 004$	$1.4825e - 004$	2.0156	2.0219
12	$3.4768e - 003$	$8.3962e - 004$	$2.0540e - 004$	2.0500	2.0313
16	$4.3743e - 003$	$1.0402e - 003$	$2.5371e - 004$	2.0722	2.0356
20	$5.1677e - 003$	$1.2153e - 003$	$2.9544e - 004$	2.0882	2.0404

Using the similar method to the estimate for $\|\tau^{n+1}\|$ and (31), we obtain

$$\begin{aligned} \|\kappa_2^{n+1}\| &\leq C\Delta t^2 \|\tilde{q}_{xttt}^h\|_{L^\infty(L^2)} \\ &\leq C\Delta t^2 (\|\rho_{xttt}\|_{L^\infty(L^2)} + \|q_{xttt}\|_{L^\infty(L^2)}) \\ &\leq C\Delta t^2 (h^r \|q_{xttt}\|_{L^\infty(H^{r+1})} + \|q_{xttt}\|_{L^\infty(L^2)}). \end{aligned} \quad (46)$$

□

3.2. Optimal Error Estimates. In this subsection, we derive the fully discrete optimal error estimates and obtain the following theorem.

Theorem 5. Assuming that $U^0, U^1 \in V_h$ and $Z^0, Z^1 \in W_h$ are given, then for $1 \leq J \leq M$, $j = 0, 1$, one has

$$\begin{aligned} \|u^{J+1} - U^{J+1}\|_j &\leq C(h^{\min(k+1-j, r+1)} + \Delta t^2), \\ \|q^{J+1} - Z^{J+1}\|_j + \|2(q^{J+1} - Z^{J+1}) - (q^J - Z^J)\|_j \\ &\leq C(h^{\min(k+1, r+1-j)} + \Delta t^2). \end{aligned} \quad (47)$$

Proof. Taking $v^h = \zeta^{n+1}$ in (33) and using Cauchy-Schwarz's inequality and Poincaré's inequality, we get

$$\|\zeta^{n+1}\| \leq C\|\zeta_x^{n+1}\| \leq C(\|\rho^{n+1}\| + \|\xi^{n+1}\|). \quad (48)$$

Set $w^h = \xi^{n+1}$ in (34) to obtain

$$\begin{aligned} &\left(\frac{3\xi^{n+1} - 4\xi^n + \xi^{n-1}}{2\Delta t}, \xi^{n+1}\right) + \beta \left(\frac{3\xi_x^{n+1} - 4\xi_x^n + \xi_x^{n-1}}{2\Delta t}, \xi_x^{n+1}\right) \\ &= -\left((1 - \beta\lambda) \frac{3\rho^{n+1} - 4\rho^n + \rho^{n-1}}{2\Delta t} + \tau^{n+1}, \xi^{n+1}\right) \\ &\quad - \beta(\kappa^{n+1}, \xi_x^{n+1}) + \gamma(2\rho^n - \rho^{n-1}, \xi_x^{n+1}) \\ &\quad + \gamma(2\xi^n - \xi^{n-1}, \xi_x^{n+1}) + \gamma(R_1^{n+1}, \xi_x^{n+1}) + \delta(R_2^{n+1}, \xi_x^{n+1}) \\ &\quad + \delta(2(u(t_n)q(t_n) - U^n Z^n) \\ &\quad - (u(t_{n-1})q(t_{n-1}) - U^{n-1} Z^{n-1}), \xi_x^{n+1}). \end{aligned} \quad (49)$$

Use (48) as well as the Cauchy-Schwarz and Young's inequalities to obtain

$$\begin{aligned} &\left(\frac{3\xi^{n+1} - 4\xi^n + \xi^{n-1}}{2\Delta t}, \xi^{n+1}\right) + \beta \left(\frac{3\xi_x^{n+1} - 4\xi_x^n + \xi_x^{n-1}}{2\Delta t}, \xi_x^{n+1}\right) \\ &\leq C(\|\xi^{n+1}\|^2 + \|\xi_x^{n+1}\|^2 + \|\xi^n\|^2 + \|2\rho^n - \rho^{n-1}\|^2 \\ &\quad + \|2\xi^n - \xi^{n-1}\|^2 + \left\|\frac{3\rho^{n+1} - 4\rho^n + \rho^{n-1}}{2\Delta t}\right\|^2 \\ &\quad + \|\tau^{n+1}\|^2 + \|\kappa^{n+1}\|^2 + \|R_1^{n+1}\|^2 + \|R_2^{n+1}\|^2) \\ &\quad + \delta|2(u(t_n)q(t_n) - U^n Z^n) \\ &\quad - (u(t_{n-1})q(t_{n-1}) - U^{n-1} Z^{n-1}), \xi_x^{n+1})|. \end{aligned} \quad (50)$$

Note that

$$\begin{aligned} (a) \quad &\left(\frac{3\xi^{n+1} - 4\xi^n + \xi^{n-1}}{2\Delta t}, \xi^{n+1}\right) \\ &= \frac{1}{4\Delta t} [(\|\xi^{n+1}\|^2 + \|2\xi^{n+1} - \xi^n\|^2) \\ &\quad - (\|\xi^n\|^2 + \|2\xi^n - \xi^{n-1}\|^2) \\ &\quad + \|\xi^{n+1} - 2\xi^n + \xi^{n-1}\|^2], \\ (b) \quad &\beta \left(\frac{3\xi_x^{n+1} - 4\xi_x^n + \xi_x^{n-1}}{2\Delta t}, \xi_x^{n+1}\right) \\ &= \frac{1}{4\Delta t} [(\|\xi_x^{n+1}\|^2 + \|2\xi_x^{n+1} - \xi_x^n\|^2) \\ &\quad - (\|\xi_x^n\|^2 + \|2\xi_x^n - \xi_x^{n-1}\|^2) \\ &\quad + \|\xi_x^{n+1} - 2\xi_x^n + \xi_x^{n-1}\|^2], \\ (c) \quad &\left\|\frac{3\rho^{n+1} - 4\rho^n + \rho^{n-1}}{2\Delta t}\right\|^2 \leq \frac{3}{2\Delta t} \int_{t_n}^{t_{n+1}} \|\rho_t\|^2 ds \\ &\quad + \frac{1}{2\Delta t} \int_{t_{n-1}}^{t_n} \|\rho_t\|^2 ds, \end{aligned} \quad (51)$$

$$\begin{aligned}
& \left| (2(u(t_n)q(t_n) - U^n Z^n) \right. \\
& \quad \left. - (u(t_{n-1})q(t_{n-1}) - U^{n-1} Z^{n-1}), \xi_x^{n+1}) \right| \\
& \leq \left| (2u(t_n)(q(t_n) - Z^n) - u(t_n)(q(t_{n-1}) - Z^{n-1}), \xi_x^{n+1}) \right| \\
& \quad + \left| ((u(t_n) - u(t_{n-1}))(q(t_{n-1}) - Z^{n-1}), \xi_x^{n+1}) \right| \\
& \quad + \left| (2(u(t_n) - U^n)Z^n - (u(t_{n-1}) - U^{n-1})Z^n, \xi_x^{n+1}) \right| \\
& \quad + \left| ((u(t_{n-1}) - U^{n-1})(Z^n - Z^{n-1}), \xi_x^{n+1}) \right| \\
& \doteq T_1 + T_2 + T_3 + T_4.
\end{aligned} \tag{52}$$

We now estimate T_1 , T_2 , T_3 , and T_4 as

$$\begin{aligned}
T_1 & \leq \left| (u(t_n)(2\xi^n - \xi^{n-1}), \xi_x^{n+1}) \right| \\
& \quad + \left| (u(t_n)(2\rho^n - \rho^{n-1}), \xi_x^{n+1}) \right| \\
& \leq \|u(t_n)\|_\infty \|2\xi^n - \xi^{n-1}\| \|\xi_x^{n+1}\| \\
& \quad + \|u(t_n)\|_\infty \|2\rho^n - \rho^{n-1}\| \|\xi_x^{n+1}\|, \\
T_2 & = \left| \left(\int_{t_{n-1}}^{t_n} u_t ds (\rho^{n-1} + \xi^{n-1}), \xi_x^{n+1} \right) \right| \\
& \leq \Delta t \|u_t\|_\infty \|\rho^{n-1} + \xi^{n-1}\| \|\xi_x^{n+1}\|, \\
T_3 & \leq \left| (Z^n(2\zeta^n - \zeta^{n-1}), \xi_x^{n+1}) \right| \\
& \quad + \left| (Z^n(2\eta^n - \eta^{n-1}), \xi_x^{n+1}) \right| \\
& \leq \|Z^n\|_\infty \|2\zeta^n - \zeta^{n-1}\| \|\xi_x^{n+1}\| \\
& \quad + \|Z^n\|_\infty \|2\eta^n - \eta^{n-1}\| \|\xi_x^{n+1}\|, \\
T_4 & \leq \left| \left(\int_{t_{n-1}}^{t_n} Z_t ds (\eta^{n-1} + \zeta^{n-1}), \xi_x^{n+1} \right) \right| \\
& \leq \Delta t \|Z_t\|_\infty \|\eta^{n-1} + \zeta^{n-1}\| \|\xi_x^{n+1}\|.
\end{aligned} \tag{53}$$

Substituting (53) into (52), we obtain

$$\begin{aligned}
& \left| (2(u(t_n)q(t_n) - U^n Z^n) \right. \\
& \quad \left. - (u(t_{n-1})q(t_{n-1}) - U^{n-1} Z^{n-1}), \xi_x^{n+1}) \right| \\
& \leq C \left(\|2\xi^n - \xi^{n-1}\|^2 + \|\xi^{n-1}\|^2 \right. \\
& \quad + \|\rho^n\|^2 + \|\rho^{n-1}\|^2 + \|\zeta^n\|^2 + \|\zeta^{n-1}\|^2 \\
& \quad \left. + \|\eta^n\|^2 + \|\eta^{n-1}\|^2 + \|\xi_x^{n+1}\|^2 \right).
\end{aligned} \tag{54}$$

Substituting (36), (51), and (54) into (50), using (48), and summing from $n = 1, 2, \dots, J$, the resulting inequality becomes

$$\begin{aligned}
& (1 - C\Delta t) \left(\|\xi^{J+1}\|^2 + \|2\xi^{J+1} - \xi^J\|^2 + \|\xi_x^{J+1}\|^2 \right. \\
& \quad \left. + \|2\xi_x^{J+1} - \xi_x^J\|^2 \right) \\
& \leq C\Delta t \sum_{n=1}^J \left(\|\eta^{n-1}\|^2 + \|\rho^{n-1}\|^2 \right) + C \int_0^{t_{J+1}} \|\rho_t\|^2 ds \\
& \quad + C\Delta t^4 \left(\|q_{ttt}\|_{L^\infty(L^2)}^2 + \|q_{ttt}\|_{L^\infty(L^2)}^2 \right. \\
& \quad + \|u_{ttt}\|_{L^\infty(L^2)} + \|u_t q_t\|_{L^\infty(L^2)} + \|u_{tt} q\|_{L^\infty(L^2)} \\
& \quad \left. + h^r \|q_{xttt}\|_{L^\infty(H^{r+1})} + \|q_{xttt}\|_{L^\infty(L^2)} \right) \\
& \quad + C\Delta t \sum_{n=1}^J \left(\|\xi^n\|^2 + \|\xi_x^n\|^2 + \|2\xi^n - \xi^{n-1}\|^2 \right. \\
& \quad \left. + \|2\xi_x^n - \xi_x^{n-1}\|^2 \right).
\end{aligned} \tag{55}$$

Choose Δt_0 in such a way that for $0 < \Delta t \leq \Delta t_0$, $(1 - C\Delta t) > 0$. Then, as an application of Gronwall's lemma, we obtain

$$\begin{aligned}
& \|\xi^{J+1}\|^2 + \|2\xi^{J+1} - \xi^J\|^2 + \|\xi_x^{J+1}\|^2 + \|2\xi_x^{J+1} - \xi_x^J\|^2 \\
& \leq C\Delta t \sum_{n=1}^J \left(\|\eta^{n-1}\|^2 + \|\rho^{n-1}\|^2 \right) + C \int_0^{t_{J+1}} \|\rho_t\|^2 ds \\
& \quad + C\Delta t^4 \left(\|q_{ttt}\|_{L^\infty(L^2)}^2 + \|q_{ttt}\|_{L^\infty(L^2)}^2 + \|u_{ttt}\|_{L^\infty(L^2)} \right. \\
& \quad + \|u_t q_t\|_{L^\infty(L^2)} + \|u_{tt} q\|_{L^\infty(L^2)} \\
& \quad \left. + h^r \|q_{xttt}\|_{L^\infty(H^{r+1})} + \|q_{xttt}\|_{L^\infty(L^2)} \right).
\end{aligned} \tag{56}$$

Combine (28), (31), and (56) with the triangle inequality to complete the L^2 and H^1 error estimates for q . Furthermore, use (48) and the triangle inequality to complete the optimal error estimates for $\|u(t_n) - U^n\|$ and $\|u(t_n) - U^n\|_1$. $\square$

Remark 6. Compared to a variety of difference methods in [17], our method is studied based on mixed element scheme (6) and (7).

Remark 7. Although some convergence proofs of multistep methods for RLW/BBM are provided in [20, 25, 29], our convergence results of multistep methods are proved based on a mixed finite element scheme. Based on the current discussion, we have to provide the detailed proofs for multistep mixed finite element methods in this paper.

The surface of the exact solution u

FIGURE 1: Surface for exact solution u .

The surface of the exact solution q

FIGURE 3: Surface for exact solution q .

The surface of the numerical solution u_h

FIGURE 2: Surface for numerical solution u_h .

The surface of the numerical solution q_h

FIGURE 4: Surface for numerical solution q_h .

4. Numerical Results

In order to test the viability of the proposed method, we consider a test problem. We write conservation laws as [3, 4]

$$\begin{aligned} Q_1 &= \int_a^b u \, dx \approx h \sum_{j=1}^N u_j^n, \\ Q_2 &= \int_a^b (u^2 + \mu(u_x)^2) \, dx \approx h \sum_{j=1}^N \left[(u_j^n)^2 + \mu [(u_x)_j^n]^2 \right], \\ Q_3 &= \int_a^b (u^3 + 3u^2) \, dx \approx h \sum_{j=1}^N \left[(u_j^n)^3 + 3[(u)_j^n]^2 \right], \end{aligned} \quad (57)$$

where Q_1 , Q_2 , and Q_3 are usually called mass, momentum, and energy, respectively, which are observed to check the conservation of the numerical scheme.

We consider RLW equation (1) and let, in (1), $\delta = \gamma = \beta = 1$. Then, the solitary wave solution of (1) is

$$u(x, t) = 3c \operatorname{sech}^2(k[x - x_0 - vt]), \quad (58)$$

where

$$k = \frac{1}{2} \sqrt{\frac{c}{1+c}}, \quad v = 1 + c. \quad (59)$$

We consider the motion of a single solitary wave and take as initial condition, with $c = 0.1$ and $x_0 = 0$,

$$u(x, 0) = 0.3 \operatorname{sech}^2\left(\frac{x}{2\sqrt{11}}\right). \quad (60)$$

The corresponding exact solution with initial condition (60) is

$$u(x, t) = 0.3 \operatorname{sech}^2\left(\frac{x - 1.1t}{2\sqrt{11}}\right). \quad (61)$$

In this procedure, we take space-time domain as $-40 \leq x \leq 60$ and $0 \leq t \leq 20$.

FIGURE 5: Comparison between u and u_h at times $t = 4, 8, 12, 16$, and 20 with $h = 0.125$ and $\Delta t = 0.2$.

FIGURE 6: Comparison between q and q_h at times $t = 4, 8, 12, 16$, and 20 with $h = 0.125$ and $\Delta t = 0.2$.

In Table 1, we take spatial mesh parameter $h = 0.125$ and time discretization parameter $\Delta t = 0.1$ and list the three invariants Q_1 , Q_2 , and Q_3 and the optimal error estimate in L^2 and L^∞ norms for u at different times $t = 0, 4, 8, 12, 16$, and 20 . At the same time, we show some numerical results at time $t = 20$ obtained by other numerical methods in Table 1. From Table 1, we find that our method is more accurate than the numerical methods in [17, 18, 28] but is less than the numerical methods in [23, 24]. From the shown results in Table 1, we can see that Q_1 , Q_2 and Q_3 keep almost constants, so the conservation for our method is very well.

In Tables 2 and 3, we take spatial mesh parameter $h = 0.125$ and obtain the optimal error results in L^2 and L^∞ norms for u at different times $t = 0, 4, 8, 12, 16$, and 20 with different time discretization parameters $\Delta t = 0.1, 0.2$, and 0.4 . From Tables 2 and 3, we see easily that the convergence rate for time is close to order 2. Similarly, the results for q are shown in Tables 4 and 5.

In Tables 6 and 7, the optimal error results in L^2 and L^∞ norms for u at different times $t = 0, 4, 8, 12, 16$, and 20 with different spatial mesh parameters $h = 0.2, 0.4$, and 0.8 and time discretization parameter $\Delta t = 0.01$ are shown. It is easy to see that the convergence rate for space is close to order 2. The similar results for q are listed in Tables 8 and 9.

Figure 1 shows the surface for the exact solution u in space-time domain $((x, t) \in [-20, 60] \times [0, 20])$, and the corresponding surface for the numerical solution u_h with $h = 2$ and $\Delta t = 0.4$ is described in Figure 2. From Figures 1 and 2, we see easily that the exact solution u is approximated very well by the numerical solution u_h . In Figures 3 and 4, we show the surface for the exact solution q and the numerical solution q_h , respectively, with $h = 2$ and $\Delta t = 0.4$ and obtain a good approximation solution q_h for the exact solution q .

The comparison between the exact solution u and the numerical solution u_h is described at different times

FIGURE 7: Comparison between u and u_h at spaces $x = 0, 10, 20$, and 30 with $h = 0.125$ and $\Delta t = 0.2$.

$t = 4, 8, 12, 16$, and 20 with $h = 0.125$ and $\Delta t = 0.2$ in Figure 5. The similar comparison for the exact solution q and the numerical solution q_h is shown in Figure 6. Figures 5 and 6 show that the solitary wave for u and q moves to the right with unchanged form and velocity, respectively. Furthermore, the exact solutions u and q are approximated well by the numerical solutions u_h and q_h , respectively.

In Figure 7, we show the comparison between u and u_h at different spaces $x = 0, 10, 20$, and 30 with $h = 0.125$ and $\Delta t = 0.2$ to verify the efficiency for the proposed scheme in this paper. Figure 8 describes a similar result for q .

FIGURE 8: Comparison between q and q_h at spaces $x = 0, 10, 20$, and 30 with $h = 0.125$ and $\Delta t = 0.2$.

From the previous analysis in Tables 1–9 and Figures 1–8, we can see that the numerical results confirm the theoretical results of Theorem 5 and our method is efficient for RLW equation.

5. Concluding Remarks

In this paper, we propose and analyze an explicit multistep mixed finite element method, which combines spatial mixed finite element method and time explicit multistep method, for RLW equation. We discuss the numerical process for our method, prove the theoretical results for the fully discrete explicit multistep mixed scheme, obtain the optimal convergence order, and compare our method's accuracy with some other numerical schemes. Compared with the numerical method in [20, 25, 26], our method can obtain the optimal error estimates in L^2 and H^1 norms for the scalar unknown u and its flux $q = u_x$ simultaneously. From our numerical results, we can see that our method is efficient for RLW equation.

Acknowledgments

This work was supported by the National Natural Science Fund of China (11061021), the Scientific Research Projection of Higher Schools of Inner Mongolia (NJZZ12011 and NJZY13199), the Natural Science Fund of Inner Mongolia Province (2012MS0108 and 2012MS0106), and the Program of Higher-Level Talents of Inner Mongolia University (125119).

References

[1] D. H. Peregrine, "Calculations of the development of an undular bore," *Journal of Fluid Mechanics*, vol. 25, no. 2, pp. 321–330, 1966.

[2] T. B. Benjamin, J. L. Bona, and J. J. Mahony, "Model equations for long waves in nonlinear dispersive systems," *Philosophical Transactions of the Royal Society of London A*, vol. 272, no. 1220, pp. 47–78, 1972.

[3] P. J. Olver, "Euler operators and conservation laws of the BBM equation," *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 85, no. 1, pp. 143–160, 1979.

[4] B. Saka and I. Dağ, "Quartic B-spline collocation algorithms for numerical solution of the RLW equation," *Numerical Methods for Partial Differential Equations*, vol. 23, no. 3, pp. 731–751, 2007.

[5] J. L. Bona and P. J. Bryant, "A mathematical model for long waves generated by wavemakers in non-linear dispersive systems," *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 73, no. 5, pp. 391–405, 1973.

[6] D. Bhardwaj and R. Shankar, "A computational method for regularized long wave equation," *Computers & Mathematics with Applications*, vol. 40, no. 12, pp. 1397–1404, 2000.

[7] X. H. Zhao, D. S. Li, and D. M. Shi, "A finite difference scheme for RLW-Burgers equation," *Journal of Applied Mathematics & Informatics*, vol. 26, no. 3–4, pp. 573–581, 2008.

[8] L. Zhang, "A finite difference scheme for generalized regularized long-wave equation," *Applied Mathematics and Computation*, vol. 168, no. 2, pp. 962–972, 2005.

[9] I. Dağ, A. Dogan, and B. Saka, "B-spline collocation methods for numerical solutions of the RLW equation," *International Journal of Computer Mathematics*, vol. 80, no. 6, pp. 743–757, 2003.

[10] A. Esen and S. Kutluay, "Application of a lumped Galerkin method to the regularized long wave equation," *Applied Mathematics and Computation*, vol. 174, no. 2, pp. 833–845, 2006.

[11] I. Dağ, B. Saka, and D. Irk, "Galerkin method for the numerical solution of the RLW equation using quintic B-splines," *Journal of Computational and Applied Mathematics*, vol. 190, no. 1–2, pp. 532–547, 2006.

[12] K. R. Raslan, "A computational method for the regularized long wave (RLW) equation," *Applied Mathematics and Computation*, vol. 167, no. 2, pp. 1101–1118, 2005.

[13] L. Wahlbin, "Error estimates for a Galerkin method for a class of model equations for long waves," *Numerische Mathematik*, vol. 23, pp. 289–303, 1975.

[14] Z. D. Luo and R. X. Liu, "Mixed finite element analysis and numerical solitary solution for the RLW equation," *SIAM Journal on Numerical Analysis*, vol. 36, no. 1, pp. 89–104, 1999.

[15] L. Guo and H. Chen, " H^1 -Galerkin mixed finite element method for the regularized long wave equation," *Computing*, vol. 77, no. 2, pp. 205–221, 2006.

[16] H. M. Gu and N. Chen, "Least-squares mixed finite element methods for the RLW equations," *Numerical Methods for Partial Differential Equations*, vol. 24, no. 3, pp. 749–758, 2008.

[17] J. L. Bona, W. G. Pritchard, and L. R. Scott, "Numerical schemes for a model for nonlinear dispersive waves," *Journal of Computational Physics*, vol. 60, no. 2, pp. 167–186, 1985.

[18] S. U. Islam, S. Haq, and A. Ali, "A meshfree method for the numerical solution of the RLW equation," *Journal of Computational and Applied Mathematics*, vol. 223, no. 2, pp. 997–1012, 2009.

[19] D. Kaya, "A numerical simulation of solitary-wave solutions of the generalized regularized long-wave equation," *Applied Mathematics and Computation*, vol. 149, no. 3, pp. 833–841, 2004.

[20] L. Q. Mei and Y. P. Chen, "Explicit multistep method for the numerical solution of RLW equation," *Applied Mathematics and Computation*, vol. 218, no. 18, pp. 9547–9554, 2012.

- [21] L. R. T. Gardner, G. A. Gardner, and A. Dogan, "A least-squares finite element scheme for the RLW equation," *Communications in Numerical Methods in Engineering*, vol. 12, no. 11, pp. 795–804, 1996.
- [22] I. Dağ, "Least squares quadratic B-spline finite element method for the regularized long wave equation," *Computer Methods in Applied Mechanics and Engineering*, vol. 182, no. 1-2, pp. 205–215, 2000.
- [23] I. Dağ and M. N. Özer, "Approximation of the RLW equation by the least square cubic B-spline finite element method," *Applied Mathematical Modelling*, vol. 25, no. 3, pp. 221–231, 2001.
- [24] A. Dogan, "Numerical solution of RLW equation using linear finite elements within Galerkin's method," *Applied Mathematical Modelling*, vol. 26, no. 7, pp. 771–783, 2002.
- [25] P. Chatzipantelidis, "Explicit multistep methods for nonstiff partial differential equations," *Applied Numerical Mathematics*, vol. 27, no. 1, pp. 13–31, 1998.
- [26] G. Akrivis, O. Karakashian, and F. Karakatsani, "Linearly implicit methods for nonlinear evolution equations," *Numerische Mathematik*, vol. 94, no. 3, pp. 403–418, 2003.
- [27] A. K. Pani, R. K. Sinha, and A. K. Otta, "An H^1 -Galerkin mixed method for second order hyperbolic equations," *International Journal of Numerical Analysis and Modeling*, vol. 1, no. 2, pp. 111–129, 2004.
- [28] M. F. Wheeler, "A priori L_2 error estimates for Galerkin approximations to parabolic partial differential equations," *SIAM Journal on Numerical Analysis*, vol. 10, pp. 723–759, 1973.
- [29] J. L. Bona, W. G. Pritchard, and L. R. Scott, "An evaluation of a model equation for water waves," *Philosophical Transactions of the Royal Society of London A*, vol. 302, no. 1471, pp. 457–510, 1981.