Research Article

Stability Analysis and Optimal Control Strategy for Prevention of Pine Wilt Disease

Kwang Sung Lee

Institute of Environmental Technology and Industry, Pusan National University, 30 Geumjeong-Gu, Busan 609-735, Republic of Korea

Correspondence should be addressed to Kwang Sung Lee; ksl@pusan.ac.kr

Received 21 March 2014; Revised 26 May 2014; Accepted 26 May 2014; Published 25 June 2014

Academic Editor: Fasma Diele

Copyright © 2014 Kwang Sung Lee. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We propose a mathematical model of pine wilt disease (PWD) which is caused by pine sawyer beetles carrying the pinewood nematode (PWN). We calculate the basic reproduction number R_0 and investigate the stability of a disease-free and endemic equilibrium in a given mathematical model. We show that the stability of the equilibrium in the proposed model can be controlled through the basic reproduction number R_0 . We then discuss effective optimal control strategies for the proposed PWD mathematical model. We demonstrate the existence of a control problem, and then we apply both analytical and numerical techniques to demonstrate effective control methods to prevent the transmission of the PWD. In order to do this, we apply two control strategies: tree-injection of nematicide and the eradication of adult beetles through aerial pesticide spraying. Optimal prevention strategies can be determined by solving the corresponding optimality system. Numerical simulations of the optimal control problem using a set of reasonable parameter values suggest that reducing the number of pine sawyer beetles is more effective than the tree-injection strategy for controlling the spread of PWD.

1. Introduction

Pine wilt disease (PWD) is caused by the pinewood nematode *Bursaphelenchus xylophilus*. The vector for this parasite is the Japanese pine sawyer beetle *Monochamus alternatus*, which disperses the nematode to healthy trees [1–8]. PWD is lethal to healthy pine trees such as *Pinus densiflora* S. et Z., *P. thunbergii* Parl., and *P. sylvestris* L. [9]. The disease (PWD), caused by the pinewood nematode (PWN), *Bursaphelenchus xylophilus*, was first observed in 1905 in Japan [10]. By the 1980s, the PWD epidemic had spread to other Asian countries such as China, Taiwan, Hong Kong, and Korea. It subsequently reached Europe (Portugal) in 1999 [11–13]. Thus, PWD is the most serious global threat to pine forest ecosystems worldwide.

Pine trees infected by PWD usually die within a few months. The first observable symptom is the lack of resin exudation from bark wounds. The foliage then becomes pale green, then yellow, and finally reddish brown as the tree succumbs to the disease. It is well known that there are three transmission pathways of PWD. One occurs during

maturation feeding, when the nematode is transferred from insect vectors to healthy pine trees via the insect feeding on wounds [4]. The second occurs during oviposition of the mature female on dead, dying, or recently cut pine trees via the oviposition wounds [6]. The third occurs during mating, that is, as mature males search for females in bark wounds, such as the oviposition wounds. This is referred to as horizontal transmission [14].

Vector-borne diseases are infectious diseases caused by viruses, bacteria, protozoa, or rickettsia. They are primarily transmitted by disease transmitting biological agents (arthropods), called vectors, who carry the disease without getting it themselves. For example, the most prevalent vector-borne disease is malaria, whose vectors are mosquitoes. Numerous cases of vector-borne disease are known for plants. Among them, there are some important wilting diseases of trees, such as PWD and the red ring disease of palms. In recent years, many mathematical models have been used to investigate the transmission dynamics of the PWD [15–18]. Mathematical models have become important tools in analyzing the spread and control of infectious diseases. Thus, we present

a mathematical model to describe the vector-host interaction between the pine trees and nematode-carrying pine sawyers. The objective of this study is to investigate global dynamics and control the spread of the PWD based on the proposed mathematical model.

In this paper, we propose a mathematical model of PWD using ordinary differential equations, with mass action incidence, and we consider a controlled PWD transmission model to prevent the spread of disease using optimal control strategies. Generally, PWD control has focused on the elimination of pine sawyer larvae inhabiting pine wilt trees (eradication of infected host pine trees), either by winter fumigation or by controlling the adult sawyers using insecticide over the summer. We apply two control parameters: the injection of nematocide to the host pine tree and the aerial spraying of pesticide to kill adult pine sawyers.

For this, we first introduce control variables representing the optimal treatment for infected hosts and vectors. We formulate an optimal system for the mathematical model of PWD caused by adult pine sawyers carrying pinewood nematodes.

Furthermore, we show the existence of an optimal control for this problem and analyze the system using optimal control techniques.

This paper is organized as follows. In Section 2, we derive the ordinary differential equations that describe the interactions between the host pine tree and vector forest beetles populations. In Section 3, the control problem is formulated. In Section 4, we derive the necessary conditions for an optimal control and the corresponding states using Pontryagin's Maximum Principle. Section 5 presents the results of numerical simulations, and our conclusions are given in Section 6.

2. Model Formulation

In this paper, we consider a four-dimensional model, which consists of the populations of susceptible host pine trees S_h , infected host pine trees I_h , susceptible vector beetles that do not carry the PWN S_v , and infected vector beetles that carry the PWN I_v . The total population sizes at time t for the host pine trees and vector beetles are denoted by N_h and N_v , respectively. In the host pine trees populations, Pine trees infected with the PWD never recovers from an infected states, and most infected pine trees die within a year of infection. Thus we do not consider the number of immune hosts R_h in the model. Thus $N_h = S_h + I_h$ and $N_v = S_v + I_v$. The model is given by the following system of differential equations:

$$\begin{aligned} \frac{dS_h}{dt} &= \Lambda_h - \alpha\phi S_h I_v - \beta\psi\theta S_h I_v - \mu_1 S_h, \\ \frac{dI_h}{dt} &= \alpha\phi S_h I_v + \beta\psi\theta S_h I_v - \mu_1 I_h, \\ \frac{dS_v}{dt} &= \Lambda_v - \gamma I_h S_v - \mu_2 S_v, \\ \frac{dI_v}{dt} &= \gamma I_h S_v - \mu_2 I_v. \end{aligned} \quad (1)$$

Susceptible host pine trees are recruited at a constant rate Λ_h . Parameter α represents the transmission rate per contact with infected vectors during maturation feeding. Parameter ϕ is the average number of contacts per day with vector adult beetles during the maturation feeding period. We denote the incidence of new infections via this route by the mass actions term $\alpha\phi S_h I_v$. The transmission probability that infected beetles transmit a nematode by oviposition is denoted by β , and the average number of contacts per day when adult beetles oviposit is denoted by ψ . The parameter θ is the probability that susceptible host pine trees cease oleoresin exudation without being infected by the nematode. In the model, the rate of transmission through oviposition is denoted by $\beta\psi\theta$, and the incidence of new infections via this route is given by the mass action term $\beta\psi\theta S_h I_v$. A constant emergence rate of the vector pine sawyer beetle is denoted by Λ_v , and parameter γ denotes the rate at which adult beetles carry the PWN when the beetles escape from dead trees. The incidence terms for vector populations are given by $\gamma I_h S_v$. Finally, μ_1, μ_2 denote the natural death rate of the host population and the natural death rate of the vector population, respectively.

Since the model deals with host pine trees and vector beetle populations, it can easily be seen that every state variable will remain nonnegative for nonnegative initial conditions, that is, for all $t \geq 0$, $S_h \geq 0$, $I_h \geq 0$, $S_v \geq 0$, and $I_v \geq 0$. And we make some reasonable technical assumptions on the parameters of the model, namely, $\alpha > 0$, $\beta > 0$, $\gamma > 0$, $\phi > 0$, $\theta > 0$, $\Lambda_h > 0$, $\Lambda_v > 0$, $\mu_1 > 0$, and $\mu_2 > 0$. The above systems for the host population and the vector are also equipped with initial conditions as follows: $S_h(0) = S_h^0$, $E_h(0) = E_h^0$, $I_h(0) = I_h^0$, $S_v(0) = S_v^0$, and $I_v(0) = I_v^0$. The total host population dynamics is given by $dN_h/dt = \Lambda_h - \mu_1 N_h$. The given initial conditions make sure that $N_h(0) \geq 0$. The total dynamics of vector population is $dN_v/dt = \Lambda_v - \mu_2 N_v$. We know that $N_h(t) \rightarrow \Lambda_h/\mu_1$, $N_v(t) \rightarrow \Lambda_v/\mu_2$.

Obviously,

$$\Omega = \left\{ (S_h, I_h, S_v, I_v) \in R_+^4 \mid 0 \leq S_h + I_h \leq \frac{\Lambda_h}{\mu_1}, \right. \\ \left. 0 \leq S_v + I_v \leq \frac{\Lambda_v}{\mu_2} \right\} \quad (2)$$

is positively invariant, and system (1) is dissipative and the global attractor is contained in Ω . On Ω , we have $S_v = (\Lambda_v/\mu_2) - I_v$. Hence, we will study the following three-dimensional nonlinear system:

$$\begin{aligned} \frac{dS_h}{dt} &= \Lambda_h - \alpha\phi S_h I_v - \beta\psi\theta S_h I_v - \mu_1 S_h, \\ \frac{dI_h}{dt} &= \alpha\phi S_h I_v + \beta\psi\theta S_h I_v - \mu_1 I_h, \\ \frac{dI_v}{dt} &= \gamma I_h \left(\frac{\Lambda_v}{\mu_2} - I_v \right) - \mu_2 I_v. \end{aligned} \quad (3)$$

For system (3), the region $\Gamma = \{(S_h, I_h, I_v) \in R_+^3 \mid 0 \leq S_h + I_h \leq \Lambda_h/\mu_1, 0 \leq I_v \leq \Lambda_v/\mu_2\}$ is positively invariant.

System (3) has always a disease-free equilibrium $E_0 = (\Lambda_h/\mu_1, 0, 0)$ and basic reproduction number $R_0 = \Lambda_h \Lambda_v \gamma (\alpha\phi + \beta\psi\theta) / \mu_1^2 \mu_2^2$.

If $R_0 > 1$, system (3) has a unique endemic equilibrium $E^* = (S_h^*, I_h^*, I_v^*)$ in the interior of Γ , where

$$\begin{aligned} S_h^* &= \frac{\Lambda_h}{(\alpha\phi + \beta\psi\theta) I_v^* + \mu_1}, & I_h^* &= \frac{\Lambda_h (\alpha\phi + \beta\psi\theta) I_v^*}{\mu_1 (\alpha\phi + \beta\psi\theta) I_v^* + \mu_1^2}, \\ I_v^* &= \frac{\mu_1^2 \mu_2^2 (R_0 - 1)}{\mu_2 (\alpha\phi + \beta\psi\theta) (\Lambda_h \gamma + \mu_1 \mu_2)}. \end{aligned} \quad (4)$$

Mathematical analysis of the system (3) is provided in Appendices A and B.

3. A Model for Optimal Control of PWD

In the host pine tree population, the force of infections is reduced by $1 - u_1$, where $u_1 \in [0, 1]$ measures the level of successful prevention efforts. It follows that control variable u_1 represents the use of drugs or vaccine which are alternative preventive measures to minimize or eliminate PWN infection (e.g., tree-injection of a nematicide). In system (1), we modified the recruitment rate in susceptible vector populations by including density effects. Further, we replace the previous recruitment rates with $\Lambda_v \rightarrow \Lambda_v N_v$. In the vector adult beetles population, the control variable $u_2 \in [0, 1]$ represents the level of adulticide used for vector control such as aerial spraying of pesticide. It follows that the reproduction rate of the vector population is reduced by a factor of $1 - u_2$. Further, we assumed that the mortality rate of the vector population increases at a rate proportional to u_2 , where $r_0 > 0$ is a rate constant. Considering these assumptions and extensions, it follows that the extended model is described by an initial value problem with a system of four differential equations:

$$\begin{aligned} \frac{dS_h}{dt} &= \Lambda_h - \alpha\phi(1 - u_1) I_v S_h - \beta\psi\theta(1 - u_1) I_v S_h - \mu_1 S_h, \\ \frac{dI_h}{dt} &= \alpha\phi(1 - u_1) I_v S_h + \beta\psi\theta(1 - u_1) I_v S_h - \mu_1 I_h, \\ \frac{dS_v}{dt} &= \Lambda_v N_v (1 - u_2) - \gamma I_h S_v - \mu_2 S_v - r_0 u_2 S_v, \\ \frac{dI_v}{dt} &= \gamma I_h S_v - \mu_2 I_v - r_0 u_2 I_v, \end{aligned} \quad (5)$$

with initial conditions $S_h(0) \geq 0$, $I_h(0) \geq 0$, $S_v(0) \geq 0$, and $I_v(0) \geq 0$.

4. Analysis of Optimal Control

We define the objective functional $J(u_1, u_2) = \int_0^T (A_1 I_h + A_2 N_v + (B_1/2)u_1^2 + (B_2/2)u_2^2) dt$ subject to the state system given by (5). The objective of our work is to minimize

the infected host pine tree and the total number of vector population and the cost of implementing the control by using possible minimal control variables u_1, u_2 . For the optimal control problem of the given system, we consider the control variables $u = (u_1, u_2) \in U$ relative to the state variables S_h, I_h, S_v , and I_v where both control variables are bounded and measured with

$$\begin{aligned} U &= \{(u_1, u_2) \mid u_i \text{ is Lebesgue measurable on } [0, 1], \\ &0 \leq u_i \leq 1, t \in [0, T], i = 1, 2\}. \end{aligned} \quad (6)$$

In the objective functional the quantities A_1, A_2 represent the weight constants of the infected host population and total vector population, respectively. In the objective functional B_1 and B_2 are weight factors for reduction of vector-host contacts and vector control, respectively. The terms $(B_1/2)u_1^2$ and $(B_2/2)u_2^2$ describe the costs associated with prevention of vector-host contacts and vector control, respectively.

The cost associated with the first control could come from the use of tree-injection of a nematicide. The cost associated with the second control could come from physical or chemical treatment of wilt pines to kill their larvae or eradication of vector beetles by aerial pesticide spraying.

Our aim is to find control functions u_1^*, u_2^* such that

$$J(u_1^*, u_2^*) = \min_U J(u_1, u_2) \quad (7)$$

subject to system (5), where the control set is defined as

$$U = \{(u_1(t), u_2(t)) \mid 0 \leq u_1, u_2 \leq 1, t \in [0, T]\}. \quad (8)$$

We note that the existence of an optimal control pair can be proved by using results from Fleming and Rishel [19]. We know easily that the system of equations given by (5) is bounded from above by a linear system. According to the theorem in [19], the solution exists if the following hypotheses are met.

- (H₁) The set of controls and corresponding state variables is nonempty.
- (H₂) The control set U is convex and closed.
- (H₃) Each right-hand side of the state system is bounded by a linear functional in the state and control and can be written as a linear function of u with coefficients depending on time and state.
- (H₄) There exist constants $c_1, c_2 > 0$ and $\beta > 1$ such that the integrand $L(y, u, t)$ of the objective functional J is convex and satisfies

$$L(y, u, t) \geq c_1(|u_1|^2 + |u_2|^2)^{\beta/2} - c_2. \quad (9)$$

In order to verify these conditions, we use a result by Lukes [20] to give the existence of solutions of the state system (5) with bounded coefficients, which gives (H₁). We note that the solutions are bounded. Our control set satisfies condition (H₂). Since our state system is bilinear in u_1, u_2 , the right-hand side the state system (5) satisfies condition (H₃), using the boundedness of the solutions. Note that the integrand of

our objective functional is convex. Moreover we have the last condition needed

$$A_1 I_h + A_2 N_v + \frac{B_1}{2} u_1^2 + \frac{B_2}{2} u_2^2 \geq c_1 (|u_1|^2 + |u_2|^2)^{\beta/2} - c_2, \quad (10)$$

where $c_1, c_2 > 0, A_1, A_2, B_1, B_2 > 0$, and $\beta > 1$. Thus we obtain the following.

Theorem 1. *Given the objective functional $J(u_1, u_2) = \int_0^T (A_1 I_h + A_2 N_v + (B_1/2)u_1^2 + (B_2/2)u_2^2)dt$, where $U = \{(u_1, u_2) \mid 0 \leq u_1, u_2 \leq 1, t \in [0, T]\}$ subject to the system (5) with initial conditions, then there exists an optimal control $u^* = (u_1^*, u_2^*)$ such that $J(u_1^*, u_2^*) = \min_U J(u_1, u_2)$.*

Lagrangian for a problem discusses how the techniques come and Hamiltonian helps in solving the adjoint variable, which we need to construct for the optimal control problem. In order to find an optimal solution pair, first we should construct the Lagrangian and Hamiltonian for the optimal control problem (5). In fact, the Lagrangian of the optimal problem is given by

$$L(I_h, N_v, u_1, u_2) = A_1 I_h + A_2 N_v + \frac{B_1}{2} u_1^2 + \frac{B_2}{2} u_2^2. \quad (11)$$

We seek for the minimal value of the Lagrangian. To do this, we define the Hamiltonian H for the control problem as taking $X = (S_h, I_h, S_v, I_v)$, $U = (u_1, u_2)$, and $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4)$, to obtain

$$H(X, U, \lambda)$$

$$\begin{aligned} &= L(I_h, N_v, u_1, u_2) \\ &+ \lambda_1 [\Lambda_h - \alpha\phi(1 - u_1)I_v S_h - \beta\psi\theta(1 - u_1)I_v S_h - \mu_1 S_h] \\ &+ \lambda_2 [\alpha\phi(1 - u_1)I_v S_h + \beta\psi\theta(1 - u_1)I_v S_h - \mu_1 I_h] \\ &+ \lambda_3 [\Lambda_v N_v(1 - u_2) - \gamma I_h S_v - \mu_2 S_v - r_0 u_2 S_v] \\ &+ \lambda_4 [\gamma I_h S_v - \mu_2 I_v - r_0 u_2 I_v]. \end{aligned} \quad (12)$$

In order to find the necessary conditions for this optimal control pair, we apply Pontryagin's Maximum Principle [21] as follows.

If (u_1^*, u_2^*) is an optimal solution of an optimal control problem, then there exists a nontrivial vector function $\lambda(t) = (\lambda_1(t), \lambda_2(t), \dots, \lambda_n(t))$ satisfying the following equalities. The state equation is

$$\frac{dx}{dt} = \frac{\partial H(t, u_1^*, u_2^*, \lambda(t))}{\partial \lambda}, \quad (13)$$

the optimality condition

$$0 = \frac{\partial H(t, u_1^*, u_2^*, \lambda(t))}{\partial u}, \quad (14)$$

and the adjoint equation

$$\frac{d\lambda}{dt} = -\frac{\partial H(t, u_1^*, u_2^*, \lambda(t))}{\partial x}. \quad (15)$$

Now we apply the necessary conditions to the Hamiltonian H .

Theorem 2. *Given optimal controls u_1^*, u_2^* and solutions S_h^*, I_h^*, S_v^* , and I_v^* of the corresponding state system (5), there exist adjoint variables $\lambda_1, \lambda_2, \lambda_3$, and λ_4 satisfying*

$$\begin{aligned} \frac{d\lambda_1}{dt} &= (\alpha\phi + \beta\psi\theta)(1 - u_1(t))(\lambda_1 - \lambda_2)I_v + \lambda_1\mu_1, \\ \frac{d\lambda_2}{dt} &= -A_1 + \gamma(\lambda_3 - \lambda_4)S_v + \lambda_2\mu_1, \\ \frac{d\lambda_3}{dt} &= \gamma(\lambda_3 - \lambda_4)I_h + \lambda_3(\mu_2 + r_0 u_2(t)) - \lambda_3\Lambda_v(1 - u_2), \\ \frac{d\lambda_4}{dt} &= (\alpha\phi + \beta\psi\theta)(1 - u_1(t))(\lambda_1 - \lambda_2)S_h \\ &+ \lambda_4(\mu_2 + r_0 u_2(t)) - \lambda_3\Lambda_v(1 - u_2), \end{aligned} \quad (16)$$

with transversality conditions

$$\lambda_1(T) = \lambda_2(T) = \lambda_3(T) = \lambda_4(T) = 0. \quad (17)$$

Furthermore, u_1^*, u_2^* are represented by

$$\begin{aligned} u_1^* &= \max \left\{ \min \left\{ 1, \frac{1}{B_1}(\lambda_2 - \lambda_1)(\alpha\phi + \beta\psi\theta)I_v S_h \right\}, 0 \right\}, \\ u_2^* &= \max \left\{ \min \left\{ 1, \frac{1}{B_2}[\Lambda_v \lambda_3(S_v + I_v) \right. \right. \\ &\quad \left. \left. + r_0(\lambda_3 S_v + \lambda_4 I_v)] \right\}, 0 \right\}. \end{aligned} \quad (18)$$

Proof. To determine the adjoint equations and the transversality conditions we use the Hamilton H . We differentiate the Hamiltonian H with respect to S_h, I_h, S_v, I_v . Thus we obtain

$$\begin{aligned} \frac{d\lambda_1}{dt} &= (\alpha\phi + \beta\psi\theta)(1 - u_1(t))(\lambda_1 - \lambda_2)I_v + \lambda_1\mu_1, \\ \frac{d\lambda_2}{dt} &= -A_1 + \gamma(\lambda_3 - \lambda_4)S_v + \lambda_2\mu_1, \\ \frac{d\lambda_3}{dt} &= \gamma(\lambda_3 - \lambda_4)I_h + \lambda_3(\mu_2 + r_0 u_2(t)) - \lambda_3\Lambda_v(1 - u_2), \\ \frac{d\lambda_4}{dt} &= (\alpha\phi + \beta\psi\theta)(1 - u_1(t))(\lambda_1 - \lambda_2)S_h \\ &+ \lambda_4(\mu_2 + r_0 u_2(t)) - \lambda_3\Lambda_v(1 - u_2), \end{aligned} \quad (19)$$

with transversality conditions

$$\lambda_1(T) = \lambda_2(T) = \lambda_3(T) = \lambda_4(T) = 0. \quad (20)$$

Using the optimality conditions we have and the property of the control space U we can drive

$$\begin{aligned} u_1^* &= \max \left\{ \min \left\{ 1, \frac{1}{B_1} (\lambda_2 - \lambda_1) (\alpha\phi + \beta\psi\theta) I_v S_h \right\}, 0 \right\}, \\ u_2^* &= \max \left\{ \min \left\{ 1, \frac{1}{B_2} [\Lambda_v \lambda_3 (S_v + I_v) \right. \right. \\ &\quad \left. \left. + r_0 (\lambda_3 S_v + \lambda_4 I_v)] \right\}, 0 \right\}. \end{aligned} \quad (21)$$

5. Numerical Results

In this section, we show the numerical simulations of the impacts of the optimal control strategy in the proposed the PWD model. The optimality system is solved using the Runge-Kutta fourth order scheme. The optimal strategy is obtained by solving the state and adjoint systems and the transversality conditions. First we start to solve the state (16) using the Runge-Kutta fourth order forward in time with a guess for the controls over the simulated time. Then, using the current iteration of the state equations, the adjoint equations in system (16) are solved by a backward method with the transversality conditions (17). Then the controls are updated by using a convex combination of the previous controls and the value from the characterizations (18). This process is repeated and iterations stopped if the values of the unknowns at the previous iterations are very close to the ones at the present iterations [21]. We have chosen the same set of weight factors $A_1 = 0.01$, $A_2 = 0.001$, $B_1 = 50$, and $B_2 = 200$ initial state variables $S_h(0) = 300$, $I_h(0) = 50$, $S_v(0) = 65$, and $I_v(0) = 20$ to illustrate the effect of different optimal control strategies on the spread of pine wilt disease in a population. In this section, we investigate numerically the effect of the following optimal control strategies on the spread of pine wilt disease. Three different control strategies are explored. Then we look at the following three alternatives:

- (i) strategy 1: precaution effort by tree-injection of a nematicide and eradication of vector beetles (controls u_1 and u_2);
- (ii) strategy 2: physical or chemical treatment of wilt pines to kill their larvae or eradication of vector beetles by aerial pesticide spraying (control u_2 alone);
- (iii) strategy 3: precaution effort by tree-injection of a nematicide (control u_1 alone).

Baseline values of model parameters used for simulations are presented in Table 1. Owing to the absence of data, some of the other parameters associated with given model are assumed (see Table 1).

5.1. Strategy 1: Optimal Use of Tree-Injection of a Nematicide ($u_1 \neq 0$) and Eradication of Vector Beetles ($u_2 \neq 0$). The computed results for optimal control under strategy 1

($u_1 \neq 0$, $u_2 \neq 0$) are shown in Figures 1, 2, 3, and 4. In Figures 2, 3, and 4, we can see that the control strategy results in a decrease in the number of infected hosts I_h , susceptible vectors S_v , and infected vectors I_v . With this strategy, an increase in the number of susceptible hosts S_h is observed in Figure 1.

5.2. Strategy 2: Optimal Use of Eradication of Vector Beetles ($u_2 \neq 0$). The computed results for optimal control under strategy 2 ($u_1 = 0$, $u_2 \neq 0$) are shown in Figures 5, 6, 7, and 8. Figures 6, 7, and 8 show a significant difference in the number of infected hosts, susceptible vectors, and infected vectors, respectively, compared to the situation where there is no control. Our numerical results suggest that tree-injection of a nematicide is not effective if we want to reduce the spread of PWD.

5.3. Strategy 3: Optimal Use of Tree-Injection of a Nematicide ($u_1 \neq 0$). The computed results for optimal control under strategy 3 ($u_1 \neq 0$, $u_2 = 0$) are shown in Figures 9, 10, 11, and 12. Figures 10 and 12 show no significant difference in the number of infected hosts I_h and infected vectors I_v , respectively, compared to the situation where there is no control. This shows that the eradication of vector beetles is more effective than injecting trees with nematicide. Figure 13 represents the optimal controls.

5.4. Effects of Weight Constants. A sensitivity analysis is carried out by studying the adequacy of our simulations in relation to the weight constants. The role of weight constants is explored by assessing their quantitative impact in terms of the number of infected cases. Comparative results with the implementation of strategy 1 under different weight constants on the controls u_1, u_2 are shown in Figures 14 and 16 ($B_1 = 50$, $B_1 = 5$, $B_1 = 2$, fixing $B_2 = 200$) and Figures 15 and 17 (fixing $B_1 = 50$, $B_2 = 200$, $B_2 = 100$, and $B_2 = 50$). In Figure 16, the value of the weight constant B_1 is varied. As the value of weight constants increases, control functions decrease, and increasing the cost of successful prevention efforts, the population of infected host I_h is increased. In Figure 17, the value of the weight constant B_2 is varied. As the value of weight constants increases, control functions decrease, and increasing the cost of vector control, the population of infected vector I_v is increased.

6. Conclusions

In this paper, we proposed a host vector transmission model to study the dynamics of PWD. A mathematical analysis was carried out for a model in which PWD is caused by the PWN, a parasite hosted by the pine sawyer beetle. The global dynamics of the model showed the existence and stability of disease-free and endemic equilibria. We proved that if $R_0 \leq 1$, the disease-free equilibrium E_0 is globally asymptotically stable in Γ , and thus the disease always dies out. However, if $R_0 > 1$, the unique endemic equilibrium E^* exists and is globally asymptotically stable, indicating that PWD persists at the endemic equilibrium if it is initially present. We have

TABLE 1: Parameters used for numerical simulation.

Notation	Parameter description	Estimated value	Ref.
Λ_h	The recruitment rate of the host pine population	$0.009041 \text{ day}^{-1}$	Assumed
Λ_v	A constant emergence rate of the vector pine sawyer beetle	$0.002691 \text{ day}^{-1}$	Assumed
μ_1	The natural death rate of host population	$0.0000301 \text{ day}^{-1}$	[22]
μ_2	The natural death rate of vector population	$0.011764 \text{ day}^{-1}$	[23]
α	The rate in which infected beetles transmit nematode by contact	0.00166 day^{-1}	[24]
β	The rate in which infected beetles transmit nematode by oviposition	0.0004 day^{-1}	[24]
γ	The rate in which the adult beetles have pinewood nematode when it escapes from dead trees	0.00305 day^{-1}	[25]
ϕ	The number of contacts during maturation feeding period	0.2 day	[26]
ψ	The number of contacts during the oviposition period	0.41 day^{-1}	Assumed
θ	The probability in which susceptible host pine is not infectious by nematode and ceases oleoresin exudation naturally	$0.0000301 \text{ day}^{-1}$	Assumed

FIGURE 1: Plot of the number of susceptible hosts with and without control, $u_1 \neq 0$, $u_2 \neq 0$. FIGURE 2: Plot of the number of infected hosts with and without control, $u_1 \neq 0$, $u_2 \neq 0$.

considered the optimal control strategy for preventing the spread of PWD. The conditions for optimal control of the disease were derived and analyzed, and the injection of nematicide to infected hosts was compared with the aerial spraying of insecticides to control infected vectors. From the control plots, we showed that the number of infected hosts and the total vector population decreased in the optimal system. Control programs that follow these strategies can effectively reduce the spread of PWD in forest ecosystems. Numerical simulations illustrated the effectiveness and efficiency of the proposed control problem more effective than

the tree-injection strategy for controlling the spread of PWD in the forest ecosystem.

Appendices

A. The Disease-Free Equilibrium and Its Stability

In this section, we will investigate the disease-free equilibrium and its stability of the system (3).

FIGURE 3: Plot of the number of susceptible vectors with and without control, $u_1 \neq 0, u_2 \neq 0$.

FIGURE 4: Plot of the number of infected vectors with and without control, $u_1 \neq 0, u_2 \neq 0$.

Theorem A.1. *If $R_0 < 1$, then the disease-free equilibrium E_0 is locally asymptotically stable.*

Proof. Linearize the system (3) around the disease-free equilibrium E_0 . The matrix of the linearization at E_0 is given by

$$J(E_0) = \begin{pmatrix} -\mu_1 & 0 & -\frac{\Lambda_h(\alpha\phi + \beta\psi\theta)}{\mu_1} \\ 0 & -\mu_1 & \frac{\Lambda_h(\alpha\phi + \beta\psi\theta)}{\mu_1} \\ 0 & \frac{\Lambda_v\gamma}{\mu_2} & -\mu_2 \end{pmatrix}. \quad (\text{A.1})$$

The characteristic equation of this matrix is given by $\det(\lambda I - J(E_0)) = 0$, where I is the 3×3 unit matrix. Expanding the determinant into a characteristic equation we obtain the following equation:

$$(\lambda + \mu_1) \left[\lambda^2 + (\mu_1 + \mu_2)\lambda + \mu_1\mu_2 - \frac{\Lambda_h\Lambda_v\gamma(\alpha\phi + \beta\psi\theta)}{\mu_1\mu_2} \right] = 0. \quad (\text{A.2})$$

So, $\lambda = -\mu_1$ and $\lambda^2 + a_1\lambda + a_2 = 0$, where

$$a_1 = \mu_1 + \mu_2 > 0,$$

$$a_2 = \mu_1\mu_2 - \frac{\Lambda_h\Lambda_v\gamma(\alpha\phi + \beta\psi\theta)}{\mu_1\mu_2} = \mu_1\mu_2(1 - R_0) > 0. \quad (\text{A.3})$$

FIGURE 5: Plot of the number of susceptible hosts with and without control, $u_1 = 0, u_2 \neq 0$.

Thus, according to the Routh-Hurwitz criteria [27], the quadratic equation has only roots with negative real parts. Hence, the disease-free equilibrium E_0 of the system (3) is locally asymptotically stable. $\square$

Now, we study the global behavior of the disease-free equilibrium for system (3).

FIGURE 6: Plot of the number of infected hosts with and without control, $u_1 = 0$, $u_2 \neq 0$.

FIGURE 8: Plot of the number of infected vectors with and without control, $u_1 = 0$, $u_2 \neq 0$.

FIGURE 7: Plot of the number of susceptible vectors with and without control, $u_1 = 0$, $u_2 \neq 0$.

Theorem A.2. *If $R_0 \leq 1$, then the disease-free equilibrium E_0 is globally asymptotically stable in Γ .*

Proof. We define the following Lyapunov function on Γ .

$$V = b_1 I_h + b_2 I_v, \quad (\text{A.4})$$

where

$$\begin{aligned} b_1 &= \mu_1 \mu_2, \\ b_2 &= \Lambda_h (\alpha \phi + \beta \psi \theta). \end{aligned} \quad (\text{A.5})$$

Clearly, $V \geq 0$ along the solution of the system (3) and is zero if and only if both I_h and I_v are zeros. The derivative of V along the solutions of (3) is

$$\begin{aligned} V' &= \mu_1 \mu_2 I_h' + \Lambda_h (\alpha \phi + \beta \psi \theta) I_v' \\ &= \mu_1 \mu_2 (\alpha \phi + \beta \psi \theta) S_h I_v - \mu_1^2 \mu_2 I_h \\ &\quad + \frac{\Lambda_h \Lambda_v \gamma (\alpha \phi + \beta \psi \theta)}{\mu_2} I_h \\ &\quad - \Lambda_h \gamma (\alpha \phi + \beta \psi \theta) I_v I_h - \Lambda_v \mu_2 (\alpha \phi + \beta \psi \theta) I_v. \end{aligned} \quad (\text{A.6})$$

We know that

$$\begin{aligned} &\mu_1 \mu_2 (\alpha \phi + \beta \psi \theta) S_h I_v - \Lambda_h \mu_2 (\alpha \phi + \beta \psi \theta) I_v \\ &= \mu_1 \mu_2 (\alpha \phi + \beta \psi \theta) \left(S_h - \frac{\Lambda_h}{\mu_1} \right) I_v \leq 0. \end{aligned} \quad (\text{A.7})$$

Thus

$$\begin{aligned} V' &\leq \frac{\Lambda_h \Lambda_v \gamma (\alpha \phi + \beta \psi \theta)}{\mu_2} I_h - \Lambda_h \gamma (\alpha \phi + \beta \psi \theta) I_v I_h - \mu_1^2 \mu_2 I_h \\ &= I_h [\mu_1^2 \mu_2 (R_0 - 1) - \Lambda_h \gamma (\alpha \phi + \beta \psi \theta) I_v] \leq 0. \end{aligned} \quad (\text{A.8})$$

FIGURE 9: Plot of the number of susceptible hosts with and without control, $u_1 \neq 0, u_2 = 0$.

FIGURE 11: Plot of the number of susceptible vectors with and without control, $u_1 \neq 0, u_2 = 0$.

FIGURE 10: Plot of the number of infected hosts with and without control, $u_1 \neq 0, u_2 = 0$.

FIGURE 12: Plot of the number of infected vectors with and without control, $u_1 \neq 0, u_2 = 0$.

We see that if $R_0 < 1$, the derivative $V' = 0$ if and only if $I_h = 0$, while in the case of $R_0 = 1$ the derivative $V' = 0$ if and only if $I_h = 0$ or $I_v = 0$. Consequently, the largest compact invariant set in $\{(S_h, I_h, I_v) \in \Gamma \mid V' = 0\}$, when $R_0 \leq 1$, is the singleton $\{E_0\}$. Hence by Lasalle's Invariance Principle [28], E_0 is globally asymptotically stable in Γ . This completes the proof. $\square$

B. The Endemic Equilibrium and Its Stability

In this section, we will investigate the endemic equilibrium and its stability of the system (3). We first give the following results.

FIGURE 13: Plot of the optimal controls, $u_1 \neq 0$, $u_2 \neq 0$. FIGURE 14: Plot of different weight constants on the control ($B_1 = 50$, $B_1 = 5$, and $B_1 = 2$). FIGURE 15: Plot of the number of different weight constants on the control ($B_2 = 200$, $B_2 = 100$, and $B_2 = 50$).

Lemma B.1. Let M be a 3×3 real matrix. If $\text{tr}(M)$, $\det(M)$, and $\det(M^{[2]})$ are all negative, then all eigenvalues of M have negative real part.

Proof. The proof of this lemma is given in [29]. $\square$

Theorem B.2. If $R_0 \geq 1$, then the endemic equilibrium E^* is locally asymptotically stable.

Proof. In order to investigate the local stability of the endemic equilibrium, the additive compound matrices approach as in [30, 31] is used. We will linearize system (3) about an endemic equilibrium E^* and get the following Jacobian matrix:

$$J(E^*) = \begin{pmatrix} -(\alpha\phi + \beta\psi\theta)I_v^* - \mu_1 & 0 & -(\alpha\phi + \beta\psi\theta)S_h^* \\ (\alpha\phi + \beta\psi\theta)I_v^* & -\mu_1 & (\alpha\phi + \beta\psi\theta)S_h^* \\ 0 & \frac{\Lambda_v\gamma}{\mu_2} & -\gamma I_h^* - \mu_2 \end{pmatrix}. \quad (\text{B.1})$$

FIGURE 16: Plot of different weight constants on the control (without control, $B_1 = 50$, $B_1 = 5$, and $B_1 = 2$).

FIGURE 17: Plot of the number of different weight constants on the control (without control, $B_2 = 200$, $B_2 = 100$, and $B_2 = 50$).

From the Jacobian matrix $J(E^*)$, the second additive compound matrix $J^{[2]}(E^*)$ is given by

$$J^{[2]}(E^*) = \begin{pmatrix} -(\alpha\phi + \beta\psi\theta)I_v^* - 2\mu_1 & (\alpha\phi + \beta\psi\theta)S_h^* & (\alpha\phi + \beta\psi\theta)S_h^* \\ \frac{\Lambda_v\gamma}{\mu_2} & -(\alpha\phi + \beta\psi\theta)I_v^* - \gamma I_h^* - \mu_1 - \mu_2 & 0 \\ 0 & (\alpha\phi + \beta\psi\theta)I_v^* & -\gamma I_h^* - \mu_1 - \mu_2 \end{pmatrix}. \quad (B.2)$$

From the Jacobian matrix $J(E^*)$, we have

$$\text{tr}(J(E^*)) = -[(\alpha\phi + \beta\psi\theta)I_v^* + \gamma I_h^* + 2\mu_1 + \mu_2] < 0. \quad (B.3)$$

And we know that system (3) is given

$$\begin{aligned} \mu_1 &= \frac{(\alpha\phi + \beta\psi\theta)S_h^* I_v^*}{I_h^*}, \\ \mu_2 &= \frac{\gamma I_h^* ((\Lambda_v/\mu_2) - I_v^*)}{I_v^*}. \end{aligned} \quad (B.4)$$

Thus,

$$\begin{aligned} \det(J(E^*)) &= -[(\alpha\phi + \beta\psi\theta)I_v^* + \mu_1] \\ &\quad \times \left[\mu_1(\gamma I_h^* + \mu_2) - \frac{\Lambda_v\gamma}{\mu_2}(\alpha\phi + \beta\psi\theta)S_h^* \right] \\ &\quad - \frac{\Lambda_v\gamma}{\mu_2}(\alpha\phi + \beta\psi\theta)^2 I_v^* S_h^* = -\frac{\Lambda_v\gamma}{\mu_2}(\alpha\phi + \beta\psi\theta)^2 I_v^* S_h^* < 0, \end{aligned} \quad (B.5)$$

since we know that

$$\mu_1\mu_2 = (\alpha\phi + \beta\psi\theta)\gamma \left(\frac{\Lambda_v}{\mu_2} - I_v^* \right) S_h^*. \quad (B.6)$$

So

$$\mu_1(\gamma I_h^* + \mu_2) - \frac{\Lambda_v\gamma}{\mu_2}(\alpha\phi + \beta\psi\theta)S_h^* = 0. \quad (B.7)$$

Computing directly the determinant of $J^{[2]}(E^*)$ and from (B.4) and (B.6) we can get

$$\begin{aligned} \det(J^{[2]}(E^*)) &= -(\gamma I_h^* + \mu_1 + \mu_2) \left[[(\alpha\phi + \beta\psi\theta)I_v^* + 2\mu_1] \right. \\ &\quad \times ((\alpha\phi + \beta\psi\theta)I_v^* + \gamma I_h^* + \mu_1 + \mu_2) \\ &\quad \left. - \frac{\Lambda_v\gamma}{\mu_2}(\alpha\phi + \beta\psi\theta)S_h^* \right] \\ &\quad + \frac{\Lambda_v\gamma}{\mu_2}(\alpha\phi + \beta\psi\theta)^2 S_h^* I_v^* \end{aligned}$$

$$\begin{aligned}
&= -(\gamma I_h^* + \mu_1 + \mu_2) \left[((\alpha\phi + \beta\psi\theta) I_v^* + \mu_1)^2 \right. \\
&\quad \left. + ((\alpha\phi + \beta\psi\theta) I_v^* + \mu_1) (\gamma I_h^* + \mu_2) \right. \\
&\quad \left. + \mu_1 (\alpha\phi + \beta\psi\theta) I_v^* \right] \\
&\quad + \frac{\Lambda_v \gamma}{\mu_2} (\alpha\phi + \beta\psi\theta)^2 S_h^* I_v^* \\
&= -(\gamma I_h^* + \mu_1 + \mu_2) [(\alpha\phi + \beta\psi\theta) I_v^* + \mu_1] \\
&\quad \times [(\alpha\phi + \beta\psi\theta) I_v^* + \mu_1 + \gamma I_h^* + \mu_2] \\
&\quad - \mu_1 (\gamma I_h^* + \mu_1 + \mu_2) (\alpha\phi + \beta\psi\theta) I_v^* \\
&\quad + (\alpha\phi + \beta\psi\theta)^2 \frac{\Lambda_v \gamma}{\mu_2} S_h^* I_v^* \\
&= -(\gamma I_h^* + \mu_1 + \mu_2) [(\alpha\phi + \beta\psi\theta) I_v^* + \mu_1] \\
&\quad \times [(\alpha\phi + \beta\psi\theta) I_v^* + \mu_1 + \gamma I_h^* + \mu_2] \\
&\quad - (\alpha\phi + \beta\psi\theta) I_v^* \mu_1^2 < 0.
\end{aligned} \tag{B.8}$$

Hence by Lemma B.1, the endemic equilibrium E^* of the model (3) is locally asymptotically stable in Γ . $\square$

Now we prove the global stability of the endemic equilibrium E^* , when the reproduction number R_0 is greater than the unity. For this, we will prove the following result.

Theorem B.3. *If $R_0 > 1$, then system (3) is uniformly persistent; that is, there exists $c > 0$ (independent of initial conditions), such that $\liminf_{t \rightarrow \infty} E_h(t) \geq c$, $\liminf_{t \rightarrow \infty} I_h(t) \geq c$, and $\liminf_{t \rightarrow \infty} I_v(t) \geq c$.*

Proof. Let π be a semidynamical system (3) in $(R_0^+)^3$, χ a locally compact metric space, and $\Gamma_0 = \{(S_h, I_h, I_v) \in \Gamma \mid I_v = 0\}$. The set Γ_0 is a compact subset of Γ and Γ/Γ_0 is positively invariant set of system (3). Let $P : \chi \rightarrow R_0^+$ be defined by $P(E_h, S_h, I_v) = I_v$ and set $S = \{(S_h, I_h, I_v) \in \Gamma \mid P(E_h, I_h, I_v) < \rho\}$, where ρ is sufficiently small so that $R_0(1 - (\mu_2/\Lambda_v)\rho) > 1$. Assume that there is a solution $x \in S$ such that, for each $t > 0$, we have $P(\pi(x, t)) < P(x) < \rho$. Let us consider the following:

$$L(t) = \frac{\Lambda_h (\alpha\phi + \beta\psi\theta) (1 - \delta^*)}{\mu_1 \mu_2} I_v + I_h, \tag{B.9}$$

where $\delta^* > 0$ is a sufficiently small constant so that $R_0(1 - (\mu_2/\Lambda_v)\rho)(1 - \delta^*) > 1$. By a direct calculation, we have

$$\begin{aligned}
&L'(t) \\
&\geq \mu_1 \left[\frac{\Lambda_h \Lambda_v \gamma (\alpha\phi + \beta\psi\theta) (1 - \delta^*) (1 - (\mu_2 \rho / \Lambda_v))}{\mu_1^2 \mu_2^2} - 1 \right] I_h \\
&\quad + \frac{\Lambda_h (\alpha\phi + \beta\psi\theta) \delta^*}{\mu_1} I_v.
\end{aligned} \tag{B.10}$$

Let

$$\begin{aligned}
&\delta \\
&= \min \left\{ \mu_1 \left[\frac{\Lambda_h \Lambda_v \gamma (\alpha\phi + \beta\psi\theta) (1 - \delta^*) (1 - (\mu_2 \rho / \Lambda_v))}{\mu_1^2 \mu_2^2} \right. \right. \\
&\quad \left. \left. - 1 \right], \frac{\mu_2 \delta^*}{1 - \delta^*} \right\} > 0.
\end{aligned} \tag{B.11}$$

Thus, we have

$$L'(t) \geq \delta L(t). \tag{B.12}$$

The inequality (B.12) implies that $L(t) \rightarrow \infty$ as $t \rightarrow \infty$. However, $L(t)$ is bounded on the set Γ . According to Theorem 1 in [32], we complete the proof of Theorem 2. $\square$

Finally, we will investigate the global stability of the endemic equilibrium E^* in the feasible region Γ . To do this we will use the results about the three dimensionless competitive systems that live in convex sets [33–35] and powerful theory of additive compound matrix to prove asymptotic orbital stability of periodic solutions [30]. We easily can get that the system (3) is competitive in Γ , by looking at its Jacobian matrix and choosing the matrix H as

$$H = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{B.13}$$

with respect to the partial order defined by the orthant

$$\Delta = \{(S_h, I_h, I_v) \in R^3 \mid S_h \geq 0, I_h \leq 0, I_v \geq 0\}. \tag{B.14}$$

Theorem B.4. *If $R_0 > 1$, then the endemic equilibrium E^* is globally asymptotically stable in $\text{int } \Gamma$.*

Since system (3) is competitive and persistent, and E^* is locally asymptotically stable if $R_0 > 1$. Furthermore, in accordance with Theorem 2.1 and Theorem 2.2 [35], Theorem 600 would be established if we show that system (3) has the property of stability of periodic orbits. Hence we will prove the following theorem.

Theorem B.5. *If $R_0 > 1$, then system (3) has the property of stability of periodic orbits.*

Proof. In accordance with the criterion given by Li and Muldowney [31], for the asymptotic orbital stability of a periodic orbit of a general autonomous system, it is sufficient to prove that the linear nonautonomous system

$$X'(t) = J(E^{[2]}(p(t))) X(t) \tag{B.15}$$

is asymptotically stable, where $J(E^{[2]})$ is the second additive compound matrix of the Jacobian $J(E)$. The Jacobian of system (3) is given by

$$J(E) = \begin{pmatrix} -(\alpha\phi + \beta\psi\theta)I_v - \mu_1 & 0 & -(\alpha\phi + \beta\psi\theta)S_h \\ (\alpha\phi + \beta\psi\theta)I_v & -\mu_1 & (\alpha\phi + \beta\psi\theta)S_h \\ 0 & \frac{\Lambda_v\gamma}{\mu_2} & -\gamma I_h - \mu_2 \end{pmatrix}. \quad (\text{B.16})$$

For the solution $p(t)$, (B.15) becomes

$$\begin{aligned} X' &= -[(\alpha\phi + \beta\psi\theta)I_v + 2\mu_1]X \\ &\quad + (\alpha\phi + \beta\psi\theta)S_h Y + (\alpha\phi + \beta\psi\theta)S_h Z, \\ Y' &= \frac{\Lambda_v\gamma}{\mu_2}X - ((\alpha\phi + \beta\psi\theta)I_v + \gamma I_h + \mu_1 + \mu_2)Y, \\ Z' &= (\alpha\phi + \beta\psi\theta)I_v Y - (\gamma I_h + \mu_1 + \mu_2)Z. \end{aligned} \quad (\text{B.17})$$

In order to prove that (B.15) is asymptotically stable, we will use the following Lyapunov function which is similar to the one found in [31] for the SEIR model. Let

$$\begin{aligned} V(X(t), Y(t), Z(t), S_h, I_h, I_v) \\ = \left\| \left(X(t), \frac{I_h}{I_v} Y(t), \frac{I_h}{I_v} Z(t) \right) \right\|, \end{aligned} \quad (\text{B.18})$$

where $\|\cdot\|$ is the norm in R^3 defined by

$$\|(X, Y, Z)\| = \sup\{|X|, |Y + Z|\}. \quad (\text{B.19})$$

Since the given system is uniformly persistence, we obtain that the orbit of $p(t)$ remains at a positive distance from the boundary of Γ . Therefore, we have $I_h(t) \geq c$, $I_h(t) \geq c$ with $0 < c < \min\{\Lambda_h/\mu_1, \Lambda_v/\mu_2\}$. Hence, the function $V(t)$ is well defined along $p(t)$ and

$$V(X, Y, Z, S_h, I_h, I_v) \geq c \|(X, Y, Z)\|. \quad (\text{B.20})$$

Along the positive solution (X, Y, Z) of the system, $V(t)$ becomes

$$V(t) = \sup\left\{|X|, \frac{I_h}{I_v}(|Y + Z|)\right\}. \quad (\text{B.21})$$

Similarly as was done in [31], we found the following inequalities:

$$\begin{aligned} D_+ |X| &\leq -((\alpha\phi + \beta\psi\theta)I_v + 2\mu_1)|X| \\ &\quad + (\alpha\phi + \beta\psi\theta)S_h |Y + Z| \\ &= -((\alpha\phi + \beta\psi\theta)I_v + 2\mu_1)|X| \\ &\quad + (\alpha\phi + \beta\psi\theta)S_h \frac{I_v}{I_h} \left(\frac{I_h}{I_v}(|Y| + |Z|) \right), \\ D_+ |Y| &\leq \frac{\Lambda_v\gamma}{\mu_2}|X| - ((\alpha\phi + \beta\psi\theta)I_v + \gamma I_h + \mu_1 + \mu_2)|Y|, \\ D_+ |Z| &\leq (\alpha\phi + \beta\psi\theta)I_v |Y| - (\gamma I_h + \mu_1 + \mu_2)|Z|. \end{aligned} \quad (\text{B.22})$$

We know that

$$D_+ (|Y + Z|) \leq \frac{\Lambda_v\gamma}{\mu_2}|X| - (\gamma I_h + \mu_1 + \mu_2)(|Y| + |Z|). \quad (\text{B.23})$$

Therefore

$$\begin{aligned} D_+ \left(\frac{I_h}{I_v} (|Y + Z|) \right) \\ = \left(\frac{I_h'}{I_h} - \frac{I_v'}{I_v} \right) \frac{I_h}{I_v} (|Y + Z|) + \frac{I_h}{I_v} D_+ (|Y + Z|) \\ \leq \left(\frac{I_h'}{I_h} - \frac{I_v'}{I_v} \right) \frac{I_h}{I_v} (|Y + Z|) \\ + \frac{I_h}{I_v} \left[\frac{\Lambda_v\gamma}{\mu_2}|X| - (\gamma I_h + \mu_1 + \mu_2)|Y + Z| \right] \\ = \frac{I_h}{I_v} \frac{\gamma\Lambda_v}{\mu_2}|X| + \left(\frac{I_h'}{I_h} - \frac{I_v'}{I_v} - \gamma I_h - \mu_1 - \mu_2 \right) \frac{I_h}{I_v} |Y + Z|. \end{aligned} \quad (\text{B.24})$$

Thus we get

$$D_+ V(t) \leq \sup\{g_1(t), g_2(t)\} V(t), \quad (\text{B.25})$$

where

$$\begin{aligned} g_1(t) &= -((\alpha\phi + \beta\psi\theta)I_v + 2\mu_1) + (\alpha\phi + \beta\psi\theta)S_h \frac{I_v}{I_h}, \\ g_2(t) &= \frac{\Lambda_v\gamma I_h}{\mu_2 I_v} + \frac{I_h'}{I_h} - \frac{I_v'}{I_v} - \gamma I_h - \mu_1 - \mu_2. \end{aligned} \quad (\text{B.26})$$

Since

$$(\alpha\phi + \beta\psi\theta)S_h \frac{I_v}{I_h} = \frac{I_v'}{I_h} + \mu_1, \quad (\text{B.27})$$

$$\frac{\Lambda_v\gamma I_h}{\mu_2 I_v} = \frac{I_v'}{I_v} + \gamma I_h + \mu_2,$$

we have

$$g_1 \leq \frac{I_h'}{I_h} - \mu_1, \quad g_2 = \frac{I_h'}{I_h} - \mu_1. \quad (\text{B.28})$$

Hence we find

$$\sup\{g_1, g_2\} \leq \frac{I_h'}{I_h} - \mu_1. \quad (\text{B.29})$$

Therefore, by Gronwall's inequality, we obtain

$$V(t) \leq V(0) I_h e^{-\mu_1 t} \leq V(0) \frac{\Lambda_h}{\mu_1} e^{-\mu_1 t}, \quad (\text{B.30})$$

which implies that $V(t) \rightarrow 0$ as $t \rightarrow \infty$. This implies that the given linear system is asymptotically stable and the periodic solution is asymptotically orbitally stable. $\square$

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (NRF-2013RIA6A3A01060805).

References

- [1] T. Kiyohara and Y. Tokushige, "Inoculation experiments of a nematode, *Bursaphelenchus* sp., onto pine trees," *Journal of the Japanese Forestry Society*, vol. 53, pp. 210–218, 1971 (Japanese).
- [2] Y. Mamiya and T. Kiyohara, "Description of *Bursaphelenchus lignicolus* n. sp. (Nematoda: Aphelenchoididae) from pine wood and histopathology of nematode-infested trees," *Nematologica*, vol. 18, no. 1, pp. 120–124, 1972.
- [3] W. A. R. Nickle, A. M. Golden, Y. Mamiya, and W. P. Wergin, "On the taxonomy and morphology of the pine wood nematode, *Bursaphelenchus xylophilus* (Steiner and Buhner 1934) Nickle 1970," *Journal of Nematologica*, vol. 13, no. 3, pp. 385–392, 1981.
- [4] Y. Mamiya and N. Enda, "Transmission of *Bursaphelenchus lignicolus* (Nematoda: Aphelenchoididae) by *Monochamus alternatus* (Coleoptera: Cerambycidae)," *Nematologica*, vol. 18, no. 2, pp. 159–162, 1972.
- [5] K. Morimoto and A. Iwasaki, "Role of *Monochamus alternatus* (Coleoptera: Cerambycidae) as a vector of *Bursaphelenchus lignicolus* (Nematoda: Aphelenchoididae)," *Journal of the Japanese Forestry Society*, vol. 54, pp. 177–183, 1972 (Japanese).
- [6] M. J. Wingfield and R. B. Blanchette, "The pine-wood nematode, *Bursaphelenchus xylophilus*, in Minnesota and Wisconsin: insect associates and transmission studies," *Canadian Journal of Forest Research*, vol. 13, no. 6, pp. 1068–1076, 1983.
- [7] H. Sato, T. Sakuyama, and M. Kobayashi, "Transmission of *Bursaphelenchus xylophilus* (Steiner et Buhner) Nickle (Nematoda, Aphelenchoididae) by *Monochamus saltuarius* (Gebler) (Coleoptera, Cerambycidae)," *Journal of the Japanese Forestry Society*, vol. 69, pp. 492–496, 1987 (Japanese).
- [8] M. J. Linit, "Nematode-vector relationships in the pine wilt disease system," *Journal of Nematology*, vol. 20, no. 2, pp. 227–235, 1988.
- [9] K. Konda, A. Foundin, M. Linit et al., *Pine Wilt Disease-Nematological, Entomological, and Biochemical Investigations*, University of Missouri-Columbia Agricultural Experiment Station, Columbia, Mo, USA, 1982.
- [10] S. Yano, "Investigation on pine death in Nagasaki prefecture," *Sanrin-Kouhou*, vol. 4, pp. 1–14, 1913 (Japanese).
- [11] K. Futai, "Pine wilt in Japan: from first incidence to the present," in *Pine Wilt Disease*, G. B. Zhao and K. Futai, Eds., pp. 5–12, Springer, Tokyo, Japan, 2008.
- [12] M. M. Mota and P. C. Viera, "Pine wilt disease in Portugal," in *Pine Wilt Disease*, G. B. Zhao and K. Futai, Eds., pp. 33–38, Springer, Tokyo, Japan, 2008.
- [13] B. G. Zhao, "Pine wilt disease in China," in *Pine Wilt Disease*, B. G. Zhao and K. Futai, Eds., pp. 18–25, Springer, Tokyo, Japan, 2008.
- [14] Y. Arakawa and K. Togashi, "Newly discovered transmission pathway of *Bursaphelenchus xylophilus* from males of the beetle *Monochamus alternatus* to *Pinus densiflora* trees via oviposition wounds," *Journal of Nematology*, vol. 34, pp. 396–404, 2002.
- [15] K. S. Lee and D. Kim, "Global dynamics of a pine wilt disease transmission model with nonlinear incidence rates," *Applied Mathematical Modelling*, vol. 37, no. 6, pp. 4561–4569, 2013.
- [16] K. S. Lee and A. A. Lashari, "Stability analysis and optimal control of pine wilt disease with horizontal transmission in vector population," *Applied Mathematics and Computation*, vol. 226, pp. 793–804, 2014.
- [17] K. S. Lee and A. A. Lashari, "Global stability of a host-vector model for pine wilt disease with nonlinear incidence rate," *Abstract and Applied Analysis*, vol. 2014, Article ID 219173, 11 pages, 2014.
- [18] X. Shi and G. Song, "Analysis of the mathematical model for the spread of pine wilt disease," *Journal of Applied Mathematics*, vol. 2013, Article ID 184054, 10 pages, 2013.
- [19] W. H. Fleming and R. W. Rishel, *Deterministic and Stochastic Optimal Control*, Springer, New York, NY, USA, 1975.
- [20] D. L. Lukes, *Differential Equations: Classical to Controlled*, vol. 162 of *Mathematics in Science and Engineering*, Academic Press, New York, NY, USA, 1982.
- [21] S. Lenhart and J. T. Workman, *Optimal Control Applied to Biological Models*, Mathematical and Computational Biology Series, Chapman and Hall, London, UK, 2007.
- [22] R. A. Monserud and H. Sterba, "Modeling individual tree mortality for Austrian forest species," *Forest Ecology and Management*, vol. 113, no. 2–3, pp. 109–123, 1999.
- [23] K. Togashi, "Population density of *Monochamus alternatus* adults (Coleoptera: Cerambycidae) and incidence of pine wilt disease caused by *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae)," *Researches on Population Ecology*, vol. 30, no. 2, pp. 177–192, 1988.
- [24] F. Kobayashi, A. Yamane, and T. Ikeda, "The Japanese pine sawyer beetle as the vector of pine wilt disease," *Annual Review of Entomology*, vol. 29, pp. 115–135, 1984.
- [25] D. S. kim, S. M. Lee, H. S. Huh, N. C. Park, and C. G. Park, "Escape of pine wood Nematode, *Bursaphelenchus xylophilus*, through feeding and oviposition behavior of *Monochamus alternatus* and *M. saltuarius* (Coleoptera: Cerambycidae) adults," *Korean Journal of Applied Entomology*, vol. 48, no. 4, pp. 527–533, 2009.
- [26] D. S. kim, S. M. Lee, C. S. Kim, D. W. Lee, and C. G. Park, "Movement of *Monochamus alternatus* hope (Coleoptera: Cerambycidae) adults among young black pine trees in a screen cage," *Korean Journal of Applied Entomology*, vol. 50, no. 1, pp. 1–6, 2011.
- [27] L. J. S. Allen, *An Introduction to Mathematical Biology*, Pearson Education, Upper Saddle River, NJ, USA, 2007.
- [28] J. P. LaSalle, *The Stability of Dynamical Systems*, Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics, Philadelphia, Pa, USA, 1976.
- [29] C. C. McCluskey and P. van den Driessche, "Global analysis of two tuberculosis models," *Journal of Dynamics and Differential Equations*, vol. 16, no. 1, pp. 139–166, 2004.
- [30] J. S. Muldowney, "Compound matrices and ordinary differential equations," *Rocky Mountain Journal of Mathematics*, vol. 20, no. 4, pp. 857–872, 1990.
- [31] M. Y. Li and J. S. Muldowney, "Global stability for the SEIR model in epidemiology," *Mathematical Biosciences*, vol. 125, no. 2, pp. 155–164, 1995.

- [32] A. Fonda, "Uniformly persistent semidynamical systems," *Proceedings of the American Mathematical Society*, vol. 104, no. 1, pp. 111–116, 1988.
- [33] M. W. Hirsch, "Systems of differential equations that are competitive or cooperative. IV. Structural stability in three-dimensional systems," *SIAM Journal on Mathematical Analysis*, vol. 21, no. 5, pp. 1225–1234, 1990.
- [34] H. L. Smith and H. R. Thieme, "Convergence for strongly order-preserving semiflows," *SIAM Journal on Mathematical Analysis*, vol. 22, no. 4, pp. 1081–1101, 1991.
- [35] H. L. Smith, "Systems of ordinary differential equations which generate an order preserving flow: a survey of results," *SIAM Review*, vol. 30, no. 1, pp. 87–113, 1988.