Research Article

A Note on Strongly Starlike Mappings in Several Complex Variables

Hidetaka Hamada,¹ Tatsuhiko Honda,² Gabriela Kohr,³ and Kwang Ho Shon⁴

¹ Faculty of Engineering, Kyushu Sangyo University, Fukuoka 813-8503, Japan

² Hiroshima Institute of Technology, Hiroshima 731-5193, Japan

³ Faculty of Mathematics and Computer Science, Babeş-Bolyai University, 1 M. Kogălniceanu Street, 400084 Cluj-Napoca, Romania

⁴ Department of Mathematics, College of Natural Sciences, Pusan National University, Busan 609-735, Republic of Korea

Correspondence should be addressed to Kwang Ho Shon; khshon@pusan.ac.kr

Received 3 December 2013; Accepted 27 January 2014; Published 3 March 2014

Academic Editor: Junesang Choi

Copyright © 2014 Hidetaka Hamada et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Let f be a normalized biholomorphic mapping on the Euclidean unit ball $\mathbb{B}^n$ in $\mathbb{C}^n$ and let $\alpha \in (0, 1)$. In this paper, we will show that if f is strongly starlike of order α in the sense of Liczberski and Starkov, then it is also strongly starlike of order α in the sense of Kohr and Liczberski. We also give an example which shows that the converse of the above result does not hold in dimension $n \geq 2$.

1. Introduction and Preliminaries

Let $\mathbb{C}^n$ denote the space of n complex variables $z = (z_1, \dots, z_n)$ with the Euclidean inner product $\langle z, w \rangle = \sum_{j=1}^n z_j \bar{w}_j$ and the norm $\|z\| = \langle z, z \rangle^{1/2}$. The open unit ball $\{z \in \mathbb{C}^n : \|z\| < 1\}$ is denoted by $\mathbb{B}^n$. In the case of one complex variable, $\mathbb{B}^1$ is denoted by U .

If Ω is a domain in $\mathbb{C}^n$, let $H(\Omega)$ be the set of holomorphic mappings from Ω to $\mathbb{C}^n$. If Ω is a domain in $\mathbb{C}^n$ which contains the origin and $f \in H(\Omega)$, we say that f is normalized if $f(0) = 0$ and $Df(0) = I_n$, where I_n is the identity matrix.

A normalized mapping $f \in H(\mathbb{B}^n)$ is said to be *starlike* if f is biholomorphic on $\mathbb{B}^n$ and $tf(\mathbb{B}^n) \subset f(\mathbb{B}^n)$ for $t \in [0, 1]$, where the last condition says that the image $f(\mathbb{B}^n)$ is a starlike domain with respect to the origin. For a normalized locally biholomorphic mapping f on $\mathbb{B}^n$, f is starlike if and only if

$$\Re \langle [Df(z)]^{-1} f(z), z \rangle > 0, \quad z \in \mathbb{B}^n \setminus \{0\} \quad (1)$$

(see [1–4] and the references therein, cf. [5]).

Let $\alpha \in (0, 1]$. A function $f \in H(U)$, normalized by $f(0) = 0$ and $f'(0) = 1$, is said to be *strongly starlike of order α* if

$$\left| \arg \frac{zf'(z)}{f(z)} \right| < \alpha \frac{\pi}{2}, \quad z \in U. \quad (2)$$

If f is strongly starlike of order α , then f is also starlike and thus univalent on U . Stankiewicz [6] proved that if $\alpha \in (0, 1)$, then a domain $\Omega \neq \mathbb{C}$ which contains the origin is α -accessible if and only if $\Omega = f(U)$, where U is the unit disc in $\mathbb{C}$ and f is a strongly starlike function of order $1 - \alpha$ on U . For strongly starlike functions on U , see also Brannan and Kirwan [7], Ma and Minda [8], and Sugawa [9].

Kohr and Liczberski [10] introduced the following definition of strongly starlike mappings of order α on $\mathbb{B}^n$.

Definition 1. Let $0 < \alpha \leq 1$. A normalized locally biholomorphic mapping $f \in H(\mathbb{B}^n)$ is said to be *strongly starlike of order α* if

$$\left| \arg \langle [Df(z)]^{-1} f(z), z \rangle \right| < \alpha \frac{\pi}{2}, \quad z \in \mathbb{B}^n \setminus \{0\}. \quad (3)$$

Obviously, if f is strongly starlike of order α , then f is also starlike, and if $\alpha = 1$ in (3), one obtains the usual notion of starlikeness on the unit ball $\mathbb{B}^n$.

Using this definition, Hamada and Honda [11], Hamada and Kohr [12], Liczberski [13], and Liu and Li [14] obtained

various results for strongly starlike mappings of order α in several complex variables.

Recently, Liczberski and Starkov [15] gave another definition of strongly starlike mappings of order α on the Euclidean unit ball $\mathbb{B}^n$ in $\mathbb{C}^n$, where $\alpha \in (0, 1]$, and proved that a normalized biholomorphic mapping f on $\mathbb{B}^n$ is strongly starlike of order $1 - \alpha$ if and only if $f(\mathbb{B}^n)$ is an α -accessible domain in $\mathbb{C}^n$ for $\alpha \in (0, 1)$. Their definition is as follows.

Definition 2. Let $0 < \alpha \leq 1$. A normalized locally biholomorphic mapping $f \in H(\mathbb{B}^n)$ is said to be strongly starlike of order α (in the sense of Liczberski and Starkov) if

$$\begin{aligned} & \Re \langle [Df(z)]^{-1} f(z), z \rangle \\ & \geq \left\| ([Df(z)]^{-1})^* z \right\| \cdot \|f(z)\| \sin \left((1 - \alpha) \frac{\pi}{2} \right), \quad (4) \\ & z \in \mathbb{B}^n \setminus \{0\}. \end{aligned}$$

In the case $n = 1$, it is obvious that both notions of strong starlikeness of order α are equivalent. Thus, the following natural question arises in dimension $n \geq 2$.

Question 1. Let $\alpha \in (0, 1)$. Is there any relation between the above two definitions of strong starlikeness of order α ?

Let f be a normalized biholomorphic mapping on the Euclidean unit ball $\mathbb{B}^n$ in $\mathbb{C}^n$ and let $\alpha \in (0, 1)$. In this paper, we will show that if f is strongly starlike of order α in the sense of Definition 2, then it is also strongly starlike of order α in the sense of Definition 1. As a corollary, the results obtained in [11–14] for strongly starlike mappings of order α in the sense of Definition 1 also hold for strongly starlike mappings of order α in the sense of Definition 2. We also give an example which shows that the converse of the above result does not hold in dimension $n \geq 2$.

2. Main Results

Let $\angle(a, b)$ denote the angle between $a, b \in \mathbb{C}^n \setminus \{0\}$ regarding a, b as real vectors in $\mathbb{R}^{2n}$.

Lemma 3. Let $a, b \in \mathbb{C}^n \setminus \{0\}$ be such that $\langle a, b \rangle \neq 0$. If $|\arg \langle a, b \rangle| \leq \pi$ and $0 \leq \angle(a, b) < \pi/2$, then

$$|\arg \langle a, b \rangle| \leq \angle(a, b). \quad (5)$$

Proof. Let $\theta = \arg \langle a, b \rangle$, $\varphi = \angle(a, b)$. Then we have $\langle a, b \rangle = re^{i\theta}$ for some $r \geq 0$ and

$$\Re \langle a, b \rangle = \|a\| \cdot \|b\| \cos \varphi = r \cos \theta. \quad (6)$$

Since $\cos \varphi > 0$ and $r = |\langle a, b \rangle| \leq \|a\| \cdot \|b\|$, we have

$$\cos \varphi \leq \cos \theta. \quad (7)$$

Therefore, we have $|\theta| \leq \varphi$, as desired. $\square$

Theorem 4. Let f be a normalized biholomorphic mapping on the Euclidean unit ball $\mathbb{B}^n$ in $\mathbb{C}^n$ and let $\alpha \in (0, 1)$. If f is

strongly starlike of order α in the sense of Definition 2, then it is also strongly starlike of order α in the sense of Definition 1.

Proof. Assume that f is strongly starlike of order α in the sense of Definition 2. Then by (4), we have $\langle [Df(z)]^{-1} f(z), z \rangle \neq 0$ and

$$\angle \left(([Df(z)]^{-1})^* z, f(z) \right) \leq \alpha \frac{\pi}{2}, \quad z \in \mathbb{B}^n \setminus \{0\}. \quad (8)$$

Using Lemma 3, we have

$$\begin{aligned} |\arg \langle [Df(z)]^{-1} f(z), z \rangle| &= |\arg \langle f(z), ([Df(z)]^{-1})^* z \rangle| \\ &\leq \angle \left(([Df(z)]^{-1})^* z, f(z) \right) \\ &\leq \alpha \frac{\pi}{2}, \quad z \in \mathbb{B}^n \setminus \{0\}. \end{aligned} \quad (9)$$

For fixed $z \in \mathbb{B}^n \setminus \{0\}$, let $w = z/\|z\|$ and

$$p(\zeta) = \begin{cases} \frac{1}{\zeta} \langle [Df(\zeta w)]^{-1} f(\zeta w), w \rangle, & \text{for } \zeta \in U \setminus \{0\}, \\ 1, & \text{for } \zeta = 0. \end{cases} \quad (10)$$

Then p is a holomorphic function on U with $|\arg p(\zeta)| \leq \pi\alpha/2$ for $\zeta \in U$. Since $\arg p$ is a harmonic function on U and $\arg p(0) = 0$, by applying the maximum and minimum principles for harmonic functions, we obtain $|\arg p(\zeta)| < \pi\alpha/2$ for $\zeta \in U$. Thus, we have

$$|\arg \langle [Df(z)]^{-1} f(z), z \rangle| < \alpha \frac{\pi}{2}, \quad z \in \mathbb{B}^n \setminus \{0\}. \quad (11)$$

Hence f is strongly starlike of order α in the sense of Definition 1, as desired. $\square$

The following example shows that the converse of the above theorem does not hold in dimension $n \geq 2$.

Example 5. For $\alpha \in (0, 1)$, let

$$f(z) = f_\alpha(z) = (z_1 + bz_2^2, z_2), \quad z = (z_1, z_2) \in \mathbb{B}^2, \quad (12)$$

where

$$b = \frac{3\sqrt{3}}{2} \sin \left(\alpha \frac{\pi}{2} \right). \quad (13)$$

Then

$$Df(z) = \begin{bmatrix} 1 & 2bz_2 \\ 0 & 1 \end{bmatrix}, \quad [Df(z)]^{-1} = \begin{bmatrix} 1 & -2bz_2 \\ 0 & 1 \end{bmatrix}. \quad (14)$$

Therefore,

$$\begin{aligned} \langle [Df(z)]^{-1} f(z), z \rangle &= (z_1 + bz_2^2 - 2bz_2^2) \overline{z_1} \\ &\quad + |z_2|^2 = |z_1|^2 + |z_2|^2 - b\overline{z_1}z_2^2. \end{aligned} \quad (15)$$

Since $|z_1 z_2^2| \leq 2/(3\sqrt{3})$, for $z \in \partial\mathbb{B}^2$, we obtain that $|bz_1 z_2^2| \leq \sin(\alpha\pi/2)\|z\|^3$ for $z \in \mathbb{B}^2$. This implies that $\langle [Df(z)]^{-1}f(z), z \rangle$ lies in the disc of center $\|z\|^2$ and radius $\sin(\alpha\pi/2)\|z\|^2$ for each $z \in \mathbb{B}^2 \setminus \{0\}$ and thus

$$\left| \arg \langle [Df(z)]^{-1}f(z), z \rangle \right| < \alpha \frac{\pi}{2}, \quad z \in \mathbb{B}^2 \setminus \{0\}. \quad (16)$$

Therefore, $f = f_\alpha$ is strongly starlike of order α in the sense of Definition 1.

On the other hand,

$$([Df(z)]^{-1})^* z = (z_1, z_2 - 2b\bar{z}_2 z_1). \quad (17)$$

So, for $z^0 = (1/\sqrt{3}, \sqrt{2}/\sqrt{3})$, we have

$$\begin{aligned} \langle [Df(z^0)]^{-1}f(z^0), z^0 \rangle &= 1 - m, \\ \left\| ([Df(z^0)]^{-1})^* z^0 \right\|^2 &= \frac{1}{3} + \frac{2}{3}(1 - 3m)^2, \\ \|f(z^0)\|^2 &= \frac{1}{3}(1 + 3m)^2 + \frac{2}{3}, \\ \sin\left((1 - \alpha)\frac{\pi}{2}\right) &= \sqrt{1 - m^2}, \end{aligned} \quad (18)$$

where

$$m = \sin\left(\alpha\frac{\pi}{2}\right). \quad (19)$$

Then, we obtain

$$\begin{aligned} &\left\| ([Df(z^0)]^{-1})^* z^0 \right\|^2 \|f(z^0)\|^2 \sin^2\left((1 - \alpha)\frac{\pi}{2}\right) \\ &\quad - \left(\Re \langle [Df(z^0)]^{-1}f(z^0), z^0 \rangle \right)^2 \\ &= (1 - m) \left\{ \left[\frac{1}{3} + \frac{2}{3}(1 - 3m)^2 \right] \left[\frac{1}{3}(1 + 3m)^2 + \frac{2}{3} \right] \right. \\ &\quad \left. \times (1 + m) - (1 - m) \right\}. \end{aligned} \quad (20)$$

Since

$$\left[\frac{1}{3} + \frac{2}{3}(1 - 3m)^2 \right] \left[\frac{1}{3}(1 + 3m)^2 + \frac{2}{3} \right] (1 + m) - (1 - m) \quad (21)$$

is increasing on $[1/3, 1]$ and positive for $m = 1/3$, we have

$$\begin{aligned} \Re \langle [Df(z^0)]^{-1}f(z^0), z^0 \rangle &< \left\| ([Df(z^0)]^{-1})^* z^0 \right\| \\ &\quad \times \|f(z^0)\| \sin\left((1 - \alpha)\frac{\pi}{2}\right) \end{aligned} \quad (22)$$

for $m \in [1/3, 1)$.

On the other hand, for $\tilde{z}^0 = (i/\sqrt{3}, \sqrt{2}/\sqrt{3})$, we have

$$\begin{aligned} \langle [Df(\tilde{z}^0)]^{-1}f(\tilde{z}^0), \tilde{z}^0 \rangle &= 1 + mi, \\ \left\| ([Df(\tilde{z}^0)]^{-1})^* \tilde{z}^0 \right\|^2 &= \frac{1}{3} + \frac{2}{3}|1 - 3mi|^2 = 6m^2 + 1, \\ \|f(\tilde{z}^0)\|^2 &= \frac{1}{3}|i + 3m|^2 + \frac{2}{3} = 3m^2 + 1. \end{aligned} \quad (23)$$

Then, we obtain

$$\begin{aligned} &\left\| ([Df(\tilde{z}^0)]^{-1})^* \tilde{z}^0 \right\|^2 \|f(\tilde{z}^0)\|^2 \sin^2\left((1 - \alpha)\frac{\pi}{2}\right) \\ &\quad - \left(\Re \langle [Df(\tilde{z}^0)]^{-1}f(\tilde{z}^0), \tilde{z}^0 \rangle \right)^2 \\ &= (6m^2 + 1)(3m^2 + 1)(1 - m^2) - 1 \\ &= m^2(-18m^4 + 9m^2 + 8). \end{aligned} \quad (24)$$

Since $-18m^4 + 9m^2 + 8$ is positive for $m \in [0, 1/3]$, we have

$$\begin{aligned} \Re \langle [Df(\tilde{z}^0)]^{-1}f(\tilde{z}^0), \tilde{z}^0 \rangle &< \left\| ([Df(\tilde{z}^0)]^{-1})^* \tilde{z}^0 \right\| \\ &\quad \times \|f(\tilde{z}^0)\| \sin\left((1 - \alpha)\frac{\pi}{2}\right) \end{aligned} \quad (25)$$

for $m \in (0, 1/3]$.

Thus, $f = f_\alpha$ is not strongly starlike of order α in the sense of Definition 2 for $\alpha \in (0, 1)$.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

Hidetaka Hamada is supported by JSPS KAKENHI Grant no. 25400151. Tatsuhiko Honda is partially supported by Brain Korea Project, 2013. The work of Gabriela Kohr was supported by a grant of the Romanian National Authority for Scientific Research, CNCS-UEFISCDI, Project no. PN-II-ID-PCE-2011-3-0899. Kwang Ho Shon was supported by a 2-year research grant of Pusan National University.

References

- [1] T. J. Suffridge, "Starlikeness, Convexity and Other Geometric Properties of Holomorphic Maps in Higher Dimensions," in *Complex Analysis*, vol. 599 of *Lecture Notes in Mathematics*, pp. 146–159, Springer, Berlin, Germany, 1977.
- [2] K. R. Gurganus, " Φ -like holomorphic functions in $\mathbb{C}^n$ and Banach spaces," *Transactions of the American Mathematical Society*, vol. 205, pp. 389–406, 1975.
- [3] T. J. Suffridge, "Starlike and convex maps in Banach spaces," *Pacific Journal of Mathematics*, vol. 46, pp. 575–589, 1973.
- [4] S. Gong, *Convex and Starlike Mappings in Several Complex Variables*, Kluwer Academic, Dordrecht, The Netherlands, 1998.
- [5] H. Hamada and G. Kohr, " Φ -like and convex mappings in infinite dimensional spaces," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 47, no. 3, pp. 315–328, 2002.
- [6] J. Stankiewicz, "Quelques problèmes extrémaux dans les classes des fonctions α -angulairement étoilées," vol. 20, pp. 59–75, 1966 (French).
- [7] D. A. Brannan and W. E. Kirwan, "On some classes of bounded univalent functions," *Journal of the London Mathematical Society. Second Series*, vol. 1, pp. 431–443, 1969.

- [8] W. Ma and D. Minda, "An internal geometric characterization of strongly starlike functions," *Annales Universitatis Mariae Curie-Skłodowska A*, vol. 45, pp. 89–97, 1991.
- [9] T. Sugawa, "A self-duality of strong starlikeness," *Kodai Mathematical Journal*, vol. 28, no. 2, pp. 382–389, 2005.
- [10] G. Kohr and P. Liczberski, "On strongly starlikeness of order α in several complex variables," *Glasnik Matematički. Serija III*, vol. 33, no. 2, pp. 185–198, 1998.
- [11] H. Hamada and T. Honda, "Sharp growth theorems and coefficient bounds for starlike mappings in several complex variables," *Chinese Annals of Mathematics B*, vol. 29, no. 4, pp. 353–368, 2008.
- [12] H. Hamada and G. Kohr, "On some classes of bounded univalent mappings in several complex variables," *Manuscripta Mathematica*, vol. 131, no. 3-4, pp. 487–502, 2010.
- [13] P. Liczberski, "A geometric characterization of some biholomorphic mappings in $\mathbb{C}^n$," *Journal of Mathematical Analysis and Applications*, vol. 375, no. 2, pp. 538–542, 2011.
- [14] H. Liu and X. Li, "The growth theorem for strongly starlike mappings of order α on bounded starlike circular domains," *Chinese Quarterly Journal of Mathematics*, vol. 15, no. 3, pp. 28–33, 2000.
- [15] P. Liczberski and V. V. Starkov, "Domains in $\mathbb{R}^n$ with conically accessible boundary," *Journal of Mathematical Analysis and Applications*, vol. 408, no. 2, pp. 547–560, 2013.