Research Article

Hemi-Slant Warped Product Submanifolds of Nearly Kaehler Manifolds

Falleh R. Al-Solamy¹ and Meraj Ali Khan²

¹ Department of Mathematics, King Abdulaziz University, P.O. Box 80015, Jeddah 21589, Saudi Arabia

² Department of Mathematics, University of Tabuk, Tabuk, Saudi Arabia

Correspondence should be addressed to Meraj Ali Khan; meraj79@gmail.com

Received 25 January 2014; Revised 16 April 2014; Accepted 24 April 2014; Published 12 May 2014

Academic Editor: Irena Rachůnková

Copyright © 2014 F. R. Al-Solamy and M. A. Khan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hemi-slant warped product submanifolds of nearly Kaehler manifolds are studied and some interesting results are obtained. Moreover, an inequality is established for squared norm of second fundamental form and equality case is also discussed. The results obtained are also true if ambient manifold is replaced by a Kaehler manifold. These results generalize several known results in the literature.

1. Introduction

In [1] Bishop and O'Neill introduced the notion of warped product manifolds as a natural generalization of Riemannian product manifolds. For instance, a surface of revolution is a warped product manifold. So far as its applications are concerned, it has been shown that warped product manifolds provide excellent setting to model space time near black holes or bodies with large gravitational forces (see [1, 2]). Due to wide applications of warped product submanifolds, this becomes a fascinating and interesting topic for research and many articles are available in literature (see [1, 3–5]). Chen [6] initiated the study of warped product submanifolds by showing that there do not exist warped product CR-submanifolds of the type $N_{\perp} \times_f N_T$, and he considered warped product CR-submanifolds of the types $N_T \times_f N_{\perp}$ and established a relationship between the warping function f and the squared norm of the second fundamental form. Extending the study of Chen, Sahin [7] proved that there exist no semislant warped product submanifolds in a Kaehler manifold. In [8], V. A. Khan and K. A. Khan studied generic warped product submanifolds of nearly Kaehler manifolds and obtained an inequality for squared norm of second fundamental form in terms of warping function. Recently, Sahin [4] investigated hemi-slant warped product for Kaehler manifolds and obtained an inequality for squared norm of

second fundamental form for mixed totally geodesic submanifolds. In view of the interesting geometric characteristic of nearly Kaehler manifolds and the nonexistence of CR-product submanifolds in S^6 [9], it will be significant to explore hemi-slant warped product submanifolds of a nearly Kaehler manifold. In this continuation we have achieved success in extending the results of Sahin [4] and Chen [6] to the setting of nearly Kaehler manifolds.

2. Preliminaries

Let $(\overline{M}, J, g)$ be a nearly Kaehler manifold with an almost complex structure J and Hermitian metric g and a Levi-Civita connection $\overline{\nabla}$ such that

$$g(JX, JY) = g(X, Y), \quad (1)$$

$$(\overline{\nabla}_X J)Y + (\overline{\nabla}_Y J)X = 0, \quad (2)$$

for all vector fields X and Y on $\overline{M}$. Six-dimensional sphere S^6 is a classic example of a nearly Kaehler non-Kaehler manifold. It has an almost complex structure J defined by the vector cross product in the space of purely imaginary Cayley numbers which satisfies the condition $(\overline{\nabla}_X J)X = 0$. Let C be the Cayley division algebra generated by $\{e_0 = 1, e_i, (1 \leq i \leq 7)\}$ over R and C_+ the subspace of C consisting

of all purely imaginary Cayley numbers. We may identify C_+ with a 7-dimensional Euclidean space R^7 with the canonical inner product $g = (\cdot, \cdot)$. The automorphism group of C is the compact simple Lie group G_2 and furthermore the inner product g is invariant under the action of G_2 and hence, the group G_2 may be considered as a subgroup of $SO(7)$. A vector cross product for vectors in $R^7 (= C_+)$ is defined by

$$x \times y = (x, y) e_0 + xy, \quad \forall x, y \in C_+. \quad (3)$$

Then the multiplication table for $e_j \times e_k$ is given by

$\begin{smallmatrix} j \\ k \end{smallmatrix}$	1	2	3	4	5	6	7	
1	0	e_3	$-e_2$	e_5	$-e_4$	e_7	$-e_6$	
2	$-e_3$	0	e_1	e_6	$-e_7$	$-e_4$	e_5	
3	e_2	$-e_1$	0	$-e_7$	$-e_6$	e_5	e_4	.
4	$-e_5$	$-e_6$	e_7	0	e_1	e_2	$-e_3$	
5	e_4	e_7	e_6	$-e_1$	0	$-e_3$	$-e_2$	
6	$-e_7$	e_4	$-e_5$	$-e_2$	e_3	0	e_1	
7	e_6	$-e_5$	$-e_4$	e_3	e_2	$-e_1$	0	

(4)

Considering S^6 as $\{x \in C_+ : (x, x) = 1\}$, the almost complex structure J on S^6 is defined by

$$J_x(X) = x \times X, \quad (5)$$

where $x \in S^6$ and $X \in T_x S^6$. The almost complex structure defined by the above equation together with the induced metric on S^6 from g on $R^7 (= C_+)$ gives rise to a nearly Kaehler structure S^6 [10].

Let M be a submanifold of $\overline{M}$. Then the induced Riemannian metric on M is denoted by the same symbol g and the induced connection on M is denoted by the symbol ∇ . If $T\overline{M}$ and TM denote the tangent bundle on $\overline{M}$ and M , respectively, and $T^\perp M$, the normal bundle on M , then the Gauss and Weingarten formulae are given by

$$\overline{\nabla}_X Y = \nabla_X Y + h(X, Y), \quad (6)$$

$$\overline{\nabla}_X N = -A_N X + \nabla_X^\perp N, \quad (7)$$

for $X, Y \in TM$ and $N \in T^\perp M$ where $\nabla^\perp$ denotes the connection on the normal bundle $T^\perp M$. h and A_N are the second fundamental form and the shape operator of immersions of M into $\overline{M}$. Corresponding to the normal vector field V they are related as

$$g(A_V X, Y) = g(h(X, Y), V). \quad (8)$$

The mean curvature vector H of M is given by

$$H = \frac{1}{n} \sum_{i=1}^n h(e_i, e_i), \quad (9)$$

where n is the dimension of M and $\{e_1, e_2, \dots, e_n\}$ is a local orthonormal frame of vector fields on M . The squared norm of the second fundamental form is defined as

$$\|h\|^2 = \sum_{i,j=1}^n g(h(e_i, e_j), h(e_i, e_j)). \quad (10)$$

A submanifold M of $\overline{M}$ is said to be a totally geodesic submanifold if $h(X, Y) = 0$ for each $X, Y \in TM$, and totally umbilical submanifold if $h(X, Y) = g(X, Y)H$.

For $X \in TM$ and $N \in T^\perp M$ we write

$$\begin{aligned} JX &= PX + FX, \\ JN &= tN + fN, \end{aligned} \quad (11)$$

where PX and tN are tangential components of JX and JN , respectively, and FX and fN are the normal components of JX and JN .

The covariant differentiation of the tensors P, F, t , and f is defined, respectively, as

$$(\overline{\nabla}_X P)Y = \nabla_X PY - P\nabla_X Y, \quad (12)$$

$$(\overline{\nabla}_X F)Y = \nabla_X^\perp FY - F\nabla_X Y, \quad (13)$$

$$(\overline{\nabla}_X t)N = \nabla_X tN - t\nabla_X^\perp N, \quad (14)$$

$$(\overline{\nabla}_X f)N = \nabla_X^\perp fN - f\nabla_X^\perp N. \quad (15)$$

Furthermore, for any $X, Y \in TM$, the tangential and normal parts of $(\overline{\nabla}_X J)Y$ are denoted by $\mathcal{P}_X Y$ and $\mathcal{Q}_X Y$; that is,

$$(\overline{\nabla}_X J)Y = \mathcal{P}_X Y + \mathcal{Q}_X Y. \quad (16)$$

On using (6)–(13), we may obtain that

$$\mathcal{P}_X Y = (\overline{\nabla}_X P)Y - A_{FY}X - th(X, Y), \quad (17)$$

$$\mathcal{Q}_X Y = (\overline{\nabla}_X F)Y + h(X, TY) - fh(X, Y). \quad (18)$$

Similarly, for $N \in T^\perp M$, denoting by $\mathcal{P}_X N$ and $\mathcal{Q}_X N$, respectively, the tangential and normal parts of $(\overline{\nabla}_X J)N$, we find that

$$\begin{aligned} \mathcal{P}_X N &= (\overline{\nabla}_X t)N + PA_N X - A_{fN} X, \\ \mathcal{Q}_X N &= (\overline{\nabla}_X f)N + h(tN, X) + FA_N X. \end{aligned} \quad (19)$$

On a submanifold M of a nearly Kaehler manifold by (2) and (16)

$$(a) \mathcal{P}_X Y + \mathcal{P}_Y X = 0, \quad (b) \mathcal{Q}_X Y + \mathcal{Q}_Y X = 0 \quad (20)$$

for any $X, Y \in TM$.

Let $\overline{M}$ be an almost Hermitian manifold with an almost complex structure J and Hermitian metric g and let M be a submanifold of $\overline{M}$. Submanifold M is said to be CR-submanifold if there exist two orthogonal complementary distributions D and $D^\perp$ such that D is holomorphic distribution, that is, $JD \subseteq D$ and $D^\perp$ is totally real distribution, that is, $JD^\perp \subseteq T^\perp M$.

An immersed submanifold M of an almost Hermitian manifold $\overline{M}$ is said to be slant submanifold if the Wirtinger angle $\theta(X) \in [0, \pi/2]$ between JX and $T_x M$, and $X \neq 0$ (cf. [11]). Holomorphic and totally real submanifolds are slant submanifolds with wirtinger angle 0 and $\pi/2$.

A submanifold is called proper slant if it is neither holomorphic nor totally real. More generally, a distribution D_θ on $\overline{M}$ is called a slant distribution if the angle $\theta(X)$ between JX and D_θ has the same value θ for each $x \in \overline{M}$ and $X \in D_\theta$ and $X \neq 0$.

If M is a slant submanifold of an almost Hermitian manifold $\overline{M}$, then we have (cf. [11])

$$P^2X = -\cos^2\theta X, \quad (21)$$

where θ is the wirtinger angle of M in $\overline{M}$. Hence we have

$$g(PX, PY) = \cos^2\theta g(X, Y), \quad (22)$$

$$g(FX, FY) = \sin^2\theta g(X, Y), \quad (23)$$

for any $X, Y \in TM$.

Papaghiuc [12] introduced a class of submanifolds in almost Hermitian manifolds called the semislant submanifolds; this class includes the class of proper CR-submanifolds and slant submanifolds. Cabrerizo et al. [13] initiated the study of contact version of semislant submanifold and also bislant submanifolds. As a step forward, Carriazo [14] defined and studied bislant submanifolds and simultaneously gave the notion of antislant submanifolds in almost Hermitian manifolds; after that V. A. Khan and M. A. Khan [15] studied antislant submanifolds with the name pseudo-slant submanifolds in the setting of Sasakian manifolds.

Recently, Sahin [4] renamed pseudo-slant submanifolds as hemi-slant submanifolds and studied hemi-slant submanifolds for their warped product.

Definition 1. A submanifold M of an almost Hermitian manifold is called hemi-slant submanifold if it is endowed with two orthogonal complementary distributions $D^\perp$ and D_θ such that $D^\perp$ is totally real, that is, $JD^\perp \subseteq T^\perp M$ and D_θ is slant distribution with slant angle θ .

It is straight forward to see that CR-submanifolds and slant submanifolds are hemi-slant submanifolds with $\theta = 0$ and $D^\perp = \{0\}$, respectively.

If μ is the invariant subspace of the normal bundle $T^\perp M$ then in the case of hemi-slant submanifold, the normal bundle $T^\perp M$ can be decomposed as follows:

$$T^\perp M = \mu \oplus FD^\perp \oplus FD_\theta. \quad (24)$$

As $D^\perp$ and D_θ are orthogonal distributions on M , then it is easy to see that the distributions $FD^\perp$ and FD_θ are mutually perpendicular. In fact, the decomposition (24) is an orthogonal direct decomposition.

A hemi-slant submanifold M is called a hemi-slant product if the distributions $D^\perp$ and D_θ are parallel on M . In this case M is foliated by the leaves of these distributions. In particular if M is CR-submanifold with parallel distribution then it is called CR-product. In general, if M_1 and M_2 are Riemannian manifolds with Riemannian metrics g_1 and g_2 , respectively, then the product manifold $(M_1 \times M_2, g)$ is a Riemannian manifold with Riemannian metric g defined as

$$g(X, Y) = g_1(d\pi_1 X, d\pi_1 Y) + g_2(d\pi_2 X, d\pi_2 Y), \quad (25)$$

where π_i ($i = 1, 2$) are the projection maps of M onto M_1 and M_2 , respectively, and $d\pi_i$ ($i = 1, 2$) are their differentials.

As a generalization of the product manifold and in particular of a hemi-slant product submanifold, one can consider warped product of manifolds which are defined in the following.

Definition 2. Let (B, g_B) and (C, g_C) be two Riemannian manifolds with Riemannian metrics g_B and g_C , respectively, and f a positive differentiable function on B . The warped product of B and C is the Riemannian manifold $(B \times_f C, g)$, where

$$g = g_B + f^2 g_C. \quad (26)$$

For a warped product manifold $N_1 \times_f N_2$, we denote by D_1 and D_2 the distributions defined by the vectors tangent to the leaves and fibers, respectively. In other words, D_1 is obtained by the tangent vectors of N_1 via the horizontal lift and D_2 is obtained by the tangent vectors of N_2 via vertical lift. In case of semi-invariant warped product submanifolds D_1 and D_2 are replaced by D and $D^\perp$, respectively.

The warped product manifold $(B \times_f C, g)$ is denoted by $B \times_f C$. If X is the tangent vector field to $M = B \times_f C$ at (p, q) then

$$\|X\|^2 = \|d\pi_1 X\|^2 + f^2(p) \|d\pi_2 X\|^2. \quad (27)$$

Bishop and O'Neill [1] proved the following.

Theorem 3. Let $M = B \times_f C$ be warped product manifolds. If $X, Y \in TB$ and $V, W \in TC$ then

- (i) $\nabla_X Y \in TB$,
- (ii) $\nabla_X V = \nabla_V X = (Xf/f)V$,
- (iii) $\nabla_V W = (-g(V, W)/f)\nabla f$.

∇f is the gradient of f and is defined as

$$g(\nabla f, X) = Xf, \quad (28)$$

for all $X \in TM$.

Corollary 4. On a warped product manifold $M = N_1 \times_f N_2$, the following statements hold:

- (i) N_1 is totally geodesic in M
- (ii) N_2 is totally umbilical in M .

In what follows, $N_\perp$ and N_θ will denote a totally real and slant submanifold, respectively, of an almost Hermitian manifold $\overline{M}$.

A warped product manifold is said to be trivial if its warping function f is constant. More generally, a trivial warped product manifold $M = N_1 \times N_2$ is a Riemannian product $N_1 \times N_2^f$, where N_2^f is the manifold with the Riemannian metric $f^2 g_2$ which is homothetic to the original metric g_2 of N_2 . For example, a trivial CR-warped product is CR-product.

Sahin [4] extended the study of warped product hemi-slant submanifolds and hemi-slant warped product of Kaehler manifolds introducing warped product submanifolds as $N_\theta \times_{f_2} N_\perp$ and $N_\perp \times_f N_\theta$, where θ is the slant angle.

3. Hemi-Slant Warped Product Submanifolds

In [5] Uddin and Chi investigated warped product pseudo-slant (hemi-slant) submanifolds of nearly Kaehler manifolds and they only showed that there do not exist warped products of the form $N_\perp \times_f N_\theta$ in nearly Kaehler manifolds, where $N_\perp$ is totally real submanifold and N_θ is slant submanifold. In this section we study the warped products of the types $N_\theta \times_f N_\perp$.

Let $M = N_\theta \times_f N_\perp$ be a hemi-slant warped product of a nearly Kaehler manifold $\bar{M}$. Then by Theorem 3,

$$\nabla_X Z = \nabla_Z X = X \ln f Z, \quad (29)$$

for any $X \in TN_\theta, Z \in TN_\perp$.

Now by formula (12) and Theorem 3,

$$(\bar{\nabla}_Z P)W = g(Z, W)P(\nabla \ln f) - g(Z, PW)\nabla \ln f, \quad (30)$$

for each $Z, W \in TN_\theta$. Now we will investigate some interesting results of the second fundamental form.

Proposition 5. *On a hemi-slant warped product submanifold $M = N_\theta \times_f N_\perp$ of a nearly Kaehler manifold $\bar{M}$, one has*

$$(i) \ 2g(h(X, Y), FZ) = g(h(X, Z), FY) + g(h(Y, Z), FX),$$

$$(ii) \ g(h(X, Z), FX) = g(h(X, X), FZ),$$

for any $X, Y \in N_\theta$ and $Z \in N_\perp$.

Proof. As N_θ is totally geodesic in M then $(\bar{\nabla}_X P)Y \in TN_\theta$ and therefore by formula (17),

$$g(\mathcal{P}_X Y, Z) = -g(A_{FX} Y, Z) - g(th(X, Y), Z), \quad (31)$$

or

$$g(h(X, Y), FZ) = g(\mathcal{P}_X Y, Z) + g(h(Y, Z), FX). \quad (32)$$

Similarly, we have

$$g(h(X, Y), FZ) = g(\mathcal{P}_Y X, Z) + g(h(X, Z), FY). \quad (33)$$

Adding above two equations and using (20)(a), we get part (i).

Now by formula (17) and (20)(a),

$$0 = (\bar{\nabla}_X P)Z + (\bar{\nabla}_Z P)X - 2th(X, Z) - A_{FZ}X - A_{FX}Z, \quad (34)$$

and by (29) the above equation gives

$$(PX \ln f)Z = 2th(X, Z) + A_{FZ}X + A_{FX}Z. \quad (35)$$

Taking inner product of (35) with $X \in D_\theta$, we get

$$g(h(X, Z), FX) = g(h(X, X), FZ), \quad (36)$$

which proves part (ii). $\square$

Theorem 6. *For a hemi-slant warped product submanifold $N_\theta \times_f N_\perp$ of a nearly Kaehler manifold the warping function satisfies the following relation:*

$$\cos^2 \theta X \ln f \|Z\|^2 = g(h(PX, Z), FZ) - g(H, FPX) \|Z\|^2, \quad (37)$$

for any $X \in TN_\theta$ and $Z \in TN_\perp$.

Proof. If M is a hemi-slant warped product submanifold $N_\theta \times_f N_\perp$ of a nearly Kaehler manifold then $(\bar{\nabla}_X P)Z = 0$ for each $X \in TN_\theta$ and $Z \in TN_\perp$, and thus by (17),

$$\mathcal{P}_X Z = -A_{FZ}X - th(X, Z). \quad (38)$$

On the other hand

$$\mathcal{P}_Z X = \nabla_Z PX - P\nabla_Z X - A_{FX}Z - th(X, Z). \quad (39)$$

Now using (29), the above equation takes the form

$$\mathcal{P}_Z X = PX \ln f Z - A_{FX}Z - th(X, Z). \quad (40)$$

Adding (38) and (40) and using (20)(a),

$$A_{FZ}X + A_{FX}Z = PX \ln f Z - 2th(X, Z), \quad (41)$$

taking inner product with $Z \in TN_\perp$, and using the fact that $N_\perp$ is totally umbilical, one gets the following equation:

$$PX \ln f \|Z\|^2 = -g(h(X, Z), FZ) + g(H, FX) \|Z\|^2. \quad (42)$$

By replacing X by PX the required result follows. $\square$

Remark 7. In [4] Sahin proved that hemi-slant warped products of the type $N_\perp \times_f N_\theta$ do not exist in the setting of Kaehler manifolds. Therefore, in the following Corollary we discuss the warped products of the type $N_\theta \times_f N_\perp$.

Corollary 8. *For a hemi-slant warped product submanifold $N_\theta \times_f N_\perp$ of a Kaehler manifold the warping function satisfies the following relation:*

$$\cos^2 \theta X \ln f \|Z\|^2 = g(h(PX, Z), FZ) - g(H, FPX) \|Z\|^2, \quad (43)$$

for any $X \in TN_\theta$ and $Z \in TN_\perp$.

Proof. Since M is a hemi-slant warped product submanifold of a Kaehler manifold, then by tensorial equation of Kaehler manifold, it is easy to see that $\mathcal{P}_Z X = 0$, for any $X \in TN_\theta$ and $Z \in TN_\perp$, and using this fact in (40) and taking inner product with $Z \in TN_\perp$, we get the required result. $\square$

Let us denote by D_θ and $D^\perp$ the tangent bundles on N_θ and $N_\perp$, respectively, and let $\{X_1, X_2, \dots, X_p, X_{p+1} = PX_1, \dots, X_{2p} = PX_p\}$ and $\{Z_1, Z_2, \dots, Z_q\}$ be local orthonormal frames of vector fields on N_θ and $N_\perp$, respectively, with $2p$ and q being their real dimensions; then

$$\begin{aligned} \|h\|^2 &= \sum_{i,j=1}^{2p} g(h(X_i, X_j), h(X_i, X_j)) \\ &+ \sum_{i=1}^{2p} \sum_{r=1}^q g(h(X_i, Z_r), h(X_i, Z_r)) \\ &+ \sum_{r,s=1}^q g(h(Z_r, Z_s), h(Z_r, Z_s)). \end{aligned} \quad (44)$$

Now we calculate the inequality for the squared norm of second fundamental form in the following theorem.

Theorem 9. Let $M = N_\theta \times_f N_\perp$ be a hemi-slant warped product of a nearly Kaehler manifold $\overline{M}$ with N_θ and $N_\perp$ slant and totally real submanifolds, respectively, of $\overline{M}$. If $H \in \mu$, then

(i) the second fundamental form satisfies the following inequality:

$$\|h\|^2 \geq 4q\cos^2\theta \|\nabla \ln f\|^2, \quad (45)$$

(ii) equality holds if $h(D, D) = 0$, $h(D^\perp, D^\perp) = 0$, $h(PX_i, Z_r)$ is normal to FZ_t and $h(PX_i, Z_r)$ is normal to FX_i and FX_s where $i, s = 1, 2, \dots, 2p$ and $r, t = 1, \dots, q$ ($i \neq s$ and $r \neq t$),

where $\nabla \ln f$ is the gradient of $\ln f$ and q is dimension of $N_\perp$.

Proof. In view of (24) the second fundamental form can be decomposed as follows:

$$h(U, V) = h_{FD_\theta}(U, V) + h_{FD_\perp}(U, V) + h_\mu(U, V), \quad (46)$$

for each $U, V \in TM$, where $h_{FD_\theta}(U, V) \in FD_\theta$, $h_{FD_\perp}(U, V) \in FD_\perp$ and $h_\mu(U, V) \in \mu$ with

$$h_{FD_\theta}(U, V) = \sum_{i=1}^{2p} h^i(U, V) FX_i, \quad (47)$$

where,

$$h^i(U, V) = \csc^2\theta g(h(U, V), FX_i), \quad (48)$$

$$h_{FD_\perp}(U, V) = \sum_{r=1}^q h^r(U, V) FZ_r, \quad (49)$$

where,

$$h^r(U, V) = g(h(U, V), FZ_r), \quad (50)$$

for each $X_i \in D_\theta$ and $Z_r \in D_\perp$.

Now, making use of (37) with assumption $H \in \mu$ and formulae (48) and (49), we obtain

$$\begin{aligned} &g(h_{FD_\perp}(PX_i, Z_r), h_{FD_\perp}(PX_i, Z_r)) \\ &= \cos^2\theta h^r(PX_i, Z_r)(X_i \ln f) + \sum_{t \neq r} (h^t(PX_i, Z_r))^2. \end{aligned} \quad (51)$$

Summing over $i = 1, \dots, 2p$ and $r = 1, \dots, q$ and (37) and assumption $H \in \mu$ with formula (50) the above equation gives

$$\begin{aligned} &g(h_{FD_\perp}(PX_i, Z_r), h_{FD_\perp}(PX_i, Z_r)) \\ &= 2q\cos^4\theta \|\nabla \ln f\|^2 + \sum_{i=1}^{2p} \sum_{r,t=1, r \neq t}^q (h^t(PX_i, Z_r))^2. \end{aligned} \quad (52)$$

Let us consider the orthonormal frame of vector fields on $D_\perp$ as $\{Z_1, \dots, Z_q\}$, and the second term in the right hand side of last equation on using (50) can be written as

$$\begin{aligned} &\sum_{i=1}^{2p} \left[\sum_{r=1}^q (g(h(PX_i, Z_r), FZ_r))^2 \right. \\ &\quad \left. + \sum_{r=1}^q \sum_{t=1, t \neq r}^q (g(h(PX_i, Z_r), FZ_t))^2 \right]. \end{aligned} \quad (53)$$

On applying (37), the first part of above expression reduced to

$$\sum_{i=1}^{2p} \sum_{r=1}^q (g(h(PX_i, Z_r), FZ_r))^2 = 2q\cos^4\theta \|\nabla \ln f\|^2. \quad (54)$$

Taking account of the above equation into (52), we obtain

$$g(h_{FD_\perp}(PX_i, Z_r), h_{FD_\perp}(PX_i, Z_r)) \geq 4q\cos^4\theta \|\nabla \ln f\|^2. \quad (55)$$

The inequality follows from (44) and (55).

To discuss the equality case we will explore the expression $\|h_{FD_\theta}(PX_i, Z_r)\|^2$ as follows.

Making use of (47), (48), and (23) and summing over $r = 1, \dots, q$ and $i = 1, \dots, 2p$ we find

$$\begin{aligned} &g(h_{FD_\theta}(PX_i, Z_r), h_{FD_\theta}(PX_i, Z_r)) \\ &= \sin^2\theta \sum_{r=1}^q \left[\left\{ \sum_{i=1}^{2p} (h^i(PX_i, Z_r))^2 \right. \right. \\ &\quad \left. \left. + \sum_{i,s=1, s \neq i}^{2p} (h^s(PX_i, Z_r))^2 \right\} \right]. \end{aligned} \quad (56)$$

Let us choose the orthonormal frame of vectors fields on D_θ as $\{X_1, \dots, X_p, PX_1, PX_2, \dots, PX_p = X_{2p}\}$. Then the right hand side of the above equation with the help of (48) can be written as

$$\begin{aligned} &g(h_{FD_\theta}(PX_i, Z_r), h_{FD_\theta}(PX_i, Z_r)) \\ &= \csc^2\theta \sum_{r=1}^q \left[2 \sum_{i=1}^{2p} (g(h(PX_i, Z_r), FX_i))^2 \right. \\ &\quad \left. + \sum_{i=1}^{2p} \sum_{s=1, s \neq i}^{2p} (g(h(PX_i, Z_r), FX_s))^2 \right]. \end{aligned} \quad (57)$$

From (44), (50), and expression (53), it is clear that equality holds if $h(D, D) = 0$, $h(D^\perp, D^\perp) = 0$, $h(PX_i, Z_r)$ is normal to FZ_t and $h(PX_i, Z_r)$ is normal to FX_i and FX_s , $i, s = 1, 2, \dots, 2p$, $r, t = 1, \dots, q$, where $i \neq s$ and $r \neq t$. $\square$

Remark 10. Since (37) is the key result of the paper which helps to get the inequality in Theorem 9 and moreover (37) is also true for the Kaehler manifolds, hence the results in Theorem 9 are also true for hemi-slant warped product submanifolds of a Kaehler manifold.

Now we compile some results of [16] and give the following example of a warped product submanifold in S^6 .

Example 11. Let $\{e_0, e_i \ (1 \leq i \leq 7)\}$ be the canonical basis of the Cayley division algebra on R^8 over R and R^7 the subspace of R^8 generated by the purely imaginary Cayley numbers $e_i \ (1 \leq i \leq 7)$. Then

$$S^6 = \{x_1 e_1 + x_2 e_2 + \cdots + x_7 e_7 : x_1^2 + x_2^2 + \cdots + x_7^2 = 1\} \subset R^7 \quad (58)$$

is a unit sphere admitting a nearly Kaehler structure $(J, g, \bar{\nabla})$ as has been specified earlier. Now suppose that $S^2 = \{x = (x_2, x_4, x_6) \in R^3 : x_2^2 + x_4^2 + x_6^2 = 1\}$ is a unit 2-sphere. For a real triple $p = (p_1, p_2, p_3)$ with $p_1 + p_2 + p_3 = 0$ and $p_1 p_2 p_3 \neq 0$, define a mapping ψ_p of $S^2 \times R$ to $S^5 \subset S^6$ as follows:

$$\begin{aligned} \psi_p(x_1, x_2, x_3, t) = & x_1 (\cos(tp_1)e_1 + \sin(tp_1)e_5) \\ & + x - 2(\cos(tp_2)e_2 + \sin(tp_2)e_6) \quad (59) \\ & + x_3 (\cos(tp_3)e_3 + \sin(tp_3)e_7), \end{aligned}$$

where $x_1^2 + x_2^2 + x_3^2 = 1$ and $t \in R$. Then ψ_p gives rise to an isometric immersion from warped product Riemannian manifold $S^2 \times_f R$ into S^6 . Moreover, induced warped product metric $\bar{g}$ on $S^2 \times_f R$ is given by

$$\bar{g} = \pi_1^* g_0 + \left(\sum_{i=1}^3 (x_i p_i)^2 \right) \pi_2^* dt^2, \quad (60)$$

where $\pi_1 : S^2 \times_f R \rightarrow S^2$ and $\pi_2 : S^2 \times R \rightarrow R$ are natural projections and g_0 is the Riemannian metric on S^2 and from (27) it is evident that warping function f is as follows:

$$f = \sqrt{\sum_{i=1}^3 (x_i p_i)^2}. \quad (61)$$

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

The authors are highly thankful to anonymous referees for their valuable suggestions and comments which have improved the paper.

References

- [1] R. L. Bishop and B. O'Neill, "Manifolds of negative curvature," *Transactions of the American Mathematical Society*, vol. 145, pp. 1–49, 1969.
- [2] S.-T. Hong, "Warped products and black holes," *Il Nuovo Cimento della Società Italiana di Fisica B*, vol. 120, no. 10–11, pp. 1227–1234, 2005.
- [3] B.-Y. Chen, "Geometry of warped product CR-submanifolds in Kaehler manifolds," *Monatshefte für Mathematik*, vol. 133, no. 3, pp. 177–195, 2001.
- [4] B. Sahin, "Warped product submanifolds of Kaehler manifolds with a slant factor," *Annales Polonici Mathematici*, vol. 95, no. 3, pp. 207–226, 2009.
- [5] S. Uddin and A. Y. M. Chi, "Warped product pseudo-slant submanifolds of nearly Kaehler manifolds," *Analele Stiintifice ale Universitatii Ovidius Constanta Seria Matematica*, vol. 19, no. 3, pp. 195–204, 2011.
- [6] B.-Y. Chen, "CR-submanifolds of a Kaehler manifold. I," *Journal of Differential Geometry*, vol. 16, no. 2, pp. 305–322, 1981.
- [7] B. Sahin, "Non existence of warped product semi-slant submanifolds of Kaehler manifolds," *Geometriae Dedicata*, vol. 117, pp. 195–202, 2006.
- [8] V. A. Khan and K. A. Khan, "Generic warped product submanifolds in nearly Kaehler manifolds," *Contributions to Algebra and Geometry*, vol. 50, no. 2, pp. 337–352, 2009.
- [9] K. Sekigawa, "Some CR-submanifolds in a S^6 -dimensional sphere," *Tensor. New Series*, vol. 41, no. 1, pp. 13–20, 1984.
- [10] T. Fukami and S. Ishihara, "Almost Hermitian structure on S^6 ," *The Tohoku Mathematical Journal*, vol. 7, pp. 151–156, 1955.
- [11] B.-Y. Chen, "Slant immersions," *Bulletin of the Australian Mathematical Society*, vol. 41, no. 1, pp. 135–147, 1990.
- [12] N. Papaghiuc, "Semi-slant submanifolds of a Kaehlerian manifold," *Analele Universitatii "Al. I. Cuza" Iasi*, vol. 40, no. 1, pp. 55–61, 1994.
- [13] J. L. Cabrerizo, A. Carriazo, L. M. Fernández, and M. Fernández, "Semi-slant submanifolds of a Sasakian manifold," *Geometriae Dedicata*, vol. 78, no. 2, pp. 183–199, 1999.
- [14] A. Carriazo, *New Developments in Slant Submanifolds Theory*, Narosa Publishing House, New Delhi, India, 2002.
- [15] V. A. Khan and M. A. Khan, "Pseudo-slant submanifolds of a Sasakian manifold," *Indian Journal of Pure and Applied Mathematics*, vol. 38, no. 1, pp. 31–42, 2007.
- [16] H. Hashimoto and K. Mashimo, "On some 3-dimensional CR submanifolds in S^6 ," *Nagoya Mathematical Journal*, vol. 156, pp. 171–185, 1999.