Research Article

A New Legendre Collocation Method for Solving a Two-Dimensional Fractional Diffusion Equation

A. H. Bhrawy^{1,2}

¹ Department of Mathematics, Faculty of Science, King Abdulaziz University, Jeddah 21589, Saudi Arabia

² Department of Mathematics, Faculty of Science, Beni-Suef University, Beni-Suef 62511, Egypt

Correspondence should be addressed to A. H. Bhrawy; alibhrawy@yahoo.co.uk

Received 11 March 2014; Accepted 5 April 2014; Published 4 May 2014

Academic Editor: Dumitru Baleanu

Copyright © 2014 A. H. Bhrawy. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A new spectral shifted Legendre Gauss-Lobatto collocation (SL-GL-C) method is developed and analyzed to solve a class of two-dimensional initial-boundary fractional diffusion equations with variable coefficients. The method depends basically on the fact that an expansion in a series of shifted Legendre polynomials $P_{L,n}(x)P_{L,m}(y)$, for the function and its space-fractional derivatives occurring in the partial fractional differential equation (PFDE), is assumed; the expansion coefficients are then determined by reducing the PFDE with its boundary and initial conditions to a system of ordinary differential equations (SODEs) for these coefficients. This system may be solved numerically by using the fourth-order implicit Runge-Kutta (IRK) method. This method, in contrast to common finite-difference and finite-element methods, has the exponential rate of convergence for the two spatial discretizations. Numerical examples are presented in the form of tables and graphs to make comparisons with the results obtained by other methods and with the exact solutions more easier.

1. Introduction

In recent years there has been a high level of interest of employing spectral methods for numerically solving many types of integral and differential equations, due to their ease of applying them for finite and infinite domains [1–8]. The speed of convergence is one of the great advantages of spectral method. Besides, spectral methods have exponential rates of convergence; they also have high level of accuracy. From the overview of spectral approximation to differential equations, the spectral methods have been divided to four types, namely, collocation [9–11], tau [12, 13], Galerkin [14, 15], and Petrov Galerkin [16, 17] methods. The main idea of all versions of spectral methods is to express the spectral solution of the problem as a finite sum of certain basis functions (combination of orthogonal polynomials or functions) and then to choose the coefficients in order to minimize the difference between the exact and numerical solutions as possible. The spectral collocation method is a specific type of spectral methods, which is more applicable and widely used to solve almost types of differential equations [18–21].

Fractional differential equations [22–31] model many phenomena in several fields such as biology, viscoelasticity, finance, fluid mechanics, physics, chemistry, and engineering. Most fractional differential equations do not have exact solutions, so approximation and numerical techniques must be used. Here, we focus on the application of SL-GL-C scheme for the numerical solution of the time dependent fractional diffusion equation in two dimensions. This problem models phenomena exhibiting [32–34] anomalous diffusion that cannot be modeled accurately by the classical two-dimensional diffusion equations with variable coefficients. For the previous numerical methods used for time dependent fractional diffusion equation, see, for example, the papers in [35–40].

The solution u of this equation is approximated as $u_{N,M}$ which can be expressed as a finite expansion of shifted Legendre polynomials for the space variables and then evaluate the spatial fractional partial derivatives of the finite expansion of $u_{N,M}$ at the shifted Legendre-Gauss-Lobatto (SL-GL) quadrature points. Substituting these approximations in the underlined PFDE provides a SODEs in time. This system

may be solved by IRK scheme. This scheme is one of the suitable schemes for solving SODEs of first order. Since the IRK scheme has excellent stability properties, then it allows us to reduce computational costs. Indeed, this is the first work for employing SL-GL-C scheme to solve two-dimensional fractional diffusion equations.

This paper is organized as follows. We present few relevant properties of fractional integration in the Riemann-Liouville and shifted Legendre polynomials in the coming section. Section 3 presents the numerical solution of the two-dimensional fractional diffusion equations with initial-boundary conditions. Section 4 is devoted to solve two test examples. Finally, some concluding remarks are given in the last section.

2. Preliminaries and Notation

2.1. The Fractional Integration in the Riemann-Liouville Sense. There are several definitions of the fractional integration of order $\nu > 0$ and not necessarily equivalent to each other; see [41, 42]. The most used definition is due to Riemann-Liouville, which is defined as

$$J^\nu f(x) = \frac{1}{\Gamma(\nu)} \int_0^x (x-t)^{\nu-1} f(t) dt, \quad \nu > 0, \quad x > 0, \quad (1)$$

$$J^0 f(x) = f(x).$$

One of the basic properties of the operator J^ν is

$$J^\nu x^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta+1+\nu)} x^{\beta+\nu}. \quad (2)$$

The Riemann-Liouville fractional derivative of order ν will be denoted by D^ν . The next relation defines the Riemann-Liouville fractional derivative of order ν as

$$D^\nu f(x) = \frac{d^m}{dx^m} (J^{m-\nu} f(x)), \quad (3)$$

where $m-1 < \nu \leq m$, $m \in \mathbb{N}$, and m is the smallest integer greater than ν .

Lemma 1. If $m-1 < \nu \leq m$, $m \in \mathbb{N}$, then

$$D^\nu J^\nu f(x) = f(x),$$

$$J^\nu D^\nu f(x) = f(x) - \sum_{i=0}^{m-1} f^{(i)}(0^+) \frac{x^i}{i!}, \quad x > 0. \quad (4)$$

2.2. Properties of Shifted Legendre Polynomials. The well-known Legendre polynomials $P_i(x)$ are defined on the interval $(-1, 1)$. Some properties about the standard Legendre polynomials will be recalled in this section. The Legendre polynomials $P_k(x)$ ($k = 0, 1, \dots$) satisfy the following Rodrigue's formula:

$$P_k(x) = \frac{(-1)^k}{2^k k!} D^k \left((1-x^2)^k \right). \quad (5)$$

We recall also that $P_k(x)$ is a polynomial of degree k , and therefore $P_k^{(q)}(x)$ (the q th derivative of $P_k(x)$) is

$$P_k^{(q)}(x) = \sum_{i=0(k+i=\text{even})}^{k-q} C_q(k, i) P_i(x), \quad (6)$$

where

$$C_q(k, i) = 2^{q-1} (2i+1) \Gamma\left(\frac{q+k-i}{2}\right) \Gamma\left(\frac{q+k+i+1}{2}\right) \times \left(\Gamma(q) \Gamma\left(\frac{2-q+k-i}{2}\right) \Gamma\left(\frac{3-q+k+i}{2}\right) \right)^{-1}. \quad (7)$$

The Legendre polynomials satisfy the following relations:

$$P_0(x) = 1, \quad P_1(x) = x,$$

$$P_{k+2}(x) = \frac{2k+3}{k+2} x P_{k+1}(x) - \frac{k+1}{k+2} P_k(x), \quad (8)$$

and the orthogonality relation

$$(P_k(x), P_l(x)) = \int_{-1}^1 P_k(x) P_l(x) w(x) dx = h_k \delta_{lk}, \quad (9)$$

where $w(x) \equiv 1$, $h_k = 2/(2k+1)$, and δ_{lk} is the Kronecker delta function. The Legendre-Gauss-Lobatto quadrature is used to evaluate the previous integral accurately. For any $\phi \in S_{2N-1}[-1, 1]$, we get

$$\int_{-1}^1 \phi(x) dx = \sum_{j=0}^N \omega_{N,j} \phi(x_{N,j}). \quad (10)$$

We introduce the following discrete inner product:

$$(u, v)_N = \sum_{j=0}^N u(x_{N,j}) v(x_{N,j}) \omega_{N,j}, \quad (11)$$

where $x_{N,j}$ ($0 \leq j \leq N$) and $\omega_{N,j}$ ($0 \leq j \leq N$) are used as usual nodes and the corresponding Christoffel numbers in the interval $[-1, 1]$, respectively. Let the shifted Legendre polynomials $P_i((2x/L) - 1)$ be denoted by $P_{L,i}(x)$; $x \in (0, L)$. Then $P_{L,i}(x)$ can be obtained with the aid of the following recurrence formula:

$$(i+1) P_{L,i+1}(x) = (2i+1) \left(\frac{2x}{L} - 1 \right) P_{L,i}(x) - i P_{L,i-1}(x), \quad i = 1, 2, \dots \quad (12)$$

The analytic form of the shifted Legendre polynomials $P_{L,i}(x)$ of degree i is given by

$$P_{L,i}(x) = \sum_{k=0}^i (-1)^{i+k} \frac{(i+k)!}{(i-k)!(k!)^2 L^k} x^k. \quad (13)$$

The Riemann-Liouville fractional integration of shifted Legendre polynomials $P_{L,i}(x)$ of degree i is given by

$$\begin{aligned} J^\nu P_{L,i}(x) &= \sum_{k=0}^i (-1)^{i+k} \frac{(i+k)!}{(i-k)!(k!)^2 L^k} J^\nu x^k \\ &= \sum_{k=0}^i (-1)^{i+k} \frac{(i+k)!k!}{(i-k)!(k!)^2 L^k (k+\nu+1)} x^{k+\nu}, \\ &\quad i = 0, 1, \dots, N, \\ D^\nu P_{L,i}(x) &= \sum_{k=0}^i (-1)^{i+k} \frac{(i+k)!}{(i-k)!(k!)^2 L^k} J^\nu x^k \\ &= \sum_{k=0}^i (-1)^{i+k} \frac{(i+k)!\Gamma(k+1)}{(i-k)!(k!)^2 L^k \Gamma(k+1-\nu)} x^{k-\nu}, \\ &\quad i = 0, 1, \dots, N, \end{aligned} \quad (14)$$

where $P_{L,i}(0) = (-1)^i$. The orthogonality condition is

$$\int_0^L P_{L,j}(x) P_{L,k}(x) w_L(x) dx = h_{L,k} \delta_{jk}, \quad (15)$$

where $w_L(x) \equiv 1$ and $h_{L,k} = L/(2k+1)$.

The function $u(x)$, square integrable in $(0, L)$, may be expressed in terms of shifted Legendre polynomials as

$$u(x) = \sum_{j=0}^{\infty} c_j P_{L,j}(x), \quad (16)$$

where the coefficients c_j are given by

$$c_j = \frac{1}{h_{L,j}} \int_0^L u(x) P_{L,j}(x) dx, \quad j = 0, 1, 2, \dots \quad (17)$$

In practice, only the first $(N+1)$ -terms shifted Legendre polynomials are considered. Hence $u(x)$ can be expressed in the form

$$u_N(x) \approx \sum_{j=0}^N c_j P_{L,j}(x). \quad (18)$$

3. Shifted Legendre Spectral Collocation Method

In this section, we derive the SL-GL-C method for solving two-dimensional fractional diffusion equation of the form

$$\begin{aligned} &\frac{\partial v(x, y, t)}{\partial t} \\ &= g_1(x, y) \frac{\partial^{\nu_1} v(x, y, t)}{\partial x^{\nu_1}} + g_2(x, y) \frac{\partial^{\nu_2} v(x, y, t)}{\partial y^{\nu_2}} \\ &\quad + f(x, y, t), \quad (x, y, t) \in [0, L_1] \times [0, L_2] \times (0, T], \end{aligned} \quad (19)$$

which is subject to the initial condition

$$v(x, y, 0) = g_3(x, y), \quad (x, y) \in [0, L_1] \times [0, L_2] \quad (20)$$

and boundary conditions

$$\begin{aligned} v(0, y, t) &= g_4(y, t), & v(L_1, y, t) &= g_5(y, t), \\ (y, t) &\in [0, L_2] \times [0, T], \\ v(x, 0, t) &= g_6(x, t), & v(x, L_2, t) &= g_7(x, t) \\ (x, t) &\in [0, L_1] \times [0, T]. \end{aligned} \quad (21)$$

Denote by $x_{N,j}(x_{L,N,j})$, $0 \leq j \leq N$, and $\omega_{N,j}(\omega_{L,N,j})$, $0 \leq i \leq N$, the nodes and Christoffel numbers of the standard (shifted) Legendre-Gauss-Lobatto quadrature on the intervals $(-1, 1)$, $(0, L)$, respectively. Then one can clearly deduce that

$$\begin{aligned} x_{L,N,j} &= \frac{L}{2} (x_{N,j} + 1), \quad 0 \leq j \leq N, \\ \omega_{L,N,j} &= \left(\frac{L}{2}\right)^{\alpha+\beta+1} \omega_{N,j}, \quad 0 \leq j \leq N, \end{aligned} \quad (22)$$

and if $S_N(0, L)$ denotes the set of all polynomials of degree at most N , then it follows for any $\phi \in S_{2N-1}(0, L)$ that

$$\begin{aligned} \int_0^L w_L(x) \phi(x) dx &= \left(\frac{L}{2}\right) \int_{-1}^1 \phi\left(\frac{L}{2}(x+1)\right) dx \\ &= \left(\frac{L}{2}\right) \sum_{j=0}^N \omega_{N,j} \phi\left(\frac{L}{2}(x_{N,j}+1)\right) \\ &= \sum_{j=0}^N \omega_{L,N,j} \phi(x_{L,N,j}). \end{aligned} \quad (23)$$

Here, the SL-GL-C method will be employed to transform the previous two-dimensional fractional diffusion equation into SODEs. Let us expand the dependent variables in the form

$$v(x, y, t) = \sum_{i=0}^N \sum_{j=0}^M a_{i,j}(t) P_{L_1,i}(x) P_{L_2,j}(y). \quad (24)$$

We observe from (15) that the coefficients $a_{i,j}(t)$ can be approximated by

$$\begin{aligned} a_{i,j}(t) &= \frac{1}{h_{L_1,i} h_{L_2,j}} \\ &\times \sum_{l=0}^N \sum_{k=0}^M (P_{L_2,j}(y_{L_2,M,k}) \omega_{M,k} P_{L_1,i}(x_{L_1,N,l}) \omega_{N,l}) \\ &\times v(x_{L_1,N,l}, y_{L_2,M,k}, t). \end{aligned} \quad (25)$$

TABLE 1: Maximum absolute errors for problem (40) for $T = 1$.

$N = M$	SL-GL-C method		$\Delta x = \Delta y = \Delta t$	[43]
	M_E^t	M_E		
4	3.41785×10^{-3}	9.11427×10^{-3}	—	—
6	2.13706×10^{-5}	6.83861×10^{-5}	$\frac{1}{5}$	1.05564×10^{-3}
8	1.32866×10^{-6}	4.85909×10^{-6}	$\frac{1}{10}$	2.61694×10^{-4}
10	1.74023×10^{-7}	7.07142×10^{-7}	$\frac{1}{20}$	6.57067×10^{-5}
12	3.43827×10^{-8}	1.52415×10^{-7}	$\frac{1}{40}$	1.55866×10^{-5}

TABLE 2: Maximum absolute errors for problem (40) for $T = 1$ and $N = M = 12$.

x	y	t	$E(x, y, t)$	x	y	t	$E(x, y, t)$	x	y	t	$E(x, y, t)$
0.1	0.1	1	2.74515×10^{-11}	0.1	0.5	1	9.17923×10^{-12}	0.1	1	1	4.41521×10^{-11}
0.2			2.13241×10^{-10}	0.2			7.57278×10^{-11}	0.2			3.58585×10^{-10}
0.3			7.22081×10^{-10}	0.3			2.61927×10^{-10}	0.3			1.20622×10^{-9}
0.4			1.71192×10^{-9}	0.4			6.43406×10^{-10}	0.4			2.93613×10^{-9}
0.5			3.35365×10^{-9}	0.5			1.29893×10^{-9}	0.5			5.8445×10^{-9}
0.6			5.80344×10^{-9}	0.6			2.33485×10^{-9}	0.6			1.02404×10^{-8}
0.7			9.20721×10^{-9}	0.7			3.82976×10^{-9}	0.7			1.65193×10^{-8}
0.8			1.39884×10^{-8}	0.8			5.84963×10^{-9}	0.8			2.50569×10^{-8}
0.9			2.06847×10^{-8}	0.9			8.5766×10^{-9}	0.9			3.64012×10^{-8}
1			2.75218×10^{-8}	1			1.26484×10^{-8}	1			5.60815×10^{-8}

TABLE 3: Maximum absolute errors for problem (46) for $T = 1$ and $v_1 = v_2 = 1.5$.

$N = M$	M_E^t	M_E
6	2.57399×10^{-8}	1.02853×10^{-7}
8	1.51431×10^{-8}	1.21098×10^{-7}
10	8.24341×10^{-14}	4.13336×10^{-13}

The expression (24) can be rewritten as

$$\begin{aligned}
 v(x, y, t) &= \sum_{i=0}^N \sum_{j=0}^M \sum_{l=0}^N \sum_{k=0}^M \frac{(P_{L_2,j}(y_{L_2,M,k}) \omega_{M,k} P_{L_1,i}(x_{L_1,N,l}) \omega_{N,l})}{h_{L_1,i} h_{L_2,j}} \\
 &\quad \times P_{L_1,i}(x) P_{L_2,j}(y) v_{l,k}(t), \quad (26)
 \end{aligned}$$

where $v(x_{L_1,N,l}, y_{L_2,M,k}, t) = v_{l,k}(t)$. In what follows, the approximation of the fractional spatial partial derivative

$\partial^{v_1} v(x, y, t) / \partial x^{v_1}$ with respect to x for $v(x, y, t)$ can be computed as

$$\begin{aligned}
 &\frac{\partial^{v_1} v(x, y, t)}{\partial x^{v_1}} \\
 &= \sum_{i=0}^N \sum_{j=0}^M \sum_{l=0}^N \sum_{k=0}^M \frac{(P_{L_2,j}(y_{L_2,M,k}) \omega_{M,k} P_{L_1,i}(x_{L_1,N,l}) \omega_{N,l})}{h_{L_1,i} h_{L_2,j}} \\
 &\quad \times D^{(v_1)}(P_{L_1,i}(x)) P_{L_2,j}(y) v_{l,k}(t). \quad (27)
 \end{aligned}$$

Also, the fractional spatial partial derivative of fractional order $\partial^{v_2} v(x, y, t) / \partial y^{v_2}$ with respect to y of the approximate solution is

$$\begin{aligned}
 &\frac{\partial^{v_2} v(x, y, t)}{\partial y^{v_2}} \\
 &= \sum_{l=0}^N \sum_{k=0}^M \sum_{i=0}^N \sum_{j=0}^M \frac{(P_{L_2,j}(y_{L_2,M,k}) \omega_{M,k} P_{L_1,i}(x_{L_1,N,l}) \omega_{N,l})}{h_{L_1,i} h_{L_2,j}} \\
 &\quad \times P_{L_1,i}(x) D^{(v_2)}(P_{L_2,j}(y)) v_{l,k}(t). \quad (28)
 \end{aligned}$$

TABLE 4: Maximum absolute errors for problem (46) for $T = 1$ and $\nu_1 = \nu_2 = 1.5$.

x	y	t	$E(x, y, t)$	x	y	t	$E(x, y, t)$	x	y	t	$E(x, y, t)$
0.1	0.1	1	4.06522×10^{-16}	0.1	0.5	1	7.05989×10^{-15}	0.1	1	1	2.35055×10^{-15}
0.2			3.12066×10^{-15}	0.2			7.445×10^{-15}	0.2		1	6.63792×10^{-15}
0.3			4.22817×10^{-15}	0.3			1.73386×10^{-15}	0.3		1	6.92588×10^{-15}
0.4			8.7365×10^{-15}	0.4			4.25354×10^{-15}	0.4		1	1.70948×10^{-14}
0.5			1.40339×10^{-14}	0.5			1.56351×10^{-14}	0.5		1	2.83974×10^{-14}
0.6			1.81075×10^{-14}	0.6			1.14631×10^{-14}	0.6		1	3.44343×10^{-14}
0.7			2.69246×10^{-14}	0.7			1.8454×10^{-14}	0.7		1	5.44946×10^{-14}
0.8			3.42964×10^{-14}	0.8			3.14054×10^{-14}	0.8		1	6.78416×10^{-14}
0.9			4.3951×10^{-14}	0.9			2.78319×10^{-14}	0.9		1	9.03513×10^{-14}
1			3.1379×10^{-15}	1			1.20095×10^{-14}	1		1	1.0103×10^{-14}

The previous fractional derivatives can be computed at the SL-GL interpolation nodes as

$$\left(\frac{\partial^{\nu_1} v(x, y, t)}{\partial x^{\nu_1}} \right)_{y=y_{L_2, M, m}}^{x=x_{L_1, N, n}} = \sum_{l=0}^N \sum_{k=0}^M \rho_{l, k}^{n, m} v_{l, k}(t), \quad n = 0, 1, \dots, N, \quad m = 0, 1, \dots, M, \quad (29)$$

$$\left(\frac{\partial^{\nu_2} v(x, y, t)}{\partial y^{\nu_2}} \right)_{y=y_{L_2, M, m}}^{x=x_{L_1, N, n}} = \sum_{l=0}^N \sum_{k=0}^M \lambda_{l, k}^{n, m} v_{l, k}(t), \quad n = 0, 1, \dots, N, \quad m = 0, 1, \dots, M,$$

In the proposed SL-GL-C method the residual of (19) is set to zero at $(N-1) \times (M-1)$ of SL-GL points. Moreover, the values of $v_{0, k}(t)$, $v_{N, k}(t)$, $v_{l, 0}(t)$, and $v_{l, M}(t)$ are given by

$$v_{0, k}(t) = g_4(y_{L_2, M, k}, t), \quad v_{N, k}(t) = g_5(y_{L_2, M, k}, t), \quad k = 0, \dots, M, \quad (31)$$

$$v_{l, 0}(t) = g_6(x_{L_1, N, l}, t), \quad v_{l, M}(t) = g_7(x_{L_1, N, l}, t), \quad l = 0, \dots, N.$$

Therefore, adapting (26)–(31) enables one to write (19)–(21) as a $(N-1) \times (M-1)$ SODEs:

$$\dot{v}_{n, m}(t) = g_1^{n, m} \sum_{l=0}^N \sum_{k=0}^M \rho_{l, k}^{n, m} v_{l, k}(t) + g_2^{n, m} \sum_{l=0}^N \sum_{k=0}^M \lambda_{l, k}^{n, m} v_{l, k}(t)$$

where

$$+ f^{n, m}(t), \quad n = 1, \dots, N-1, \quad m = 1, \dots, M-1, \quad (32)$$

which is subject to the initial conditions:

$$v_{n, m}(0) = g_3^{n, m} \left(x_{L_1, N, n}, y_{L_2, M, m} \right) = g_1^{n, m}, \quad n = 1, \dots, N-1, \quad m = 1, \dots, M-1, \quad (33)$$

where

$$g_2(x_{L_1, N, n}, y_{L_2, M, m}) = g_2^{n, m}, \quad (34)$$

$$g_3(x_{L_1, N, n}, y_{L_2, M, m}) = g_3^{n, m},$$

$$f(x_{L_1, N, n}, y_{L_2, M, m}, t) = f^{n, m}(t).$$

$$\rho_{l, k}^{n, m} = \sum_{i=0}^N \sum_{j=0}^M \frac{(P_{L_2, j}(y_{L_2, M, k}) \omega_{M, k} P_{L_1, i}(x_{L_1, N, l}) \omega_{N, l})}{h_i h_j} \times D^{(\nu_1)} P_{L_1, i}(x_{L_1, N, n}) P_{L_2, j}(y_{L_2, M, m}), \quad (30)$$

$$\lambda_{l, k}^{n, m} = \sum_{i=0}^N \sum_{j=0}^M \frac{(P_{L_2, j}(y_{L_2, M, k}) \omega_{M, k} P_{L_1, i}(x_{L_1, N, l}) \omega_{N, l})}{h_i h_j} \times D^{(\nu_2)} P_{L_2, j}(y_{L_2, M, m}) P_{L_1, i}(x_{L_1, N, n}).$$

Finally, we can arrange (32) and (33) to their matrix formulation, which can be solved by IRK scheme:

$$\begin{aligned}
 & \begin{pmatrix} \dot{v}_{1,1}(t) & \cdots & \dot{v}_{1,M-1}(t) \\ \dot{v}_{2,1}(t) & \vdots & \dot{v}_{2,M-1}(t) \\ \cdots & \ddots & \cdots \\ \dot{v}_{N-2,1}(t) & \vdots & \dot{v}_{N-2,M-1}(t) \\ \dot{v}_{N-1,1}(t) & \cdots & \dot{v}_{N-1,M-1}(t) \end{pmatrix} \\
 &= \begin{pmatrix} H_{1,1}(t, v_1, \dots, v_{N-1}) & \cdots & H_{1,M-1}(t, v_1, \dots, v_{N-1}) \\ H_{2,1}(t, v_1, \dots, v_{N-1}) & \vdots & H_{2,M-1}(t, v_1, \dots, v_{N-1}) \\ \cdots & \ddots & \cdots \\ H_{N-2,1}(t, v_1, \dots, v_{N-1}) & \vdots & H_{N-2,M-1}(t, v_1, \dots, v_{N-1}) \\ H_{N-1,1}(t, v_1, \dots, v_{N-1}) & \cdots & H_{N-1,M-1}(t, v_1, \dots, v_{N-1}) \end{pmatrix}, \\
 & \begin{pmatrix} v_{1,1}(0) & \cdots & v_{1,M-1}(0) \\ v_{2,1}(0) & \vdots & v_{2,M-1}(0) \\ \cdots & \ddots & \cdots \\ v_{N-2,1}(0) & \vdots & v_{N-2,M-1}(0) \\ v_{N-1,1}(0) & \cdots & v_{N-1,M-1}(0) \end{pmatrix} \\
 &= \begin{pmatrix} g_3^{1,1} & \cdots & g_3^{1,M-1} \\ g_3^{2,1} & \vdots & g_3^{2,M-1} \\ \cdots & \ddots & \cdots \\ g_3^{N-2,1} & \vdots & g_3^{N-2,M-1} \\ g_3^{N-1,1} & \cdots & g_3^{N-1,M-1} \end{pmatrix}, \quad t \in [-\tau, 0],
 \end{aligned} \tag{35}$$

where

$$\begin{aligned}
 & H_{n,m}(t, v_{1,1}, \dots, v_{N-1,M-1}) \\
 &= g_1^{n,m} \sum_{l=0}^N \sum_{k=0}^M \rho_{l,k}^{n,m} v_{l,k}(t) + g_2^{n,m} \sum_{l=0}^N \sum_{k=0}^M \lambda_{l,k}^{n,m} v_{l,k}(t) \\
 &+ f^{n,m}(t), \quad n = 1, \dots, N-1, \quad m = 1, \dots, M-1.
 \end{aligned} \tag{36}$$

The Runge-Kutta method can be expressed as one of powerful numerical integrations tools used for initial value SODEs of first order. The IRK method presents a subclass of the well-known family of Runge-Kutta methods and has many applications in the numerical solution of systems of ordinary differential equations.

4. Numerical Results

In this section, we present two numerical examples to show the accuracy, robustness, and applicability of the proposed

FIGURE 1: Space graph of numerical solution of problem (40).

FIGURE 2: Space graph of absolute error of problem (40).

method. We compare the results obtained by our method with those obtained in [43].

The difference between the measured value of the approximate solution and the exact solution (absolute error) is given by

$$E(\mathbf{x}, \mathbf{y}, t) = |u(\mathbf{x}, \mathbf{y}, t) - u_N(\mathbf{x}, \mathbf{y}, t)|, \tag{37}$$

where $\mathbf{x}$, $\mathbf{y}$, t , $u(\mathbf{x}, \mathbf{y}, t)$, and $u_N(\mathbf{x}, \mathbf{y}, t)$ are the space vectors, time, exact, and numerical solutions, respectively. Moreover, the maximum absolute error (MAEs) is given by

$$M_E = \text{Max} \{E(\mathbf{x}, \mathbf{y}, t) : \text{over all domains}\}. \tag{38}$$

Also, we define the norm infinity as

$$M_E^t = \text{Max} \{E(\mathbf{x}, \mathbf{y}, t_k) : \text{over all space domains}\}. \tag{39}$$

Example 2. Consider the two-dimensional problem [43] is

$$\begin{aligned}
 & \frac{\partial u}{\partial t}(x, y, t) \\
 &= g_1(x, t) \frac{\partial^{1.8} u(x, y, t)}{\partial x^{1.8}} + g_2(x, t) \frac{\partial^{1.6} u(x, y, t)}{\partial x^{1.6}} \\
 &+ f(x, y, t), \quad (x, y, t) \in [0, 1] \times [0, 1] \times [0, T],
 \end{aligned} \tag{40}$$

FIGURE 3: x -direction curves of numerical and exact solutions of problem (40).

FIGURE 4: y -direction curves of numerical and exact solutions of problem (40).

with the diffusion coefficients

$$g_1(x, t) = \frac{\Gamma(2.2) x^{2.8} y}{6}, \quad g_2(x, t) = \frac{2xy^{2.6}}{\Gamma(4.6)}, \quad (41)$$

and consider the forcing function is

$$f(x, y, t) = -(1 + 2xy) e^{-t} x^3 y^{3.6}, \quad (42)$$

with the initial condition

$$u(x, y, 0) = x^3 y^{3.6} \quad (43)$$

and Dirichlet boundary conditions

$$\begin{aligned} u(0, y, t) = u(x, 0, t) = 0, \quad u(1, y, t) = e^{-t} y^{3.6}, \\ u(x, 1, t) = e^{-t} x^3 \quad \forall t \geq 0. \end{aligned} \quad (44)$$

The exact solution of this two-dimensional fractional diffusion equation is given by

$$u(x, y, t) = e^{-t} x^3 y^{3.6}. \quad (45)$$

From the first look on Table 1, we see high accurate results based on MAEs at different choices of N and M , compared with the results given in [43]. Moreover, results listed in Table 2 confirm that the proposed method is very accurate.

In Figure 1, we display the numerical solution of problem (40) computed by the present method at $N = M = 12$, while we plot the space graph of absolute error of problem (40) computed by the present method at $N = M = 12$ in Figure 2. As shown in Figures 3 and 4, we see the agreement of numerical and exact solutions curves.

Example 3. Consider the two-dimensional fractional diffusion equation is

$$\begin{aligned} \frac{\partial u}{\partial t}(x, y, t) \\ = g_1(x, t) \frac{\partial^{v_1} u(x, y, t)}{\partial x^{v_1}} + g_2(x, t) \frac{\partial^{v_2} u(x, y, t)}{\partial y^{v_2}} \end{aligned} \quad (46)$$

$$+ f(x, y, t), \quad (x, y, t) \in [0, 1] \times [0, 1] \times [0, T],$$

with the diffusion coefficients

$$\begin{aligned} g_1(x, y) &= \frac{(3 - 2x) \Gamma(3 - v_1)}{2}, \\ g_2(x, y) &= \frac{(4 - y) \Gamma(4 - v_2)}{6}, \end{aligned} \quad (47)$$

and consider the forcing function is

$$f(x, y, t) = e^{-t} (x^2 (-y^{3/2} + y - 4) y^{3/2} + \sqrt{x} (2x - 3) y^3), \quad (48)$$

with the initial condition

$$u(x, y, 0) = x^2 y^3 \quad (49)$$

and the Dirichlet boundary conditions

$$\begin{aligned} u(0, y, t) = u(x, 0, t) = 0, \quad u(1, y, t) = e^{-t} y^3, \\ u(x, 1, t) = e^{-t} x^2 \quad \forall t \geq 0. \end{aligned} \quad (50)$$

The exact solution of this two-dimensional fractional diffusion equation is given by

$$u(x, y, t) = e^{-t} x^2 y^3. \quad (51)$$

More accurate results for maximum absolute and absolute errors of (46) have been obtained in Tables 3 and 4, respectively, with values of parameters listed in their captions, the approximate solution $u_{N,M}$ at $v_1 = v_2 = 1.5$ and $N = M = 10$ is plotted in Figure 5. Moreover, we plot the absolute error for this problem at $v_1 = v_2 = 1.5$ and $N = M = 10$ in Figure 6. Finally, the x - and y -directions curves of absolute error at $v_1 = v_2 = 1.5$ and $N = M = 10$ are displayed in Figures 7 and 8.

FIGURE 5: Space graph of numerical solution of problem (46) at $\nu_1 = \nu_2 = 1.5$ and $N = M = 10$.

FIGURE 6: Space graph of absolute error of problem (46) at $\nu_1 = \nu_2 = 1.5$ and $N = M = 10$.

5. Concluding Remarks and Future Work

The SL-GL-C method was investigated successfully in spatial discretizations to get accurate approximate solution of the two-dimensional fractional diffusion equations with variable coefficients. This problem was transformed into SODEs in time variable which greatly simplifies the problem. The IRK scheme was then applied to the resulting system. From the numerical experiments, the obtained results demonstrated the effectiveness and the high accuracy of SL-GL-C method for solving the mentioned problem.

The technique can be extended to more sophisticated problems in two- and three-dimensional spaces. In principle, this method may be extended to related problems in mathematical physics. It is possible to use other orthogonal polynomials, say, Chebyshev polynomials or Jacobi polynomials, to solve the mentioned problem in this paper. Furthermore, the proposed spectral method might be developed by considering the Legendre pseudospectral approximation in both temporal and spatial discretizations. We should note that, as a numerical method, we are restricted to solving problem over a finite domain. Also, the pseudospectral approximation might be employed based on generalized

FIGURE 7: x -direction curve of absolute error of problem (46) at $\nu_1 = \nu_2 = 1.5$ and $N = M = 10$.

FIGURE 8: y -direction curves of absolute error of problem (46) at $\nu_1 = \nu_2 = 1.5$ and $N = M = 10$.

Laguerre or modified generalized Laguerre polynomials [44] to solve similar problems in a semi-infinite spatial intervals.

Conflict of Interests

The author declares that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

This paper was funded by the Deanship of Scientific Research DSR, King Abdulaziz University, Jeddah, Grant no. 130-53-D1435. The author, therefore, acknowledges with thanks DSR for the technical and financial support.

References

- [1] C. Canuto, M. Y. Hussaini, A. Quarteroni, and T. A. Zang, *Spectral Methods: Fundamentals in Single Domains*, Springer, New York, NY, USA, 2006.
- [2] H. Schamel and K. Elsaesser, "The application of the spectral method to nonlinear wave propagation," *Journal of Computational Physics*, vol. 22, no. 4, pp. 501–516, 1976.
- [3] W. M. Abd-Elhameed, E. H. Doha, and Y. H. Youssri, "Efficient spectral-Petrov-Galerkin methods for third- and fifth-order differential equations using general parameters generalized

- Jacobi polynomials," *Quaestiones Mathematicae*, vol. 36, pp. 15–38, 2013.
- [4] E. H. Doha, A. H. Bhrawy, M. A. Abdelkawy, and R. M. Hafez, "A Jacobi collocation approximation for nonlinear coupled viscous Burgers' equation," *Central European Journal of Physics*, vol. 12, pp. 111–122, 2014.
 - [5] H. Adibi and A. M. Rismani, "On using a modified Legendre-spectral method for solving singular IVPs of Lane-Emden type," *Computers and Mathematics with Applications*, vol. 60, no. 7, pp. 2126–2130, 2010.
 - [6] E. H. Doha, A. H. Bhrawy, M. A. Abdelkawy, and R. A. van Gorder, "Jacobi-Gauss-Lobatto collocation method for the numerical solution of 1 + 1 nonlinear Schrodinger equations," *Journal of Computational Physics*, vol. 261, pp. 244–255, 2014.
 - [7] E. H. Doha, A. H. Bhrawy, D. Baleanu, and R. M. Hafez, "A new Jacobi rational-Gauss collocation method for numerical solution of generalized Pantograph equations," *Applied Numerical Mathematics*, vol. 77, pp. 43–54, 2014.
 - [8] F. M. Mahfouz, "Numerical simulation of free convection within an eccentric annulus filled with micropolar fluid using spectral method," *Applied Mathematics and Computation*, vol. 219, pp. 5397–5409, 2013.
 - [9] J. Ma, B.-W. Li, and J. R. Howell, "Thermal radiation heat transfer in one- and two-dimensional enclosures using the spectral collocation method with full spectrum k-distribution model," *International Journal of Heat and Mass Transfer*, vol. 71, pp. 35–43, 2014.
 - [10] X. Ma and C. Huang, "Spectral collocation method for linear fractional integro-differential equations," *Applied Mathematical Modelling*, vol. 38, pp. 1434–1448, 2014.
 - [11] W. M. Abd-Elhameed, E. H. Doha, and Y. H. Youssri, "New wavelets collocation method for solving second-order multi-point boundary value problems using Chebyshev polynomials of third and fourth," *Abstract and Applied Analysis*, vol. 2013, Article ID 542839, 9 pages, 2013.
 - [12] S. R. Lau and R. H. Price, "Sparse spectral-tau method for the three-dimensional helically reduced wave equation on two-center domains," *Journal of Computational Physics*, vol. 231, pp. 7695–7714, 2012.
 - [13] F. Ghoreishi and S. Yazdani, "An extension of the spectral Tau method for numerical solution of multi-order fractional differential equations with convergence analysis," *Computers and Mathematics with Applications*, vol. 61, no. 1, pp. 30–43, 2011.
 - [14] E. H. Doha and A. H. Bhrawy, "An efficient direct solver for multidimensional elliptic Robin boundary value problems using a Legendre spectral-Galerkin method," *Computers and Mathematics with Applications*, vol. 64, pp. 558–571, 2012.
 - [15] T. Boaca and I. Boaca, "Spectral galerkin method in the study of mass transfer in laminar and turbulent flows," *Computer Aided Chemical Engineering*, vol. 24, pp. 99–104, 2007.
 - [16] E. H. Doha, A. H. Bhrawy, and R. M. Hafez, "A Jacobi-Jacobi dual-Petrov-Galerkin method for third- and fifth-order differential equations," *Mathematical and Computer Modelling*, vol. 53, no. 9–10, pp. 1820–1832, 2011.
 - [17] W. M. Abd-Elhameed, E. H. Doha, and M. A. Bassuony, "Two Legendre-Dual-Petrov-Galerkin algorithms for solving the integrated forms of high odd-order boundary value problems," *The Scientific World Journal*, vol. 2013, Article ID 309264, 11 pages, 2013.
 - [18] L. Wang, Y. Ma, and Z. Meng, "Haar wavelet method for solving fractional partial differential equations numerically," *Applied Mathematics and Computation*, vol. 227, pp. 66–76, 2014.
 - [19] A. Golbabai and M. Javidi, "A numerical solution for non-classical parabolic problem based on Chebyshev spectral collocation method," *Applied Mathematics and Computation*, vol. 190, no. 1, pp. 179–185, 2007.
 - [20] A. H. Bhrawy, "A Jacobi-Gauss-Lobatto collocation method for solving generalized Fitzhugh-Nagumo equation with time-dependent coefficients," *Applied Mathematics and Computation*, vol. 222, pp. 255–264, 2013.
 - [21] E. H. Doha, A. H. Bhrawy, and R. M. Hafez, "On shifted Jacobi spectral method for high-order multi-point boundary value problems," *Communications in Nonlinear Science and Numerical Simulation*, vol. 17, no. 10, pp. 3802–3810, 2012.
 - [22] R. Garrappa and M. Popolizio, "On the use of matrix functions for fractional partial differential equations," *Mathematics and Computers in Simulation*, vol. 81, no. 5, pp. 1045–1056, 2011.
 - [23] A. A. Pedas and E. Tamme, "Numerical solution of nonlinear fractional differential equations by spline collocation methods," *Journal of Computational and Applied Mathematics*, vol. 255, pp. 216–230, 2014.
 - [24] F. Gao, X. Lee, F. Fei, H. Tong, Y. Deng, and H. Zhao, "Identification time-delayed fractional order chaos with functional extrema model via differential evolution," *Expert Systems With Applications*, vol. 41, pp. 1601–1608, 2014.
 - [25] Y. Zhao, D. F. Cheng, and X. J. Yang, "Approximation solutions for local fractional Schrödinger equation in the one-dimensional Cantorian system," *Advances in Mathematical Physics*, vol. 2013, Article ID 291386, 5 pages, 2013.
 - [26] A. H. Bhrawy and M. M. Al-Shomrani, "A shifted Legendre spectral method for fractional-order multi-point boundary value problems," *Advances in Difference Equations*, vol. 2012, article 8, 2012.
 - [27] J. W. Kirchner, X. Feng, and C. Neal, "Frail chemistry and its implications for contaminant transport in catchments," *Nature*, vol. 403, no. 6769, pp. 524–526, 2000.
 - [28] A. Pedas and E. Tamme, "Piecewise polynomial collocation for linear boundary value problems of fractional differential equations," *Journal of Computational and Applied Mathematics*, vol. 236, no. 13, pp. 3349–3359, 2012.
 - [29] A. Ahmadian, M. Suleiman, S. Salahshour, and D. Baleanu, "A Jacobi operational matrix for solving a fuzzy linear fractional differential equation," *Advances in Difference Equations*, vol. 2013, article 104, 29 pages, 2013.
 - [30] A. H. Bhrawy and M. A. Alghamdi, "A shifted Jacobi-Gauss-Lobatto collocation method for solving nonlinear fractional Langevin equation involving two fractional orders in different intervals," *Boundary Value Problems*, vol. 2012, article 62, 2012.
 - [31] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
 - [32] E. Kotomin and V. Kuzovkov, *Modern Aspects of Diffusion-Controlled Reactions: Cooperative Phenomena in Bimolecular Processes*, Comprehensive Chemical Kinetics, Elsevier, 1996.
 - [33] D. Baleanu, K. Diethelm, E. Scalas, and J. J. Trujillo, *Fractional Calculus Models and Numerical Methods*, Series on Complexity, Nonlinearity and Chaos, World Scientific, Singapore, 2012.
 - [34] R. S. Cantrell and C. Cosner, *Spatial Ecology via Reaction Diffusion Equations*, Wiley, 2004.
 - [35] H. Wang and N. Du, "Fast solution methods for space-fractional diffusion equations," *Journal of Computational and Applied Mathematics*, vol. 255, pp. 376–383, 2014.
 - [36] A. H. Bhrawy and D. Baleanu, "A spectral Legendre-Gauss-Lobatto collocation method for a space-fractional advection

- diffusion equations with variable coefficients," *Reports on Mathematical Physics*, vol. 72, pp. 219–233, 2013.
- [37] E. H. Doha, A. H. Bhrawy, and S. S. Ezz-Eldien, "Numerical approximations for fractional diffusion equations via a Chebyshev spectral-tau method," *Central European Journal of Physics*, vol. 11, pp. 1494–1503, 2013.
 - [38] F. Liu, P. Zhuang, I. Turner, K. Burrage, and V. Anh, "A new fractional finite volume method for solving the fractional diffusion equation," *Applied Mathematical Modelling*, 2013.
 - [39] S. B. Yuste and J. Quintana-Murillo, "A finite difference method with non-uniform timesteps for fractional diffusion equations," *Computer Physics Communications*, vol. 183, pp. 2594–2600, 2012.
 - [40] K. Wang and H. Wang, "A fast characteristic finite difference method for fractional advection-diffusion equations," *Advances in Water Resources*, vol. 34, no. 7, pp. 810–816, 2011.
 - [41] K. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, New York, NY, USA, 1993.
 - [42] I. Podluny, *Fractional Differential Equations*, Academic Press, San Diego, Calif, USA, 1999.
 - [43] C. Tadjeran and M. M. Meerschaert, "A second-order accurate numerical method for the two-dimensional fractional diffusion equation," *Journal of Computational Physics*, vol. 220, no. 2, pp. 813–823, 2007.
 - [44] D. Baleanu, A. H. Bhrawy, and T. M. Taha, "Two efficient generalized Laguerre spectral algorithms for fractional initial value problems," *Abstract and Applied Analysis*, vol. 2013, Article ID 546502, 10 pages, 2013.